

Symplectomorphisms with positive metric entropy

Artur Avila^{1,2} | Sylvain Crovisier³ | Amie Wilkinson⁴

¹Institut für Mathematik, Universität
Zürich, Zürich, Switzerland

²IMPA, Estrada Dona Castorina 110, Rio
de Janeiro, Brazil

³CNRS - Laboratoire de Mathématiques
d'Orsay, Université Paris-Saclay, Orsay,
France

⁴Department of Mathematics, University
of Chicago, Illinois, USA

Correspondence

A. Wilkinson, Department of
Mathematics, University of Chicago, 5734
S. University Avenue Chicago, IL 60637,
USA.

Email: wilkinso@math.uchicago.edu

Funding information

SNSF; ERC, Grant/Award Number:
692925; NSF, Grant/Award Number:
DMS-1402852

Abstract

We obtain a dichotomy for C^1 -generic symplectomorphisms: either all the Lyapunov exponents of almost every point vanish, or the map is partially hyperbolic and ergodic with respect to volume. This completes a program first put forth by Ricardo Mañé. A key ingredient is an analysis of partially hyperbolic sets of positive volume. We generalize to partially hyperbolic invariant sets the main result in Dolgopyat and Wilkinson (Astérisque **287** (2003), 33–60) that stable accessibility is C^1 -dense among partially hyperbolic diffeomorphisms.

MSC (2020)

37C20, 37D30, 37J11, 37C05 (primary)

INTRODUCTION

A measurable map $f : M \rightarrow M$ is *ergodic* with respect to an invariant probability measure μ if it is indecomposable: $f^{-1}(A) = A$ implies $\mu(A) = 0$ or 1, for every measurable $A \subset M$. In the context of this paper, where M is a closed connected manifold, f is a homeomorphism, and $\mu = m$ is a normalized volume, ergodicity is equivalent to the equidistribution of almost every orbit: for m -almost every $x \in M$ and every continuous $\phi : M \rightarrow \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \phi(f^j(x)) = \int_M \phi \, dm.$$

© 2022 The Authors. *Proceedings of the London Mathematical Society* is copyright © London Mathematical Society. This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

In his 1983 ICM address [14], Mañé announced the following result, whose proof was completed in 2002 by Bochi [7].

Theorem (Mañé–Bochi). *C^1 -generically, an area-preserving diffeomorphism f of a closed, connected surface M^2 is either Anosov and ergodic or has volume entropy 0.*

The C^1 -genericity means that the statement holds for any f in a dense G_δ subset of the space of C^1 area-preserving diffeomorphisms. The volume entropy $h(f, m)$ measures how orbits separate. By the Pesin formula, it vanishes if and only if for m -a.e. $x \in M$ and every $v \in T_x M \setminus \{0\}$

$$\lim_{n \rightarrow \pm\infty} \frac{1}{n} \log \|D_x f^n v\| = 0.$$

The focus of Mañé's discussion in [14] was in fact the C^1 -generic behavior of *symplectomorphisms*, which in dimension 2 coincide with the area-preserving diffeomorphisms. If $f : M^{2n} \rightarrow M^{2n}$ preserves a symplectic form ω , then it preserves the normalized volume m induced by the form ω^n . The question of whether f is typically ergodic with respect to this volume has a long history going back to the ergodic hypothesis for Hamiltonian systems.

As a consequence of the main result in this paper we obtain the following optimal generalization of Mañé–Bochi theorem to symplectomorphisms in any dimension.

Theorem A. *C^1 -generically among the symplectomorphisms of a closed, connected symplectic manifold (M, ω) , positive volume entropy implies partial hyperbolicity and ergodicity.*

Here, *partial hyperbolicity* of $f : M \rightarrow M$ means that there exists a Df -invariant splitting $TM = E^u \oplus E^c \oplus E^s$, with E^u, E^s both nontrivial, and $N \geq 1$ such that for all unit vectors $v^u \in E_x^u, v^c \in E_x^c$, and $v^s \in E_x^s$, we have

$$2\|D_x f^N(v^s)\| \leq \|D_x f^N(v^c)\| \leq \frac{1}{2}\|D_x f^N(v^u)\|, \quad (1)$$

$$\text{and } \max\{\|D_x f^N(v^s)\|, \|D_x f^{-N}(v^u)\|\} < \frac{1}{2}. \quad (2)$$

An f -invariant set $\Lambda \subset M$ is *partially hyperbolic* if there is a Df -invariant splitting $T_\Lambda M = E^u \oplus E^c \oplus E^s$, with E^u, E^s both nontrivial, and $N \geq 1$ such that (1) and (2) hold for all $x \in \Lambda$. A partially hyperbolic set Λ is *hyperbolic* if E^c is trivial, and f is *Anosov* if M is hyperbolic. In dimension 2, partially hyperbolic sets are hyperbolic, and partially hyperbolic diffeomorphisms are Anosov.

Note that there are obstructions to partial hyperbolicity on certain symplectic manifolds (see [13] for a discussion); for example, surfaces different from \mathbb{T}^2 do not carry an Anosov diffeomorphism, and $\mathbb{C}\mathbb{P}^n$ does not carry a partially hyperbolic symplectomorphism. For these manifolds, Theorem A implies that the C^1 -generic symplectomorphism has volume entropy 0. We also remark that the assumption “positive volume entropy” cannot be replaced by “positive topological entropy:” on any symplectic manifold there exist symplectic horseshoes with positive topological entropy. These horseshoes persist under C^1 -small perturbation.

The proof of the Mañé–Bochi theorem in [7] uses C^1 ergodic-theoretic perturbative techniques to show that C^1 -generically, an area-preserving diffeomorphism f with positive entropy has an invariant hyperbolic set with positive area. The final step of the proof is a well-known and simple folklore result that states that for a C^2 area-preserving diffeomorphism, any invariant

hyperbolic set with positive measure must equal the whole manifold, and the diffeomorphism must be Anosov (for a proof, see, e.g., [11]). This in particular implies that C^1 -generically, any diffeomorphism with an invariant set with positive measure must be Anosov (it is not known whether the C^1 -generic condition can be removed for area-preserving diffeomorphisms, although there exist C^1 -diffeomorphisms with hyperbolic sets of positive volume in the dissipative setting [10]).

The analog to the folklore result is decidedly false in the partially hyperbolic setting; that is, there exist C^2 symplectomorphisms containing proper partially hyperbolic sets of positive volume (such examples are easy to construct as a nonpartially-hyperbolic product of an area-preserving Anosov diffeomorphism (\mathbb{T}^2, f_1) with another area-preserving diffeomorphism (\mathbb{T}^2, f_2) preserving a compact disc D : the invariant proper set is then $\mathbb{T}^2 \times D$).

Question. Suppose that f is a C^2 symplectomorphism that has a partially hyperbolic set of positive volume. What additional conditions imply that f is (globally) partially hyperbolic?

While this question remains open, we show that C^1 -generically the folklore result holds for partially hyperbolic sets in any dimension.

Theorem B. *C^1 -generically among the symplectomorphisms of a closed, connected symplectic manifold (M, ω) , if a diffeomorphism f has a partially hyperbolic set of positive volume, then f is globally partially hyperbolic.*

Ten years ago, building on intermediate work with Viana, Bochi introduced substantially new techniques to prove that for the C^1 -generic symplectomorphism, in any dimension, positive volume entropy implies the existence of a partially hyperbolic invariant set of positive volume [8]. Using this result, we prove Theorem A assuming Theorem B.

Proof of Theorem A. Let (M, ω) be a closed symplectic manifold. Bochi proved [8] that there are two disjoint open sets, \mathcal{Z} and \mathcal{P} in the space $\text{Symp}^1(M, \omega)$ of C^1 symplectomorphisms of M , such that

- $\mathcal{Z} \cup \mathcal{P}$ is a dense G_δ subset of $\text{Symp}^1(M, \omega)$;
- for f in \mathcal{Z} , the volume entropy is zero;
- for $f \in \mathcal{P}$, there exists a positive volume, partially hyperbolic f -invariant set Δ_f .

Theorem B implies that the diffeomorphisms $f \in \mathcal{P}$ are partially hyperbolic and [4, Theorem A] that they are ergodic, completing the proof. \square

Symplectic versus volume-preserving

In a previous paper, we proved the optimal generalization of Mañé–Bochi theorem to *volume-preserving* diffeomorphisms in any dimension.

Theorem [5]. *C^1 -generically among the volume-preserving diffeomorphisms of a closed, connected manifold M , a diffeomorphism with positive volume entropy is nonuniformly Anosov and ergodic.* \square

Despite being optimal generalizations of the same result, the conclusions (nonuniformly Anosov vs. partially hyperbolic) of Theorem A and of [5] are truly different.

In the case of volume-preserving diffeomorphisms f , there exist C^1 open sets of volume-preserving diffeomorphisms which admits a dominated splitting (hence have positive volume entropy) but which are not partially hyperbolic [18], so that the statement of Theorem A is not satisfied. Note that the “nonuniformly Anosov” condition implies in particular that there is a constant $c > 0$ such that for almost every $x \in M$ and every $v \in T_x M \setminus \{0\}$, we have

$$\liminf_{n \rightarrow \pm\infty} \left| \frac{1}{n} \log \|D_x f^n v\| \right| > c.$$

For symplectomorphisms, the exact conclusion of [5] does not hold. Indeed there exists an open set of *partially hyperbolic* C^1 symplectomorphisms that are not Anosov, obtained near the product of an Anosov diffeomorphism and of a surface diffeomorphism with an elliptic fixed point; these systems have positive entropy but by [8, Theorem 1.3] at almost every point the Lyapunov exponents along the center direction vanish and the nonuniformly Anosov condition is not satisfied.

Needless to say, the techniques behind the proof of Theorem A are essentially disjoint from those in the volume-preserving setting of [5]. In the volume-preserving setting, the positive entropy condition implies the existence of nonzero Lyapunov exponents on the phase space, and the proof in [5] harnesses the presence of some nonzero exponents to eliminate all zero Lyapunov exponents throughout large parts of the phase space. A Baire argument completes the proof. In the symplectic setting, we prove that C^1 -generically, the partially hyperbolic set provided by [8] in the presence of positive entropy is the entire manifold.

Partial hyperbolicity and accessibility

For partially hyperbolic diffeomorphisms, a conjecture of Pugh and Shub [16] asserts that the ergodicity should be a consequence of a geometrical property called *accessibility*. We extend this definition to some proper invariant sets that are partially hyperbolic.

Let Λ be a compact partially hyperbolic set for f . Through each $x \in \Lambda$ are unique local stable and unstable manifolds $\mathcal{W}_f^s(x, \text{loc})$ and $\mathcal{W}_f^u(x, \text{loc})$, respectively, which are given by a graph transform argument in a suitable neighborhood of Λ . The local stable and unstable manifolds determine global manifolds by

$$\mathcal{W}_f^u(x) = \bigcup_{n \geq 0} f^n(\mathcal{W}_f^u(f^{-n}(x), \text{loc})), \quad \text{and} \quad \mathcal{W}_f^s(x) = \bigcup_{n \geq 0} f^{-n}(\mathcal{W}_f^s(f^n(x), \text{loc})).$$

We say that Λ is *u-saturated* if for any $x \in \Lambda$, $\mathcal{W}_f^u(x) \subset \Lambda$ and *s-saturated* if for any $x \in \Lambda$, $\mathcal{W}_f^s(x) \subset \Lambda$. We say that Λ is *bisaturated* if it is both *s*- and *u*-saturated. The bisaturated set Λ is *accessible* if for every $p, q \in \Lambda$ there is an *su-path* for f in Λ — that is, a piecewise C^1 path with finitely many segments, each of them contained in a single leaf of \mathcal{W}_f^s or a single leaf of \mathcal{W}_f^u — from p to q .

Note that if f is partially hyperbolic, then M is automatically bisaturated. In this case, we say that f is *accessible* if for every $p, q \in M$ there is an *su-path* from p to q . Dolgopyat and Wilkinson proved in [12] that accessibility holds for a C^1 -open and dense set of partially hyperbolic

diffeomorphisms, volume-preserving diffeomorphisms, and symplectomorphisms of a closed, connected manifold M .

The key ingredient of our proof of Theorem B is a local version of the main result in [12] that implies both Theorem B and the results of [12].[†] Let us denote by $\text{Diff}^k(M)$ the space of C^k diffeomorphisms endowed with the C^k topology. If m is a volume form on M , we denote by $\text{Diff}_m^k(M)$ the subspace of C^k diffeomorphisms that preserve m ; if (M^{2n}, ω) is a symplectic manifold, we denote by $\text{Symp}^k(M, \omega)$ the subspace of C^k symplectomorphisms.

Theorem C. *Let M be a closed manifold, let Λ be a partially hyperbolic set of a diffeomorphism $f : M \rightarrow M$, and let \mathcal{U} be a neighborhood of f in $\text{Diff}^1(M)$. Then there exists a neighborhood U of Λ and a nonempty open set $\mathcal{O} \subset \mathcal{U}$ such that: for any $g \in \mathcal{O}$, any bisaturated partially hyperbolic set $\Delta \subset U$ for g has nonempty interior and is accessible.*

The same result holds in $\text{Diff}_m^1(M)$ and in $\text{Symp}^1(M, \omega)$, if (M^{2n}, ω) is a symplectic manifold.

Using Theorem C, we give a proof of Theorem B.

Proof of Theorem B. Theorem C in [3] shows that for any diffeomorphism f in a dense G_δ subset $\mathcal{R}_1 \subset \text{Symp}^1(M)$, for any f -invariant compact set Λ with positive volume, and for any neighborhood U of Λ , any diffeomorphism $g \in \text{Symp}^1(M)$ close to f also admits a compact g -invariant set $\Delta_g \subset U$ with positive volume. Moreover Δ_g converges to Λ in the Hausdorff topology, and $m(\Delta_g) \rightarrow m(\Lambda)$ as $g \rightarrow f$.

Note that if Λ is partially hyperbolic and U small enough, then Δ_g is partially hyperbolic as well. For $g \in C^2$, the set Δ_g is the union of a bi-saturated set and of a set with zero volume, see [19, Corollary B]. Passing to the limit as $g \rightarrow f$, one deduces that any partially hyperbolic set of f is the union of a bi-saturated set and of a set with zero volume.

[2, Theorem 1] implies that any diffeomorphism in a dense G_δ subset $\mathcal{R}_2 \subset \text{Symp}^1(M)$ is transitive.

Let \mathcal{P} be the set of diffeomorphisms $f \in \mathcal{R}_1 \cap \mathcal{R}_2$ admitting a partially hyperbolic set Λ with positive volume. As we have explained, we can assume with no loss of generality that Λ is bisaturated. For any $f \in \mathcal{P}$, Theorem C above implies that there exists an open set \mathcal{O} containing f in its closure such that for $g \in \mathcal{O} \cap \mathcal{R}_1 \cap \mathcal{R}_2$, the set Δ_g has nonempty interior; but transitivity implies $\Delta_g = M$, so that M is partially hyperbolic. Hence any g in an open and dense subset of \mathcal{P} is partially hyperbolic. \square

The boundary of a bisaturated partially hyperbolic set is also bisaturated (see Lemma 1.3 in Section 1.4). When M is connected, this has the following consequence, which generalizes the main result in [12].

Corollary D. *Let M be a closed, connected manifold, let Λ be a partially hyperbolic set of a diffeomorphism $f : M \rightarrow M$, and let \mathcal{U} be a neighborhood of f in $\text{Diff}^1(M)$.*

There exists a neighborhood U of Λ and a nonempty open set $\mathcal{O} \subset \mathcal{U}$ such that for any $g \in \mathcal{O}$, there is no proper bisaturated subset of U .

In particular, if f is partially hyperbolic, then there exists a nonempty $\mathcal{O} \subset \mathcal{U}$ such that every $g \in \mathcal{O}$ is accessible.

[†] Our proof of Theorem C below also corrects some omissions in the proof in [12]. We will indicate there.

Proof. Let Λ be a partially hyperbolic set for f , and let \mathcal{U} be a neighborhood of f in $\text{Diff}^1(M)$. Applying Theorem C, we obtain a neighborhood U of Λ and a nonempty open set $\mathcal{O} \subset \mathcal{U}$ such that for $g \in \mathcal{O}$, any bisaturated set Δ for g in U has empty interior and is accessible.

Thus, if U contained a proper bisaturated set for a diffeomorphism $g \in \mathcal{O}$, then its boundary would be a bisaturated set with empty interior, a contradiction.

If f is partially hyperbolic, then applying this argument to $\Lambda = M$ gives that any $g \in \mathcal{O}$ is accessible. \square

We remark that in the dissipative setting an earlier version of Theorem C was proved for bi-Lyapunov homoclinic classes by Abdenur–Bonatti–Díaz [1]. Theorem C has the following direct consequence.

Corollary E. *Let M be closed and connected. Then the C^1 -generic f in $\text{Diff}^1(M)$ has no proper, partially hyperbolic, bisaturated invariant compact set.*

The same result holds in $\text{Diff}_m^1(M)$ and in $\text{Symp}^1(M, \omega)$, if (M^{2n}, ω) is a symplectic manifold.

Proof. Let \mathcal{B} be a countable collection of proper open sets of M , such that any compact proper subset of M is contained in an element of \mathcal{B} . For U in \mathcal{B} , let C_U be the set of diffeomorphisms $f \in \text{Diff}^1(M)$ whose maximal invariant set in \bar{U} is partially hyperbolic and let $D_U = \text{Diff}^1(M) \setminus C_U$. Clearly $C_U \cup D_U$ is open and dense in $\text{Diff}^1(M)$.

By Theorem C, there exists a dense open subset $\mathcal{G}_U \subset C_U$ such that U does not contain any proper bisaturated set. Now let

$$\mathcal{R} := \bigcap_{U \in \mathcal{B}} (D_U \cup \mathcal{G}_U).$$

The set \mathcal{R} is residual in $\text{Diff}^1(M)$, and $g \in \mathcal{R}$ implies that g has no proper bisaturated partially hyperbolic subsets. \square

In the case of one-dimensional center, a related result was proved earlier by J. Rodríguez–Hertz [17]: the C^r -generic volume-preserving diffeomorphism has no proper partially hyperbolic invariant compact set with one-dimensional center and positive Lebesgue measure.

Another application of our results is to the Gibbs su -states of a partially hyperbolic diffeomorphism. Let f be partially hyperbolic. Recall that a Gibbs u -state (respectively, s -state) is an f -invariant probability measure μ such that the disintegration of μ along leaves of the \mathcal{W}^u (respectively, \mathcal{W}^s) foliation is absolutely continuous with respect to volume on \mathcal{W}^u (respectively, \mathcal{W}^s) leaves. A Gibbs su -state is an f -invariant probability measure that is both a Gibbs u -state and s -state.

Corollary F. *Let f be partially hyperbolic, and let \mathcal{U} be a neighborhood of f in $\text{Diff}^1(M)$. Then there exists a nonempty open set $\mathcal{O} \subset \mathcal{U}$ such that for every $g \in \mathcal{O}$, if μ is a Gibbs su -state for f , then μ has full support, that is, $\text{supp}(\mu) = M$.*

Proof. The Corollary D implies that there exists a nonempty open set $\mathcal{O} \subset \mathcal{U}$ such that every $g \in \mathcal{O}$ is accessible. Continuity of the foliations \mathcal{W}^u and \mathcal{W}^s implies that the support of a Gibbs su -state is bisaturated. \square

The proof of Theorem C broadly follows the lines of the proof of the main result in [12]. A particular difficulty is the absence of a global partially hyperbolic structure, since holonomy maps are not well defined.

Discussion about stable ergodicity

For $k \leq r$, we say that a diffeomorphism f is C^k -stably ergodic in $\text{Symp}^r(M, \omega)$ if any diffeomorphism g that is C^k -close to f in $\text{Symp}^r(M, \omega)$ is ergodic. Since we do not know examples of stably ergodic diffeomorphisms in $\text{Symp}^1(M, \omega)$, a higher smoothness is usually required. In [6] we have proved that C^1 -stable ergodicity is C^1 -dense among partially hyperbolic diffeomorphisms in $\text{Diff}_m^r(M)$ for any $r > 1$ and an important step was the theorem of [5] stated above. One can ask if the same result holds in $\text{Symp}^r(M, \omega)$.

Question. Is C^1 -stable ergodicity C^1 -dense among $\text{Symp}^r(M, \omega)$, $r > 1$?

Again the strategy for addressing this question should be completely different from the volume-preserving case.

1 | NOTATION AND OUTLINE OF THE PROOF OF THEOREM C

Throughout, M denotes a closed connected Riemannian manifold and m denotes a smooth volume on M , normalized so that $m(M) = 1$. When $M = M^{2n}$ is equipped with a symplectic structure ω , we will indicate so.

1.1 | Outline of the proof

In order to prove that a bisaturated set is accessible, as in the statement of Theorem C, one has to show that sliding alternatively along stable and unstable manifolds also gives progress in the center direction. This is checked by introducing a family of disjoint discs D transverse to the planes $E^u \oplus E^s$ and whose union meets any bisaturated set (Proposition 1.5). Instead of proving accessibility inside the whole bisaturated set Λ , it is enough to prove it locally: for each disc D and any points $x, y \in \Lambda \cap D$, there exists an su -path connecting x to y . The discs are chosen small and satisfy for some integer $J > 0$ large:

$$f^i(D) \cap f^j(D') = \emptyset \text{ when } D \neq D' \text{ and } 0 \leq i < j \leq J.$$

This property ensures that one can perform independent perturbations in the neighborhood of the discs $f^i(D)$ for $0 \leq j \leq J$ to obtain local accessibility on each disc D (Proposition 1.4). This is achieved by building Brin quadrilaterals that produce displacement in each center direction, see Figure 1 in Section 2.2. Since at small scale the bundles are almost constant, the two laminations are almost jointly integrable.

The perturbation then rotates the unstable direction while almost preserving the stable one: the control of the directions is enabled by a control of stable and unstable cones, which requires some time J between perturbations along an orbit. Note that in our setting, since the




FIGURE 1 θ -accessibility

diffeomorphism is a priori not globally partially hyperbolic, the support of the stable and unstable laminations changes when the diffeomorphism is perturbed, which complicates the control of their holonomies.

1.2 | Charts

We introduce for each point $p \in M$ a chart

$$\varphi_p : B(0, 1) \subset T_p M \rightarrow M, \quad \text{with } \varphi_p(0) = p,$$

which has the following properties:

- (1) The map $p \mapsto \varphi_p$ is “piecewise continuous in the C^1 topology.”[†] More precisely, there exist open sets $U_1, \dots, U_\ell \subset M$ and:
 - compact sets K_1, \dots, K_ℓ covering M with $K_i \subset U_i$,
 - trivialisations $\psi_i : U_i \times \mathbb{R}^d \rightarrow T_{U_i} M$ such that $\psi_i(\{p\} \times B(0, 2))$ contains the unit ball in $T_p M$ for each $p \in U_i$, and
 - smooth maps $\Phi_i : U_i \times B(0, 2) \rightarrow M$,
 such that each $p \in M$ belongs to some K_i , with

$$\varphi_p = \Phi_i \circ \psi_i^{-1} \text{ on } B(0, 1) \subset T_p M.$$

- (2) When a volume or symplectic form has been fixed on M , it pulls back under φ_p to a constant form on $T_p M$.

The construction can be done as follows. Assume that a Riemannian metric on M has been fixed. We first choose a cover of TM by trivialisations $\psi_i : U_i \times \mathbb{R}^d \rightarrow T_{U_i} M$ such that $\psi_i(p, 0) = 0 \in T_p M$. By taking a large number of charts with small supports, the maps $u \mapsto \psi(p, u)$ can be chosen close to isometries in the C^1 topology.

Denoting by $B^d(0, R)$ the standard open ball in \mathbb{R}^d with radius R , we then construct finitely many charts

$$\Phi : B^d(0, 3) \rightarrow M$$

[†] In [12] similar charts are constructed, but it is claimed that these can be defined on a fixed domain in \mathbb{R}^d , depending continuously on $p \in M$. This continuity is not possible (when TM is not trivialisable) and also unnecessary.

such that (after replacing the U_i by smaller open sets, if necessary) for any U_i , there exists one such chart Φ satisfying $U_i \subset \Phi(B(0, 1))$. We can thus define

$$\Phi_i(p, u) := \Phi(\Phi^{-1}(p) + u) \text{ on } U_i \times B(0, 2),$$

and for each $p \in M$, choose K_i containing p and set $\varphi_p = \Phi_i \circ \psi_i^{-1}$ on $B(0, 1) \subset T_p M$.

When M is equipped with a volume form, one can require (by Moser’s theorem [15]) that φ_p send divergence-free vector fields to divergence-free vector fields. When M is equipped with a symplectic form ω , one can require (by Darboux’s theorem) that $(\varphi_p)_* \omega$ coincide (up to multiplication by a constant) with the standard symplectic form $\Sigma dp_i \wedge dq_i$ of $\mathbb{R}^d = \mathbb{R}^{2n}$. This concludes the construction of the charts.

Remark 1.1. Given a compact set X with a continuous splitting $T_X M = E \oplus F$ and $\gamma > 0$, the construction below shows that one can choose a Riemannian metric on M and the charts φ_p such that for each $p \in \Lambda$ and $z \in \varphi_p^{-1}(X) \cap B(0, 1)$, the norm of the orthogonal projection of $D\varphi_p(z)^{-1}(F_{\varphi_p(z)})$ onto $D\varphi_p(z)^{-1}(E_{\varphi_p(z)})$ is less than γ .

Indeed, one can first choose a Riemannian metric such that the norm of the orthogonal projection from F to E is arbitrarily small. One then chooses the charts Φ in such a way that the bundles E and F lifted in $B^d(0, 3)$ are close to constant bundles (this is possible by the continuity of E and F). Since the ψ_i are close to isometries, this shows that for $p \in X$, the bundles E and F lifted by φ_p in $B(0, 1) \subset T_p M$ are close to E_p and F_p , respectively, which are close to orthogonal.

1.3 | Conefields

A k -conefield C over a subset $U \subset M$ is a subset of the tangent bundle $T_U M$ satisfying:

- $tv \in C$ for any $v \in C$ and $t \in \mathbb{R}$; and
- there is a continuous subbundle $E \subset T_U M$ with k -dimensional fibers such that $\{v \in C : \|v\| = 1\}$ is a neighborhood of $\{v \in E : \|v\| = 1\}$.

We denote $C(x) := C \cap T_x M$. The conefield C is *forward invariant* under a diffeomorphism f if for any $x \in U \cap f^{-1}(U)$, the image of $C(x)$ is contained in $\text{Interior}(C(f(x))) \cup \{0\}$. It is *contracted* if there exists $N \geq 1$ such that $\|Df.v\| \leq \frac{1}{2}\|v\|$ for any $v \in C(x)$. It is *transverse* to a bundle E if $C(x) \cap E(x) = \{0\}$ at any point $x \in U$.

A conefield C' over U is a δ -perturbation of C with support in $V \subset U$ if there exists a diffeomorphism h that is δ -close to the identity in the C^1 topology such that $h(U) = U$, h coincides with the identity on $U \setminus V$ and $h^*(C) = C'$. A k -conefield C is δ -close to a subbundle E of $T_U M$ with k -dimensional fibers if $\{v \in C : \|v\| = 1\}$ is δ -close to $\{v \in E : \|v\| = 1\}$ in the Hausdorff distance.

Let $f : M \rightarrow M$ be a diffeomorphism, and let Λ be a compact f -invariant set with a partially hyperbolic splitting $T_\Lambda M = E^u \oplus E^c \oplus E^s$.

A neighborhood U of Λ is *admissible* if there exist continuous conefields C^u, C^s, C^{cu}, C^{cs} over U containing E^u, E^s, E^{cu}, E^{cs} on $T_\Lambda M$ that are transverse to E^{cs}, E^{cu}, E^s, E^u , respectively, and such that C^u, C^{cu} are forward invariant, C^s, C^{cs} backward invariant, C^s (respectively, C^u) is contracted by f (respectively by f^{-1}). The following proposition is standard (see [9, Appendix B.1]).

Proposition 1.2. *For every partially hyperbolic set Λ for f , there exist neighborhoods \mathcal{U}_0 and U_0 of Λ and f , and cone fields $C_0^u, C_0^s, C_0^{cu}, C_0^{cs}$ on U_0 with the following property. If $\Delta \subset U_0$ is a compact g -invariant set for $g \in \mathcal{U}_0$, then it is partially hyperbolic and U_0 is an admissible neighborhood of Δ with respect the cone fields $C_0^u, C_0^s, C_0^{cu}, C_0^{cs}$ and the diffeomorphism g .*

1.4 | Bi-saturated partially hyperbolic sets: Accessibility

Consider a diffeomorphism f and a partially hyperbolic set Λ . We denote by $U_0(f, \Lambda)$ and $\mathcal{U}_0(f, \Lambda)$ the neighborhoods given by Proposition 1.2.

Let M be a manifold of dimension $d \geq 2$, and let K be a compact subset of M . A k -dimensional topological lamination \mathcal{L} of K is a decomposition of K into path-connected subsets

$$K = \bigcup_{x \in K} \mathcal{L}(x)$$

called *leaves*, where $x \in \mathcal{L}(x)$, and two leaves $\mathcal{L}(x)$ and $\mathcal{L}(y)$ are either disjoint or equal, and a covering of K by coordinate neighborhoods $\{U_\alpha\}$ with local coordinates $(x_\alpha^1, \dots, x_\alpha^d)$ with the following property. For $x \in U_\alpha \cap K$, denote by $\mathcal{L}_{U_\alpha}(x)$ the connected component of $\mathcal{L}(x) \cap U_\alpha$ containing x . Then in coordinates on U_α the local leaf $\mathcal{L}_{U_\alpha}(x)$ is given by a set of equations of the form $x_\alpha^{k+1} = \dots = x_\alpha^d = cst$. If the local coordinates $(x_\alpha^1, \dots, x_\alpha^d)$ can be chosen uniformly C^r along the local leaves (i.e., to have uniformly C^r overlaps on the sets $x_\alpha^{k+1} = \dots = x_\alpha^d = cst$), then we say that \mathcal{L} has C^r leaves.

Note that the leaves of a lamination with C^r leaves are C^r , injectively immersed submanifolds of M . A lamination of M is called a foliation.

Let Λ be a partially hyperbolic set for f . Continuity and invariance of the partially hyperbolic splitting implies that Λ is u - (respectively, s -) saturated if and only if $\{\mathcal{W}_f^u(x) : x \in \Lambda\}$ (respectively, $\{\mathcal{W}_f^s(x) : x \in \Lambda\}$) is a lamination of Λ .

Lemma 1.3. *Let Λ be a bisaturated partially hyperbolic set for f . Then the boundary $\partial\Lambda$ is also bisaturated.*

Proof. A set is bisaturated if and only if it is laminated by \mathcal{W}^s leaves and by \mathcal{W}^u leaves. If x, y belong to the same leaf of a compact lamination $\mathcal{W} \subset M$, then there exist neighborhoods V_x and V_y in \mathcal{W} , of x and y , respectively, that are homeomorphic; thus, x belongs to the interior of \mathcal{W} if and only if y does. \square

Let $\mathcal{P}(M)$ be the collection of all subsets of M . We say that (f, Λ) is *accessible on* $X \in \mathcal{P}(M)$ if for every $p \in X \cap \Lambda$, and every $q \in X$, there is an su -path for f in Λ from p to q . In particular, if $X \cap \Lambda \neq \emptyset$, and f is accessible on X , then $\Lambda \supset X$.

We say that (f, Λ) is *stably accessible on* $X \in \mathcal{P}(M)$ if there exists a neighborhood $U \subset U_0(f, \Lambda)$ of Λ with $X \subset U$, and a neighborhood $\mathcal{U} \subset \mathcal{U}_0(f, \Lambda)$ of f such that for every $\tilde{f} \in \mathcal{U}$ and every \tilde{f} -invariant bisaturated compact set $\tilde{\Lambda} \subset U$, we have that $(\tilde{f}, \tilde{\Lambda})$ is accessible on X .

We say that a set $X \in \mathcal{P}(M)$ is a *c-section for* (f, Λ) if for every bisaturated subset $\Delta \subset \Lambda$, we have $X \cap \Delta \neq \emptyset$ (the name comes from the fact that it meets each class, once one quotients along the \mathcal{W}^s and \mathcal{W}^u leaves, which are transverse to E^c).

1.5 | Admissible families of disks

Since we do not assume that E^c is tangent to a foliation, we will work with approximate center manifolds.

For $\rho > 0$ small and $p \in \Lambda$, we denote by $B^c(0, \rho)$ the ball inside E_p^c of radius ρ and set

$$V_\rho(p) := \varphi_p(B^c(0, \rho)).$$

We refer to $V_\rho(p)$ as a *c-admissible disk* (with respect to (f, Λ)) with center p and radius ρ and write $r(V_\rho(p)) = \rho$. If D is a *c-admissible disk* with center p and radius ρ , then for $\beta \in (0, 1)$, we denote by βD the *c-admissible disk* with center p and radius $\beta\rho$. A *c-admissible family* (with respect to (f, Λ)) is a finite collection of pairwise disjoint, *c-admissible disks*.

Define the return time $R : \mathcal{P}(M) \rightarrow \mathbf{N} \cup \{\infty\}$ as follows. For $X \in \mathcal{P}(M)$, let $R(X)$ be the smallest $J \in \mathbf{N} \cup \{\infty\}$ satisfying:

$$f^i(X) \cap X \neq \emptyset, \text{ with } |i| = J + 1. \tag{3}$$

Note that $R(B_\rho(p)) \rightarrow \text{per}(p)$, as $\rho \rightarrow 0$, where $\text{per}(p)$ denotes the period if p is periodic and $\text{per}(p) = \infty$ if p is not periodic.

For \mathcal{D} a *c-admissible family* and $\beta \in (0, 1)$, we introduce the following notations:

$$\begin{aligned} \beta \mathcal{D} &= \{\beta D \mid D \in \mathcal{D}\}, \\ |\mathcal{D}| &= \bigcup_{D \in \mathcal{D}} D, \\ r(\mathcal{D}) &= \sup_{D \in \mathcal{D}} r(D), \text{ and} \\ R(\mathcal{D}) &= R(|\mathcal{D}|), \end{aligned}$$

where as before $r(D)$ denotes the radius and $R(D)$ the smallest return time.

1.6 | Two propositions

Our first proposition is the counterpart to [12, Lemma 1.1].[†]

Proposition 1.4 (Stable accessibility on center disks). *Let Λ be partially hyperbolic for f , and let $\delta > 0$ be given. Then there exist $J \geq 1$ and a neighborhood U of Λ satisfying $\overline{U} \subset U_0(f, \Lambda)$ and the following property.*

If \mathcal{D} is a c-admissible family with respect to (f, Λ) with $r(\mathcal{D}) < J^{-1}$ and $R(\mathcal{D}) > J$, then for all $\sigma > 0$ there exists $g \in \mathcal{U}_0(f, \Lambda)$ such that:

- (1) $d_{C^1}(f, g) < \delta$,
- (2) $d_{C^0}(f, g) < \sigma$,
- (3) *for each $D \in \mathcal{D}$, and every bisaturated partially hyperbolic set $\Delta \subset U$ for g , we have that (g, Δ) is stably accessible on D .*

[†] While Lemma 1.1 is stated correctly in [12], its proof has an error. In particular, in Lemma 3.3, $r(\mathcal{D})$ and $R(\mathcal{D})$ are chosen after θ is given, when they should be chosen before. A correct proof is given here.

The second proposition is the counterpart to [12, Lemmas 1.2 and 1.3].

Proposition 1.5 (Stable c -sections exist). *Let Λ be a partially hyperbolic set for f . Then there exists $\delta > 0$ with the following property.*

Let U be a neighborhood of Λ satisfying $\overline{U} \subset U_0(f, \Lambda)$. For any $J \geq 1$ there exists a c -admissible disk family \mathcal{D} and $\sigma > 0$ such that:

- (1) $r(\mathcal{D}) < J^{-1}$,
- (2) $R(\mathcal{D}) > J$, and
- (3) *if g satisfies $d_{C^1}(f, g) < \delta$ and $d_{C^0}(f, g) < \sigma$, then for any bisaturated partially hyperbolic set $\Delta \subset U$ for g , the set $|\mathcal{D}|$ is a c -section for (g, Δ) .*

1.7 | Proof of Theorem C from Propositions 1.4 and 1.5

Let f , Λ , and \mathcal{U} be given as in the statement of the theorem. Let $\delta > 0$ be given by Proposition 1.5. By shrinking the value of δ if necessary, we may assume that $d_{C^1}(f, g) < \delta$ implies $g \in \mathcal{U}$.

Let the neighborhood U of Λ such that $\overline{U} \subset U_0(f, \Lambda)$ and $J \geq 1$ be given by Proposition 1.4, using the value $\delta/2$. Let \mathcal{D} and σ be given by Proposition 1.5. Applying Proposition 1.4 to f , Λ , $\delta/2$, \mathcal{D} , $\sigma/2$ we associate a perturbation g_0 of f satisfying:

- (1) $d_{C^1}(f, g_0) < \delta/2$,
- (2) $d_{C^0}(f, g_0) < \sigma/2$,
- (3) for each $D \in \mathcal{D}$, and every bisaturated partially hyperbolic set $\Delta \subset U$ for g_0 , we have that (g_0, Δ) is stably accessible on D .

By compactness of the Hausdorff topology, there exists a neighborhood $\mathcal{O} \subset \mathcal{U}$ of g_0 in the $\delta/2$ -neighborhood of g_0 such that accessibility holds on each D , for bisaturated sets of any $g \in \mathcal{O}$. Then for any $g \in \mathcal{O}$, we have $d_{C^1}(f, g) < \delta$ and $d_{C^0}(f, g) < \sigma$. Let $\Delta \subset U$ be a bisaturated set for such a g .

On the one hand, Proposition 1.5 implies that $|\mathcal{D}|$ is a c -section for Δ , and so there exists $D \in \mathcal{D}$ such that $\Delta \cap D \neq \emptyset$. On the other hand, Proposition 1.4 then implies that $\Delta \supset D$. By saturating D by local stable and unstable manifolds for Δ and using again the bisaturation of Δ , we see that Δ has nonempty interior.

Consider any point $p \in \Delta$ and its accessibility class $C(p)$, that is, the set of points $p' \in \Delta$ that can be connected to p by a su-path in Δ . Note that the closure of $C(p)$ is a bisaturated set and hence meets the c -section $|\mathcal{D}|$ at a point z . This point belongs to a disc $\varphi_{x_i}(B^c(0, \rho_i)) \subset |\mathcal{D}|$. Any point y close to z can be joined by a su-path with two legs to a point in $\varphi_{x_i}(B^c(0, \rho_i))$: this proves that $C(p)$ intersects $|\mathcal{D}|$ at a point z_p . If q is another point in Δ , its accessibility class $C(q)$ meets $|\mathcal{D}|$ as well at a point z_q and the stable accessibility relative to $|\mathcal{D}|$ implies that the two points z_p, z_q can be connected by a su-path. We have thus proved that p and q belong to the same accessibility class, hence that Λ is accessible, completing the proof of Theorem C. \square

2 | PROOF OF PROPOSITION 1.4

Fix f , Λ , δ as in the statement of Proposition 1.4. We denote by c the dimension of the center bundle. The proof follows closely the proof of Lemma 1.1 in [12]. The main adaptation is that we

work inside bisaturated sets Δ for g in a small neighborhood of Λ and consider unstable and stable holonomies in restriction to Δ .

The partially hyperbolic splitting for f at a point z will be denoted by $E_z^u \oplus E_z^c \oplus E_z^s$, whereas the splitting for another diffeomorphism g will be denoted by $E_{g,z}^u \oplus E_{g,z}^c \oplus E_{g,z}^s$. As before $d = \dim(M)$.

For any $p \in M$ we have defined a chart $\varphi_p : B(0, 1) \subset T_pM \rightarrow M$. From Remark 1.1, we can assume that for any $p \in \Lambda$ and $z \in B(0, 1) \cap \varphi_p^{-1}(\Lambda)$, the orthogonal projections of E_z^s on $E_z^u \oplus E_z^c$ and of E_z^u on $E_z^c \oplus E_z^s$ have norms smaller than 10^{-1} .

In the following, we will reduce the C^1 -size δ of the perturbation, the size of the neighborhood U of Λ , and the size ρ of the c -admissible discs.

2.1 | A center covering

We will need to replace c -admissible discs by families of disjoint smaller balls, which will be the support of independent perturbations.

Lemma 2.1. *There exist $\delta_1, \rho_1 > 0$, $K > 1$ and a neighborhood U_1 of Λ such that for any $\rho \in (0, \rho_1)$, any c -admissible disc D with radius ρ , centered $p \in \Lambda$, and for any $\varepsilon \in (0, K^{-1}\rho)$, there exist $z_1, \dots, z_\ell \in T_pM$ such that:*

- (1) *the balls $B(z_i, 100d^2\varepsilon)$ are in the $K\varepsilon$ -neighborhood of $\varphi_p^{-1}(D)$;*
- (2) *the balls $B(z_i, 100d^2\varepsilon)$ are pairwise disjoint;*
- (3) *for any $x \in D$, there exists z_i such that for any g that is δ_1 -close to f in the C^1 distance and for any bisaturated set $\Delta \subset U_1$ for g :*
 - (a) *if $x \in \Delta$ then there is a su-path for g between x and $\varphi_p(B(z_i, \varepsilon))$,*
 - (b) *if $\varphi_p(B(z_i, \varepsilon)) \subset \Delta$, then any point $y \in \varphi_p(B(\varphi_p^{-1}(x), \varepsilon/2))$ belongs to an su-path that intersects $\varphi_p(B(z_i, \varepsilon))$.*

Proof. There exists $K_0 > 1$ such that for any $\varepsilon > 0$, the unit ball $B^c(0, 1) \subset \mathbb{R}^c$ can be covered by balls $B(x_1, \varepsilon/4), \dots, B(x_\ell, \varepsilon/4)$ in \mathbb{R}^d with the property that any ball $B(x_i, 200d^2\varepsilon)$ intersects at most $K_0 - 1$ others.

We introduce a local flow along the unstable leaves W_g^u of Δ for g . Fix $p \in \Lambda$, and for $x \in B(0, \rho_1) \cap \varphi_p^{-1}(\Delta) \subset T_pM$, denote by π_x^u the projection along $(E_p^u)^\perp$ of $B(0, 2\rho_1)$ onto the connected component of $\varphi_p^{-1}(W_{g, \varphi_p(x)}^u) \cap B(0, 2\rho_1)$ containing x .

Given a unit vector $v^u \in E_p^u$, we define a vector field X_{v^u} along the local leaves of $\varphi_p^{-1}(W_g^u)$ as follows: for $x \in B(0, \rho_1) \cap \varphi_p^{-1}(\Delta) \subset T_pM$, let

$$X_{v^u}(x) = D\pi_x^u(x + v^u).$$

The vector field X_{v^u} induces a local flow Φ^u on the set $B(0, 2\rho_1) \cap \varphi_p^{-1}(\Delta)$, for $|t| < \rho_1$: the orbit of x is the projection by π_x^u of the curve $t \mapsto x + tv^u$. The orbits are C^1 curves with a tangent space arbitrarily close to $\mathbb{R}v^u$ if ρ_1, δ_1 and U_1 have been chosen small enough.

Let D be a c -admissible disk centered at p , with radius $\rho < \rho_1$. From the property above, one can choose points $x_1, \dots, x_\ell \in E_p^c$ such that the balls $B(x_1, \varepsilon/4), \dots, B(x_\ell, \varepsilon/4)$ cover $\varphi_p^{-1}(D) \subset E_p^c$ and choose integers k_1, \dots, k_ℓ in $\{1, \dots, K_0\}$ such that the balls $B(z_1, 100d^2\varepsilon), \dots, B(z_\ell, 100d^2\varepsilon)$,

centered at points $z_i := x_i + 500d^2k_i\varepsilon v^u$ are pairwise disjoint. The two first items are satisfied with $K = 1000d^2K_0$.

Since the flow lines under Φ^u are C^1 -close to lines parallel to v^u , for each point $x \in D \cap \Delta$, there exists $|t| < 10K_0$ such that $\Phi_t(\varphi_p^{-1}(x))$ belongs to one of the balls $B(z_i, \varepsilon/2)$. Hence the unstable manifold of x intersects $\varphi_p(B(z_i, \varepsilon/2))$.

If $\varphi_p(B(z_i, \varepsilon)) \subset \Delta$, then the continuous map $H : (t, y) \mapsto \Phi_t^u(y)$ defined on $[-K\varepsilon, K\varepsilon] \times B(z_i, \varepsilon)$ is $\varepsilon/2$ -close in the C^0 metric to the map $(t, y) \mapsto y + tv^u$. Hence $B(z_i, \varepsilon/2) + [-(K - 1)\varepsilon, (K - 1)\varepsilon]v^u$ is contained in the image of H . By construction $B(\varphi_p^{-1}(x), \varepsilon/2)$ is contained in this image. We have thus proved that any point in $\varphi_p(B(\varphi_p^{-1}(x), \varepsilon/2))$ belongs to the unstable manifold of some point in $\varphi_p(B(z_i, \varepsilon))$.

The lemma is proved. □

2.2 | A center accessibility criterion

Let $\theta \in (0, 1)$, $p \in \Lambda$ and $z \in T_pM$.

We say that the pair (g, Δ) is θ -accessible on the ball $\varphi_p(B(z, 2d\varepsilon))$ if there exist an orthonormal basis w_1, \dots, w_c of E_p^c and for each $j \in \{1, \dots, c\}$, a continuous map

$$H^j : [-1, 1] \times [0, 1] \times \varphi_p^{-1}(\Delta) \cap B(z, 2d\varepsilon) \rightarrow \varphi_p^{-1}(\Delta) \cap B(0, 2\rho)$$

such that for any $x \in \varphi_p^{-1}(\Delta) \cap B(z, 2d\varepsilon)$ and $s \in [-1, 1]$,

- (a) $H^j(s, 0, x) = x$,
- (b) the map $\varphi_p \circ H^j(s, \cdot, x) : [0, 1] \rightarrow \Delta$ is a 4-legged su-path, that is, the concatenation of 4 curves, each contained in a stable or unstable leaf,
- (c) $\|H^j(s, 1, x) - x\| < \frac{\varepsilon}{10d}$, and
- (d) $\|H^j(\pm 1, 1, x) - (x \pm \theta\varepsilon w_j)\| < \theta \frac{\varepsilon}{10d}$.

See the Figure 1.

The following replaces [12, Lemma 3.2] in our setting.[†]

Lemma 2.2. *For any $\theta > 0$, there exist $\delta_2, \rho_2 > 0$ and a neighborhood U_2 of Λ such that*

- for any $p \in \Lambda$, any z in the ball $B(0, \rho_2) \subset T_pM$ and $\varepsilon \in (0, \rho_2)$,
- for any diffeomorphism g which is δ_2 -close to f in the C^1 topology,
- for any bi-saturated set $\Delta \subset U_2$ such that (g, Δ) is θ -accessible on the ball $\varphi_p(B(z, 2d\varepsilon))$,

the pair (g, Δ) is accessible on $\varphi_p(B(z, \varepsilon))$.

Proof. Let u, s be the dimensions of the bundles E^u, E^s . Hence $d = u + c + s$. Let v_1, \dots, v_u and v_{u+c+1}, \dots, v_d be orthonormal bases of E_p^u and E_p^s , respectively. We define local flows (Φ_i^t) (for $i \in \{1, \dots, u\} \cup \{u + c + 1, \dots, d\}$) on $\varphi_p^{-1}(\Delta)$, as in the proof of Lemma 2.1.

[†] The proof of [12, Lemma 3.2] contains a subtle error, in the sentence beginning: “By a standard argument . . .” The argument described there is indeed standard in the dynamically coherent setting, but it is not clear in the more general setting. We bypass this argument here by removing the requirement that the su -paths end in the c -admissible disk, establishing local accessibility directly.

As in the proof of Lemma 2.1, we define for each $j \in \{1, \dots, u\}$ a local flow Φ^j : at any $x \in B(z, 2d\varepsilon) \cap \varphi_p^{-1}(\Delta)$ it is tangent to the vector field obtained by projecting the vector $\theta\varepsilon v_j$ at x orthogonally to E^u on the tangent space $\varphi_p^{-1}(W^u(\varphi_p(x)))$. The point $\Phi_t^j(x)$ is the projection of $x + t\theta\varepsilon v_j$ orthogonally onto $\varphi_p^{-1}(W^u(\varphi_p(x)))$. Similarly, for $j \in \{1, \dots, s\}$, we define a flow $\Phi_t^{u+c+j}(x)$ in the direction of v_{u+c+j} , along the stable leaves of $\varphi_p^{-1}(\Delta)$. Choosing ρ_2, δ_2, U_2 small, the tangent spaces of the unstable and stable leaves of g inside $\varphi_p^{-1}(\Delta)$ in $B(p, \rho_2)$ are close to E_p^u and E_p^s . This gives

$$\|\Phi_t^j(x) - (x + t\theta\varepsilon v_j)\| < |t|\theta \frac{\varepsilon}{10d}. \tag{4}$$

We also define maps in the center direction. Let us set $v_{u+j} = w_j$. For $j \in \{1 \dots, c\}$, we introduce inductively $\Phi_t^{u+j}(x)$ (while it can be defined) by

$$\begin{aligned} \Phi_t^{u+j}(x) &= H^j(t, 1, x) \text{ when } t \in [0, 1], \\ \Phi_t^{u+j}(x) &= \Phi_{t-1}^{u+j} \circ \Phi_1^{u+j}(x) \text{ when } t > 1, \\ \Phi_t^{u+j}(x) &= \Phi_{t+1}^{u+j} \circ (\Phi_{-1}^{u+j})^{-1}(x) \text{ when } t < 0, \end{aligned}$$

Let us consider a point $x_0 \in \varphi_p^{-1}(\Delta) \cap B(z, \varepsilon)$. From (c), (d), and (4), one can define for each $(t_1, \dots, t_d) \in [-3\theta^{-1}, 3\theta^{-1}]^d$

$$P(t_1, \dots, t_d) = \Phi_{t_1}^1 \dots \Phi_{t_d}^d(x_0).$$

This induces a continuous map satisfying

$$\|P(t_1, \dots, t_d) - (x_0 + \sum_i t_i \theta \varepsilon v_i)\| < \frac{2\varepsilon}{10}.$$

It follows that the degree of the map $x_0 + \sum_i t_i \theta \varepsilon v_i \mapsto P(t_1, \dots, t_d)$ over any point that is $(3 - \frac{1}{2})\varepsilon$ -close to x_0 is equal to 1. Hence the image of P contains the ball centered at x_0 of radius $(3 - \frac{1}{2})\varepsilon$, and the ball $B(z, \varepsilon)$. By construction, the points in the image of $\varphi_p \circ P$ are be connected to z_0 by an su-path in Δ . We have thus shown that (g, Δ) is accessible on $B(z, \varepsilon)$. □

2.3 | Elementary perturbations

The perturbation will be built from the following.

Lemma 2.3. *There exists $\eta, \alpha_0 > 0$ small with the following properties.*

For any $\alpha \in (0, \alpha_0)$, $p \in \Lambda$, $z \in B(0, 1/4) \subset T_p M$, $r \in (0, 1/4)$ and any unit vector $v \in E_p^c$, there exists a diffeomorphism T of $T_p M$:

- which is supported on $B(z, 3r)$,

– whose restriction to $B(z, 2r)$ coincides with

$$x \mapsto x + \alpha\eta rv,$$

- whose tangent map $DT(y)$ is α -close to id for any $y \in T_p M$,
- which is $\frac{r}{100d^2}$ -close to the identity in the C^0 distance.

Moreover, if f preserves a volume m or a symplectic form ω , then such a T can be constructed so that the maps $\varphi_p \circ T \circ \varphi_p^{-1}$ preserve m or ω as well.

Proof. The construction is standard. One first notices that it can be done in the case $r = 1/4$. One then reduces r by conjugating by an homothety.

With a bump function, one builds a vector field which takes the constant value v on $B(z, 2r)$ and which vanishes outside $B(z, 3r)$. There exists $\eta > 0$ such that the time t of the flow is at distance $\eta^{-1} \cdot t$ from the identity in the C^1 topology. For $\alpha > 0$ small, the map T is the time $\alpha\eta$ of the flow.

In the volume-preserving case, the lift of the volume form is constant in the domain of the charts. Choosing a divergence free vector field, the map T preserves the volume.

In the symplectic case, the symplectic form in the chart is constant. The constant vector field is hamiltonian. Using a bump function, one can extend the hamiltonian to a function which vanishes outside $B^c(z, 3r)$. The associated vector field is then symplectic as required. \square

The diffeomorphism g of Proposition 1.4 will be obtained from f as a composition of some number L of diffeomorphisms obtained from Lemma 2.3:

$$g := \Psi_L \circ \dots \circ \Psi_1 \circ f, \quad \text{with } \Psi_i := \varphi_{p_i} \circ T_i \circ \varphi_{p_i}^{-1}, \quad (5)$$

where the points p_i belong to Λ and where the maps T_i are diffeomorphisms of $T_{p_i} M$ given by Lemma 2.3 which coincide with the identity outside some sets Ω_i contained in $B(0, 2\rho) \subset T_{p_i} M$, for some $\rho > 0$ small which will be chosen later. The supports $\varphi_{p_i}(\Omega_i)$ will be chosen pairwise disjoint so that the maps Ψ_i commute.

We consider cone fields C_0^u, C_0^s on U_0 , respectively, invariant by f and f^{-1} as in Proposition 1.2. If $\delta_3 > 0$ is small, for any g that is δ_3 -close to f in the C^1 topology, the same cone fields C_0^u, C_0^s are still invariant by g and g^{-1} .

We may assume without loss of generality that $\delta < \min(\delta_1, \delta_2, \delta_3)$. Recall that the charts φ_p depend continuously on p in the C^1 topology when p belong to the atoms of a finite compact covering of M . Consequently, there exists $\rho_3 > 0$ and $\alpha \in (0, \alpha_0)$ small such that if the support Ω_i of each map T_i is contained in $B(0, 2\rho) \subset T_{p_i} M$ for some $\rho \in (0, \rho_3)$ and has a tangent map DT which is α -close to the identity, then the diffeomorphism g is δ -close to f in the C^1 distance.

2.4 | Choice of J and U

For $J \geq 1$, let us introduce the iterates $C^u = Df^J(C_0^u)$ and $C^s = Df^{-J}(C_0^s)$ on a neighborhood U of Λ satisfying

$$\bar{U} \subset U_1 \cap U_2 \cap \bigcap_{|k| \leq J} f^k(U_0). \quad (6)$$

The contraction of the cone fields ensures that C^u and C^s get arbitrarily close to the bundles E^u and E^s (defined on the maximal invariant set of \bar{U}) as $J \rightarrow +\infty$. Hence, there exist $J_1 \geq 1$ and $\rho_4 > 0$ such that if $J \geq J_1$ and if $\rho < \rho_4$, then for any $p \in \Lambda$,

- $\varphi_p(B(0, 2\rho)) \subset U$,
- the cone fields $D\varphi_p^{-1}(C^s)$ and $D\varphi_p^{-1}(C^u)$ on $B(0, 2\rho)$ are γ -close to the spaces E_p^s and E_p^u in T_pM for some $\gamma > 0$ much smaller than $\alpha\eta$.

In particular, from the choice of the Riemannian metric, for any point $z \in B(0, 2\rho) \subset T_pM$, the orthogonal projection of any unit vector $u \in D\varphi_p(z)^{-1}(C^s(\varphi_p(z)))$ on E_p^c has norm smaller than $1/2$.

We now fix:

- J_1 and the neighborhood U to satisfy (6),
- $\rho > 0$ smaller than $\min(\rho_1, \rho_2, \rho_3, \rho_4)$,
- $J \geq J_1$ large enough so that any c -admissible disk D with center $p \in \Lambda$ and radius $r(D) < J^{-1}$ lifts by φ_p as a subset of $B(0, \rho) \subset T_pM$,
- a c -admissible family \mathcal{D} of disks as in the statement of Proposition 1.4,
- $\sigma > 0$ as in the statement of Proposition 1.4.

The construction also depends on a number $\varepsilon > 0$ smaller than σ, ρ_2 and $K^{-1}r(D)$ for any $D \in \mathcal{D}$ (where K is the constant in Lemma 2.1). We will specify later the value of ε .

2.5 | Construction of the diffeomorphism g

We associate to each c -admissible disc $D \in \mathcal{D}$ a set of balls as given by Lemma 2.1. The union of these sets defines a family \mathcal{B} of balls $B_i := \varphi_{p_i}(B(z_i, 100d^2\varepsilon))$, $1 \leq i \leq L$, inside the tangent spaces of points $p_i \in \Lambda$. Since the discs $D \in \mathcal{D}$ are disjoint, by choosing $\varepsilon > 0$ small enough the items (a) and (b) in Lemma 2.1 ensure that the balls B_i are pairwise disjoint.

We now define $g \circ f^{-1}$ in each B_i separately. The choice of ρ gives $\varphi_{p_i}(B(z_i, 100d^2\varepsilon)) \subset U$, and one can choose two spaces $\mathcal{E}^u, \mathcal{E}^s \subset T_{p_i}M$ with the same dimension as $E_{p_i}^u$ and $E_{p_i}^s$ and satisfying

$$D\varphi_{p_i}(z_i) \cdot \mathcal{E}^s \subset C^s(\varphi_{p_i}(z_i)), \quad D\varphi_{p_i}(z_i) \cdot \mathcal{E}^u \subset C^u(\varphi_{p_i}(z_i)).$$

We choose two unit vectors $e^s \in \mathcal{E}^s$, $e^u \in \mathcal{E}^u$ and we also fix an orthonormal basis w_1, \dots, w_c of $E_{p_i}^c$.

For each $j \in \{1, \dots, c\}$, the Lemma 2.3 provides us with:

- diffeomorphisms $T_{i,j}$ of $T_{p_i}M$, whose restriction to $B(z_i + 10jd\varepsilon e^s, 2d\varepsilon)$ coincides with the translation by $\alpha\eta d\varepsilon w_j$,
- diffeomorphisms $T_{i,-j}$ of $T_{p_i}M$, whose restriction to $B(z_i - 10jd\varepsilon e^s, 2d\varepsilon)$ coincides with the translation by $-\alpha\eta d\varepsilon w_j$,

Moreover $DT_{i,\pm j}$ is α -close to the identity and $T_{i,\pm j}$ coincides with the identity outside $B(z_i \pm 10jd\varepsilon e^s, 3d\varepsilon)$.

Since the norm of the orthogonal projection of \mathcal{E}^s to E^c is less than $1/2$, the supports of the $T_{i,j}$ and $T_{i,-j}$ for $j \in \{1, \dots, c\}$, are pairwise disjoint, and also disjoint from $B(z_i, 2d\varepsilon)$. Also the union of the supports is contained in $B(z_i, 100d^2\varepsilon) \subset T_{p_i}M$. Since the balls B_i are disjoint, the composition

of f with all the $\Psi_{i,\pm j} := \varphi_{p_i} \circ T_{i,\pm j} \circ \varphi_{p_i}^{-1}$ as in (5) defines a diffeomorphism g , which is δ -close to f in the C^1 topology. Since the diameters of these balls have been chosen small, g is also σ -close to f in the C^0 -topology.

It remains to check the item (3) of Proposition 1.4.

2.6 | θ -accessibility

We set $\theta = \alpha\eta d$ and check the criterion in Section 2.2.

Lemma 2.4. *If $\varepsilon > 0$ is small enough, then for any diffeomorphism \tilde{g} that belongs to a small C^1 -neighborhood of g and for any bisaturated set $\tilde{\Delta} \subset U$ for \tilde{g} , then the pair $(\tilde{g}, \tilde{\Delta})$ is θ -accessible on each ball $\varphi_{p_i}(B(z_i, 2d\varepsilon))$.*

Proof. Let us describe the holonomies in each ball $B(z_i, 100d^2\varepsilon) \subset T_{p_i}M$. We introduce the affine foliations $\mathcal{F}_0^u, \mathcal{F}_0^s$ of $T_{p_i}M$ by leaves parallel to \mathcal{E}^u and \mathcal{E}^s and the perturbed foliation

$$\mathcal{F}^u := (T_{i,-c} \circ \dots \circ T_{i,-1} \circ T_{i,1} \circ \dots \circ T_{i,c})(\mathcal{F}_0^u).$$

We next introduce flows Φ^s, Φ^u along the leaves of \mathcal{F}_0^s and \mathcal{F}^u in the directions e^s and e^u as in proof of Lemma 2.1. For the linear foliation \mathcal{F}_0^s , Φ^s simply coincides with the linear flow $(x, t) \mapsto x + te^s$. For \mathcal{F}^u , one defines $\Phi^u(x, t)$ by projecting the flow $(x, t) \mapsto x + te^u$ on the leaves of \mathcal{F}^u , along the space $\mathcal{E}^s + E_{p_i}^c$.

Claim. For each $j \in \{1, \dots, c\}$, the composition

$$\Phi_{-10jd\varepsilon}^s \circ \Phi_{-10jd\varepsilon}^u \circ \Phi_{10jd\varepsilon}^s \circ \Phi_{10jd\varepsilon}^u$$

coincides on $B(z_i, 2d\varepsilon)$ with the translation by $\theta\varepsilon w_j$.

Similarly, the composition $\Phi_{10jd\varepsilon}^s \circ \Phi_{-10jd\varepsilon}^u \circ \Phi_{-10jd\varepsilon}^s \circ \Phi_{10jd\varepsilon}^u$ coincides on $B(z_i, 2d\varepsilon)$ with the translation by $-\theta\varepsilon w_j$. \square

Proof. Indeed on $B(z_i, 2d\varepsilon)$ the map $\Phi_{10jd\varepsilon}^u$ coincides with $x \mapsto x + (10jd\varepsilon)e^u$ (by construction the support of the maps $T_{i,j}$ does not intersect $B(z_i, 2d\varepsilon) + \mathbb{R}e^u$). On $B(z_i, 2d\varepsilon) + (10jd\varepsilon)e^u + 10jd\varepsilon e^s$, the map $\Phi_{-10jd\varepsilon}^s$ coincides with the composition of $x \mapsto x - (10jd\varepsilon)e^u$ with the translation by $\alpha\eta d\varepsilon w_j = \theta\varepsilon w_j$. The first part of the claim follows. The second is obtained analogously.

Arguing in a similar way and using the fact that the C^0 size of the perturbations $T_{i,j}$ is smaller than $\frac{\varepsilon}{100d}$, one gets the following.

Claim. For each $j \in \{1, \dots, c\}$ and $s \in [0, 1]$ the compositions

$$\begin{aligned} &\Phi_{-s10jd\varepsilon}^s \circ \Phi_{-s10jd\varepsilon}^u \circ \Phi_{s10jd\varepsilon}^s \circ \Phi_{s10jd\varepsilon}^u \\ &\text{and } \Phi_{s10jd\varepsilon}^s \circ \Phi_{-s10jd\varepsilon}^u \circ \Phi_{-s10jd\varepsilon}^s \circ \Phi_{s10jd\varepsilon}^u \end{aligned}$$

are at a C^0 -distance smaller than $\frac{\varepsilon}{10d}$ from the identity. \square

Let \tilde{g} be a diffeomorphism C^1 -close to g and $\tilde{\Delta} \subset U$ be a bisaturated set for \tilde{g} . At each x in $B(0, 2\rho) \cap \varphi_p^{-1}(\Delta)$, the connected components of the leaves $\varphi_p^{-1}(W_{\tilde{g}}^{s/u}(\varphi_p(x))) \cap B(0, 2\rho)$ containing x define leaves $W_{\tilde{g},loc}^{s/u}(x)$. One can define flows $\tilde{\Phi}^s$ and $\tilde{\Phi}^u$ in $B(0, 2\rho) \cap \varphi_{z_i}^{-1}(\tilde{\Delta})$ as before: $\tilde{\Phi}^u(x, t)$ is obtained by projecting $(x, t) \mapsto x + te^u$ on the leaves $W_{\tilde{g},loc}^u(x)$ along the space $\mathcal{E}^u + E_{p_i}^c$; $\tilde{\Phi}^s(x, t)$ is obtained by projecting $(x, t) \mapsto x + te^s$ on the leaves $W_{\tilde{g},loc}^s(x)$ along the space $\mathcal{E}^s + E_{p_i}^c$.

We then define the map

$$H^j : [-1, 1] \times [0, 1] \times (\varphi_p^{-1}(\Delta) \cap B(z_i, 2d\varepsilon)) \rightarrow \varphi_p^{-1}(\Delta) \cap B(0, 2\rho).$$

For $s \in [-1, 1]$ and x in $\varphi_p^{-1}(\Delta) \cap B(0, 2d\varepsilon)$, the arc $t \mapsto H^j(s, t, x)$ is the concatenation of the four arcs (defined on $[0, 1/4]$)

$$\begin{aligned} t &\mapsto \Phi_{|s|t40jd\varepsilon}^u(x), \\ t &\mapsto \Phi_{st40jd\varepsilon}^s(\Phi_{|s|10jd\varepsilon}^u(x)), \\ t &\mapsto \Phi_{-|s|t40jd\varepsilon}^u(\Phi_{s10jd\varepsilon}^s \circ \Phi_{|s|10jd\varepsilon}^u(x)), \\ t &\mapsto \Phi_{-st40jd\varepsilon}^s(\Phi_{-|s|10jd\varepsilon}^u \circ \Phi_{s10jd\varepsilon}^s \circ \Phi_{|s|10jd\varepsilon}^u(x)). \end{aligned}$$

The items (a) and (b) of the definition of the θ -accessibility hold by construction.

The stable leaves $W_{\tilde{g},loc}^s(x)$ are C^1 -close to the leaves of the foliation \mathcal{F}_0^s . Indeed the J first iterates of g and f coincide on B_i (since $D(D) > J$) so that the tangent spaces of the unstable leaves of g are tangent to the cone field C^s and its preimage by φ_{p_i} is γ -close to the direction \mathcal{E}^s on $B(z_i, 100d^2\varepsilon)$. The same holds for \tilde{g} which is C^1 -close to g .

Similarly the unstable leaves of g and \tilde{g} are tangent to the cone field C^u on $f^{-1}(B_i)$. Hence on each ball $\varphi_{p_i}(B(z_i \pm 10jd\varepsilon^s, 3d\varepsilon))$, they are tangent to the cone field $D\Psi_{i,\pm j}(C^u)$. Their preimages by φ_{p_i} are thus γ -close to the leaves of \mathcal{F}^u on $B(z_i, 100d^2\varepsilon)$. This implies that the trajectories of the flows $\tilde{\Phi}^{s/u}$ and $\tilde{\Phi}^{s/u}$ are C^1 -close. Together with the two previous claims, it gives the items (c) and (d) of the definition of the θ -accessibility.

2.7 | Conclusion of the proof of Proposition 1.4

Let $\Delta \subset U$ be a bisaturated set for g . Let us consider any diffeomorphism \tilde{g} that is C^1 -close to g and any bisaturated set $\tilde{\Delta} \subset U$ for \tilde{g} (in particular, any bisaturated $\tilde{\Delta}$ for \tilde{g} contained in a small neighborhood of Δ). The pair $(\tilde{g}, \tilde{\Delta})$ is θ -accessible on each ball $\varphi_{p_i}(B(z_i, 2d\varepsilon))$ by Lemma 2.4. By Lemma 2.2, the pair $(\tilde{g}, \tilde{\Delta})$ is accessible on each ball $\varphi_{p_i}(B(z_i, \varepsilon))$.

Let D be any c -admissible disk in the family \mathcal{D} , which intersects $\tilde{\Delta}$ at a point z . By Lemma 2.1(3a), there exists $\varphi_{p_i}(B(z_i, \varepsilon))$ and a su-path for \tilde{g} between z and a point $y \in \varphi_{p_i}(B(z_i, \varepsilon))$. By accessibility, $\varphi_{p_i}(B(z_i, \varepsilon)) \subset \tilde{\Delta}$. By Lemma 2.1(3b), any point x in the $\varepsilon/2$ -neighborhood of z in D can be connected by a su-path to a point $y' \in \varphi_{p_i}(B(z_i, \varepsilon))$. By accessibility, $y, y' \in \varphi_{p_i}(B(z_i, \varepsilon))$ belong to a su-path. This shows that any point in the $\varepsilon/2$ -neighborhood of z in D can be connected to z by a su-path. Since D is connected, any two points in D belong to a su-path in $\tilde{\Delta}$, showing that $(\tilde{g}, \tilde{\Delta})$

is accessible on D . This concludes the proof of the stable accessibility of (g, Δ) on any disk $D \in \mathcal{D}$, and of the last item of Proposition 1.4.

3 | PROOF OF PROPOSITION 1.5

For Λ a partially hyperbolic set for f and U a neighborhood of Λ with $\bar{U} \subset U_0(f, \Lambda)$, we denote the maximal f -invariant set in \bar{U} by

$$\Lambda_U(f) := \bigcap_{j \in \mathbb{Z}} f^j(\bar{U}).$$

For $J \geq 1$, denote by $\text{Per}_{\leq J}(f, U)$ the set of points

$$\text{Per}_{\leq J}(f, U) := \{p \in \bar{U} \mid \text{per}(p) \leq J\}.$$

3.1 | Existence of c -admissible families

We restate [12, Lemma 2.3] in our setting.

Lemma 3.1. *Let Λ be a partially hyperbolic set for f , let $\beta \in (0, 1)$, let U be neighborhood of Λ with $\bar{U} \subset U_0(f, \Lambda)$ and let $J \geq 1$.*

Then for every $\rho > 0$ sufficiently small and denoting by B_ρ the ρ -neighborhood of $\text{Per}_{\leq J}(f, U)$, there exist $q_1, \dots, q_k \in \Lambda_U(f) \setminus B_\rho$ such that

- (1) *the balls $B_{\beta\rho}(q_1), \dots, B_{\beta\rho}(q_k)$ cover $\Lambda_U(f) \setminus B_\rho$,*
- (2) *$V_\rho(q_i) \cap V_\rho(q_j) = \emptyset$, for all $i \neq j$, and*
- (3) *$V_\rho(q_i) \cap f^m(V_\rho(q_j)) = \emptyset$, for all i, j and $0 < |m| \leq J$.*

In particular $\mathcal{D} := \{V_\rho(q_1), \dots, V_\rho(q_k)\}$ is a c -admissible disk family with respect to $(f, \Lambda_U(f))$ satisfying $R(\mathcal{D}) > J$.

Remark. There exists $\sigma > 0$ such that for any diffeomorphism g with $d_{C^0}(g, f) < \sigma$, the set $\Lambda_U(g)$ is contained in an arbitrarily small neighborhood of $\Lambda_U(f)$ so that we still have $\Lambda_U(g) \setminus B_\rho \subset \cup_i B_{\beta\rho}(q_i)$.

3.2 | From c -admissible families to stable c -sections

The conclusions of Proposition 1.5 follow from the next lemma, choosing $\rho < J^{-1}$.

Lemma 3.2. *Given a partially hyperbolic set Λ for f , there exist $\delta, \beta > 0$ with the following property.*

Let U be neighborhood of Λ with $\bar{U} \subset U_0(f, \Lambda)$, let $J \geq 1$ and, given $\rho > 0$ sufficiently small, let \mathcal{D} be the c -admissible disk family given by Lemma 3.1.

Then there exists $\sigma > 0$ such that

- for any diffeomorphism g satisfying $d_{C^1}(f, g) < \delta$ and $d_{C^0}(f, g) < \sigma$,
- for any bisaturated partially hyperbolic set $\Delta \subset U$ for g ,

the set $|D|$ is a c -section for g .

Proof. Consider f, Λ, U, J satisfying the hypotheses. The set $\text{Per}_{\leq J}(f, U)$ is compact, partially hyperbolic and satisfies for every $p \in \text{Per}_{\leq J}(f, U)$:

$$(\mathcal{W}^s(p) \cup \mathcal{W}^u(p)) \cap \text{Per}_{\leq J}(f, U) = \{p\}. \tag{7}$$

The following property follows easily from (7) by a compactness argument.

Claim. There exist $\rho_0, \delta_0 > 0$ such that the ρ_0 -neighborhood B_0 of $\text{Per}_{\leq J}(f, U)$ has the following properties. For every diffeomorphism g with $d_{C^1}(f, g) < \delta_0$, every $p \in B_0 \cap \Lambda_U(g)$ can be connected to a point in $\bar{U} \setminus B_0$ by an su -path for g with one leg. □

There is also a projection onto admissible discs by su -paths.

Claim. There exist $\beta \in (0, 1)$ and $\rho_1, \delta_1 > 0$ such that for every diffeomorphism g with $d_{C^1}(f, g) < \delta_1$, every $\rho \in (0, \rho_1)$, every bisaturated set $\Delta \subset U$ for g , every $q \in \Lambda_U(g)$ and $p \in B_{\beta\rho}(q) \cap \Delta$, there is an su -path for g with 2 legs from p to some point in $V_\rho(q)$.

Proof. The proof follows the same argument as in the proofs of Lemmas 2.1 and 2.2. Let $B^u(0, 1)$ and $B^s(0, 1)$ be the unit balls in the spaces E_q^u and E_q^s . Working in the chart ϕ_q , define a continuous map Φ from $B^u(0, 1) \times B^s(0, 1)$ to $\varphi_q^{-1}(\Delta) \subset T_q M$ in the following way. For $v, w \in B^u(0, 1) \times B^s(0, 1)$, first project orthogonally along the space $E_q^c \oplus E_q^s$ the point $\varphi_q^{-1}(p) + \rho v$ onto the unstable leaf of $\varphi_q^{-1}(p)$ lifted in the chart: this defines a point $y \in \varphi_q^{-1}(\Delta)$. Then project orthogonally along the space $E_q^u \oplus E_q^c$ the point $y + \rho w$ onto the stable leaf of y in the chart: this defines the point $\varphi_p^{-1}(\Phi(v, w))$. It belongs to the bisaturated set Δ .

The restriction of Φ to the boundary of $B^u(0, 1) \times B^s(0, 1)$ is disjoint from E_q^c , and the Brouwer fixed-point theorem ensures that the image of $\varphi_p^{-1} \circ \Phi$ intersects E_q^c . Since the stable and unstable leaves of g lifted in the chart are close to the directions E_q^u and E_q^s , the intersection point belongs to $B^c(0, \rho) \subset E_q^c$.

Fix $\delta = \min(\delta_0, \delta_1)$, $\rho < \min\{\rho_0, \rho_1\}$ and let \mathcal{D} be the c -admissible disk family with centers $q_1, \dots, q_k \in \Lambda_U(f) \setminus B_\rho$ given by Lemma 3.1. Let $\sigma > 0$ be given by the remark following Lemma 3.1. Consider a diffeomorphism g satisfying $d_{C^1}(f, g) < \delta$ and $d_{C^0}(f, g) < \sigma$, and let $\Delta \subset U$ be a bisaturated partially hyperbolic set for g .

Claim 3.2 gives that every $p \in B_\rho \cap \Delta$ can be connected to a point in $\Lambda_U(g) \setminus B_\rho$ by an su -path for g . Since Δ is bisaturated, it intersects $\Lambda_U(g) \setminus B_\rho$.

By Lemma 3.1 and the remark that follows, the balls $B_{\beta\rho}(q_1), \dots, B_{\beta\rho}(q_k)$ cover $\Lambda_U(g) \setminus B_\rho$. Hence, there exists i such that $\Delta \cap B_{\beta\rho}(q_i) \neq \emptyset$. Now the Claim 3.2 and the bisaturation of Δ imply that $\Delta \cap V_\rho(q_i) \neq \emptyset$.

We have thus shown that Δ intersects $|D|$, as required. □

ACKNOWLEDGEMENTS

We thank Rohil Prasad for helpful comments on a first version of the text. A.A. was supported by the [SNSF](#) grant “Dynamical Systems.” S.C. was partially supported by the [ERC](#) project 692925 NUHGD. A.W. was supported by [NSF](#) grant DMS-1402852.

JOURNAL INFORMATION

The *Proceedings of the London Mathematical Society* is wholly owned and managed by the London Mathematical Society, a not-for-profit Charity registered with the UK Charity Commission. All surplus income from its publishing programme is used to support mathematicians and mathematics research in the form of research grants, conference grants, prizes, initiatives for early career researchers and the promotion of mathematics.

REFERENCES

1. F. Abdenur, C. Bonatti, and L. J. Díaz, *Non-wandering sets with non-empty interiors*, *Nonlinearity* **17** (2004), 175–191.
2. M.-C. Arnaud, C. Bonatti, and S. Crovisier, *Dynamiques symplectiques génériques*, *Ergodic Theory Dynam. Systems* **25** (2005), 1401–1436.
3. A. Avila and J. Bochi, *Nonuniform hyperbolicity, global dominated splittings and generic properties of volume-preserving diffeomorphisms*, *Trans. Amer. Math. Soc.* **364** (2012), 2883–2907.
4. A. Avila, J. Bochi, and A. Wilkinson, *Nonuniform center bunching and the genericity of ergodicity among C^1 partially hyperbolic symplectomorphisms*, *Ann. Sci. Éc. Norm. Supér.* **42** (2009), 931–979.
5. A. Avila, S. Crovisier, and A. Wilkinson, *Diffeomorphisms with positive metric entropy*, *Publ. Math. Inst. Hautes Études Sci.* **124** (2016), 319–347.
6. A. Avila, S. Crovisier, and A. Wilkinson, *C^1 density of stable ergodicity*, *Adv. Math.* **379** (2021), 107496, 68 pp.
7. J. Bochi, *Genericity of zero Lyapunov exponents*, *Ergodic Theory Dynam. Systems* **22** (2002), no. 6, 1667–1696.
8. J. Bochi, *C^1 -generic symplectic diffeomorphisms: partial hyperbolicity and zero centre Lyapunov exponents*, *J. Inst. Math. Jussieu* **9** (2010), 49–93.
9. C. Bonatti, L. Díaz, and M. Viana, *Dynamics beyond uniform hyperbolicity: a global geometric and probabilistic perspective*, *Encyclopaedia of Mathematical Sciences*, vol. 102, Springer, Berlin, 2005.
10. R. Bowen, *A horseshoe with positive measure*, *Invent. Math.* **29** (1975), 203–204.
11. R. Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*, *Lecture Notes in Mathematics*, vol. 470, Springer, Berlin, 1975.
12. D. Dolgopyat and A. Wilkinson, *Stable accessibility is C^1 dense, geometric methods in dynamics. II*, *Astérisque* **287** (2003), 33–60.
13. A. Kisilev, *Hofer growth of C^1 -generic Hamiltonian flows*, *Comment. Math. Helv.* **91** (2016), 21–38.
14. R. Mañé, *Oseledec’s theorem from the generic viewpoint*, *Proceedings of International Congress of Mathematicians (Warszawa 1983)*, vol. 2, pp. 1259–1276.
15. J. Moser, *On volume elements on manifolds*, *Trans. Amer. Math. Soc.* **120** (1965), 285–294.
16. C. Pugh and M. Shub, *Stable ergodicity and julienne quasi-conformality*, *J. Eur. Math. Soc.* **2** (2000), 1–52.
17. J. Rodríguez-Hertz, *Genericity of nonuniform hyperbolicity in dimension 3*, *J. Mod. Dyn.* **6** (2012), 121–138.
18. A. Tahzibi, *Stably ergodic diffeomorphisms which are not partially hyperbolic*, *Israel J. Math.* **142** (2004), 315–344.
19. P. Zhang, *Partially hyperbolic sets with positive measure and ACIP for partially hyperbolic systems*, *Discrete Contin. Dyn. Syst.* **32** (2012), 1435–1447.