



US 20210079063A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2021/0079063 A1**

KOIDE et al. (43) **Pub. Date: Mar. 18, 2021**

(54) **METHODS AND COMPOSITION INVOLVING THERMOPHILIC FIBRONECTIN TYPE III (FN3) MONOBODIES**

Publication Classification

(51) **Int. Cl.**
C07K 14/78 (2006.01)
C40B 40/10 (2006.01)
(52) **U.S. Cl.**
CPC *C07K 14/78* (2013.01); *C40B 40/10* (2013.01)

(71) Applicant: **The University of Chicago**, Chicago, IL (US)

(72) Inventors: **Shohei KOIDE**, Chicago, IL (US);
Shun-Ichi TANAKA, Chicago, IL (US); **Akiko KOIDE**, Chicago, IL (US)

(57) **ABSTRACT**

(21) Appl. No.: **15/733,530**

(22) PCT Filed: **Feb. 21, 2019**

(86) PCT No.: **PCT/US2019/018866**

§ 371 (c)(1),

(2) Date: **Aug. 21, 2020**

Related U.S. Application Data

(60) Provisional application No. 62/634,616, filed on Feb. 23, 2018.

The current application describes various compositions and methods for the production of FN3-based binding proteins with improved stability properties. Aspects of the disclosure relate to polypeptides comprising a variant fibronectin type III (FN3) domain from *Sulfolobus tokodaii* or *Pyrococcus horikoshii* comprising one or more amino acid substitutions or insertions in a loop region of FN3, in a non-loop region of FN3, or in both.

Specification includes a Sequence Listing.

(a)


STOFN3-1	315	PP--PKPQIASIASGNE--TITVKWYDTNASG---YYITYWSNFSQKVT	356
STOFN3-2	399	PSSVPNPPIIKVKIGNL--NATLTWYDTFNGG---YPIEGYYLYVNGKG	442
STOFN3-3	488	-----ANLTVTVYKIN--GFLVSWNSTSKAK---YIL--TVSKENVVL	524
STOFN3-4	568	----PASPTVNWSITLN--TVSLNWSKVSAGAE---Y----YLIYDNGKL	603
PDB ID: 2DJU	1	----PKPPIDLVTETTATSVTLTWDSGNSEPVYYGIQYRAAGTEGPF	45




STOFN3-1	357	INVGNV--TSYTIKHLKDGVTYYIQIVPYNSLGNGT-PSDIISAT---	398
STOFN3-2	443	INVGNI--TSYVLTNLTAGELYTIELIAYNKIGNSS-ISSVSFIAASK	484
STOFN3-3	525	LNVSTT--NTSYFVKVPEGV-YNISLEAVNIVGITKYAFILIIYYIQ--	566
STOFN3-4	604	ITNTTN--TAFTFN-LTIGQN-EIEVYAANAYYKSA-PYIINDVR---	643
PDB ID: 2DJU	46	QEV DGVATTRYSIGGLSPFSEYAFRVLAVNSIGRGPPEAVRARTGEQ	106



(b)



(c)






FIG. 1A-C




FIG. 2A-C




FIG. 3A-D




FIG. 4A

(b)

	AB	BC	CD	DE	EF	FG	T_M (°C)
STOFN3-1	ASGNETIT	WYDTNASG	WSNFSQK	VGNVTSY	IKHLKDGVT	PYNSLGNGTPSDII	80.0
STOFN3-1/ABSer	ASSSSSIT	WYDTNASG	WSNFSQK	VGNVTSY	IKHLKDGVT	PYNSLGNGTPSDII	79.0
STOFN3-1/BCSer	ASGNETIT	WYSSSSSG	WSNFSQK	VGNVTSY	IKHLKDGVT	PYNSLGNGTPSDII	74.7
STOFN3-1/CDSer	ASGNETIT	WYDTNASG	WSSSSQK	VGNVTSY	IKHLKDGVT	PYNSLGNGTPSDII	82.2
STOFN3-1/DESer1	ASGNETIT	WYDTNASG	WSNFSQK	VGSSSSY	IKHLKDGVT	PYNSLGNGTPSDII	81.7
STOFN3-1/DESer2	ASGNETIT	WYDTNASG	WSNFSQK	VSSSSSY	IKHLKDGVT	PYNSLGNGTPSDII	54.3
STOFN3-1/EFSer	ASGNETIT	WYDTNASG	WSNFSQK	VGNVTSY	IKSSSSSVT	PYNSLGNGTPSDII	64.0
STOFN3-1/FGSer1	ASGNETIT	WYDTNASG	WSNFSQK	VGNVTSY	IKHLKDGVT	PYSSSSSSSSPSDII	66.7
STOFN3-1/FGSer2	ASGNETIT	WYDTNASG	WSNFSQK	VGNVTSY	IKHLKDGVT	PSSSSSSSSSSPSDII	60.0
STOFN3-1/FGSer3	ASGNETIT	WYDTNASG	WSNFSQK	VGNVTSY	IKHLKDGVT	PYSSSSSSSSSSSDII	63.2
STOFN3-1/FGSer4	ASGNETIT	WYDTNASG	WSNFSQK	VGNVTSY	IKHLKDGVT	PSSSSSSSSSSSSSII	62.0
STOTEMP3*	ASGPETIT	WYDTNASG	WSSSSQK	VGNVTSY	IKHLKPGVT	PYNSLGNGTPSDII	>95.0
STOTEMP4*	ASGPETIT	WYDTNASY	WSSSSQK	VGNVTSY	IKHLKPGVT	PVNSLGNGTPSDII	>95.0
STOTEMP4*/FGSer5	ASGPETIT	WYDTNASY	WSSSSQK	VGNVTSY	IKHLKPGVT	PVSSSSSSSSPSDII	89.3
STOTEMP4*/FGSer6	ASGPETIT	WYDTNASY	WSSSSQK	VGNVTSY	IKHLKPGVT	PYSSSSSSSSPSDII	85.8
STOTEMP4*/FGSer7	ASGPETIT	WYDTNASY	WSSSSQK	VGNVTSY	IKHLKPGVT	PSSSSSSSSSSPSDII	86.5

*C-terminal extented.

FIG. 4B

(c)

	BC	CD	DE	FG	T_M (°C)
STOFN3-1	WYDT-----NASG	WSN-----FSQ	VGN-----VTSY	PYNS-----LGNGTPSDII	80.0
STOFN3-1/BC+2	WYDTSS-----NASG	WSN-----FSQ	VGN-----VTSY	PYNS-----LGNGTPSDII	80.3
STOFN3-1/BC+4	WYDTSSSS---NASG	WSN-----FSQ	VGN-----VTSY	PYNS-----LGNGTPSDII	76.7
STOFN3-1/BC+8	WYDTSSSSSSSSSNASG	WSN-----FSQ	VGN-----VTSY	PYNS-----LGNGTPSDII	77.7
STOFN3-1/CD+2	WYDT-----NASG	WSNSS-----FSQ	VGN-----VTSY	PYNS-----LGNGTPSDII	71.7
STOFN3-1/CD+4	WYDT-----NASG	WSNSSSS---FSQ	VGN-----VTSY	PYNS-----LGNGTPSDII	70.0
STOFN3-1/CD+8	WYDT-----NASG	WSNSSSSSSSSFSQ	VGN-----VTSY	PYNS-----LGNGTPSDII	67.7
STOFN3-1/DE+2	WYDT-----NASG	WSN-----FSQ	VGNSS-----VTSY	PYNS-----LGNGTPSDII	55.8
STOFN3-1/DE+4	WYDT-----NASG	WSN-----FSQ	VGNSSSS---VTSY	PYNS-----LGNGTPSDII	55.0
STOFN3-1/DE+8	WYDT-----NASG	WSN-----FSQ	VGNSSSSSSSSVTSY	PYNS-----LGNGTPSDII	54.8
STOFN3-1/FG+2	WYDT-----NASG	WSN-----FSQ	VGN-----VTSY	PYNSSS-----LGNGTPSDII	68.2
STOFN3-1/FG+4	WYDT-----NASG	WSN-----FSQ	VGN-----VTSY	PYNSSSSS---LGNGTPSDII	68.0
STOFN3-1/FG+8	WYDT-----NASG	WSN-----FSQ	VGN-----VTSY	PYNSSSSSSSSSSLGNGTPSDII	67.2

FIG. 4C






FIG. 5



Target	Clone	K_D (nM)	Amino acid sequence									
			315	320	330	BC	350	DE	370	380	FG	400
	Library	-	sSSSVQIASIASGPETITVKWY (X_{3-9}) ---- YYYITWSSSSQKVIINVG (X_3) SYTIKHLKPGVYYYIQIVP (X_{8-12}) -----PSDIIATPSSs									
ySUMO	SUMOMbL03	63 ±18	sSSSVQIASIASGPETITVKWYDTNAS----YYYITWSSSSQKVTINVGKRVSYTIKHLKPGVYYYIQIVPYYEPSEDFWGI PSDIIATPSSs									
ySUMO	SUMOMbL14	44 ±15	sSSSVQIASIASGPETITVKWYDTNAS----YYYITWSSSSQKVTINVGSPISYTIKHLKPGVYYYIQIVPYYITIGPEYYYI--PSDIIATPSSs									
MBP	MBPMbL08	12 ±3	sSSSVQIASIASGPETITVKWYSSYYWSPFYIITWSSSSQKVTINVGNVTSYTIKHLKPGVYYYIQIVPYYQYFPRYGY--PSDIIATPSSs									
MBP	MBPMbL17	41±14	sSSSVQIASIAVGPETITVKWYRI PHAY---YYYITWSSSSQKVTINVGNVTSYTIKHLKPGVYYYIQIVPYYPSWYLSYGY--PSDIIATPSSs									
EGFP	EGFPMbL24	58 ±9	sSSSVQIASIASGPETITVKWYHHSNY---YYYITWSSSSQKVTINVGNVTSYTIKHLKPGVYYYIQIVPDAGGYDKG----PSDIIATPSSs									
EGFP	EGFPMbL26	23 ±7	sSSSVQIASIASGPETITVKWYDATYTVI--YYYITWSSSSQKVTINVGNVTSYTIKHLKPGVYYYIQIVPESGWYSRW----PSDIIATPSSs									
AbSH2	AbSH2MbL03	105 ±11	sSSSVQIASIASGPETITVKWYFWAYGD---YYYITWSSSSQKVTINVGRRYSYTIKHLKPGVYYYIQIVPVNSLNGMT----PSDIIATPSSs									
AbSH2	AbSH2MbL04	78 ±18	sSSSVQIASIASGPETITVKWYYSIDV---YYYITWSSSSQKVTINVGQPTSYTIKHLKPGVYYYIQIVPYSFYSDSDYGY--PSDIIATPSSs									
NSH2	NSH2MbL06	81 ±24	sSSSVQIASIASGPETITVKWYPTGY----YYYITWSSSSQKVTINVGYNRSYTIKHLKPGVYYYIQIVPYYLSKSGATP--PSDIIATPSSs									
NSH2	NSH2MbL10	158 ±54	sSSSVQIASIASGPETITVKWYPTGY----YYYITWSSSSQKVTINVGYNRSYTIKHLKPGVYYYIQIVPASYVWVYDT----PSDIIATPSSs									
CSH2	CSH2MbL01	67 ±17	sSSSVQIASIASGPETITVKWYDTNAS----YYYITWSSSSQKVTINVGNRHSYTIKHLKPGVYYYIQIVPGWADYAWLWY--PSDIIATPSSs									
CSH2	CSH2MbL03	103 ±53	sSSSVQIASIASGPETITVKWYDTNAS----YYYITWSSSSQKVTINVGNRYSYTIKHLKPGVYYYIQIVPGWSSSETYTWY--PSDIIATPSSs									



Target	Clone	K_D (nM)	Amino acid sequence									
			315	320	330	β C	CD	β D	360	370	380	FG
	Library	-	sSSSVQIASIASGPETITVKWYDTNASYYYIYBYUS (X_{3-3}) ZZVZIZVGNVTSYTIKHLKPGVYYYIQIVP (X_{8-12}) -----PSDIIATPSSs									
ySUMO	SUMOMbS08	65 ±16	sSSSVQIASIASGPETITVKWYDTNASYYYIYWSLQYREAVKIEVGNVTSYTIKHLKPGVYYYIQIVPFEEFPGFWYSGDPSDIIATPSSs									
ySUMO	SUMOMbS20	95 ±26	sSSSVQIASIAVGPETITVKWYDTNASYYYIYWSKHP--NKVKIEVGNVTSYTIKHLKPGVYYYIQIVPFISYYDLYYSGDPSDIIATPSSs									
ySUMO	SUMOMbS34	20 ±8	sSSSVQIASIASGPETITVKWYDTNASYYYIYWSGPPYKVTIIVGNVTSYTIKHLKPGVYYYIQIVPVHVEDWGYYYY--PSDIIATPSSs									
MBP	MBPMbS01	33 ±13	sSSSVQIASIASGPETITVKWYDTNASYYYISYWSADYDSEAVAIAVGNVTSYTIKHLKPGVYYYIQIVPYKANSYYPGYGYPDIIATPSSs									
MBP	MBPMbS09	9 ±1	sSSSVQIASIASGPETITVKWYDTNASYYYISYWSYST--ATVAIAVGNVTSYTIKHLKPGVYYYIQIVPYYGYRFPYPSYGYPSDIIATPSSs									
MBP	MBPMbS12	484 ±66	sSSSVQIASIASGPETITVKWYDTNASYYYISYWSGQS--EIVAEVGNVTSYTIKHLKPGVYYYIQIVPYYPGYYPVHYV--PSDIIATPSSs									
EGFP	EGFPMbS04	7 ±2	sSSSVQIASIAVGPETITVKWYDTNASYYYIDYWSHGE--EIVAEVGNVTSYTIKHLKPGVYYYIQIVPYYGDLGLG---PSDIIATPSSs									
EGFP	EGFPMbS05	8 ±1	sSSSVQIASIASGPETITVKWYDTNASYYYIDYWSYSK--EIVTIEVGNVTSYTIKHLKPGVYYYIQIVPYYGSLGQ---PSDIIATPSSs									
EGFP	EGFPMbS12	4 ±2	sSSSVQIASIASGPETITVKWYDTNASYYYIDYWSYGT--EIVTINVGNTSYTIKHLKPGVYYYIQIVPYYGVLEHQ---PSDIIATPSSs									
AbSH2	AbSH2MbS01	71 ±10	sSSSVQIASIASGPETITVKWYDTNASYYYIYWSYIWPVSEVEIVGNVTSYTIKHLKPGVYYYIQIVPVNSLNGMT----PSDIIATPSSs									
AbSH2	AbSH2MbS02	13 ±6	sSSSVQIASIASGPETITVKWYDTNASYYYIYWSYIWPVTEVEIVGNVTSYTIKHLKPGVYYYIQIVPQYWYLGAGY---PSDIIATPSSs									
AbSH2	AbSH2MbS07	827 ±140	sSSSVQIASIASGPETITVKWYDTNASYYYIYWSYIWPVTEVEIVGNVTSYTIKHLKPGVYYYIQIVPGYYSYFYLS--PSDIIATPSSs									

FIG. 6A-B




FIG. 6C




FIG. 7A

(b)




FIG. 7B

(c)

	Oligomerization state	Thermal stability (T_M , °C)
STOTEMP5	Monomeric	>95.0
SUMOMbL03	Monomeric	63.5
SUMOMbL14	Monomeric	63.5
MBPMbL08	Large aggregate	-
MBPMbL17	Late elution*	73.3
EGFPMbL24	Monomeric	54.2
EGFPMbL26	Late elution*	53.5
AbISH2MbL03	Monomeric	74.0
AbISH2MbL04	Monomeric	69.5
NSH2MbL06	Monomeric	86.3
NSH2MbL10	Monomeric	80.5
CSH2MbL01	Late elution*	64.8
CSH2MbL03	Late elution*	66.5
	Mean	68.1 ±10.1
SUMOMbS08	Large aggregate	-
SUMOMbS20	Monomeric	53.5
SUMOMbS34	Large aggregate	-
MBPMbS01	Late elution*	53.7
MBPMbS09	Late elution*	53.2
MBPMbS12	Monomeric	53.5
EGFPMbS04	Monomeric	85.5
EGFPMbS05	Monomeric	70.2
EGFPMbS12	Monomeric	66.0
AbISH2MbS01	Monomeric	71.0
AbISH2MbS02	Monomeric	53.5
AbISH2MbS07	Monomeric	50.5
	Mean	61.1 ±11.6


*Mono-dispersed single peak

FIG. 7C

(a)

Clone	K_D (nM)	Amino acid sequece
Clamp(Ptpn11_pY580) cpGrb2 SH2 Monobody	2.5*	VKFNSLNELVDYHRSTSVSRNQIFLRDIGGGGGHPWYKGIKIPRAKAEMLSKQRHDGAFLIRESEAPGDFSLSVKFGNDVQHEKVLKRDGAGKYFLWV GGSGGSVSSVPTKLEVVAATPTSLLSWD AW SGSDWPVSYRITYYGE TGGNSPVQEFYVPGSSYTATISGLSPGVDTITIVYAGYDGKYYYQSPI-SINVRT
STOTEMP4		SSSKPQIASI-ASGPETITVKWYD-----TNASYYYITYWSSSS--QKVTINVG-NVTSYTIKHLKPGVTYYYIQIVPVNSLNG--TPSDIISATPSSs
ClampSTOTEMP4_L0	-	GSSKPQIASI-ASGPETITVKWYAWSGSDWPVSYRITYWSSSS--QKVTINVG-NVTSYTIKHLKPGVTYYYIQIYAGYDGKYYYQTPSDIISATPSSs
ClampSTOTEMP4_L2	-	GSSSSKPQIASI-ASGPETITVKWYAWSGSDWPVSYRITYWSSSS--QKVTINVG-NVTSYTIKHLKPGVTYYYIQIYAGYDGKYYYQTPSDIISATPSSs
ClampSTOTEMP4_L5	-	GGSGSSSKPQIASI-ASGPETITVKWYAWSGSDWPVSYRITYWSSSS--QKVTINVG-NVTSYTIKHLKPGVTYYYIQIYAGYDGKYYYQTPSDIISATPSSs
ClampSTOTEMP5_L0	>2000	GSSVPQIASI-ASGPETITVKWYAWSGSDWPVSYRITYWSSSS--QKVTINVG-NVTSYTIKHLKPGVTYYYIQIYAGYDGKYYYQTPSDIISATPSSs
ClampSTOTEMP5_L2	156 ±42	GSSSSVPQIASI-ASGPETITVKWYAWSGSDWPVSYRITYWSSSS--QKVTINVG-NVTSYTIKHLKPGVTYYYIQIYAGYDGKYYYQTPSDIISATPSSs
ClampSTOTEMP5_L5	121 ±21	GGSGSSSVVPQIASI-ASGPETITVKWYAWSGSDWPVSYRITYWSSSS--QKVTINVG-NVTSYTIKHLKPGVTYYYIQIYAGYDGKYYYQTPSDIISATPSSs

(b)



(c)




FIG. 8A-C




FIG. 9





FIG. 10



strand




FIG. 12




FIG. 13

**METHODS AND COMPOSITION INVOLVING
THERMOPHILIC FIBRONECTIN TYPE III
(FN3) MONOBODIES**

CROSS-REFERENCE TO RELATED
APPLICATIONS

[0001] This application claims the benefit of priority of U.S. Provisional Patent Application No. 62/634,616 filed Feb. 23, 2018, which is hereby incorporated by reference in its entirety.

STATEMENT OF GOVERNMENT SUPPORT

[0002] This invention was made with government support under GM090324 awarded by the National Institutes of Health. The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

II. Field of the Invention

[0003] Embodiments are directed generally to biology, medicine, and protein engineering.

III. Background

[0004] The fibronectin type III domain (FN3) has been established as an effective non-antibody, "alternative" scaffold for the generation of novel binding proteins. A member of the immunoglobulin superfamily, FN3 has three surface exposed loops at one end of the molecule which are analogous to antibody CDRs. Engineering strategies using this scaffold are based on combinatorial libraries created by diversifying both the length and amino acid sequence of these surface loops. From such libraries, FN3 variants capable of binding to a target of interest can be isolated using various selection methods. The utility of the FN3 scaffold has been demonstrated in producing high-affinity binding proteins to a number of different protein targets (reviewed in Bloom and Calabro, 2009). These binding proteins generated from this scaffold are referred to as monobodies. The FN3 scaffold offers many advantages compared to conventional antibodies or fragments thereof because it lacks disulfide bonds, can be readily and highly expressed in bacterial systems, and is relatively small.

[0005] The FN3 scaffold has produced high-affinity binding proteins to a number of distinct targets, achieving dissociation constants in the low- to mid-nanomolar range and as low as 1.1 picomolar (Bloom and Calabro, 2009; Hackel et al., 2008). Binding proteins based on the FN3 domain have also been used in a number of applications including, conformation specific purification of a target protein (Huang et al., 2006), probing protein conformational changes in cells (Koide et al., 2002), antagonism of a growth factor receptor for therapeutic use (Getmanova et al., 2006), inhibition of virus replication (Liao et al., 2009), and specifically inhibiting a protein-protein interaction in cells (Wojcik et al., 2010).

[0006] The ability to generate novel binding proteins capable of interacting with other proteins with high-affinity and/or specificity is important in biotechnology, medicine and molecular biology. Such designer binding proteins can be used in numerous applications. They can be used to label a protein of interest for detection and visualization, to purify a target protein from a complex mixture or to functionally perturb a target by blocking a functional site. Because of

their broad utility, there is a need to develop strategies for producing novel binding proteins quickly and effectively and with improved stability properties. To date, FN3-based binding proteins have been constructed by selections from combinatorial libraries in which loop regions are diversified (Bloom and Calabro, 2009).

SUMMARY OF THE INVENTION

[0007] The present application describes various compositions and methods for the production and use of FN3-based binding proteins with improved stability properties. Aspects of the disclosure relate to polypeptides comprising a variant fibronectin type III (FN3) domain from *Sulfolobus tokodaii* that includes one or more amino acid substitutions or insertions in a loop region of FN3, in a non-loop region of FN3, or in both.

[0008] In some embodiments, a polypeptide comprises a non-variant FN3 domain. In particular embodiments, the non-variant FN3 domain comprises at least 70% identity to SEQ ID NO:1. In some embodiments, the non-variant FN3 domain comprises SEQ ID NO:1. In further embodiments, the non-variant FN3 domain comprises a polypeptide with at least 70% identity to SEQ ID NO:2 (STOFN3-2), 3 (STOFN3-3), or 4 (STOFN3-4). In some embodiments, the non-variant FN3 domain comprises SEQ ID NO:2, 3, or 4. In some embodiments, the amino acid substitution comprises one or more substitutions of the FN3 domain corresponding to amino acid positions 1, 2, and 3 of SEQ ID NO:1. In some embodiments, the amino acid substitutions at one or more positions corresponding to amino acid positions 1, 2, and/or 3 of SEQ ID NO:1 are a substitution with a hydrophilic amino acid or combinations thereof. In some embodiments, the amino acid insertion comprises one or more amino acid insertions of the FN3 domain after the amino acid corresponding to amino acid position 84 of SEQ ID NO:1. In some embodiments, the amino acid insertion comprises 1-3 amino acids. In some embodiments, the inserted amino acids are proline and serine or combinations thereof. In some embodiments, the substitution comprises one or more substitutions of the FN3 domain corresponding to amino acid positions 4, 14, 28, and 58 of SEQ ID NO:1. In some embodiments, the substitution corresponding to amino acid position 4 of SEQ ID NO:1 is with a valine. In some embodiments, the substitution corresponding to amino acid position 14 of SEQ ID NO:1 is with a proline. In some embodiments, the substitution corresponding to amino acid position 28 of SEQ ID NO:1 is with a tyrosine. In some embodiments, the substitution corresponding to amino acid position 58 of SEQ ID NO:1 is with a proline. In some embodiments, the one or more amino acid substitutions or insertions in the non-loop segment is in one or more of beta strand C or beta strand D. In some embodiments, the one or more amino acid substitutions or insertions correspond to or after positions 28, 29, 30, 31, 32, 33, 34, or 35 of SEQ ID NO:1. In some embodiments, the one or more amino acid substitutions or insertions correspond to or after positions 38, 39, 40, 41, 42, 43, 44, 45, or 46 of SEQ ID NO:1. In some embodiments, the one or more amino acid substitutions or insertions in the loop region of FN3 are in the BC loop, DE loop, FG loop and/or CD loop. In some embodiments, the DE loop does not comprise any inserted amino acids. In some embodiments, the EF loop comprises less than 3 substitutions or insertions. In some embodiments, the FN3 domain does not have substitutions at amino acid

positions corresponding to positions 69 and 77 of SEQ ID NO:1. In some embodiments, the polypeptide further comprises a non-FN3 polypeptide that enhances the FN3 polypeptide binding affinity for a target molecule. In some embodiments, the polypeptide comprises or further comprises an insertion or deletion of at least 1 amino acids in at least one loop region of FN3. In some embodiments, at least one loop region of FN3 comprises an insertion of at least 2 amino acids. In some embodiments, at least one region of FN3 comprises an insertion of 2 to 25 amino acids in at least one loop region. In some embodiments, at least two loop regions comprise an insertion. In some embodiments, at least one loop region of FN3 comprises a deletion of at least 1 amino acid. In some embodiments, at least one loop region of FN3 comprises a deletion of 2 to 10 amino acids. In some embodiments, at least two loop regions comprise a deletion of at least 1 amino acid. In some embodiments, the polypeptide comprises at least 1 amino acid insertion and 1 amino acid deletion in at least one loop region. In some embodiments, the polypeptide comprises an insertion and deletion of at least 1 amino acid in the same loop region. In some embodiments, the polypeptide further comprises one or more serine residues immediately before amino acid 1 or immediately after amino acid 84 of SEQ ID NO:1.

[0009] In certain aspects, the insertion, deletion, or substitution, when described with respect to an amino acid at a position in SEQ ID NO:1 may be the corresponding amino acid at a corresponding position in SEQ ID NO:2, 3, and/or 4. The corresponding positions are those that are aligned with the amino acid in SEQ ID NO:1 in FIG. 1.

[0010] In some embodiments, the AB, BC, CD, DE, EF, and/or FG loop or A, B, C, D, E, F, or G beta-strand may have at least, at most, or exactly 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 amino acid substitutions, insertions and/or deletions (or combinations thereof). In certain aspects, beta strand variations can be used in conjunction with variations in the AB loop, the BC loop, the CD loop, the DE loop, and/or the FG loop of FN3 to generate a polypeptide library or a nucleic acid library encoding the same. FN3 polypeptides can be modified by inserting or deleting 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20 or more amino acids, or any range derivable therein, in a FN3 loop. In certain aspects, variations in loops AB, CD, and EF may be specifically excluded from invention, either individually or in various combinations. In some embodiments it is contemplated that an FN3 variant does not include a substitution, insertion and/or deletion in a bottom loop (loop AB, CD, EF). In further embodiments modifications in the bottom loop(s) is limited to 1, 2, 3, 4, or 5 or fewer substitutions, insertions, and/or deletions.

[0011] In certain embodiments, polypeptides comprise a variant fibronectin type III (FN3) domain comprising one or more amino acid substitutions in both a loop region of FN3 and in a non-loop region of FN3. In certain aspects, the one or more amino acid substitution in the non-loop segment is one or more substitution in beta strand C, beta strand D, beta strand F, and/or beta strand G.

[0012] In certain embodiments, the polypeptide can comprise 1, 2, 3, 4 or more insertions and/or deletions of amino acids corresponding to amino acids of SEQ ID NO:1. Insertions can include, but are not limited to stretches of poly-serine, poly-alanine, poly-valine, poly-threonine, or polymers of any other of the 20 amino acids, that is

subsequently mutagenized or diversified for generating a combinatorial polypeptide library. Diversification of these inserted residues can include alteration to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 of the other natural amino acids. In certain aspects 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more contiguous amino acids are inserted into one or more of the AB, BC, CD, DE, EF, FG loops of a FN3 domain polypeptide. In a further aspect, the polypeptide can comprise an insertion, a deletion, or both an insertion and a deletion. The insertion and/or deletion can be at the beginning and/or end of the polypeptide. The insertion and deletion need not be located at the same position and may be located at sites distal or proximal to each other. The insertion and/or deletion can be in a loop or non-loop portion of the FN3 domain polypeptide. In certain aspects, at least one loop region of FN3 comprises an insertion of at least 2 amino acids. In a further aspect, at least one region of FN3 comprises an insertion of 2 to 25 amino acids in at least one loop region. In certain aspects at least 2, 3, or more loop regions comprise an insertion. In certain aspects, the polypeptide has at least one loop region of FN3 comprises a deletion of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids, including all values and ranges there between. In certain aspects, at least 2, 3, or 4 loop or non-loop segments, portions, or regions comprise a deletion of at least 1 amino acid. In certain aspects, the polypeptide comprises at least one insertion and one deletion in at least one loop or non-loop region. In a further aspect, the polypeptide comprises an insertion and a deletion in the same loop or non-loop region. The term region indicates the amino acids of a particular structural segment of the polypeptide as defined by secondary structure and/or crystal structure corresponding to the amino acids of SEQ ID NO:1 or its variants.

[0013] Further aspects relate to a polypeptide comprising a variant fibronectin type III (FN3) domain from *Pyrococcus horikoshii* comprising one or more amino acid substitutions or insertions in a loop region of FN3, in a non-loop region of FN3, or in both. In some embodiments, the non-variant FN3 domain comprises at least 70% identity to SEQ ID NO:6. In some embodiments, the non-variant FN3 domain comprises SEQ ID NO:6. In some embodiments, the one or more amino acid substitutions or insertions in the non-loop segment is in one or more of beta strand C or beta strand D. In some embodiments, the one or more amino acid substitutions or insertions correspond to or after positions 13 and/or 14 of SEQ ID NO:6. In some embodiments, the one or more amino acid substitutions or insertions correspond to or after positions 20, 21, 22, 23, 24, 25, 26, and/or 27 of SEQ ID NO:6. In some embodiments, the one or more amino acid substitutions or insertions correspond to or after positions 36 and/or 37 of SEQ ID NO:6. In some embodiments, the one or more amino acid substitutions or insertions correspond to or after positions 40, 41, 42, 43, 44, 45, 46, 47, and/or 48 of SEQ ID NO:6. In some embodiments, the one or more amino acid substitutions or insertions correspond to or after positions 54, 55, 56, 57, 58, and/or 59 of SEQ ID NO:6. In some embodiments, the one or more amino acid substitutions or insertions correspond to or after positions 67, 68, 69, 70, 71, 72, 73, and/or 74 of SEQ ID NO:6. In some embodiments, the one or more amino acid substitutions or insertions in the loop region of FN3 are in the AB loop, BC loop, CD loop, DE loop, EF loop, and/or FG loop. In some

embodiments, the polypeptide further comprises a non-FN3 polypeptide that enhances the FN3 polypeptide binding affinity for a target molecule. In some embodiments, the polypeptide further comprises an insertion or deletion of at least 1 amino acids in at least one loop region of FN3. In some embodiments, at least one loop region of FN3 comprises an insertion of at least 2 amino acids. In some embodiments, at least one region of FN3 comprises an insertion of 2 to 25 amino acids in at least one loop region. In some embodiments, at least two loop regions comprise an insertion. In some embodiments, at least one loop region of FN3 comprises a deletion of at least 1 amino acid. In some embodiments, at least one loop region of FN3 comprises a deletion of 2 to 10 amino acids. In some embodiments, at least two loop regions comprise a deletion of at least 1 amino acid. In some embodiments, the polypeptide comprises at least 1 amino acid insertion and 1 amino acid deletion in at least one loop region. In some embodiments, the polypeptide comprises an insertion and deletion of at least 1 amino acid in the same loop region.

[0014] In certain aspects, the polypeptide is at least or at most 50, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identical (or any derivable range therein) to a polypeptide sequence of the disclosure, including any of SEQ ID NOS:1-4 or 6.

[0015] In some embodiments, the polypeptide is isolated, recombinant, a non-natural polypeptide, engineered, and/or synthetic.

[0016] In some embodiments, the polypeptides of the disclosure bind specifically to a site on a target motif. In some embodiments, the polypeptides of the disclosure specifically bind a target motif. In some embodiments, the polypeptide further comprises a polypeptide comprising a biorecognition module including a molecular recognition domain; wherein the polypeptide comprising the biorecognition molecule and the polypeptide comprising the variant FN3 domain are operatively linked together either directly or indirectly via the linker, and are spatially oriented to bind the same, overlapping, or distinct sites on the target motif. In some embodiments, the polypeptide comprising the biorecognition molecule and the polypeptide comprising the variant FN3 domain bind overlapping sites on the target motif at the same time and are capable of forming a complex with the target motif. In some embodiments, the K_d for the polypeptide being equal to or lower than one μM. In some embodiments, the target motif is a peptide, a phosphorylated peptide or a methylated peptide. In some embodiments, the peptide is present within a protein. In some embodiments, the molecular recognition domain comprises an interaction domain or mutants of interaction domains. In some embodiments, the interaction domain is selected from the group consisting of PDZ, WW, SH2, PTB, SH3, Bromo, Chromo, PHD, Polo-box and FHA domains. In further embodiments, the polypeptide comprising the variant FN3 domain further comprises a first signaling moiety and the polypeptide comprising the biorecognition molecule further comprises a second signaling moiety and wherein the first and second signaling moieties are capable of interacting to produce a detectable signal. In some embodiments, the signaling moiety is a dye, a quencher, a reporter protein, or a quantum dot. In some embodiments, the first and second signaling moieties comprise a fluorescent resonance energy transfer (FRET) donor group and a FRET acceptor group, respectively, and binding of the first and second molecular recog-

nition domains to the target motif results in a change in the FRET efficiency between the FRET donor and FRET acceptor groups.

[0017] The polypeptides of the invention can be comprised in a polypeptide library or encoded in a polynucleotide library that can be screened for particular polypeptide characteristics, e.g., binding affinity. One or more members of the library can then be isolated from other members of the library and analyzed. In certain aspects the library comprises or encodes a plurality of those polypeptides described herein. In certain aspects, the polypeptide library is pre-selected to bind a target and those preselected members are then further diversified in selected amino acid position to generate a targeted library that is subsequently screened for a particular characteristic or property.

[0018] Certain aspects relate to a polypeptide library comprising a plurality of modified FN3 domain polypeptides comprising one or more amino acid substitutions, insertions, or deletions a loop region of FN3, in a non-loop region of FN3, or in both; wherein the unmodified FN3 domain comprises a polypeptide comprising the amino acid sequence of SEQ ID NOS:1-4 or 6. In some embodiments, the FN3 domain polypeptides comprises one or more amino acid substitutions corresponding to amino acid positions corresponding to positions 1, 2, 3, 4, 14, 28, and/or 58 of SEQ ID NO:1.

[0019] Certain aspects are directed to polynucleotides encoding one or more polypeptide described herein. In certain embodiments the polynucleotide is an expression cassette or an expression construct. The expression construct can be capable of expressing the encoded polypeptide in a host cell, such as a prokaryotic or eukaryotic cell line or strain. In certain aspects the expression construct is functional in one or more polypeptide expression systems known in the art. In a further aspect, the expression construct is functional in bacteria, yeast, insect cells or the like.

[0020] The polypeptide can further comprise a second FN3 domain that may or may not have been selected for affinity to a particular target. The second FN3 domain may or may not contain additional amino acid variations or diversification. In other aspects, the polypeptide can further comprise a non-FN3 polypeptide that enhances the FN3 polypeptide binding affinity for a target molecule. The non-FN3 polypeptide can include, but is not limited to domains involved in phospho-tyrosine binding (e.g., SH2, PTB), phospho-serine binding (e.g., UIM, GAT, CUE, BTB/POZ, VHS, UBA, RING, HECT, WW, 14-3-3, Polo-box), phospho-threonine binding (e.g., FHA, WW, Polo-box), proline-rich region binding (e.g., EVH1, SH3, GYF), acetylated lysine binding (e.g., Bromo), methylated lysine binding (e.g., Chromo, PHD), apoptosis (e.g., BIR, TRAF, DED, Death, CARD, BH), cytoskeleton modulation (e.g., ADF, GEL, DH, CH, FH2), or other cellular functions (e.g., EH, CC, VHL, TUDOR, PUF Repeat, PAS, MH1, LRR1 IQ, HEAT, GRIP, TUBBY, SNARE, TPR, TIR, START, SOCS Box, SAM, RGS, PDZ, PB1, LIM, F-BOX, ENTH, EF-Hand, SHADOW, ARM, ANK).

[0021] In certain aspects, variants in any one or more of positions that correspond with amino acid position 13, 14, 15, 16, 22, 23, 24, 25, 26, 27, 36, 37, 47, 48, 49, 54, 55, 56, 57, 58, 59, 60, 71, 72, 73, 74, 75, 76, 77, 78, 79 and/or 80 of SEQ ID NO:1, including all ranges there between, can be specifically included in the claimed embodiments.

[0022] In other embodiments, variants in any one or more of positions that correspond with amino acid positions 13, 14, 20, 21, 22, 23, 24, 25, 26, 27, 36, 37, 40, 41, 42, 43, 44, 45, 46, 47, 48, 54, 55, 56, 57, 58, 59, 67, 68, 69, 70, 71, 72, 73 and 74 of SEQ ID NO:6, including all ranges there between, can be specifically included in the claimed embodiments.

[0023] In some embodiments, the polypeptide library comprises domain polypeptides are at least 50%, 60%, 70%, 80%, or 90% identical to SEQ ID NO:1, 2, 3, 4, or 6.

[0024] In some embodiments, the length of the polypeptide is at least, at most, or exactly about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, 265, 270, 275, 280, 285, 290, 295, 300, 305, 310, 315, 320, 325, 330, 335, 340, 345, 350, 355, 360, 365, 370, 375, 380, 385, 390, 395, 400, 405, 410, 415, 420, 425, 430, 435, 440, 445, 450, 455, 460, 465, 470, 475, 480, 485, 490, 495, 500, 505, 510, 515, 520, 525, 530, 535, 540, 545, 550, 555, 560, 565, 570, 575, 580, 585, 590, 595, 600, 625, 650, 675, 700, 750, 800, or 900 amino acids (or any derivable range therein). In some embodiments, the polypeptide is truncated or not full length.

[0025] In some embodiments, the polypeptide library further comprises a non-FN3 polypeptide that enhances the FN3 polypeptide binding affinity for a target molecule. In some embodiments, the polypeptide library further comprises an insertion or deletion of at least 1 amino acids in at least one loop region of the FN3 domain polypeptides. In some embodiments, at least one loop region of FN3 comprises an insertion of at least 2 amino acids. In some embodiments, at least one region of FN3 comprises an insertion of 2 to 25 amino acids in at least one loop region. In some embodiments, at least two loop regions comprise an insertion. In some embodiments, at least one loop region of FN3 comprises a deletion of at least 1 amino acid. In some embodiments, at least one loop region of FN3 comprises a deletion of 2 to 10 amino acids. In some embodiments, at least two loop regions comprise a deletion of at least 1 amino acid. In some embodiments, the polypeptide comprises at least 1 amino acid insertion and 1 amino acid deletion in at least one loop region. In some embodiments, the FN3 domain polypeptides comprise an insertion and deletion of at least 1 amino acid in the same loop region. In some embodiments, the library is pre-selected to bind a target.

[0026] Certain embodiments are directed to a monobody library comprising a plurality of polypeptides having a variant fibronectin type III (FN3) domain, compared to wildtype (SEQ ID NO:1-4 or 6), comprising one or more alterations or variants in a beta strand or a loop region. In some embodiments, the alterations are in one or more of beta strand C, beta strand D, BC loop, CD loop, DE loop, or and/or FG loop. In some embodiments, the monobody comprises one or more of the variant amino acids corresponding to position 1, 2, 3, 4, 14, 28, and/or 58 of SEQ ID NO:1. In some embodiments, the variant FN3 domains further comprise an insertion, substitution, or deletion of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 amino acids in at least one loop region of FN3.

[0027] The monobody library can also comprise variant FN3 domains comprising an amino acid insertion in loop FG. The monobody library can also comprise a plurality of those polypeptides described above.

[0028] Certain embodiments include methods of making a polypeptide or polynucleotide library comprising a plurality of FN3 variants. In certain aspects the library can contain 10, 100, 1000, 104, 105, 106, 107, 108, 109, 1010, 1011, 1012, 1013, 1014, 1015 or more different polypeptide or polynucleotide variants, including all values and ranges there between, though it will be understood that there may be duplicate variants. The methods of making such a polypeptide or polynucleotide include the engineering of various amino acids substitutions, deletions, and/or insertion described herein.

[0029] Certain embodiments include methods of selecting one or more FN3 variants comprising conducting one or more binding assays using a FN3 library having a plurality of different FN3 variants. In certain aspects the library can comprise FN3 polypeptides having amino acid variations in the FN3 loops, FN3 beta-strands, or both FN3 loops and beta-strands. After conducting the binding assay(s) one or more FN3 variants are selected that have a particular property, such as binding specificity and/or binding affinity to a target. In certain aspects, the amino acid or nucleic acid sequence of one or more of the selected library members can be determined using conventional methods. The sequence of the selected FN3 polypeptide(s) can then be used to produce a second library that introduces further variation of the selected sequences. The second library can then be screened for FN3 polypeptides having a particular property. The process can be repeated 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more times. Additional iterations would enrich the library as well as potentially include other variants.

[0030] In certain aspects the method for selecting a protein binding domain specific for a target comprises (a) detecting target specific binding of one or more members of a polypeptide library comprising a plurality of FN3 domain polypeptides as described herein; and (b) selecting the protein binding domain that specifically binds the target. The method can further comprise first preparing the plurality of FN3 domain polypeptide variants described herein. A polypeptide identified as exhibiting a particular characteristic can be isolated. In certain aspects, the method can further comprise determining the nucleic acid and/or the amino acid of sequence of the selected protein binding domain. The selected protein binding domain can then be synthesized or expressed.

[0031] Methods can further comprise conducting a first screen of a library having amino acid variations in only FN3 loops or only FN3 beta strands and conducting a second screen using variations in only FN3 loops or only FN3 beta strands. In certain aspects the first screen uses only variations in the FN3 loops and the second screen only uses variations in the FN3 beta-strands. In a further aspect, the second screen can use variations in both FN3 loops and beta-strands. In certain aspects, the FN3 amino acid residues varied in the first screen are or are not varied in the second screen.

[0032] Further aspects include methods of identifying a polypeptide that specifically binds a target comprising detecting specific binding of one or more polypeptides of a polypeptide library, the library comprising a plurality of fibronectin type III (FN3) polypeptides as described herein.

[0033] Further aspects include methods of detecting a target molecule comprising contacting a sample containing the target with a fibronectin type III (FN3) binding domain that specifically binds the target.

[0034] Yet further aspects relate to a method of making a polypeptide or polypeptide library according to the disclosure, the method comprising expressing a polynucleotide of the disclosure or a polynucleotide encoding such polypeptide in a host cell.

[0035] Certain aspects include methods of producing a fibronectin type III (FN3) variant comprising: (a) expressing a polypeptide comprising an amino acid sequence; and (b) isolating and/or purifying the expressed variant FN3 domain from a host cell expressing the variant FN3.

[0036] Certain embodiments are directed to kits. In certain aspects, a kit can comprise a plurality of polypeptides as described herein. In a further aspect, a kit can comprise a plurality of polynucleotides encoding FN3 domain variants as described herein.

[0037] Further aspects of the disclosure relate to methods for testing a polypeptide according to the embodiments described herein, the method comprising contacting the polypeptide with a target molecule and testing for binding activity between the polypeptide and the target molecule.

[0038] The term “fibronectin type III domain” or “FN3 domain” refers to a domain (region) from a wild-type fibronectin from any organism.

[0039] The term “fibronectin type III domain variant” or “FN3 variant domain” refers to a polypeptide region in which one or more amino acid substitutions, deletions, and/or insertions are present as compared to the amino acid sequence of a wildtype FN3 domain. In certain embodiments, the FN3 variant or FN3 variant domain has an alteration with respect to specifically the fibronectin type III domain of SEQ ID NO:1-4 or 6. The term “substitutional variant” includes the replacement of one or more amino acids in a peptide sequence with a conservative or non-conservative amino acid(s). In some embodiments, the FN3 domain variant has increased binding properties compared to the wildtype FN3 domain relative to a particular target.

[0040] The term “FN3-domain polypeptide” refers to a polypeptide that includes at least one FN3 domain. A “variant FN3 domain polypeptide” refers to a polypeptide that includes at least one variant FN3 domain. It is contemplated that such polypeptides are capable of specifically binding a polypeptide or protein.

[0041] A “non-FN3 binding sequence” refers to an amino acid sequence of more than 15 contiguous amino acid residues that is not present in an FN3 domain or an FN3 domain variant and that specifically binds to a protein or polypeptide. In some embodiments, a non-FN3 binding sequence is specifically a non-tenth module fibronectin type III domain binding sequence.

[0042] The β sheet is a form of regular secondary structure in proteins. Beta sheets consist of beta strands connected laterally by at least two or three backbone hydrogen bonds, forming a generally twisted, pleated sheet. A beta strand (also β strand) is a stretch of polypeptide chain typically 3 to 10 amino acids long with backbone in an almost fully extended conformation.

[0043] A loop is a less ordered, flexible stretch of amino acids (as compared to alpha helices and beta sheets) that typically connect other structural elements of a protein. In the context of FN3, the loops are designated by the beta-strands they connect, for example the loop connecting beta-strand A and beta-strand B is the AB loop.

[0044] Beta strand A refers to the amino acids preceding the AB loop

[0045] Beta strand B refers to the amino acids connecting the AB and BC loops

[0046] Beta strand C refers to the amino acids connecting the BC and CD loops.

[0047] Beta strand D refers to the amino acids connecting the CD and DE loops.

[0048] Beta strand E refers to the amino acids connecting the DE and EF loops.

[0049] Beta strand F refers to the amino acids connecting the EF and FG loops.

[0050] Beta strand G refers to the amino acids after the FG loop.

[0051] The term “binding protein” refers to a polypeptide that specifically binds another compound, such as a polypeptide through non-covalent chemical interactions.

[0052] As used herein, “monobody” is intended to mean a polypeptide having a sequence and structure related to the tenth module of the fibronectin type III domain (FN3) that includes a beta-strand domain lacking in disulfide bonds and containing a plurality of beta-strands, two or more loop regions each connecting one beta-strand to another beta-strand, and optionally an N-terminal tail, a C-terminal tail, or both, wherein at least one of the two or more loop regions, the N-terminal tail, and/or the C-terminal tail is characterized by activity in binding a target protein or molecule. More specifically, in some embodiments such monobodies can include three or more loop regions or, even more specifically, four or more loop regions. The size of such polypeptide monobodies is preferably less than about 30 kDa, more preferably less than about 20 kDa.

[0053] The term “library” refers to a collection (e.g., to a plurality) of polypeptides having different amino acid sequences and different protein binding properties. In some embodiments there is a variant FN3 domain library comprising polypeptides having different variations of the FN3 domain. Unless otherwise noted, the library is an actual physical library of polypeptides or nucleic acids encoding the polypeptides. In further embodiments, there is a database that comprises information about a library that has been generated or a theoretical library that can be generated. This information may be a compound database comprising descriptions or structures of a plurality of potential variant FN3 domains. “FN3-based molecule” refers to a molecule having an amino acid sequence of an FN3 domain or FN3 variant domain.

[0054] The term “specifically binds” or “specific binding” refers to the measurable and reproducible ability of an FN3-based molecule to bind another molecule (such as a target), that is determinative of the presence of the target molecule in the presence of a heterogeneous population of molecules including biological molecules. For example, an FN3-based molecule that specifically or preferentially binds to a target is a polypeptide that binds this target with greater affinity, avidity, more readily, and/or with greater duration than it binds to most or all other molecules. “Specific binding” does not necessarily require (although it can include) exclusive binding.

[0055] An polypeptide that specifically binds to a target with an affinity of at least 1×10^{-6} M at room temperature under physiological salt and pH conditions, as measured by surface plasmon resonance.

[0056] The term “non-natural amino acid residue” refers to an amino acid residue that is not present in the naturally occurring FN3 domain in a mammal, such as a human. The

term “non-natural polypeptide” refers to a polypeptide that is not wild-type and/or not found in nature.

[0057] The terms “tag”, “epitope tag” or “affinity tag” are used interchangeably herein, and usually refers to a molecule or domain of a molecule that is specifically recognized by an antibody or other binding partner. The term also refers to the binding partner complex as well. Thus, for example, biotin or a biotin/avidin complex are both regarded as an affinity tag. In addition to epitopes recognized in epitope/antibody interactions, affinity tags also comprise “epitopes” recognized by other binding molecules (e.g., ligands bound by receptors), ligands bound by other ligands to form heterodimers or homodimers, His6 bound by Ni-NTA, biotin bound by avidin, streptavidin, or anti-biotin antibodies, and the like.

[0058] Epitope tags are well known to those of skill in the art. Moreover, antibodies specific to a wide variety of epitope tags are commercially available. These include but are not limited to antibodies against the DYKDDDDK epitope, c-myc antibodies (available from Sigma, St. Louis), the HNK-1 carbohydrate epitope, the HA epitope, the HSV epitope, the His4, His5, and His6 epitopes that are recognized by the His epitope specific antibodies (see, e.g., Qiagen), and the like. In addition, vectors for epitope tagging proteins are commercially available. A polypeptide can be tagged with the FLAG® epitope (N-terminal, C-terminal or internal tagging), the c-myc epitope (C-terminal) or both the FLAG (N-terminal) and c-myc (C-terminal) epitopes.

[0059] The term “conjugate” in the context of an FN3-based molecule refers to a chemical linkage between the FN3-based molecule and a non-FN3-based molecule. It is specifically contemplated that this excludes a regular peptide bond found between amino acid residues under physiologic conditions in some embodiments.

[0060] Other embodiments of the invention are discussed throughout this application. Any embodiment discussed with respect to one aspect of the invention applies to other aspects of the invention as well and vice versa. The embodiments in the Example section are understood to be embodiments of the invention that are applicable to all aspects of the invention.

[0061] “Biorecognition module” and “recognition module” (used interchangeably), as used herein, refer to a biomolecule which makes up one module of the modular molecular affinity clamp embodying the principles of the invention. A biorecognition module contains a molecular recognition domain that has affinity for a target motif of interest.

[0062] “Molecular recognition domain” and “recognition domain”, (used interchangeably), as used herein, refer to a binding domain within a biorecognition module that demonstrates an ability to bind to a target motif, i.e., has binding affinity for a target motif.

[0063] The terms “target” and “target molecule,” as used herein, refer to a peptide, antigen or epitope that specifically binds to an FN3-based binding molecule or monobody described herein, or any biomolecule of interest for which a molecular affinity clamp is sought. Targets include, but are not limited to, epitopes present on proteins, peptides, carbohydrates, and/or lipids. Exemplary targets include, but are not limited to, secreted peptide growth factors, pharmaceutical agents, cell signaling molecules, blood proteins, portions of cell surface receptor molecules, portions of nuclear

receptors, steroid molecules, viral proteins, carbohydrates, enzymes, active sites of enzymes, binding sites of enzymes, portions of enzymes, small molecule drugs, cells, bacterial cells, proteins, epitopes of proteins, surfaces of proteins involved in protein-protein interactions, cell surface epitopes, diagnostic proteins, diagnostic markers, plant proteins, peptides involved in protein-protein interactions, and foods. The target may be associated with a biological state, such as a disease or disorder in a plant or animal as well as the presence of a pathogen. When a target is “associated with” a certain biological state, the presence or absence of the target or the presence of a certain amount of target can identify the biological state.

[0064] A “target motif”, as used herein, refers to any portion or sequence of a target of interest for which a molecular affinity clamp is sought, e.g., refers to a pattern of amino acid residues which is recognized by particular recognition domains. In accordance with the invention, the target motif can bind more than one recognition domain. In other words, a target motif is one to which an affinity clamp embodying the principles of the invention can bind with high affinity and specificity. Of particular importance are target motifs that are short peptides of about 2-100 amino acid residues, especially those of 3-10 amino acid residues. A target motif may be, be at least, or be at most 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 residues in length (or any range derivable therein), which may or may not be contiguous. A target motif may include 1, 2, 3, 4, 5 or more noncontiguous residues and/or regions of residues that may be, be at least, or be at most 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 residues in length (or any range derivable therein).

[0065] As used herein, the term “binds” in connection with the interaction between a target motif and a recognition domain indicates that the recognition domain associates with (e.g., interacts with or complexes with) the target motif to a statistically significant degree as compared to association with proteins generally (i.e., non-specific binding). Thus, the term “molecular recognition domain” is also understood to refer to a domain that has a statistically significant association or binding with a target motif.

[0066] In the context of a recognition domain binding to a target motif, the term “greater affinity” indicates that an affinity clamp binds more tightly than a reference domain, or than the same domain in a reference condition, i.e., with a lower dissociation constant. In particular embodiments, the greater affinity is at least 2-fold.

[0067] Also in the context of recognition domain binding to a target motif, the term “altered specificity” indicates that relative binding affinity of an affinity clamp is different from that exhibited by a biorecognition module alone. In other words, “altered binding specificity” may refer to an increased binding constant of the affinity clamp for one target motif without the same level of increase for another

target, an unchanged binding constant for one target with a decreased binding constant for another, target or a combination thereof.

[0068] The term “linked” refers to any method of functionally connecting peptides, particularly the two modules of the modular affinity clamps embodying the principles of the invention. “Linked” may also refer to non-covalent physical association. The biomolecular modules making up the biorecognition modules of the affinity clamps may be linked directly covalently, e.g., via a peptide linkage, or non-covalently, or indirectly via a linker.

[0069] A “linker” or “linker moiety,” (used interchangeably) may refer to a peptide sequence of about 30 or more amino acid residues that is configured to associate two biorecognition modules in an orientation that facilitates binding of each module to a target motif. The linker, generally, is bifunctional in that it includes a functionality for linking the first biorecognition module and a functionality for linking the second biorecognition module.

[0070] By “binding site” is meant an area or region within a recognition domain where a biomolecule can bind non-covalently, i.e., interact with higher affinity than background interactions between molecules. Binding sites embody particular shapes and often can contain multiple binding pockets present within the binding site. The particular shapes are often conserved within a class of molecules, such as a protein family. Binding sites within a class also can contain conserved structures such as, for example, chemical moieties, the presence of a binding pocket, and/or an electrostatic charge at the binding site or some portion of the binding site, all of which can influence the shape of the binding site. It is noted that a molecular affinity clamp is distinguishable from other protein-based compositions that are multivalent—i.e., bind multiple but separate target motifs that are the same. Moreover, in some embodiments a molecular affinity clamp has one component that specifically binds to a target motif or an amino acid sequence that is distinct, overlapping, or the same as another target motif or amino acid sequence specifically bound by another component in the affinity clamp. The different binding components of a molecular affinity clamp do not compete against one another for binding to a target motif.

[0071] By “binding pocket” is meant a specific volume of space within a binding site that is available for occupation by a biomolecule. A binding pocket can often be a particular shape, indentation, groove, or cavity in the binding site. Binding pockets can contain particular chemical groups or structures that are important in the non-covalent binding of another molecule such as, for example, groups that contribute to ionic, hydrogen bonding, or van der Waals interactions between the molecules.

[0072] By “orientation” or “oriented,” in reference to a biorecognition module bound to a target motif, is meant the spatial relationship of the biorecognition module, and at least some of its constituent atoms, to the atoms of the target motif.

[0073] By “assaying” is meant the creation of experimental conditions and the gathering of data regarding a particular result of the experimental conditions. For example, enzymes can be assayed based on their ability to act upon a detectable substrate. A particular target motif, in a test sample, can be assayed based on its ability to bind to a monobody or molecular affinity clamp.

[0074] As used herein, the terms “peptide,” “polypeptide,” and “protein” are used interchangeably and mean polymers of amino acid monomers linked by peptide linkages between carboxyl (COOH) groups and amine (NH₂) groups. A peptide may consist entirely of naturally occurring amino acid monomers, non-naturally occurring amino acids, or mixtures thereof. Unless denoted otherwise, whenever an amino acid sequence is represented, it will be understood that the amino acids are in N-terminal to C-terminal order from left to right. The term “polypeptide” may refer to small peptides, larger polypeptides, proteins containing single polypeptide chains, proteins containing multiple polypeptide chains, and multi-subunit proteins.

[0075] The term “amino acid”, as used herein, refers to any amino acid, natural or non-natural, that may be incorporated, either enzymatically or synthetically, into a polypeptide or protein. Amino acids may also be altered. The term thus encompasses amino acids that have been modified naturally or by interaction. Examples may include, but are not limited to, phosphorylation, glycosylation, methylation, biotinylation, and any covalent and non-covalent additions to a protein that do not result in a change in amino acid sequence.

[0076] The term “label” as used herein refers to any tag, marker, or identifiable moiety. The skilled artisan will appreciate that many labels may be used in the methods of the invention. For example, labels include, but are not limited to, affinity tags, fluorophores, radioisotopes, chromogens, dyes, magnetic probes, magnetic particles, paramagnetic particles, electrophoretic molecules and particles, dielectrophoretic particles, phosphorescence groups, chemiluminescent, mobility modifiers, and particles that confer a dielectrophoretic change.

[0077] As used herein, the term “modulating” or “modulate” refers to an effect of altering a biological activity, especially a biological activity associated with a particular biomolecule. For example, an agonist or antagonist of a particular biomolecule modulates the activity of that biomolecule, e.g., an enzyme.

[0078] As used herein, the term “library” refers to any collection of two or more different polypeptides or proteins. In certain embodiments, a library may be a collection of polypeptides that have been modified to favor the inclusion of certain amino acid residues, or polypeptides of certain lengths.

[0079] As used herein, the term “variant” is meant to refer to a polypeptide differing from another polypeptide by one or more amino acid substitutions resulting from engineered mutations in the gene coding the polypeptide.

[0080] As used herein in connection with numerical values, the terms “approximately” and “about” are meant to encompass variations of $\pm 20\%$ to $\pm 10\%$ or less of the indicated value.

[0081] The terms “inhibiting,” “reducing,” or “preventing,” or any variation of these terms, when used in the claims and/or the specification includes any measurable decrease or complete inhibition to achieve a desired result.

[0082] The use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.”

[0083] It is contemplated that any embodiment discussed herein can be implemented with respect to any other

embodiment discussed herein, and vice versa. Furthermore, compositions and kits can be used to achieve recited methods.

[0084] Throughout this application, the term “about” is used to indicate that a value includes the standard deviation of error for the device or method being employed to determine the value.

[0085] The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” It is also contemplated that anything listed using the term “or” may also be specifically excluded.

[0086] As used in this specification and claim(s), the words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, unrecited elements or method steps.

[0087] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

DESCRIPTION OF THE DRAWINGS

[0088] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

[0089] FIG. 1A-C. Sequence alignment and biophysical properties of predicted STOFN3 domains. (a) Alignment of four predicted STOFN3 domains and the homologue FN3 domain of human receptor-type tyrosine-protein phosphatase F (PDB ID: 2DJU). Gaps are denoted as dashes. The human FN3 domain shows the sequence identities of 27, 29, 19 and 18% to STOFN3-1, STOFN3-2, STOFN3-3 and STOFN3-4, respectively. The ranges of the secondary structure of the human FN3 domain are shown below the sequences. Dots indicate the highly conserved residues of FN3 domains reported by Main et al. and Dickinson et al. (Main et al., 1992; Dickinson et al., 1994). (b) Size-exclusion chromatograms of STOFN 3-1 and STOFN3-2. The chromatographs are shown with vertical offsets for clarity. Both STOFN3 domains exhibited a mono-dispersed peak with the calculated molecular weight of ~12 kDa based on the calibration standards (not yet done, will do it). (c) Thermal stability of purified STOFN3-1 and STOFN3-2 monitored by DSF. One representative trace of three technical replicates is shown for each STOFN3 domains. The TM value is the mean of three replicates. The melting curves were measured by heating the samples at a rate of 0.5° C. per 30 seconds.

[0090] FIG. 2A-C. The crystal structures of STOFN3-1 (a) and STOFN3-2 (b). For panel (a) and (b), the β -strands are

labeled A-G. (c) Superposition of STOFN3-1, STOFN3-2 and FN3fn10 (PDBID: 1FNA). STOFN3-1, STOFN3-2 and FN3fn10 are colored green, cyan and gray, respectively.

[0091] FIG. 3A-D. The structures of STOFN3-1 (STOTEMP1) and STOTEMP4 around the N-terminal (a-b) and C-terminal (c) regions. The 2Fo-Fc maps around the N-terminal regions of STOTEMP1 (a) and STOTEMP4 (b) are shown as mesh at the 1.5 σ level. The N-terminal residues are indicated by green stick models, in which the oxygen and nitrogen atoms are colored red and blue, respectively. The BC-, DE- and FG-loops are colored yellow, pink and red. The low electron density of the segment N-terminal to P316S of STOTEMP4 indicates that this segment is conformationally disordered and dislodged from the folded portion. (c) Superposition of STOTEMP1 and STOTEMP4. STOTEMP1 is colored gray and the C-terminal residues are indicated as gray stick models. The C-terminal residues and A- and F-strands of STOTEMP4 are colored green, and the C-terminal residues are indicated as green stick models. The label of position 399 for STOTEMP1 is shown in parentheses. (d) Structure based alignment of STOFN3-1 with FN3fn10 (PDB ID: 1FNA) and amino acid sequences and melting temperatures of STOFN3-1 mutants. The ranges of the secondary structure of FN3fn10 are shown above the sequences. Mutated residues are colored in red. The mean TM values from three replicates are indicated.

[0092] FIG. 4A-C. Schematic drawing of the amino acid sequence of STOFN3-1 in its secondary structure context (a). Loop residues as assigned by the program DSSP are shown in yellow. G342 and Y383 subjected to mutation in this work are also shown in cyan. An arrow marks the site at which poly-Serine residues were inserted. Residues of the β -strands whose side chain forms the hydrophobic core are enclosed in circles with thicker ring. (b)-(c) Amino acid sequences and melting temperatures of STOFN3-1 variants. Mutated residues are colored in red. The mean TM values from three replicates are indicated.

[0093] FIG. 5. Optimization of the signal sequence and linker length for the efficient display of STOTEMP5. The amino acid sequences corresponding to OmpT leader, linker and STOTEMP5 are shown. Mutated residues are colored in red. Deleted residues are denoted as dashes. The display of STOTEMP5 on phage particle was measured by phage ELISA using an anti-V5 tag antibody and a HRP conjugated anti-M13 phage antibody. The absorbance changes at 405 nm after the reaction with 1-step Ultra TMB ELISA for 10 min at 25° C. are shown. For this measurement, aliquots of 50 μ L of phage particles with normalized titers of 2 \times 10⁶ cfu/mL pre-blocked in 0.5% BSA/TBS were added per well.

[0094] FIG. 6A-C. therMonobody library designs and generated clones. Amino acid sequences of therMonobodies generated from the loop-only library (a) and the side-and-loop library (b). The locations of diversified residues in the libraries are shown as spheres on the STOTEMP4 structure. “X” denotes a mixture of 30% Tyr, 13% Ser, 10% Gly, 5% Asp, 5% Leu, 5% Pro, 5% Trp and 2.5% each of all pther amino acids except for Cys and Met; “B”, a mixture of Ala, Ser, Thr, Asn, Asp and Tyr; “U”, a mixture of Ser, Thr, Asn and Tyr; “Z”, a mixture of Ala, Glu, Lys and Thr. (c) Binding measurements by yeast surface display of representative therMonobodies for each combination of library and target. The mean fluorescence intensities of yeast cells displaying therMonobodies are plotted as a function of the target

concentration. The error indicated are the standard deviations from curve fitting of the 1:1 binding model.

[0095] FIG. 7A-C. Oligomerization state and thermal stability of generated therMonobodies. (a) Size-exclusion chromatograms of therMonobodies. The chromatographs are shown with vertical offsets for clarity. The labels show the identities of analyzed samples. MBPMbL17 and MBPMbS09 exhibited a mono-dispersed peak but appeared to interact with the chromatography media, resulting in late elution. SUMOMbS34 was eluted at the void volume fraction, indicating the large aggregate. (b) Thermal stability of therMonobodies monitored by DSF. The traces of three technical replicates are shown for each therMonobody. The T_m value is the mean of three replicates. The melting curves were measured by heating the samples at a rate of 0.5° C. per 30 seconds. (c) Summary of oligomerization state and thermal stability measurements of 24 therMonobodies.

[0096] FIG. 8A-C. Affinity clamp using therMonobody. (a) Amino acid sequences of Clamp(Ptpn11_pY580) and STOTEMP4 variants. The monobody segment of Clamp (Ptpn11_pY580) and STOTEMP4 variants are aligned based on structure based alignment of the monobody segment and STOTEMP4. The ranges of the secondary structure of the monobody segment of Clamp(Ptpn11_pY580) are shown below its sequence. Residues grafted to the structurally equivalent positions of STOTEMP4 are shaded in yellow. Mutated residues in STOTEMP4 are colored in red. Linkers are indicated by underlining. (b)-(c) pY-peptide binding properties of designed pY-clamps measured by yeast surface display. In panel (b), binding of the designed pY-clamps to the target pY-peptide (Ptpn11_pY580) at the concentration of 0, 100 and 500 nM is shown. In panel (c), binding titration curves and the dissociation constants (KD) of the designed pY-clamps are shown. The mean fluorescence intensities of yeast cells displaying the designed pY-clamp are plotted as a function of the target peptide concentration. The errors shown are the standard deviations from curve fitting of the 1:1 binding model.

[0097] FIG. 9. Sequence alignment of predicted PHOFN3 and the bacterial FN3 domain in *Clostridium Perfringens* Glycoside Hydrolase Gh84c (PDB ID: 2W1N). Gaps are denoted as dashes. The bacterial FN3 domain shows the sequence identities of 25% to PHOFN3, which is the highest among PDB entries. The locations of beta-strands in the bacterial FN3 domain are shown as the green arrows. Red dots indicate the highly conserved residues of FN3 domains reported by Main et al. and Dickinson et al. (Main et al., 1992; Dickinson et al., 1994).

[0098] FIG. 10. Thermal stability of purified PHOFN3 and its truncated variants monitored by DSF. Triplicate measurements are shown for each sample. The melting curves were measured by heating the samples at a rate of 0.5° C. per 30 seconds. Only PHOFN3□C35 exhibited an inflection point indicative of thermal denaturation. The T_m value for PHOFN3□C35 is the mean of the inflection points from triplicates.

[0099] FIG. 11A-B. A) The crystal structures of SeMet-labeled PHOFN3AC25. The seven β -strands are colored and labeled A-G. N- and C-terminus are labeled. B) Superposition of PHOFN3AC25 and FN3fn10 (PDBID: 1FNA). PHOFN3AC25 and FN3fn10 are colored green and gray, respectively.

[0100] FIG. 12. Schematic drawing of the amino acid sequence of PHOFN3AC25 in its secondary structure con-

text. Loop residues as assigned by the program DSSP are shown in yellow. Residues of the β -strands whose side chain forms the hydrophobic core are enclosed in circles with thicker ring.

[0101] FIG. 13. Phage ELISA analysis for surface display of PHOFN3 on phage particles. Binding of phage particles to an anti-V5 tag antibody. The V5 tag is located in the linker between PHOFN3 and the phage coat protein, P3, in the phage display vector. The absorbance changes at 405 nm after the HRP reaction with 1-step Ultra TMB ELISA for 10 min for phages produced with the M13KO7 helper phage or 1 min for hyperphage-produced phages are shown. For these measurements, 50 μ L of 0.5% BSA/TBS containing 2.3×10^6 cfu/mL M13KO7-produced phages or 7.4×10^6 cfu/mL hyperphage-produced phages were added to each well and bound phages were detected with a HRP conjugated anti-M13 phage antibody.

DETAILED DESCRIPTION OF THE INVENTION

[0102] The fibronectin type III domain (FN3) has been particularly successful as a protein scaffold for generating synthetic binding proteins. Since the pioneering work by inventors on the tenth FN3 of human fibronectin (FNfn10), numerous binding proteins, termed monobodies, have been generated to diverse target molecules. To date, all FN3-based scaffolds have been derived from human proteins, primarily because of the prediction that molecules engineered from a human protein may have low immunogenicity, an important consideration in biotherapeutic development. However, immunogenicity concerns are less important in applications where synthetic binding proteins are not exposed directly to patients or consumers including chemical processing and research tools. Therefore, this current application is directed toward the development of a molecular scaffold for industrial applications using FN3's from non-human origins, in particular thermophiles.

IV. Fibronectin Type III (FN3) Domain

[0103] A. *Sulfolobus tokodaii*

[0104] The inventors utilized the SMART database to explore FN3 domains from hyperthermophiles. The database predicted many FN3 domains in hyperthermophilic archaea and bacteria such as *Thermococcus kodakaraensis*, *Sulfolobus tokodaii*, *Pyrococcus horikoshii* and *Thermotoga lettingae*. The inventors first eliminated predicted domains that were shorter than the length of the shortest FN3 domains that had been structurally characterized (75 amino acids). Then, four predicted FN3 domains in the sequence of Kelch domain-containing protein ST0939 from the hyperthermophilic archaeon *Sulfolobus tokodaii* DSM 16993 were chosen as the candidate proteins, because of their detectably homology to a human FN3. In the predicted constructs, termed STOFN3-1, -2, -3 and -4.

[0105] STOFN3-1 comprises 84 amino acids at positions 315-398:

(SEQ ID NO: 1)
 PPPKPQIASIASGNETITVKWYDTNASGGYITYWNSFSQKVTINVGNTVS
 YTIKHLKDGVTYYIQIVPYNSLGNCTPSDIIISAT.

[0106] STOFN3-2 comprises 86 amino acids at positions 399-484:

(SEQ ID NO: 2)
PSSVPNPPIIKVKIGNLNATLTWYDTFNGGYPYIEGYLYVNGKGINVGNL
TSYVLTNLTAGELYTIELIAYNKIGNSSISSVSFIA.

[0107] STOFN3-3 comprises 79 amino acids at positions 488-566:

(SEQ ID NO: 3)
ANLTVTVYKINGFLVSWNSTKAKYILTVSKENVLLNVSTTNTSYFVK
VPGVYINISLEAVNIVGITYAFILIIYI.

[0108] STOFN3-4 comprises 76 amino acids at positions 568-643:

(SEQ ID NO: 4)
PASPTVNWISITLNTVSLNWSKVGAEYLIYDNGKLI TTTNTAFTFNLT
IGQNEIEVYAANAYYKSAPYIINDVR.

[0109] Each construct maintains at least two of the three highly conserved hydrophobic amino acids of FN3 domains (FIG. 1A).

[0110] The full-length ST0939 comprises the sequence:

(SEQ ID NO: 5)
MKRNTLLALVVLILIFPTLSTAYIEFTTSINQAI PDSLVIYATSAYYDGI
FLIGGENLYSTPVNSVYVYENGSWYLGPSLPFSLSSAGATVCNNTLYVVG
GANSTSIFGGILEFIGNGWKVIITNSMPVIVYGAIVFSYDYKIYVIGGMNY
SGNSLVPPVNYIQVYNLKTNSWQIIGNAPLRLAYSAYYFNGSALFVVGGF
TQSATLTSVVFVYYPENNTWISLPSLPVVEAGGVLGYNGMYLVGLGLY
VSGAYQLGEILYNYNGTWRNTNIQEIQIPQFSTSVQIGNKLIILGGFGPG
NIPSNAMQTVSIYLPKPKQIASIASGNETITVWKYDTNAGYYITYWSN
FSQKVTINVGNTSYTIKHLKDGVTYIYQIVPYNLGNSTPSDIISATPS
SVPNPPIIKVKIGNLNATLTWYDTFNGGYPYIEGYLYVNGKGINVGNITS
VYLTNLTAGELYTIELIAYNKIGNSSISSVSFIAASKANLTVTVYKING

-continued

FLVSWNSTKAKYILTVSKENVLLNVSTTNTSYFVKVPGVYINISLEAV
NIVGITYAFILIIYIYQASPVTVNWISITLNTVSLNWSKVGAEYLIYDN
GKLI TTTNTAFTFNLTIGQNEIEVYAANAYYKSAPYIINDVRNIVVNV
STAISISVPOIKVSGENTDAPLQTNIDLKSALIVITVFVIALLMILVI
LRERSDNYW.

[0111] In certain aspects, the FN3 domain comprises beta strand A, beta strand B, beta strand C, beta strand D, beta strand E, beta strand F, and beta strand G. Connecting beta strands A, B, C, D, E, F, and G are loop regions AB, BC, CD, DE, EF, and FG. Beta strand A precedes the AB loop and beta strand G follows the FG loop. The loop regions correspond to the following amino acid positions in SEQ ID NO:1—AB (13-16), BC (22-27), CD (36-37), DE (47-49), EF (54-60), and FG (71-80). The loop regions in STOFN3-2, STOFN3-3, and STOFN3-4 are the corresponding amino acids in SEQ ID NO:2, 3, and 4, respectively. The corresponding amino acids can be determined from the alignment of FIG. 1

[0112] FN3 polypeptides can be modified by inserting or deleting 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20 or more amino acids, or any range derivable therein, in a FN3 loop. Variants are discussed in U.S. Pat. No. 6,673,901, which is hereby incorporated by reference with respect to embodiments regarding FN3 monobodies.

[0113] *B. Pyrococcus horikoshii*

[0114] The inventors also characterized a FN3 domain in the sequence of putative uncharacterized protein PH0954 from the hyperthermophilic archaeon *Pyrococcus horikoshii* OT3, termed PHOFN3, as the candidate protein, because of its detectably homology to a bacterial FN3 domain in the *Clostridium Perfringens* Glycoside Hydrolase Gh84c whose FN3 fold has been experimentally confirmed. The SMART database predicted PHOFN3 with 108 (P1873-A1980) amino acid residues, but the C-terminal 26 residues (E1955-A1980) did not have detectable homology to the sequence of the bacterial homologue (FIG. 9) and shown below:

(SEQ ID NO: 6)
PSPPSGVTMLNGSYVELSWLPSDSDVAGYFIYKDGKRLNEVPIEKPNF
RDIYSGTLNYSISAI DFGFSEKTEVFPVKLEVDENLTAGYPGAVKVK
VENLDGEA.

[0115] The full-length PH0954 comprises the sequence:

(SEQ ID NO: 7)
MINIKGLILTLILFISLIPPWALGEGSKDTKVFADYLAGDSVVINATLYDAGSCNLT
SVFSPIEAPNVSEISFTWMNLSEYIES ATEATYGEYLRDGNVIMREDDGYFIYELPFS
NYFGREIKKIAVNTNGLIELLEEEYEPRIEDYYGIHEEGEFYSDVIFGLDEDLVTYD
YLLLVNLQDKIVIEWLASTYEDYSEIVDNINPQVIINSNGTITWSYKSLSEYHYDYL
PSGYYSKVGSDVKGFTKGEKSFQVPLGTPKLYTYQVRESGSYLLTLPLSNYHVE
VFANCMDDPDLNLAEVGVWPGDYWVENASINNLIPGEFASINFKVRTTSKIPSAK
VKLLRNGVEEKIEYLSFYNGIAEGEISWLVQGGNYTLALLVEGKGDINSNNIYLLGN
YNFPLPNFEVGNYSIDLPTCVDSTGEVRVNVSTANWSIPVRLTLVYEEGNRSYTRYI
STKGEESSEVIFTPMIKAGTLEKVVIEIDPWNVEESNESDNKVEVPYHIIIEKPDFTK

-continued

SLNIPGNVSIIGNLYEVNVTLDNLGGCYGRNVLVKLYENGTSKDWRVRINNETNVT
 LTKWPGNAGLVNLTVVVDPYSYVDEINEGNRLSRLIFVNAPDFKISKVELLSFDGIA
 GSKAKFNVTVKNEGEDYSGYFSIAVYGLRSSIAYLRGIKSSEEKWTIISLPINGGNST
 LIFVVDPHNVISSETNEGNVIFYNMGYIPKPNFVVKIEISLPNNTVGYIPLNITIGNVGAP
 YNATSYQVPVKIKTEYGWKVSYLRGBIIRDNYTISIDSLAMLPPGSTINVTVNYNMKV
 NETSYSDNSLIINYTTGYPDLELGIIPPSGELSAGKDVKITFLVKVNGNATLRIDRSSW
 YSPYLGPLYVTLEDENGKHTLGRYELAPATLSPGANISQVVWITLNGGTNKIMGRIV
 DEYENIYENNDTLILITLEKPDFAILNYSIPDEILNGTAYLYKAYPIVLNINSLGGNFSD
 GIRVDFLFDNGIIKTSTSVYGLESGASRDVTLRYLPSSGKHNSIVLDPYNRWIEEENEEN
 NNLTFSLSFGKPDLEKVEGITWAPYNFTSGENVLFITYVKNLQOPFLKSFTVRAEIIWNG
 TRKIYSTNAYPRNWSFGKGETKEFNWRWYNAKPGNLTVKIVVDYNSIPEGNESNN
 EFSAPLGNVGTPDFKLENLSVEDLAYGKFRINATVKNLGDSEIYRPI TVLFPVSGERY
 YRTVYGIKENESKSVTLWPYVDRVGEVVRVKEVDPGNRIVEGNESNNIERTYYVES
 PELMLSGYEWLEEEVRRGYLAYKVNVTNTGGDVYRGFVYQMFVDGEPKSSVWINK
 LLHGETAERTLRWRPSSGGRKEVRI VVDPQDYIPESNEDNNAIVENVTI VLPDIEVLSL
 NIPSMHANSYFKVNATIKNSSGQDVKRIFYVSLYQDGKLLGSAPVYSLASGEVKEVT
 LTIRPYPGNSTFKVVDPTNAVVELNEDNNEISVRSYVKAPDIVVVSADLGNFTYPGE
 MVNAKVRIRNSGDYKSGVYLLIRNKRRLGSAYVDSITPGEEIEVNVVPLVDSGDY
 NVSVIADPYNVREWEENNKL DIEVSVSPDLTVENITHSGKEVAGEEIIKVTVKNI
 GESSKLPFYIVLYANSSFVGINRVTKIDKGESITLFEKWRASYGEYALRAIVDPYDEV
 YEENESNNEGMVKVFIEDEEPPVLKLTYPENGTFTNKPYIGAYLRDEGSGVKFGEIEV
 YREGTSVPGSTKFSGGWLI FQNSTPLLDGKYTVTVKAVDRAGNEITYSWNFTLDREP
 PRIVCNLTDGTYLNGTVVPGVQVIDDNLWDYKVKVNGREFSGPIKLDGTYTLNVT
 KDKAGNLAEKIIRFTVNGVPSPPSGVTLMLNGSYVELSWLPSDSDVAGYFIYKDGK
 RLNEVP IEKPNFRDIYSGTLNYSISAIDFSGFSEKTEVFPVKLEVDEENLTAGYPGAV
 KVKVENLDGEANGTLSIILIDFEGNEIEKLSRKVEVPRGRSSHEFVFMVPRGLTLIRGE
 LKVGNSTARIHRAKVRAGENPEIRVGKLLAGFPGLVEVEIRNCGIVELNTSETLMKL
 DNSSGELIEAPLTI PPGKKTVLRKIVPPKKSYNLTFRIADVEVRKIVNVSESVLNPIT
 ISTENFIKGGKAKIYVSPRNI GSAPIFVKSIELNGMSKRLSIELPPNLSVEESFEYLI SEEN
 VEINATVNTDVGKFRKSLTLTAEQPEYNADVSVSVYEVGKEILITGVAYNESGMLS
 NVPVKVSIARGGFVREYIVTTNENGYFNLTFRPPFKGESGHFIVSATHPKIELLERDAEF
 DVVGIIEVIPSPLYLLTVPVEFNGTVRRLIN YWRASDVSVSVKAPPEYEVSI PKVLHLK
 PGSNIINIGLSSKNAVNGSIMITFKARQLGLNITRSLTLKLVLPAPAIVTS PNFLDVG
 VLTNETASAEVVVRNLGPTALRNVSIRSSIPVVKVSNFTEVDPKDNESISLYIEPPRN
 VTGTFKGEITISSSNYNPIKVPMRIRVTPNATGGVKVTVMDPNATRL ENVKLTLYNG
 YFHFEGYTNKNGTLEVENVPIGEYKLFASLEGYGYSTSI TIEVGVKEKNSIILTPSILE
 VEWEVVPVTIQDVIYIIKHEMWYSTHVPAPEIRMEGGDLEVYVDYEKLAEEGMLEFR
 GQVIVRNTHQYISVYNVTFESGGSHYIDVEFGINRIDE LKPGEAIVPYVVR IYYSRSP
 PINPCLHETKVKFLKAGVVVCEEAGKITLKAQR IHQIVVKPTCKGCWESVFPVAGKL

-continued

AFMAIAQKVGQALGNIDDTGVLSTLAGEALNNLESFDAYNAYKANPTKENMENY
 VKTFNSVKANLASLFMFDPVAYQEINSLQLTLIKTPKGDIAGFAVSRATATPVYALGM
 GVGKIENGQLKVDYKKAJNIANGIVLVNMSKMGALGGIASGVGLLQLLDDKAAED
 LPPYIAQLFLNCAICLMRNDCTLPEGEEIRPIQIIASGSLGGYPSMIPSGLAGGGGGTA
 VGRFTCGGLPTVKKSSSTSMSCSTCSSEDEVVKERVCRLPRESETHDEEPEPSNTLHMCV
 DLVLTIEQRLTFERQAFRASLKFTNTNRNYSLENVSVRVIFFDEEENRVDKFFVRLD
 EKAGLSGSLEPEKTAEMKWLII PKVGAEEKFRARYYVMANITARVGSTKLVYETW
 PAMIEVEVPVQLVLDYVLPVSYVFGDDPYTPEKELPIPIFGVVRVKNVGYGTARKLRIA
 SAQPKIERSNYPGVYIDFKIIGTLVNGKKVNSLTIDFGDLNPGESSTAAWLMIAEVSG
 KFLQYNATFKHSDELGGNETSLIKEVRTHFLIRAFNNTENDDGMDFLVDDDDGDKPK
 EKIIDSRGFDYNVLLLNFTVEEESMRKIIPEMKTPEFWVYFTVPFKGSVVRSDGKNPM
 DQWMENGLHVLDLGTPEFYILKSNQPPRIYVKEPVIANETVVLDDGSLSYDPDGSII
 AYTWKIGNESFVGDKVSYVREPGTYNVTLRDCKGTESSTMEIKVYLGPKFNES
 LKVEPQWGI VPPNLSITFNVTNVGDVSGEYSYIIKLGNSTIAEGSEIIESGRWKVINSTV
 EIRKEGNYTVTANNLSKTVTAYRKVYGNLTENYIKEKDFGHYKSFYWNFKRDFEG
 WVVEALSTIELPKVNFVNLNYPFGNWSLLNYSEMLNITKGWGINATYARRVREGE
 LEEFKYLI VNVTLQVLLGNATHELDESPTLNVTSSGIYSEIPKIQVRTCDEGTITLV
 WGA VGN YTKEFTEVESNGTCSWEGIVPLNIGNNTVAIYAEDEFGNRGNVSLWIIYLN
 PEAPVYIIESPEEKVYNSREVMINYTVVNHDLVGVVAYLNGELISSNASYSGFIKLDY
 GWHNFTIYAWDVSYNVSKSVIFRVNEPPSVDFSWEVNDLTVKFEANASDEDGISKYL
 WDFGDNESSLLVNPHTHYRKGGRYNVTLTVWDSYNLSSISKEVVVFGSSTLTMVK
 EYSYTKDFGPFYNTTSWKDFLKDPEVWVNLTLRNVTLPLEYFEEIIEVNVENWSLISVE
 KNLKNDIGEMSAEYERNATIVGIMNYTRVTLKLTQEVILSGRARKVEDKIPPLVEILFP
 RNMTYNETIREIKVRATDESGIANVTATINGESLSLEKVNETHWIGRVELDDGKYELNV
 FASDKWGNVGCSTVNFTINRSVKVRIINGTEIVTIPGDIKTRVYFEGDIIIVEIVKESLRF
 KIPSGGTLVIDERGRKDPWLLARINSTIENISKTSRIFEENGKKVHEIRYRISISRGYAIL
 VVPLEGMKVSSIRIINKGTVTREDEKHGNYKLSKGYLFIPLSEDPIVEVTLTKIEKKDIF
 RVLYYAGI IWERNYRLRKEEFIMKMSNETSQEAIRLHEEAKEYYKGREYYPRIPSPS
 AIYWYAVYMRKAYLTERKALELLSIS.

[0116] In certain aspects, the FN3 domain comprises beta strand A, beta strand B, beta strand C, beta strand D, beta strand E, beta strand F, and beta strand G. Connecting beta strands A, B, C, D, E, F, and G are loop regions AB, BC, CD, DE, EF, and FG. Beta strand A precedes the AB loop and beta strand G follows the FG loop. The loop regions correspond to the following amino acid positions in SEQ ID NO:6—AB (13-14) BC (20-27), CD (36-37), DE (40-48), EF (54-59), and FG (67-74).

[0117] FN3 polypeptides can be modified by inserting or deleting 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20 or more amino acids, or any range derivable therein, in a FN3 loop. Variants are discussed in U.S. Pat. No. 6,673,901, which is hereby incorporated by reference with respect to embodiments regarding FN3 monobodies.

[0118] C. FN3 Library

[0119] A combinatorial library is a collection of diverse compounds generated by either chemical synthesis or biological synthesis by combining a number of chemical “building blocks.” For example, a linear combinatorial chemical library such as a polypeptide (e.g., mutein or variant) library is formed by combining a set of chemical building blocks called amino acids in every possible way for a given compound length. Millions of compounds can be synthesized through such combinatorial mixing of chemical building blocks. For example, one commentator has observed that the systematic, combinatorial mixing of 100 interchangeable chemical building blocks results in the theoretical synthesis of 100 million tetrameric compounds or 10 billion pentameric compounds (Gallop et al., 1994).

[0120] Embodiments of the disclosure are directed to a combinatorial library of FN3 domains. In certain aspects, polypeptides of the library include variations of amino acid sequence in one or more of the beta strands or body of the FN3 domains. In certain aspects, the library includes variations of amino acid sequences in one or more loops of the FN3 domains. In still further aspects, the library includes variation in both loops and beta strands of the FN3 domain.

[0121] FN3 variants can include alanine substitutions at one or more of amino acid positions. In certain aspects, any of the 19 other amino acids can be substituted for one or more amino acid of SEQ ID NO:1-4 or 6. Substitutions include, but are not limited to conservative substitutions that have little or no effect on the overall net charge, polarity, or hydrophobicity of the protein.

[0122] In certain aspects, FN3 domains will have, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 amino acid substitutions that include, but are not limited to the following FN3 residue substitutions (corresponding to SEQ ID NO:1): K4V, N14P, D58P, G28Y, Y22A, Y22C, Y22D, Y22E, Y22F, Y22G, Y22H, Y22I, Y22K, Y22L, Y22M, Y22N, Y22P, Y22Q, Y22R, Y22S, Y22T, Y22V, Y22W, D23A, D23C, D23E, D23F, D23G, D23H, D23I, D23K, D23L, D23M, D23N, D23P, D23Q, D23R, D23S, D23T, D23V, D23W, D23Y, T24A, T24C, T24D, T24E, T24F, T24G, T24H, T24I, T24K, T24L, T24M, T24N, T24P, T24Q, T24R, T24S, T24V, T24W, T24Y, N25A, N25C, N25D, N25E, N25F, N25G, N25H, N25I, N25K, N25L, N25M, N25P, N25Q, N25R, N25S, N25T, N25V, N25W, N25Y, A26C, A26D, A26E, A26F, A26G, A26H, A26I, A26K, A26L, A26M, A26N, A26P, A26Q, A26R, A26S, A26T, A26V, A26W, A26Y, S27A, S27C, S27D, S27E, S27F, S27G, S27H, S27I, S27K, S27L, S27M, S27N, S27P, S27Q, S27R, S27T, S27V, S27W, S27Y, N47A, N47C, N47D, N47E, N47F, N47G, N47H, N47I, N47K, N47L, N47M, N47P, N47Q, N47R, N47S, N47T, N47V, N47W, N47Y, V48A, V48C, V48D, V48E, V48F, V48G, V48H, V48I, V48K, V48L, V48M, V48N, V48P, V48Q, V48R, V48S, V48T, V48W, V48Y, T49A, T49C, T49D, T49E, T49F, T49G, T49H, T49I, T49K, T49L, T49M, T49N, T49P, T49Q, T49R, T49S, T49V, T49W, T49Y, S71A, S71C, S71D, S71E, S71F, S71G, S71H, S71I, S71K, S71L, S71M, S71N, S71P, S71Q, S71R, S71T, S71V, S71W, S71Y, L72A, L72C, L72D, L72E, L72F, L72G, L72H, L72I, L72K, L72M, L72N, L72P, L72Q, L72R, L72S, L72T, L72V, L72W, L72Y, G73A, G73C, G73D, G73E, G73F, G73H, G73I, G73K, G73L, G73M, G73N, G73P, G73Q, G73R, G73S, G73T, G73V, G73W, G73Y, N74A, N74C, N74D, N74E, N74F, N74G, N74H, N74I, N74K, N74L, N74M, N74P, N74Q, N74R, N74S, N74T, N74V, N74W, N74Y, G75A, G75C, G75D, G75E, G75F, G75H, G75I, G75K, G75L, G75M, G75N, G75P, G75Q, G75R, G75S, G75T, G75V, G75W, G75Y, T76A, T76C, T76D, T76E, T76F, T76G, T76H, T76I, T76K, T76L, T76M, T76N, T76P, T76Q, T76R, T76S, T76V, T76W, T76Y, P77A, P77C, P77D, P77E, P77F, P77G, P77H, P77I, P77K, P77L, P77M, P77N, P77Q, P77R, P77S, P77T, P77V, P77W, P77Y, S78A, S78C, S78D, S78E, S78F, S78G, S78H, S78I, S78K, S78L, S78M, S78N, S78P, S78Q, S78R, S78T, S78V, S78W, S78Y, D79A, D79C, D79E, D79F, D79G, D79H, D79I, D79K, D79L, D79M, D79N, D79P, D79Q, D79R, D79S, D79T, D79V, D79W, D79Y, I80A, I80C, I80D, I80E, I80F, I80G,

I80H, I80K, I80L, I80M, I80N, I80P, I80Q, I80R, I80S, I80T, I80V, I80W, I80Y, N36A, N36C, N36D, N36E, N36F, N36G, N36H, N36I, N36K, N36L, N36M, N36P, N36Q, N36R, N36S, N36T, N36V, N36W, N36Y, F37A, F37C, F37D, F37E, F37G, F37H, F37I, F37K, F37L, F37M, F37N, F37P, F37Q, F37R, F37S, F37T, F37V, F37W, F37Y, G28A, G28C, G28D, G28E, G28F, G28H, G28I, G28K, G28L, G28M, G28N, G28P, G28Q, G28R, G28S, G28T, G28V, G28W, G28Y, Y29A, Y29C, Y29D, Y29E, Y29F, Y29G, Y29H, Y29I, Y29K, Y29L, Y29M, Y29N, Y29P, Y29Q, Y29R, Y29S, Y29T, Y29V, Y29W, Y30A, Y30C, Y30D, Y30E, Y30F, Y30G, Y30H, Y30I, Y30K, Y30L, Y30M, Y30N, Y30P, Y30Q, Y30R, Y30S, Y30T, Y30V, Y30W, I31A, I31C, I31D, I31E, I31F, I31G, I31H, I31K, I31L, I31M, I31N, I31P, I31Q, I31R, I31S, I31T, I31V, I31W, I31Y, T32A, T32C, T32D, T32E, T32F, T32G, T32H, T32I, T32K, T32L, T32M, T32N, T32P, T32Q, T32R, T32S, T32V, T32W, T32Y, Y33A, Y33C, Y33D, Y33E, Y33F, Y33G, Y33H, Y33I, Y33K, Y33L, Y33M, Y33N, Y33P, Y33Q, Y33R, Y33S, Y33T, Y33V, Y33W, W34A, W34C, W34D, W34E, W34F, W34G, W34H, W34I, W34K, W34L, W34M, W34N, W34P, W34Q, W34R, W34S, W34T, W34V, W34Y, S35A, S35C, S35D, S35E, S35F, S35G, S35H, S35I, S35K, S35L, S35M, S35N, S35P, S35Q, S35R, S35T, S35V, S35W, S35Y, S38A, S38C, S38D, S38E, S38F, S38G, S38H, S38I, S38K, S38L, S38M, S38N, S38P, S38Q, S38R, S38T, S38V, S38W, S38Y, Q39A, Q39C, Q39D, Q39E, Q39F, Q39G, Q39H, Q39I, Q39K, Q39L, Q39M, Q39N, Q39P, Q39R, Q39S, Q39T, Q39V, Q39W, Q39Y, K40A, K40C, K40D, K40E, K40F, K40G, K40H, K40I, K40L, K40M, K40N, K40P, K40Q, K40R, K40S, K40T, K40V, K40W, K40Y, V41A, V41C, V41D, V41E, V41F, V41G, V41H, V41I, V41K, V41L, V41M, V41N, V41P, V41Q, V41R, V41S, V41T, V41W, V41Y, T42A, T42C, T42D, T42E, T42F, T42G, T42H, T42I, T42K, T42L, T42M, T42N, T42P, T42Q, T42R, T42S, T42V, T42W, T42Y, I43A, I43C, I43D, I43E, I43F, I43G, I43H, I43K, I43L, I43M, I43N, I43P, I43Q, I43R, I43S, I43T, I43V, I43W, I43Y, N44A, N44C, N44D, N44E, N44F, N44G, N44H, N44I, N44K, N44L, N44M, N44P, N44Q, N44R, N44S, N44T, N44V, N44W, N44Y, V45A, V45C, V45D, V45E, V45F, V45G, V45H, V45I, V45K, V45L, V45M, V45N, V45P, V45Q, V45R, V45S, V45T, V45W, V45Y, G46A, G46C, G46D, G46E, G46F, G46H, G46I, G46K, G46L, G46M, G46N, G46P, G46Q, G46R, G46S, G46T, G46V, G46W, and/or G46Y, and combinations thereof. It is contemplated that one or more of these substitutions may be specifically excluded in embodiments described herein.

[0123] In still further embodiments other amino acid substitutions can be introduced before, during, or after introduction of those amino acid substitutions listed above. Further substitutions (corresponding to SEQ ID NO:1) include, but is not limited to 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 of P1A, P1C, P1D, P1E, P1F, P1G, P1H, P1I, P1K, P1L, P1M, P1N, P1Q, P1R, P1S, P1T, P1V, P1W, P1Y, P2A, P2C, P2D, P2E, P2F, P2G, P2H, P2I, P2K, P2L, P2M, P2N, P2Q, P2R, P2S, P2T, P2V, P2W, P2Y, P3A, P3C, P3D, P3E, P3F, P3G, P3H, P3I, P3K, P3L, P3M, P3N, P3Q, P3R, P3S, P3T, P3V, P3W, P3Y, K4A, K4C, K4D, K4E, K4F, K4G, K4H, K4I, K4L, K4M, K4N, K4P, K4Q, K4R, K4S, K4T, K4V, K4W, K4Y, P5A, P5C, P5D, P5E, P5F, P5G, P5H, P5I, P5K, P5L, P5M, P5N, P5Q, P5R, P5S,

P5T, P5V, P5W, P5Y, Q6A, Q6C, Q6D, Q6E, Q6F, Q6G, Q6H, Q6I, Q6K, Q6L, Q6M, Q6N, Q6P, Q6R, Q6S, Q6T, Q6V, Q6W, Q6Y, I7A, I7C, I7D, I7E, I7F, I7G, I7H, I7K, I7L, I7M, I7N, I7P, I7Q, I7R, I7S, I7T, I7V, I7W, I7Y, ABC, A8D, A8E, A8F, A8G, A8H, A8I, A8K, A8L, A8M, A8N, A8P, A8Q, A8R, A8S, A8T, A8V, A8W, A8Y, S9A, S9C, S9D, S9E, S9F, S9G, S9H, S9I, S9K, S9L, S9M, S9N, S9P, S9Q, S9R, S9T, S9V, S9W, S9Y, I10A, I10C, I10D, I10E, I10F, I10G, I10H, I10K, I10L, I10M, I10N, I10P, I10Q, I10R, I10S, I10T, I10V, I10W, I10Y, A11C, A11D, A11E, A11F, A11G, A11H, A11I, A11K, A11L, A11M, A11N, A11P, A11Q, A11R, A11S, A11T, A11V, A11W, A11Y, S12A, S12C, S12D, S12E, S12F, S12G, S12H, S12I, S12K, S12L, S12M, S12N, S12P, S12Q, S12R, S12T, S12V, S12W, S12Y, G13A, G13C, G13D, G13E, G13F, G13H, G13I, G13K, G13L, G13M, G13N, G13P, G13Q, G13R, G13S, G13T, G13V, G13W, G13Y, N14A, N14C, N14D, N14E, N14F, N14G, N14H, N14I, N14K, N14L, N14M, N14P, N14Q, N14R, N14S, N14T, N14V, N14W, N14Y, E15A, E15C, E15D, E15F, E15G, E15H, E15I, E15K, E15L, E15M, E15N, E15P, E15Q, E15R, E15S, E15T, E15V, E15W, E15Y, T16A, T16C, T16D, T16E, T16F, T16G, T16H, T16I, T16K, T16L, T16M, T16N, T16P, T16Q, T16R, T16S, T16V, T16W, T16Y, I17A, I17C, I17D, I17E, I17F, I17G, I17H, I17K, I17L, I17M, I17N, I17P, I17Q, I17R, I17S, I17T, I17V, I17W, I17Y, T18A, T18C, T18D, T18E, T18F, T18G, T18H, T18I, T18K, T18L, T18M, T18N, T18P, T18Q, T18R, T18S, T18V, T18W, T18Y, V19A, V19C, V19D, V19E, V19F, V19G, V19H, V19I, V19K, V19L, V19M, V19N, V19P, V19Q, V19R, V19S, V19T, V19W, V19Y, K20A, K20C, K20D, K20E, K20F, K20G, K20H, K20I, K20L, K20M, K20N, K20P, K20Q, K20R, K20S, K20T, K20V, K20W, K20Y, W21A, W21C, W21D, W21E, W21F, W21G, W21H, W21I, W21K, W21L, W21M, W21N, W21P, W21Q, W21R, W21S, W21T, W21V, W21Y, S50A, S50C, S50D, S50E, S50F, S50G, S50H, S50I, S50K, S50L, S50M, S50N, S50P, S50Q, S50R, S50T, S50V, S50W, S50Y, Y51A, Y51C, Y51D, Y51E, Y51F, Y51G, Y51H, Y51I, Y51K, Y51L, Y51M, Y51N, Y51P, Y51Q, Y51R, Y51S, Y51T, Y51V, Y51W, T52A, T52C, T52D, T52E, T52F, T52G, T52H, T52I, T52K, T52L, T52M, T52N, T52P, T52Q, T52R, T52S, T52V, T52W, T52Y, I53A, I53C, I53D, I53E, I53F, I53G, I53H, I53K, I53L, I53M, I53N, I53P, I53Q, I53R, I53S, I53T, I53V, I53W, I53Y, K54A, K54C, K54D, K54E, K54F, K54G, K54H, K54I, K54L, K54M, K54N, K54P, K54Q, K54R, K54S, K54T, K54V, K54W, K54Y, H55A, H55C, H55D, H55E, H55F, H55G, H55I, H55K, H55L, H55M, H55N, H55P, H55Q, H55R, H55S, H55T, H55V, H55W, H55Y, L56A, L56C, L56D, L56E, L56F, L56G, L56H, L56I, L56K, L56M, L56N, L56P, L56Q, L56R, L56S, L56T, L56V, L56W, L56Y, K57A, K57C, K57D, K57E, K57F, K57G, K57H, K57I, K57L, K57M, K57N, K57P, K57Q, K57R, K57S, K57T, K57V, K57W, K57Y, D58A, D58C, D58E, D58F, D58G, D58H, D58I, D58K, D58L, D58M, D58N, D58P, D58Q, D58R, D58S, D58T, D58V, D58W, D58Y, G59A, G59C, G59D, G59E, G59F, G59H, G59I, G59K, G59L, G59M, G59N, G59P, G59Q, G59R, G59S, G59T, G59V, G59W, G59Y, V60A, V60C, V60D, V60E, V60F, V60G, V60H, V60I, V60K, V60L, V60M, V60N, V60P, V60Q, V60R, V60S, V60T, V60W, V60Y, T61A, T61C, T61D, T61E, T61F, T61G, T61H, T61I, T61K, T61L, T61M, T61N, T61P, T61Q, T61R, T61S, T61V, T61W, T61Y, Y62A, Y62C, Y62D, Y62E, Y62F, Y62G, Y62H, Y62I,

Y62K, Y62L, Y62M, Y62N, Y62P, Y62Q, Y62R, Y62S, Y62T, Y62V, Y62W, Y63A, Y63C, Y63D, Y63E, Y63F, Y63G, Y63H, Y63I, Y63K, Y63L, Y63M, Y63N, Y63P, Y63Q, Y63R, Y63S, Y63T, Y63V, Y63W, I64A, I64C, I64D, I64E, I64F, I64G, I64H, I64K, I64L, I64M, I64N, I64P, I64Q, I64R, I64S, I64T, I64V, I64W, I64Y, Q65A, Q65C, Q65D, Q65E, Q65F, Q65G, Q65H, Q65I, Q65K, Q65L, Q65M, Q65N, Q65P, Q65R, Q65S, Q65T, Q65V, Q65W, Q65Y, I66A, I66C, I66D, I66E, I66F, I66G, I66H, I66K, I66L, I66M, I66N, I66P, I66Q, I66R, I66S, I66T, I66V, I66W, I66Y, V67A, V67C, V67D, V67E, V67F, V67G, V67H, V67I, V67K, V67L, V67M, V67N, V67P, V67Q, V67R, V67S, V67T, V67W, V67Y, P68A, P68C, P68D, P68E, P68F, P68G, P68H, P68I, P68K, P68L, P68M, P68N, P68Q, P68R, P68S, P68T, P68V, P68W, P68Y, Y69A, Y69C, Y69D, Y69E, Y69F, Y69G, Y69H, Y69I, Y69K, Y69L, Y69M, Y69N, Y69P, Y69Q, Y69R, Y69S, Y69T, Y69V, Y69W, N70A, N70C, N70D, N70E, N70F, N70G, N70H, N70I, N70K, N70L, N70M, N70P, N70Q, N70R, N70S, N70T, N70V, N70W, N70Y, I81A, I81C, I81D, I81E, I81F, I81G, I81H, I81K, I81L, I81M, I81N, I81P, I81Q, I81R, I81S, I81T, I81V, I81W, I81Y, S82A, S82C, S82D, S82E, S82F, S82G, S82H, S82I, S82K, S82L, S82M, S82N, S82P, S82Q, S82R, S82T, S82V, S82W, S82Y, A83C, A83D, A83E, A83F, A83G, A83H, A83I, A83K, A83L, A83M, A83N, A83P, A83Q, A83R, A83S, A83T, A83V, A83W, A83Y, T84A, T84C, T84D, T84E, T84F, T84G, T84H, T84I, T84K, T84L, T84M, T84N, T84P, T84Q, T84R, T84S, T84T, T84V, T84W, or T84Y or combinations thereof. It is contemplated that one or more of these substitutions may be specifically excluded in embodiments described herein.

[0124] In certain aspects, FN3 domains will have 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 amino acid substitutions that include, but are not limited to the following FN3 residue substitutions (corresponding to SEQ ID NO:6): G13A, G13C, G13D, G13E, G13F, G13H, G13I, G13K, G13L, G13M, G13N, G13P, G13Q, G13R, G13S, G13T, G13V, G13W, G13Y, S14A, S14C, S14D, S14E, S14F, S14G, S14H, S14I, S14K, S14L, S14M, S14N, S14P, S14Q, S14R, S14T, S14V, S14W, S14Y, Y15A, W20A, W20C, W20D, W20E, W20F, W20G, W20H, W20I, W20K, W20L, W20M, W20N, W20P, W20Q, W20R, W20S, W20T, W20V, W20Y, L21A, L21C, L21D, L21E, L21F, L21G, L21H, L21I, L21K, L21M, L21N, L21P, L21Q, L21R, L21S, L21T, L21V, L21W, L21Y, P22A, P22C, P22D, P22E, P22F, P22G, P22H, P22I, P22K, P22L, P22M, P22N, P22Q, P22R, P22S, P22T, P22V, P22W, P22Y, S23A, S23C, S23D, S23E, S23F, S23G, S23H, S23I, S23K, S23L, S23M, S23N, S23P, S23Q, S23R, S23T, S23V, S23W, S23Y, P24A, P24C, P24D, P24E, P24F, P24G, P24H, P24I, P24K, P24L, P24M, P24N, P24Q, P24R, P24S, P24T, P24V, P24W, P24Y, D25A, D25C, D25E, D25F, D25G, D25H, D25I, D25K, D25L, D25M, D25N, D25P, D25Q, D25R, D25S, D25T, D25V, D25W, D25Y, S26A, S26C, S26D, S26E, S26F, S26G, S26H, S26I, S26K, S26L, S26M, S26N, S26P, S26Q, S26R, S26T, S26V, S26W, S26Y, D27A, D27C, D27E, D27F, D27G, D27H, D27I, D27K, D27L, D27M, D27N, D27P, D27Q, D27R, D27S, D27T, D27V, D27W, D27Y, D36A, D36C, D36E, D36F, D36G, D36H, D36I, D36K, D36L, D36M, D36N, D36P, D36Q, D36R, D36S, D36T, D36V, D36W, D36Y, G37A, G37C, G37D, G37E, G37F, G37H, G37I, G37K, G37L, G37M, G37N, G37P, G37Q, G37R,

G37S, G37T, G37V, G37W, G37Y, L40A, L40C, L40D, L40E, L40F, L40G, L40H, L40I, L40K, L40M, L40N, L40P, L40Q, L40R, L40S, L40T, L40V, L40W, L40Y, N41A, N41C, N41D, N41E, N41F, N41G, N41H, N41I, N41K, N41L, N41M, N41P, N41Q, N41R, N41S, N41T, N41V, N41W, N41Y, E42A, E42C, E42D, E42F, E42G, E42H, E42I, E42K, E42L, E42M, E42N, E42P, E42Q, E42R, E42S, E42T, E42V, E42W, E42Y, V43A, V43C, V43D, V43E, V43F, V43G, V43H, V43I, V43K, V43L, V43M, V43N, V43P, V43Q, V43R, V43S, V43T, V43W, V43Y, P44A, P44C, P44D, P44E, P44F, P44G, P44H, P44I, P44K, P44L, P44M, P44N, P44Q, P44R, P44S, P44T, P44V, P44W, P44Y, I45A, I45C, I45D, I45E, I45F, I45G, I45H, I45K, I45L, I45M, I45N, I45P, I45Q, I45R, I45S, I45T, I45V, I45W, I45Y, E46A, E46C, E46D, E46F, E46G, E46H, E46I, E46K, E46L, E46M, E46N, E46P, E46Q, E46R, E46S, E46T, E46V, E46W, E46Y, K47A, K47C, K47D, K47E, K47F, K47G, K47H, K47I, K47L, K47M, K47N, K47P, K47Q, K47R, K47S, K47T, K47V, K47W, K47Y, P48A, P48C, P48D, P48E, P48F, P48G, P48H, P48I, P48K, P48L, P48M, P48N, P48Q, P48R, P48S, P48T, P48V, P48W, P48Y, Y54A, Y54C, Y54D, Y54E, Y54F, Y54G, Y54H, Y54I, Y54K, Y54L, Y54M, Y54N, Y54P, Y54Q, Y54R, Y54S, Y54T, Y54V, Y54W, S55A, S55C, S55D, S55E, S55F, S55G, S55H, S55I, S55K, S55L, S55M, S55N, S55P, S55Q, S55R, S55T, S55V, S55W, S55Y, G56A, G56C, G56D, G56E, G56F, G56H, G56I, G56K, G56L, G56M, G56N, G56P, G56Q, G56R, G56S, G56T, G56V, G56W, G56Y, T57A, T57C, T57D, T57E, T57F, T57G, T57H, T57I, T57K, T57L, T57M, T57N, T57P, T57Q, T57R, T57S, T57V, T57W, T57Y, L58A, L58C, L58D, L58E, L58F, L58G, L58H, L58I, L58K, L58M, L58N, L58P, L58Q, L58R, L58S, L58T, L58V, L58W, L58Y, N59A, N59C, N59D, N59E, N59F, N59G, N59H, N59I, N59K, N59L, N59M, N59P, N59Q, N59R, N59S, N59T, N59V, N59W, N59Y, F67A, F67C, F67D, F67E, F67G, F67H, F67I, F67K, F67L, F67M, F67N, F67P, F67Q, F67R, F67S, F67T, F67V, F67W, F67Y, S68A, S68C, S68D, S68E, S68F, S68G, S68H, S68I, S68K, S68L, S68M, S68N, S68P, S68Q, S68R, S68T, S68V, S68W, S68Y, G69A, G69C, G69D, G69E, G69F, G69H, G69I, G69K, G69L, G69M, G69N, G69P, G69Q, G69R, G69S, G69T, G69V, G69W, G69Y, F70A, F70C, F70D, F70E, F70G, F70H, F70I, F70K, F70L, F70M, F70N, F70P, F70Q, F70R, F70S, F70T, F70V, F70W, F70Y, E71A, E71C, E71D, E71F, E71G, E71H, E71I, E71K, E71L, E71M, E71N, E71P, E71Q, E71R, E71S, E71T, E71V, E71W, E71Y, S72A, S72C, S72D, S72E, S72F, S72G, S72H, S72I, S72K, S72L, S72M, S72N, S72P, S72Q, S72R, S72T, S72V, S72W, S72Y, E73A, E73C, E73D, E73F, E73G, E73H, E73I, E73K, E73L, E73M, E73N, E73P, E73Q, E73R, E73S, E73T, E73V, E73W, E73Y, K74A, K74C, K74D, K74E, K74F, K74G, K74H, K74I, K74L, K74M, K74N, K74P, K74Q, K74R, K74S, K74T, K74V, K74W, and/or K74Y or combinations thereof. It is contemplated that one or more of these substitutions may be specifically excluded in embodiments described herein.

[0125] In still further embodiments other amino acid substitutions can be introduced before, during, or after introduction of those amino acid substitutions listed above. Further substitutions (corresponding to SEQ ID NO:6) include, but is not limited to 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 of P1A, P1C, P1D, P1E, P1F, P1G,

P1H, P1I, P1K, P1L, P1M, P1N, P1Q, P1R, P1S, P1T, P1V, P1W, P1Y, S3A, S3C, S3D, S3E, S3F, S3G, S3H, S3I, S3K, S3L, S3M, S3N, S3P, S3Q, S3R, S3T, S3V, S3W, S3Y, P4A, P4C, P4D, P4E, P4F, P4G, P4H, P4I, P4K, P4L, P4M, P4N, P4Q, P4R, P4S, P4T, P4V, P4W, P4Y, S5A, S5C, S5D, S5E, S5F, S5G, S5H, S5I, S5K, S5L, S5M, S5N, S5P, S5Q, S5R, S5T, S5V, S5W, S5Y, G6A, G6C, G6D, G6E, G6F, G6H, G6I, G6K, G6L, G6M, G6N, G6P, G6Q, G6R, G6S, G6T, G6V, G6W, G6Y, V7A, V7C, V7D, V7E, V7F, V7G, V7H, V7I, V7K, V7L, V7M, V7N, V7P, V7Q, V7R, V7S, V7T, V7W, V7Y, T8A, T8C, T8D, T8E, T8F, T8G, T8H, T8I, T8K, T8L, T8M, T8N, T8P, T8Q, T8R, T8S, T8V, T8W, T8Y, L9A, L9C, L9D, L9E, L9F, L9G, L9H, L9I, L9K, L9M, L9N, L9P, L9Q, L9R, L9S, L9T, L9V, L9W, L9Y, M10A, M10C, M10D, M10E, M10F, M10G, M10H, M10I, M10K, M10L, M10N, M10P, M10Q, M10R, M10S, M10T, M10V, M10W, M10Y, L11A, L11C, L11D, L11E, L11F, L11G, L11H, L11I, L11K, L11M, L11N, L11P, L11Q, L11R, L11S, L11T, L11V, L11W, L11Y, N12A, N12C, N12D, N12E, N12F, N12G, N12H, N12I, N12K, N12L, N12M, N12P, N12Q, N12R, N12S, N12T, N12V, N12W, N12Y, Y15C, Y15D, Y15E, Y15F, Y15G, Y15H, Y15I, Y15K, Y15L, Y15M, Y15N, Y15P, Y15Q, Y15R, Y15S, Y15T, Y15V, Y15W, V16A, V16C, V16D, V16E, V16F, V16G, V16H, V16I, V16K, V16L, V16M, V16N, V16P, V16Q, V16R, V16S, V16T, V16W, V16Y, E17A, E17C, E17D, E17F, E17G, E17H, E17I, E17K, E17L, E17M, E17N, E17P, E17Q, E17R, E17S, E17T, E17V, E17W, E17Y, L18A, L18C, L18D, L18E, L18F, L18G, L18H, L18I, L18K, L18M, L18N, L18P, L18Q, L18R, L18S, L18T, L18V, L18W, L18Y, S19A, S19C, S19D, S19E, S19F, S19G, S19H, S19I, S19K, S19L, S19M, S19N, S19P, S19Q, S19R, S19T, S19V, S19W, S19Y, V28A, V28C, V28D, V28E, V28F, V28G, V28H, V28I, V28K, V28L, V28M, V28N, V28P, V28Q, V28R, V28S, V28T, V28V, V28Y, A29C, A29D, A29E, A29F, A29G, A29H, A29I, A29K, A29L, A29M, A29N, A29P, A29Q, A29R, A29S, A29T, A29V, A29W, A29Y, G30A, G30C, G30D, G30E, G30F, G30G, G30H, G30I, G30K, G30L, G30M, G30N, G30P, G30Q, G30R, G30S, G30T, G30V, G30W, G30Y, Y31A, Y31C, Y31D, Y31E, Y31F, Y31G, Y31H, Y31I, Y31K, Y31L, Y31M, Y31N, Y31P, Y31Q, Y31R, Y31S, Y31T, Y31V, Y31W, F32A, F32C, F32D, F32E, F32G, F32H, F32I, F32K, F32L, F32M, F32N, F32P, F32Q, F32R, F32S, F32T, F32V, F32W, F32Y, I33A, I33C, I33D, I33E, I33F, I33G, I33H, I33K, I33L, I33M, I33N, I33P, I33Q, I33R, I33S, I33T, I33V, I33W, I33Y, Y34A, Y34C, Y34D, Y34E, Y34F, Y34G, Y34H, Y34I, Y34K, Y34L, Y34M, Y34N, Y34P, Y34Q, Y34R, Y34S, Y34T, Y34V, Y34W, K35A, K35C, K35D, K35E, K35F, K35G, K35H, K35I, K35L, K35M, K35N, K35P, K35Q, K35R, K35S, K35T, K35V, K35W, K35Y, K38A, K38C, K38D, K38E, K38F, K38G, K38H, K38I, K38L, K38M, K38N, K38P, K38Q, K38R, K38S, K38T, K38V, K38W, K38Y, R39A, R39C, R39D, R39E, R39F, R39G, R39H, R39I, R39K, R39L, R39M, R39N, R39P, R39Q, R39S, R39T, R39V, R39W, R39Y, N49A, N49C, N49D, N49E, N49F, N49G, N49H, N49I, N49K, N49L, N49M, N49P, N49Q, N49R, N49S, N49T, N49V, N49W, N49Y, F50A, F50C, F50D, F50E, F50G, F50H, F50I, F50K, F50L, F50M, F50N, F50P, F50Q, F50R, F50S, F50T, F50V, F50W, F50Y, R51A, R51C, R51D, R51E, R51F, R51G, R51H, R51I, R51K, R51L, R51M, R51N, R51P, R51Q, R51S, R51T, R51V, R51W, R51Y, D52A, D52C, D52E, D52F, D52G, D52H, D52I, D52K, D52L,

D52M, D52N, D52P, D52Q, D52R, D52S, D52T, D52V, D52W, D52Y, I53A, I53C, I53D, I53E, I53F, I53G, I53H, I53K, I53L, I53M, I53N, I53P, I53Q, I53R, I53S, I53T, I53V, I53W, I53Y, Y60A, Y60C, Y60D, Y60E, Y60F, Y60G, Y60H, Y60I, Y60K, Y60L, Y60M, Y60N, Y60P, Y60Q, Y60R, Y60S, Y60T, Y60V, Y60W, S61A, S61C, S61D, S61E, S61F, S61G, S61H, S61I, S61K, S61L, S61M, S61N, S61P, S61Q, S61R, S61T, S61V, S61W, S61Y, I62A, I62C, I62D, I62E, I62F, I62G, I62H, I62K, I62L, I62M, I62N, I62P, I62Q, I62R, I62S, I62T, I62V, I62W, I62Y, S63A, S63C, S63D, S63E, S63F, S63G, S63H, S63I, S63K, S63L, S63M, S63N, S63P, S63Q, S63R, S63T, S63V, S63W, S63Y, A64C, A64D, A64E, A64F, A64G, A64H, A64I, A64K, A64L, A64M, A64N, A64P, A64Q, A64R, A64S, A64T, A64V, A64W, A64Y, I65A, I65C, I65D, I65E, I65F, I65G, I65H, I65K, I65L, I65M, I65N, I65P, I65Q, I65R, I65S, I65T, I65V, I65W, I65Y, D66A, D66C, D66E, D66F, D66G, D66H, D66I, D66K, D66L, D66M, D66N, D66P, D66Q, D66R, D66S, D66T, D66V, D66W, D66Y, T75A, T75C, T75D, T75E, T75F, T75G, T75H, T75I, T75K, T75L, T75M, T75N, T75P, T75Q, T75R, T75S, T75V, T75W, T75Y, E76A, E76C, E76D, E76F, E76G, E76H, E76I, E76K, E76L, E76M, E76N, E76P, E76Q, E76R, E76S, E76T, E76V, E76W, E76Y, V77A, V77C, V77D, V77E, V77F, V77G, V77H, V77I, V77K, V77L, V77M, V77N, V77P, V77Q, V77R, V77S, V77T, V77W, V77Y, F78A, F78C, F78D, F78E, F78G, F78H, F78I, F78K, F78L, F78M, F78N, F78P, F78Q, F78R, F78S, F78T, F78V, F78W, F78Y, P79A, P79C, P79D, P79E, P79F, P79G, P79H, P79I, P79K, P79L, P79M, P79N, P79Q, P79R, P79S, P79T, P79V, P79W, P79Y, V80A, V80C, V80D, V80E, V80F, V80G, V80H, V80I, V80K, V80L, V80M, V80N, V80P, V80Q, V80R, V80S, V80T, V80W, V80Y, K81A, K81C, K81D, K81E, K81F, K81G, K81H, K81I, K81L, K81M, K81N, K81P, K81Q, K81R, K81S, K81T, K81V, K81W, K81Y, L82A, L82C, L82D, L82E, L82F, L82G, L82H, L82I, L82K, L82M, L82N, L82P, L82Q, L82R, L82S, L82T, L82V, L82W, L82Y, E83A, E83C, E83D, E83F, E83G, E83H, E83I, E83K, E83L, E83M, E83N, E83P, E83Q, E83R, E83S, E83T, E83V, E83W, E83Y, V84A, V84C, V84D, V84E, V84F, V84G, V84H, V84I, V84K, V84L, V84M, V84N, V84P, V84Q, V84R, V84S, V84T, V84V, V84Y, D85A, D85C, D85E, D85F, D85G, D85H, D85I, D85K, D85L, D85M, D85N, D85P, D85Q, D85R, D85S, D85T, D85V, D85W, D85Y, E86A, E86C, E86D, E86F, E86G, E86H, E86I, E86K, E86L, E86M, E86N, E86P, E86Q, E86R, E86S, E86T, E86V, E86W, E86Y, E87A, E87C, E87D, E87F, E87G, E87H, E87I, E87K, E87L, E87M, E87N, E87P, E87Q, E87R, E87S, E87T, E87V, E87W, E87Y, N88A, N88C, N88D, N88E, N88F, N88G, N88H, N88I, N88K, N88L, N88M, N88P, N88Q, N88R, N88S, N88T, N88V, N88W, N88Y, L89A, L89C, L89D, L89E, L89F, L89G, L89H, L89I, L89K, L89M, L89N, L89P, L89Q, L89R, L89S, L89T, L89V, L89W, L89Y, T90A, T90C, T90D, T90E, T90F, T90G, T90H, T90I, T90K, T90L, T90M, T90N, T90P, T90Q, T90R, T90S, T90V, T90W, T90Y, A91C, A91D, A91E, A91F, A91G, A91H, A91I, A91K, A91L, A91M, A91N, A91P, A91Q, A91R, A91S, A91T, A91V, A91W, A91Y, G92A, G92C, G92D, G92E, G92F, G92H, G92I, G92K, G92L, G92M, G92N, G92P, G92Q, G92R, G92S, G92T, G92V, G92W, G92Y, Y93A, Y93C, Y93D, Y93E, Y93F, Y93G, Y93H, Y93I, Y93K, Y93L, Y93M, Y93N, Y93P, Y93Q, Y93R, Y93S, Y93T, Y93V, Y93W, P94A, P94C, P94D, P94E, P94F, P94G,

P94H, P94I, P94K, P94L, P94M, P94N, P94Q, P94R, P94S, P94T, P94V, P94W, P94Y, G95A, G95C, G95D, G95E, G95F, G95H, G95I, G95K, G95L, G95M, G95N, G95P, G95Q, G95R, G95S, G95T, G95V, G95W, G95Y, A96C, A96D, A96E, A96F, A96G, A96H, A96I, A96K, A96L, A96M, A96N, A96P, A96Q, A96R, A96S, A96T, A96V, A96W, A96Y, V97A, V97C, V97D, V97E, V97F, V97G, V97H, V97I, V97K, V97L, V97M, V97N, V97P, V97Q, V97R, V97S, V97T, V97W, V97Y, K98A, K98C, K98D, K98E, K98F, K98G, K98H, K98I, K98L, K98M, K98N, K98P, K98Q, K98R, K98S, K98T, K98V, K98W, K98Y, V99A, V99C, V99D, V99E, V99F, V99G, V99H, V99I, V99K, V99L, V99M, V99N, V99P, V99Q, V99R, V99S, V99T, V99W, V99Y, K100A, K100C, K100D, K100E, K100F, K100G, K100H, K100I, K100L, K100M, K100N, K100P, K100Q, K100R, K100S, K100T, K100V, K100W, V101Y, V101A, V101C, V101D, V101E, V101F, V101G, V101H, V101I, V101K, V101L, V101M, V101N, V101P, V101Q, V101R, V101S, V101T, V101W, V101Y, E102A, E102C, E102D, E102F, E102G, E102H, E102I, E102K, E102L, E102M, E102N, E102P, E102Q, E102R, E102S, E102T, E102V, E102W, E102Y, N103A, N103C, N103D, N103E, N103F, N103G, N103H, N103I, N103K, N103L, N103M, N103P, N103Q, N103R, N103S, N103T, N103V, N103W, N103Y, L104A, L104C, L104D, L104E, L104F, L104G, L104H, L104I, L104K, L104M, L104N, L104P, L104Q, L104R, L104S, L104T, L104V, L104W, L104Y, D105A, D105C, D105E, D105F, D105G, D105H, D105I, D105K, D105L, D105M, D105N, D105P, D105Q, D105R, D105S, D105T, D105V, D105W, D105Y, G106A, G106C, G106D, G106E, G106F, G106H, G106I, G106K, G106L, G106M, G106N, G106P, G106Q, G106R, G106S, G106T, G106V, G106W, G106Y, E107A, E107C, E107D, E107F, E107G, E107H, E107I, E107K, E107L, E107M, E107N, E107P, E107Q, E107R, E107S, E107T, E107V, E107W, E107Y, A108C, A108D, A108E, A108F, A108G, A108H, A108I, A108K, A108L, A108M, A108N, A108P, A108Q, A108R, A108S, A108T, A108V, A108W, and/or A108Y and combinations thereof. It is contemplated that one or more of these substitutions may be specifically excluded in embodiments described herein.

[0126] In certain aspects, the library comprises a variation in an amino acid corresponding to amino acid 1 of SEQ ID NO:1 in combination with one or more residue corresponding to amino acid 2, 3, 4, 14, 28, and 58.

[0127] In a further aspect, the library comprises a variation in an amino acid corresponding to amino acid 2 of SEQ ID NO:1 in combination with one or more residue corresponding to amino acid 1, 3, 4, 14, 28, and/or 58.

[0128] In a still a further aspect, the library comprises a variation in an amino acid corresponding to amino acid 3 of SEQ ID NO:1 in combination with one or more residue corresponding to amino acid 1, 2, 4, 14, 28, and/or 58.

[0129] In a further aspect, the library comprises a variation in an amino acid corresponding to amino acid 4 of SEQ ID NO:1 in combination with one or more residue corresponding to amino acid 1, 2, 3, 14, 28, and/or 58.

[0130] In a certain aspect, the library comprises a variation in an amino acid corresponding to amino acid 14 of SEQ ID NO:1 in combination with one or more residue corresponding to amino acid 1, 2, 3, 4, 28, and/or 58.

[0131] In a further aspect, the library comprises a variation in an amino acid corresponding to amino acid 28 of SEQ ID

NO:1 in combination with one or more residue corresponding to amino acid 1, 2, 3, 4, 14, and/or 58.

[0132] In a further aspect, the library comprises a variation in an amino acid corresponding to amino acid 58 of SEQ ID NO:1 in combination with one or more residue corresponding to amino acid 1, 2, 3, 4, 14, and/or 28.

[0133] D. Library Screening

[0134] Library screening can be conducted in order to select FN3 variants that bind to specific ligands or targets. Combinatorial screening can easily produce and screen a large number of variants, which is not feasible with specific mutagenesis ("rational design") approaches. Amino acid variant at various amino acid positions in FN3 can be generated using a degenerate nucleotide sequence. FN3 variants with desired binding capabilities can be selected in vitro, recovered and amplified. The amino acid sequence of a selected clone can be identified readily by sequencing the nucleic acid encoding the selected FN3.

[0135] In some embodiments, a particular FN3-based molecule has an affinity for a target that is at least 2-fold greater than the affinity of the polypeptide prior to substitutions discussed herein. In some embodiments, the affinity is, at least, or is at most about 2-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 15-, 20-, 25-, 30-, 35-, 40-, 45-, 50-, 60-, 70-, 80-, 90-, 100-fold increased (or any range derivable therein) compared to another FN3-based molecule.

[0136] Phage Display Library and Selection. A FN3 polypeptide library can be created using a "shaved" template containing polyserine sequence at locations to be diversified (Koide et al., 2007 and Wojcik et al., 2010). A synthetic DNA fragment that encodes signal sequence of DsbA (Steiner et al., 2006) can be fused to the gene for the template, and the fusion gene can be cloned into a phage display vector (Koide et al., 1998). A phage-display combinatorial library can be constructed by introducing codons for amino acid variation into the FN3 polypeptide. Library construction procedures have previously been described (Koide and Koide, 2007).

[0137] Phagemid particles can be prepared by growing XL1-Blue cells transfected with the phagemid library in the presence of IPTG and helper phage (Lo Conte et al., 1999; Fellouse et al., 2005). Phagemid library selection can be performed as follows. In the first round, 0.5 μ M of a target protein modified with EZ-Link Sulfo-NHS—SS-Biotin (Sulfosuccinimidyl 2(biotinamido)-ethyl-1,3-dithiopropionate; Pierce) can be mixed with a sufficient amount of streptavidin-conjugated magnetic beads (Streptavidin MagneSphere Pramagnetic Particles; Promega, Z5481/2) in TBS (50 mM Tris HCl buffer pH 7.5 150 mM NaCl) containing 0.5% Tween20 (TBST). To this target solution, 1012-13 phagemids suspended in 1 ml TBST plus 0.5% BSA can be added, and the solution can then be mixed and incubated for 15 min at room temperature. After washing the beads twice with TBST, the beads suspension containing bound phagemids can be added to fresh *E. coli* culture. Phagemids were amplified as described before (Fellouse et al., 2005). In a second round, phagemids can be incubated with 0.1 μ M target in TBST plus 0.5% BSA, and then captured by streptavidin-conjugated magnetic beads. Phagemids bound to the target protein can be eluted from the beads by cleaving the linker within the biotinylation reagent with 100 mM DTT in TBST. The phagemids can then be washed and recovered as described above. After amplification, the third round of selection may be performed using 0.02 μ M target.

Phage display is an established technique for generating binding members and has been described in detail in many publications such as Kontermann & Dubel (2001) and WO92/01047, each of which is incorporated herein by reference in its entirety.

[0138] Yeast Surface Display. Yeast surface experiments can be performed according to Boder and Wittrup (2000) with minor modifications. The Express-tag in the yeast display vector, pYD1, (Invitrogen) may be removed, since it can cross-react with anti-FLAG antibodies (Sigma). The genes for monobodies in the phagemid library after three rounds of selection can be amplified using PCR and mixed with the modified pYD1 cut with EcoRI and XhoI, and yeast EBY100 cells can be transformed with this mixture. The transformed yeast cells can be grown in the SD-CAA media at 30 $^{\circ}$ C for two days, and then monobody expression can be induced by growing the cells in the SG-CAA media at 30 $^{\circ}$ C for 24 h.

[0139] Sorting of monobody-displaying yeast cells may be performed as follows. The yeast cells may be incubated with a biotinylated target (50 nM) and mouse anti-V5 antibody (Sigma), then after washing incubated with anti-mouse antibody-FITC conjugate (Sigma) and neutravidin-PE conjugate (Invitrogen). The stained cells can be sorted based on the FITC and PE intensities. Typically, cells exhibiting the top 1% PE intensity and top 10% FITC intensity are recovered.

[0140] After FACS sorting, individual clones can be analyzed. Approximate K_d values can be determined from a titration curve by FACS analysis (Boder and Wittrup, 2000). Amino acid sequences can be deduced from DNA sequencing.

[0141] Effects of *E. coli* lysate on monobody-target interaction can be tested by comparing binding in the presence and absence of *E. coli* lysate prepared from cell suspension with OD₆₀₀ of 50.

[0142] Protein Expression and Purification. The nucleic acid encoding any targets can be cloned in the appropriate expression vector. In one example, genes for monobodies can be cloned in the expression vector, pHFT2, which is a derivative of pHFT1 (Huang et al., 2006) in which the His-6 tag had been replaced with a His-10 tag. Protein expression and purification can be performed as described previously (Huang et al., 2006).

[0143] An expression vector comprising cDNA encoding a FN3 polypeptide or a target molecule is introduced into *Escherichia coli*, yeast, an insect cell, an animal cell or the like for expression to obtain the polypeptide. Polypeptides used in the methods and compositions of the disclosure can be produced, for example, by expressing a DNA encoding it in a host cell using a method described in Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press (1989), Current Protocols in Molecular Biology, John Wiley & Sons (1987-1997) or the like. A recombinant vector is produced by inserting a cDNA downstream of a promoter in an appropriate expression vector. The vector is then introduced into a host cell suitable for the expression vector. The host cell can be any cell so long as it can express the gene of interest, and includes bacteria (e.g., *Escherichia coli*), an animal cell and the like. Expression vector can replicate autonomously in the host cell to be used or vectors which can be integrated into a

chromosome comprising an appropriate promoter at such a position that the DNA encoding the polypeptide can be transcribed.

V. Affinity Claims

[0144] Embodiments of the disclosure, particularly those comprising a polypeptide comprising a biorecognition module including a molecular recognition domain relate to a polypeptide capable of forming an “affinity clamp” to a target motif.

[0145] It is informative to compare characteristics of molecular affinity clamps with those of antibodies, the gold standard of affinity reagents. Antibodies are general and versatile affinity reagents. The immune system can produce an antibody to virtually any molecule. The diversity of the immunoglobulin repertoire is 10¹⁰-12, which is similar in size to the diversity of a typical phage display library (10¹⁰). This versatility of the antibodies, however, also means that the antibody repertoire is not focused and that only a small subset of the naive repertoire is available to bind to a particular class of antigen. For example, antibodies that bind to lysozyme and those that bind to a phospho-Ser peptide are distinct subsets of the same repertoire.


[0146] Economical and scalable production is another important area of consideration for affinity reagents. As noted above, polyclonal antibodies cannot be reproduced, once the original stock is depleted. Monoclonal antibodies can be reproduced, but the maintenance and large-scale culture of hybridoma cells are cumbersome and expensive. Antibodies can also be produced by recombinant technologies, but the natural diversity throughout the antibody molecules (i.e., framework diversity in addition to the extensive diversity within the antigen binding loops) makes formatting them for different applications fundamentally low throughput.

[0147] Moreover, because of the presence of critical disulfide bonds, recombinant production of antibodies is not straightforward. For this reason, a number of alternative “molecular scaffolds” for engineering affinity reagents have been developed that are small and devoid of disulfide bonds (8-10). Although these new-generation affinity reagents generally have good affinity and specificity, developing affinity reagents for short peptide motifs remains a major challenge in the field, because of the fundamental difficulties stated hereinabove.

[0148] In contrast, molecular affinity clamps in accordance with the disclosure are affinity reagents directed to a pre-defined motif. In one aspect, molecular affinity clamps are built with a particular biorecognition module comprising an interaction domain that is specific primarily to the class of target motifs that the interaction domain recognizes. Because of this pre-defined binding specificity, repertoire diversity can then be used to enhance the properties of affinity reagents rather than to blindly search for initial hits. This distinctive feature of the invention may lead to an increased success rate of producing high-affinity reagents for a motif of interest. In another embodiment, the molecular affinity clamps are built with one or more variant FN3 domains.

[0149] The two polypeptide molecules of the affinity clamp (e.g. the first polypeptide comprising the variant FN3 domain and the second polypeptide comprising the biorecognition module, which can include an interaction domain or a second variant FN3 domain) are spatially oriented to

bind distinct, overlapping, or the same sites within target motif of a target. The configuration of the two biorecognition modules about the target motif is clamp-like or clamshell-like, i.e., the target motif is “clamped” between the two biorecognition modules. The two biorecognition modules of the affinity clamp are capable of together binding a single target motif on a single target. The first polypeptide molecule and second polypeptide molecules may bind at least overlapping portions the target motif on a target. This is unlike other binding molecules, such as an antibody, where the binding molecules recognize non-overlapping target motifs, different target motifs, or the same target motif located on more than one target. In another aspect, the affinity clamp is suitably described as a ternary complex composition of the type:



wherein M1 and M2 are independently the polypeptide comprising the variant FN3 domain and the polypeptide comprising the biorecognition module, L is a direct bond or linker moiety used for tethering the first and second biorecognition modules, and T is a target motif. M1 includes a variant FN3 domain bound to a first site of the target motif, and M2 includes a molecular recognition domain bound to a second site of the target motif (or vice versa) without disrupting the binding of the variant FN3 domain. The first and second sites can be the same, overlapping, or distinct sites within the target motif. L as a linker is selected from the group consisting of a peptide which is equal to or shorter than 30 residues, a group capable of disulfide bonding, and a chemical crosslinker.

[0150] B. Target Motif

[0151] A target motif suitable in accordance with the disclosure may be any motif which can be recognized by a biorecognition module, e.g., an interaction domain. Such target motifs include peptides and covalently modified peptides, including but not limited to peptides that are phosphorylated, methylated, acetylated, ubiquitinated, SUMOylated, ISGylated, glycosylated, acylated, prenylated, ribosylated, gammacarboxylated, or sulfated.

[0152] C. Biorecognition Module

[0153] Among the commonly occurring domains identified in signaling proteins are the so-called “interaction domains.” Interaction domains are typically small (usually less than ~100 amino acids) and autonomously folded. Many of them bind to short peptide motifs that often contain modified amino acids. It has been found that a primary binding domain, i.e., the molecular recognition domain, of the biorecognition module is suitably an interaction domain. With molecular affinity clamp technology, the interaction domains as the biorecognition modules can be engineered in such a way that the enhancer domain can be connected in a proper orientation. The bifunctional module architecture of the molecular affinity clamps in accordance with the invention, after optimization, significantly increases the surface areas of the peptide-binding interface by forming the clamshell architecture, leading to higher affinity and/or specificity. Use of interaction domains as the primary binding domain is based on the following common features of these domains: a target peptide motif binds to a shallow groove on the interaction domain surface, and the peptide is still highly

exposed; there are turns and/or loops located close to the peptide-binding site; and the N- and C-termini are juxtaposed in space so that they could be connected and a new set of termini could be created elsewhere.

[0154] In short, molecular affinity clamp technology makes it possible to define the primary specificity of affinity reagents in advance (e.g., using the specificity of the interaction domain), and then, enhance that affinity and/or specificity. This modular architecture in accordance with the disclosure transforms affinity reagent development from an “unguided fishing expedition” to a focused, rational and robust process.

[0155] Interaction domains, suitable as the recognition domain, include, but are not limited to, domains involved in phospho-tyrosine binding (e.g. SH2, PTB), phospho-serine binding (e.g. UIM, GAT, CUE, BTB/POZ, VHS, UBA, RING, HECT, WW, 14-3-3, Polo-box), phospho-threonine binding (e.g. FHA, WW, Polo-box), proline-rich region binding (e.g. EVH1, SH3, GYF), acetylated lysine binding (e.g. Bromo), methylated lysine binding (e.g. Chromo, PHD), apoptosis (e.g. BIR, TRAF, DED, Death, CARD, BH), cytoskeleton modulation (e.g. ADF, GEL, DH, CH, FH2), or other cellular functions (e.g. EH, CC, VHL, TUDOR, PUF Repeat, PAS, MH1, LRR, IQ, HEAT, GRIP, TUBBY, SNARE, TPR, TIR, START, SOCS Box, SAM, RGS, PDZ, PB1, LIM, F-BOX, ENTH, EF-Hand, SHADOW, ARM, ANK).

[0156] D. Linkers

[0157] The variant FN3 polypeptide and polypeptide comprising the biorecognition module may be linked together either directly, e.g., bound together with a peptide sequence via a tail from one of the modules, or indirectly via a linker. As to the latter, the linker generally is bifunctional in that it includes a functionality for linking the biorecognition module and a functionality for linking variant FN3 polypeptide. The linker may suitably be a specific moiety, such as an amino acid sequence of about 30 or fewer residues. It is also contemplated that the two polypeptide domains may be linked non-covalently through a high affinity binding interaction or physical association such as the interaction mediated by coiled-coil peptides.

[0158] E. Detection of Affinity Clamp Binding

[0159] In general, the different conformational states of modular affinity clamps used in accordance with the disclosure will correspond to different separation distances between the polypeptide modules, whereby changes in conformation may be conveniently monitored by means of a separation sensitive signal.

[0160] Various forms of separation sensitive signal systems may be used with the affinity clamps of the disclosure. In such embodiments, the biorecognition module includes a first signaling moiety and the variant FN3 polypeptide includes a second signaling moiety, and the first and second signaling moieties are capable of interacting to produce a detectable signal. The signaling moieties may include dyes, quenchers, reporter proteins and quantum dots. Particularly useful are embodiments in which the polypeptide domains include optical signaling pairs that can produce a detectable signal when the proximity of the modules with respect to each other changes with the binding of the polypeptide domains. Suitably, the first and second signaling modules are a fluorescence resonance energy (FRET) donor group and a receptor group, respectively. The change in proximity

of the FRET groups produces an optical signal which differs between when the target motif is present and not present.

[0161] It will also be appreciated that various other means may be used for “reading” the presence of target motif binding to a modular affinity clamp, and/or the resultant change in conformational state of the affinity clamp structure. Many different labeling systems may be used, such as fluorophore labeling (including quantum dot), radio-labeling, and redox labeling.

[0162] F. Use of Affinity Clamps as Bio Sensors

[0163] Molecular affinity clamps in accordance with the disclosure may be suitably used as a biosensor wherein the polypeptide modules are each labeled with paired signaling moieties as described above.

[0164] A plurality of affinity clamps described herein may be immobilized, directly or indirectly to a support or substrate to form an array of clamps or an array of biosensors. Supports or substrates can take a variety of forms such as polymers, glasses, metal and those with coating therein. Arrays are ordered arrangements of elements, allowing them to be displayed and examined in parallel. Arrays of immobilized affinity clamps can be used to detect the target motif and demonstrate the binding reaction. Certain array formats are sometimes referred to as “biochips.” Biochips may include a plurality of locations configured so that each location is spatially addressable. Typically, the clamp format is configured in a row and column format with regular spacing between locations, wherein each location has machine-readable (e.g., computer-readable) information to identify the location on the surface of the substrate.

[0165] The affinity clamp technology provides a method of detecting the presence and amount of a target motif in a sample by using the affinity clamp as a biosensor. Specifically, a sample is contacted under specific conditions with a biosensor. Fluorescence events are sensed with the binding of the polypeptide modules to the target motif in the sample and in the absence of the sample, and the fluorescence sensing in the absence of the target motif is correlated with a change in the FRET signal in the presence of the target motif. Thus, absence of the target motif generates a specific FRET signal in terms of the wavelength and amplitude of the emission, and the presence of the target motif generates a modulated FRET signal emission in terms of either the wavelength or amplitude or both. Samples may include blood, saliva or tissue.

[0166] Accordingly, an affinity clamp array as a biosensor array includes a plurality of affinity clamps or biosensors anchored to the surface of a substrate, each at an addressable site on the substrate.

[0167] G. Construction of a Modular Molecular Affinity Clamp

[0168] The general engineering of a molecular affinity clamp is given basically in four steps. Step 1 involves identifying the potential locations for attachment, via a linker, of the variant FN3 polypeptide to the biorecognition module by visual inspection of the interaction domain structure and/or from sequence variability among interaction domain family members, and testing the tolerance of identified locations for extensive modifications, for example, by inserting four Gly residues.

[0169] Step 2 includes two sub-steps, Step 2a and 2b. Step 2a is included if circular permutations are performed to construct new termini closer to the interaction domain binding site. In some embodiments, Step 2a is not needed.

In Step 2a, if circular permutation is performed, a domain is constructed by joining the original termini and cutting the polypeptide at a location closer to the target-binding site of the interaction domain that tolerates mutations. Then, in Step 2b, the variant FN3 polypeptide is attached to the C-terminus of the circularly permuted domain or the natural C-terminus (in the case where no circular permutation is performed.) The N-terminus of FN3 is located close to its functional loops, and thus, connecting the FN3 N-terminus to the interaction domain ensures that the FN3 binding loops are facing the target motif-binding site.

[0170] In Step 3, amino acid diversity is introduced in FN3 loops to construct a large combinatorial library of mutated polypeptides, and in Step 4, library sorting is performed to optimize the enhancer domain for a specific target motif.

[0171] Further embodiments of the affinity clamp are described in WO/2009/062170, which is herein incorporated by reference.

VI. Polypeptide Compositions

[0172] The polypeptides or polynucleotides of the disclosure may include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 or more variant amino acids or nucleic acid substitutions or be at least 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% similar, identical, or homologous with at least, or at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids or nucleic acids, or any range derivable therein, of SEQ ID NOS:1-7.

[0173] The polypeptides or polynucleotides of the disclosure may include 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156,

157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:1-7.

[0174] In some aspects there is a nucleic acid molecule or polypeptide starting at position 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, or 615 of any of SEQ ID NOS:1-7 and comprising 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47,

359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, or 615 of one of SEQ ID NO:1-7.

[0177] Embodiments include polypeptides and polynucleotides with at least, at most, or exactly 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% (or any derivable range therein) identity, similarity, or homology to one of SEQ ID NO:1-7.

[0178] Substitutional variants typically contain the exchange of one amino acid for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide, with or without the loss of other functions or properties. Substitutions may be conservative, that is, one amino acid is replaced with one of similar shape and charge. Conservative substitutions are well known in the art and include, for example, the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; and valine to isoleucine or leucine. Alternatively, substitutions may be non-conservative such that a function or activity of the polypeptide is affected. Non-conservative changes typically involve substituting a residue with one that is chemically dissimilar, such as a polar or charged amino acid for a nonpolar or uncharged amino acid, and vice versa.

[0179] The current disclosure concerns methods and compositions related to the identification and use of variants of FN3 and libraries containing the same. As used herein, a "polypeptide" generally is defined herein to refer to a peptide sequence of about 10 to about 1,000 or more amino acid residues.

[0180] The polypeptides included in the methods set forth herein are variants in that they comprise a FN3 amino acid sequence that has been altered by substitution, insertion and/or deletion of one or more amino acid. The polypeptides set forth herein may demonstrate a selective and/or specific binding affinity for particular target molecules or portions thereof.

[0181] In certain embodiments, the polypeptide is a fusion polypeptide that includes a variant FN3 amino acid sequence linked at the N- or C-terminus to a second peptide or polypeptide. In other embodiments, the polypeptide comprises a linker interposed between the FN3 amino acid sequence and the second peptide or polypeptide sequence. Linkers are discussed in greater detail in the specification below.

[0182] Furthermore, the polypeptides set forth herein may comprise a sequence of any number of additional amino acid residues at either the N-terminus or C-terminus of the amino acid sequence that includes the variant FN3 amino acid sequence. For example, there may be an amino acid sequence of about 3 to about 1,000 or more amino acid residues at either the N-terminus, the C-terminus, or both the N-terminus and C-terminus of the amino acid sequence that includes the variant FN3 amino acid sequence.

[0183] The polypeptide may include the addition of an antibody epitope or other tag, to facilitate identification, targeting, and/or purification of the polypeptide. The use of 6xHis and GST (glutathione S transferase) as tags is well known. Inclusion of a cleavage site at or near the fusion junction will facilitate removal of the extraneous polypeptide after purification. Other amino acid sequences that may be included in the polypeptide include functional domains, such as active sites from enzymes such as a hydrolase, glycosylation domains, cellular targeting signals or transmembrane regions. The polypeptide may further include one or more additional tissue-targeting moieties.

[0184] Polypeptides may possess deletions and/or substitutions of amino acids relative to the native sequence. Sequences with amino acid substitutions are contemplated, as are sequences with a deletion, and sequences with a deletion and a substitution. In some embodiments, these polypeptides may further include insertions or added amino acids.

[0185] Substitutional or replacement variants typically contain the exchange of one amino acid for another at one or more sites within the protein and may be designed to modulate one or more properties of the polypeptide, particularly to increase its efficacy or specificity. Substitutions of this kind may or may not be conservative substitutions. Conservative substitution is when one amino acid is replaced with one of similar shape and charge. Being that the libraries of variant FN3 domains serves to provide a diversity of amino acid sequences and binding selectivity conservative substitutions are not required. However, if used, conservative substitutions are well known in the art and include, for example, the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; and valine to isoleucine or leucine. Changes other than those discussed above are generally considered not to be conservative substitutions. It is specifically contemplated that one or more of the conservative substitutions above may be included as embodiments. In other embodiments, such substitutions are specifically excluded. Further-

more, in additional embodiments, substitutions that are not conservative are employed in variants.

[0186] In addition to a deletion or substitution, the polypeptides may possess an insertion of one or more residues.

[0187] The variant FN3 amino acid sequence may be structurally equivalent to the native counterparts. For example, the variant FN3 amino acid sequence forms the appropriate structure and conformation for binding targets, proteins, or peptide segments.

[0188] The following is a discussion based upon changing of the amino acids of a polypeptide to create a library of molecules or a second-generation molecule. For example, certain amino acids may be substituted for other amino acids in a polypeptide without appreciable loss of function, such as ability to interact with a target peptide sequence. Since it is the interactive capacity and nature of a polypeptide that defines that polypeptide's functional activity, certain amino acid substitutions can be made in a polypeptide sequence and nevertheless produce a polypeptide with like properties.

[0189] In making such changes, the hydrophobic index of amino acids may be considered. The importance of the hydrophobic amino acid index in conferring interactive function on a protein is generally understood in the art (Kyte and Doolittle, 1982). It is accepted that the relative hydrophobic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like.

[0190] It also is understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Pat. No. 4,554,101, incorporated herein by reference, states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein. As detailed in U.S. Pat. No. 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0±1); glutamate (+3.0±1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5±1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4).

[0191] It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still produce a biologically equivalent and immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ±2 is preferred, those that are within ±1 are particularly preferred, and those within ±0.5 are even more particularly preferred.

[0192] As outlined above, amino acid substitutions generally are based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. However, in some aspects a non-conservative substitution is contemplated. In certain aspects a random substitution is also contemplated. Exemplary substitutions that take into consideration the various foregoing characteristics are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

VII. Polynucleotides

[0193] Aspects of the disclosure relate to polypeptides and polynucleotides encoding such polypeptides. The terms "polynucleotide" and "oligonucleotide" are used interchangeably and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides or analogs thereof. Polynucleotides can have any three-dimensional structure and may perform any function, known or unknown. The following are non-limiting examples of polynucleotides: a gene or gene fragment (for example, a probe, primer, EST or SAGE tag), exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, dsRNA, siRNA, miRNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes and primers. A polynucleotide can comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. If present, modifications to the nucleotide structure can be imparted before or after assembly of the polynucleotide. The sequence of nucleotides can be interrupted by non-nucleotide components. A polynucleotide can be further modified after polymerization, such as by conjugation with a labeling component. The term also refers to both double- and single-stranded molecules. Unless otherwise specified or required, any embodiment of the disclosure that is a polynucleotide encompasses both the double-stranded form and each of two complementary single-stranded forms known or predicted to make up the double-stranded form.

[0194] The term "complementary" as used herein refers to Watson-Crick base pairing between nucleotides and specifically refers to nucleotides hydrogen bonded to one another with thymine or uracil residues linked to adenine residues by two hydrogen bonds and cytosine and guanine residues linked by three hydrogen bonds. In general, a nucleic acid includes a nucleotide sequence described as having a "percent complementarity" to a specified second nucleotide sequence. For example, a nucleotide sequence may have 80%, 90%, or 100% complementarity to a specified second nucleotide sequence, indicating that 8 of 10, 9 of 10 or 10 of 10 nucleotides of a sequence are complementary to the specified second nucleotide sequence. For instance, the nucleotide sequence 3'-TCGA-5' is 100% complementary to the nucleotide sequence 5'-AGCT-3'. Further, the nucleotide sequence 3'-TCGA- is 100% complementary to a region of the nucleotide sequence 5'-TTAGCTGG-3'. It will be recognized by one of skill in the art that two complementary nucleotide sequences include a sense strand and an antisense strand.

[0195] Polypeptides may be encoded by a nucleic acid molecule in the composition. In certain embodiments, the nucleic acid molecule can be in the form of a nucleic acid vector. The term "vector" is used to refer to a carrier nucleic acid molecule into which a heterologous nucleic acid sequence can be inserted for introduction into a cell where it can be replicated and expressed. A nucleic acid sequence can be "heterologous," which means that it is in a context foreign to the cell in which the vector is being introduced or to the nucleic acid in which is incorporated, which includes a sequence homologous to a sequence in the cell or nucleic acid but in a position within the host cell or nucleic acid where it is ordinarily not found. Vectors include DNAs, RNAs, plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (e.g.,

YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (for example Sambrook et al., 2001; Ausubel et al., 1996, both incorporated herein by reference). Vectors may be used in a host cell to produce an antibody.

[0196] The term “expression vector” refers to a vector containing a nucleic acid sequence coding for at least part of a gene product capable of being transcribed or stably integrate into a host cell’s genome and subsequently be transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. Expression vectors can contain a variety of “control sequences,” which refer to nucleic acid sequences necessary for the transcription and possibly translation of an operably linked coding sequence in a particular host organism. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described herein. It is contemplated that expression vectors that express a marker may be useful in the methods and compositions of the disclosure. In other embodiments, the marker is encoded on an mRNA and not in an expression vector.

[0197] A “promoter” is a control sequence. The promoter is typically a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind such as RNA polymerase and other transcription factors. The phrases “operatively positioned,” “operatively linked,” “under control,” and “under transcriptional control” mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and expression of that sequence. A promoter may or may not be used in conjunction with an “enhancer,” which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

[0198] The particular promoter that is employed to control the expression of a peptide or protein encoding polynucleotide is not believed to be critical, so long as it is capable of expressing the polynucleotide in a targeted cell, preferably a bacterial cell. Where a human cell is targeted, it is preferable to position the polynucleotide coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a bacterial, human or viral promoter. In some embodiments, the host cell is an eukaryotic cell. In some embodiments, using eukaryotic cells is beneficial, as it provides for secondary modifications that may not be present in certain prokaryotic systems.

[0199] A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals.

[0200] Vectors can include a multiple cloning site (MCS), which is a nucleic acid region that contains multiple restriction enzyme sites, any of which can be used in conjunction with standard recombinant technology to digest the vector. (See Carbonelli et al., 1999, Levenson et al., 1998, and Cocea, 1997, incorporated herein by reference.)

[0201] Most transcribed eukaryotic RNA molecules will undergo RNA splicing to remove introns from the primary

transcripts. Vectors containing genomic eukaryotic sequences may require donor and/or acceptor splicing sites to ensure proper processing of the transcript for protein expression. (See Chandler et al., 1997, incorporated herein by reference.)

[0202] The vectors or constructs will generally comprise at least one termination signal. A “termination signal” or “terminator” is comprised of the DNA sequences involved in specific termination of an RNA transcript by an RNA polymerase. Thus, in certain embodiments a termination signal that ends the production of an RNA transcript is contemplated. A terminator may be necessary in vivo to achieve desirable message levels. In eukaryotic systems, the terminator region may also comprise specific DNA sequences that permit site-specific cleavage of the new transcript so as to expose a polyadenylation site. This signals a specialized endogenous polymerase to add a stretch of about 200 A residues (polyA) to the 3’ end of the transcript. RNA molecules modified with this polyA tail appear to more stable and are translated more efficiently. Thus, in other embodiments involving eukaryotes, it is preferred that that terminator comprises a signal for the cleavage of the RNA, and it is more preferred that the terminator signal promotes polyadenylation of the message.

[0203] In expression, particularly eukaryotic expression, one will typically include a polyadenylation signal to effect proper polyadenylation of the transcript.

[0204] In order to propagate a vector in a host cell, it may contain one or more origins of replication sites (often termed “ori”), which is a specific nucleic acid sequence at which replication is initiated. Alternatively an autonomously replicating sequence (ARS) can be employed if the host cell is yeast.

[0205] Some vectors may employ control sequences that allow it to be replicated and/or expressed in both prokaryotic and eukaryotic cells. One of skill in the art would further understand the conditions under which to incubate all of the above described host cells to maintain them and to permit replication of a vector. Also understood and known are techniques and conditions that would allow large-scale production of vectors, as well as production of the nucleic acids encoded by vectors and their cognate polypeptides, proteins, or peptides.

[0206] The polynucleotides and polypeptides of the disclosure may be transfected or transformed into host cells or expressed in host cells. As used herein, the terms “cell,” “cell line,” and “cell culture” may be used interchangeably. All of these terms also include both freshly isolated cells and ex vivo cultured, activated or expanded cells. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. In the context of expressing a heterologous nucleic acid sequence, “host cell” refers to a prokaryotic or eukaryotic cell, and it includes any transformable organism that is capable of replicating a vector or expressing a heterologous gene encoded by a vector. A host cell can, and has been, used as a recipient for vectors or viruses. A host cell may be “transfected” or “transformed,” which refers to a process by which exogenous nucleic acid, such as a recombinant protein-encoding sequence, is transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny. Common host cells include bacteria

(such as *E. coli*, *B. subtilis*, *S. violaceoruber*), yeast (such as *S. cerevisiae*, *P. pastoris*), fungi (such as *A. oryzae*) or eukaryotic cells.

VIII. Kits

[0207] Kits are also contemplated as being made or used in certain aspects of the present disclosure. For instance, a polypeptide or nucleic acid of the disclosure can be included in a kit or in a library provided in a kit. A kit can be included in a sealed container. Non-limiting examples of containers include a microtiter plate, a bottle, a metal tube, a laminate tube, a plastic tube, a dispenser, a pressurized container, a barrier container, a package, a compartment, or other types of containers such as injection or blow-molded plastic containers into which the dispersions or compositions or desired bottles, dispensers, or packages are retained. Other examples of containers include glass or plastic vials or bottles. The kit and/or container can include indicia on its surface. The indicia, for example, can be a word, a phrase, an abbreviation, a picture, or a symbol.

[0208] The containers can dispense or contain a predetermined amount of a composition of the present disclosure. The composition can be dispensed as a liquid, a fluid, or a semi-solid. A kit can also include instructions for using the kit and/or compositions. Instructions can include an explanation of how to use and maintain the compositions.

EXAMPLES

[0209] The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion. One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the methods described herein are presently representative of particular embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses which are encompassed within the spirit of the invention as defined by the scope of the claims will occur to those skilled in the art.

Example 1: TherMonobodies, Synthetic Binding Proteins Based on a Hyperthermophilic Fibronectin Type III Domain

[0210] A. Identification of FN3 Domain from Hyperthermophiles

[0211] The inventors utilized the SMART database to explore FN3 domains from hyperthermophiles. The database predicted many FN3 domains in hyperthermophilic archaea and bacteria such as *Thermococcus kodakaraensis*, *Sulfolobus tokodaii*, *Pyrococcus horikoshii* and *Thermotoga lettingae*. The inventors first eliminated predicted domains that were shorter than the length of the shortest FN3 domains that had been structurally characterized (75 amino acids). Then, four FN3 domains in the sequence of Kelch domain-containing protein ST0939 from the hyperthermophilic archaeon *Sulfolobus tokodaii* DSM 16993 were chosen as the candidate proteins, because of their detectably homology to a human FN3. In the predicted constructs, termed STOFN3-1, -2, -3 and -4, consist of 84 (positions 315-398), 86 (399-484), 79 (488-566) and 76 (568-643) amino acids,

respectively, and each construct maintain at least two of the three highly conserved hydrophobic amino acids of FN3 domains (FIG. 1A).

[0212] Expression vectors for these constructs were constructed using synthetic genes. It was found that only the N-terminal two domains (STOFN3-1 and -2) were robustly produced in *E. coli* as soluble proteins that were predominantly monomeric (FIG. 1B). We detected little protein for the third domain, and the fourth domain was expressed mostly as insoluble proteins. These results demonstrate the challenge of predicting constructs that behave well as purified proteins, even for members of the well characterized FN3 domain. Melting temperatures obtained by differential scanning fluorimetry (DSF) were 80° C. for STOFN3-1 and 70° C. for STOFN3-2, respectively, confirming that they were indeed thermostable proteins (FIG. 1C).

[0213] To confirm that STOFN3-1 and STOFN3-2 had the FN3 fold, we determined their crystal structures at 1.28 and 2.45 Å resolutions, respectively. Both segments indeed adopt the FN3 fold consisting of seven anti-parallel β -strands and six loops (FIG. 2A, 2B). Superposition with the FN3fn10 (PDBID: 1FNA), the well characterized FN3 scaffold, demonstrated that the structures of the segments are highly similar to FN3fn10 with an average RMSD value of 1.9 Å for STOFN3-1 and 1.8 Å for STOFN3-2, respectively, for aligned backbone Ca atoms (excluding loops) (FIG. 2C). STOFN3-1 was then chosen for further experiments, because of its superior biophysical properties to STOFN3-2 in terms of expression level, thermal stability, solubility and the ease to obtain high-resolution structure.

[0214] In the crystal structure, three continuous Pro residues (P315, P316 and P317) in the N-terminal region of STOFN3-1 were highly ordered and positioned between the BC and FG loops, the loops commonly utilized for presenting diversified residues (FIG. 3A). Because this N-terminal segment may interfere with creating a target binding site by presenting BC and/or FG loops, the Pro residues were substituted with Ser residues to reduce hydrophobicity and increase flexibility. The high-resolution crystal structure showed that the segment N-terminal to P317S of the mutant had low electron density indicative of conformational disorder (FIG. 3B), strongly suggesting that this segment was dislodged from the folded portion within the construct. The apparent dissociation of this fragment only slightly decreased the stability FIG. 3D).

[0215] The crystal structure of STOFN3-1 also revealed that, surprisingly, the last residues of the FN3 domain according to the SMART database prediction (T398) was located in the middle of the last β -strand (G strand). The β -strand continued by incorporating a Ser residue we added as an artificial extension to the predicted C-terminus (FIG. 3C), suggesting that the predicted boundaries did not accurately match the structural boundary of the domain. We thus extended the C-terminus by adding four residues (P399, S400, S401 and an extra Serine residue) of ST0939, of which P399-S401 were predicted to be N-terminal residues of STOFN3-2 by the SMART database. The crystal structure of this extended construct confirmed that the added segment was properly incorporated into the β -sheet (FIG. 3C), and the construct was more thermostable than the original one (FIG. 3D).

[0216] The stability of STOFN3-1 was further improved by structure-guided design. Based on a structure-guided alignment with FN3 domains, it was found that STOFN3-1

lacked highly conserved Proline residues in FN3 domains. Replacing these residues with Pro (N328P in the AB loop and D372P in the EF loop) improved the thermostability (FIG. 3D). The variant containing all the improvements, namely, the N-terminal substitution, C-terminal extension and Pro mutations now exhibited very high thermal stability with the melting temperature exceeding 95° C. (FIG. 3D). This construct was used as the template for constructing combinatorial libraries (termed STOTEMP2).

[0217] B. Identification of Positions Permissive to Amino Acid Diversification

[0218] In parallel to the scaffold improvements described above, the inventors identified positions of STOFN3-1 that are tolerant to amino acid diversification so that library designs avoid mutating positions critical for maintaining thermal stability and high solubility. A series of point mutations and insertions were introduced to the STOFN3-1 construct and their thermal stabilities were measured (FIG. 4A-C).

[0219] On the bottom end of the molecule as depicted in FIG. 2A, the poly-Ser mutations of the AB and CD loops resulted in little destabilization. In contrast, whereas that of EF loop destabilized STOFN3-1 by 16° C. The sensitivity of the EF loop to mutations is consistent with previous reports for other FN3 domains, which can be rationalized by the presence of the structurally important “Tyrosine corner” motif in the EF loop of STOFN3-1. On the top end of the molecule, the mutation of the BC loop slightly destabilized STOFN3-1 and that of the DE loop showed no destabilizing effect. The serine mutation of G360 immediately N-terminal to the DE loop resulted in inclusion body formation in *E. coli* and also dramatically decreased the stability (T_m reduction by more than 15° C.). Four sets of poly-Serine mutations of the FG loop all showed large destabilization effect (13-20° C.), and all but FGser1 were expressed predominantly as inclusion bodies. This result suggested that both Y383 and P391, residues not mutated in the FGser1 mutant, were critical for efficient folding of STOFN3-1.

[0220] In addition to substitution mutations, insertions of 2, 4 and 8 Serine residues into the BC, CD, DE and FG loops, the loops that tolerated substitution mutations were tested (FIG. 4C). Of these loops, the BC loop highly tolerated the insertions, whereas insertions in the CD and FG loops were destabilizing but these mutants still maintained the T_m of around 70° C. A very large destabilizing effect was observed for the DE-loop elongation. Even an insertion of two residues decreased the stability by 25° C. None of these insertion mutants resulted in substantial inclusion body formation. Taken together, these systematic mutation experiments identified that DE, EF and FG loops are less permissive to mutations than the other loops, but the most destabilized mutants still had T_m higher than 50° C.

[0221] In the crystal structure, the side chain of Y383, the structurally important Tyr in the F strand, interacts with G342 in the C strand, apparently providing the “Aromatic rescue” of the destabilizing Gly residue in a β -strand (Regan ref). The impact of Y383 mutation completely disappeared when we substituted G342 with Tyr, an equivalent residue in FN3fn10 (FIG. 3D and FIG. 4B). Because it was envisioned that introducing Tyr, an amino acid particularly suitable for binding, at position 342 and the ability to diversify position 383 would both positively contribute to creating molecular recognition surfaces, the G342Y mutation was incorporated in the scaffold.

[0222] C. Phage and Yeast Display of STOFN3-1

[0223] Efficient display of a scaffold on the phage particle is a prerequisite for efficient selection of binding proteins using phage display. Highly stable, rapidly folding proteins present challenges in phage display, because phage display requires that the displayed protein fused to a phage coat protein be translocated into the periplasm of *E. coli* and highly stable proteins are not efficiently translocated across the *E. coli* inner membrane using a conventional, posttranslational secretion signal such as OmpT. An elegant solution was to use a co-translational secretion signal sequence such as the DsbA signal. Different signal sequences were examined for robust display of STOFN3-1. For this examination, the variant of STOTEMP4 with K318V mutation (STOTEMP5) was used because this mutation was crucial to function as the enhancer domain of affinity clamp (see below). Unexpectedly, it was found that an OmpT-based system achieved high levels of display of STOFN3-1 on the phage after optimization of the signal sequence and a linker length between the signal and STOFN3-1 with use of the C-terminal domain of M13 pIII. Notably, a single point mutation of the signal peptidase cleavage site from -SSFA/S- found in the vector pET12a (where the slash denotes the cleavage site) to -SSFA/A- and no additional residues between this signal sequence and the first residue of STOFN3-1 were crucial for the efficient display (FIG. 5). Unlike in a previous report, the R2K mutation of the OmpT signal did not improve surface display. For yeast surface display of STOFN3-1, a standard Aga2-mediated system as previously described that robustly displayed the protein as confirmed using flow cytometric analysis was used (data not shown).

[0224] D. Design and Evaluation of Combinatorial Libraries

[0225] Following previous designs of combinatorial libraries of FNfn10, two distinct libraries were constructed. One library is “loop only” library where positions in the BC, DE, and FG loops are utilized (FIG. 6A). The other library is “side-and-loop library” where residues in the C strand (residues 346 and 348) and the D strand (residues 352-354, 356 and 358) as well as residues in the CD and FG loops are utilized (FIG. 6B). Both libraries used highly biased amino acid diversity that emphasized Tyr, Ser, Gly and excluded Cys and Met, similar to designs used for FNfn10 and Fab libraries. In both libraries, the length of the FG loop was varied. In the “loop only” library, the length of the BC loop was varied. In the “side-and-loop” library, the same design of diversification as previous studies was applied to residues in the C and D strands and the CD loop. For diversifying the position of W348 on the C-strand, two sets of oligos were used for each length of CD-loop (3, 4, and 5 residues). One contains a codon for Tryptophan and the other contains codons for Serine, Threonine, Asparagine and Tyrosine so that this position could be diversified to five amino acids. Both libraries were constructed using the STOTEMP5 as a template scaffold in the phage-display format described above with estimated numbers of independent sequences of 2.5×10^9 and 1.7×10^9 for the loop only and side-and-loop libraries, respectively.

[0226] The inventors evaluated the performance of the two libraries using a total of six target proteins, yeast small ubiquitin-like modifier (γ SUMO), maltose-binding protein (MBP), enhanced green fluorescent protein (GFP), Abl SH2, SHP2 N- and C-SH2 domains. For each combination of

target and library, the inventors first enriched binding clones from the phage-display library, performed gene shuffling among the enriched population and identified high-affinity clones using yeast surface display. The inventors successfully generated binding proteins, called therMonobodies (thermophilic Monobodies), to ySUMO, MBP, EGFP and AblSH2 from both libraries and SHP2 N- and C-SH2 from the loop-only library (FIG. 6A, 6B). Most therMonobodies had Kd values in the low nanomolar range as measured in the yeast-display format (FIG. 6C). Residues in the FG loop were mutated in almost all the therMonobodies derived from either library, suggesting the importance of residues in the FG loop in target binding. Only exceptions were tMb (ABLSH2_L03) and tMb(ABLSH2_S01) that had no mutations in the FG loop, suggesting the possibility of achieving high affinity without utilizing the FG loop. We identified loop lengths that were not encoded in our designs. tMb (SUMO_L03) had 13 residues in the FG loop, one residue longer than the longest design, and tMb(NSH2_L06) and tMb(NSH2_L10) had four residues in the BC loop, one residue shorter than the shortest design. These sequences probably arose from errors in DNA synthesis and/or PCR errors, and their high functionality indicates that we could potentially expand the range of loop lengths in our library designs. For position 348, only Trp was selected, even though the position was diversified to a combination of five amino acids, Trp, Ser, Thr, Asn and Tyr. Subsequent mutation analysis showed that the replacement of W348 with Tyr or Ser destabilized STOTEMP5 by $>18^{\circ}$ C. and $>25^{\circ}$ C., respectively, indicating the importance of W348. In the crystal structure, the indole ring of W348 interacts with Y377 on the adjacent F-strand and also form a cation- π interaction with K354 on the adjacent D-strand. Replacement of W348 may well disrupt or weaken these interactions, and this destabilization effect may lead to the low occurrence frequencies.

[0227] These results clearly demonstrate that the combinatorial libraries constructed on the newly developed scaffold and the selection strategy can generate high affinity binding proteins.

[0228] E. Biophysical Characterization of therMonobodies

[0229] The inventors characterized the oligomerization state and thermal stability of purified therMonobodies (FIG. 7). Of 24 therMonobodies tested, 15 were predominantly monomeric as assayed using size-exclusion chromatography (SEC). Six eluted from the SEC as a monodispersed peak at the volume corresponding to a molecular weight much smaller than expected, suggesting that they were predominantly monomeric but weakly interacted with the column. The remaining three proteins eluted at the void volume, indicative of large aggregates. Thus, the vast majority (21/24) of these therMonobodies were produced as soluble, monomeric species. The thermal stability of the monomeric therMonobodies was assessed using DSF (FIG. 7B, C). The therMonobodies derived from the loop only library had a mean Tm value of $68 \pm 10^{\circ}$ C. with the highest and lowest of 86.3 and 53.5° C., respectively. Those generated from the side-and-loop library had a mean Tm value of $59.2 \pm 11.3^{\circ}$ C. with the highest and lowest of 85.5 and 50.5° C., respectively. Interestingly, the most stable clones, tMb(NSH2_L06) and tMb(NSH2_L10), had four residues in the BC loop, which is one residue shorter than the designed range of

the BC loop length. This finding suggests a new library design that better maintain the high stability of the underlying STOFN3-1 scaffold.

[0230] The highest Tm value of 86° C. and the lowest Tm value of 50° C. are both 13° C. higher than those of previously generated monobodies built from the FNfn10 scaffold (ref. Vazquez-Lombardi, 2015, Drug Discov Today). Thus, these results suggest that the therMonobodies system is better suited for generating thermo stable binding proteins.

[0231] F. therMonobody Functions as the Enhancer Domain of Affinity Clamps

[0232] The affinity clamping technology involves connecting a FN3 domain to a peptide-binding domain (such as an interaction domain or another FN3 domain) and subsequent optimization of the FN3 domain via directed evolution so as to create clamshell architecture that “clamps” a target motif in the newly generated interface between the two domains. The enlarged ligand interaction interface relative to that afforded by either domain achieves high specificity and high affinity. In this context, the FN3 portion is termed as the enhancer domain. To determine whether therMonobody can be used as the enhancer domain for affinity clamping technology, the FNfn10-based monobody was replaced in an affinity clamp directed to a phosphotyrosine (pY)-containing peptide with a therMonobody. The pY-clamp consists of an engineered Grb2 SH2 domain linked, via short linker, to a FNfn10 monobody. STOFN3-1 was structurally aligned with the FNfn10 monobody segment in the pY-clamp, clamp(Ptpn11_pY580). Residues of the monobody segment that were located within 5 \AA from the Grb2 SH2 domain or 6 \AA from the target pY-peptide were grafted to the structurally equivalent positions of STOTEMP4 (FIG. 8A). Because V4 located in the N-terminal tail of the monobody segment participates in the interaction with the Grb2 SH2 domain in the pY-clamp, the inventors also prepared a construct in which structurally equivalent K318 of STOTEMP4 was mutated to Val.

[0233] The inventors first examined biophysical properties of the designed therMonobody segments of the designed pY-clamps in isolation, i.e. not linked to the engineered SH2 domain. The therMonobody segments with and without the K318V mutation were expressed as $>50\%$ of soluble proteins and showed monodispersed size-exclusion chromatography profiles with an elution time consistent with that of their parent templates. The Tm values obtained from DSF were 68.7 and 63.5° C. for the variants without and with K318V mutation, respectively, indicating that both variants retained high stability. Together with the results mentioned in the above section, these results demonstrate the high tolerance of therMonobody template to extensive mutations.

[0234] Next the inventors examined binding properties of the designed pY-clamps to the target pY-peptide (Ptpn11_pY580). A total of six constructs were tested by combining the presence and absence of the K318V mutation with linker lengths of 0, 2 and 5 amino acids between the Grb2 SH2 domain and the therMonobody segment. Two of them, with the K318V mutation and linker lengths of 2 or 5 residues showed significantly higher binding to the pY-peptide than the original Grb2 SH2 domain alone (FIG. 8B), with KD values of 156 ± 42 and 121 ± 21 nM for the 2- and 5-residue linkers, respectively (FIG. 8C). For comparison, clamp (Ptpn11_pY580) with the optimized enhancer domain based on FNfn10 had a 6 amino acids linker between the Grb2 SH2

domain and the monobody segment and a KD value of 2.5 nM. Although the affinity of the therMonobody versions of the pY-clamp was lower, these results clearly indicates that therMonobody can readily be used as a building block for generating affinity clamps.

Example 2: Hyperthermophilic Fn3 Domain from
Pyrococcus horikoshii OT3

[0235] The inventors utilized the SMART database to explore FN3 domains from hyperthermophiles. The database predicted many FN3 domains in hyperthermophilic archaea and bacteria such as *Thermococcus kodakaraensis*, *Sulfolobus tokodaii*, *Pyrococcus horikoshii* and *Thermotoga lettingae*. Predicted domains that were shorter than the length of the shortest FN3 domains that had been structurally characterized (75 amino acids) were first eliminated. Then, a FN3 domain in the sequence of putative uncharacterized protein PH0954 from the hyperthermophilic archaeon *Pyrococcus horikoshii* OT3, termed PHOFN3 was chosen as the candidate protein, because of its detectable homology to a bacterial FN3 domain in the *Clostridium Perfringens* Glycoside Hydrolase Gh84c whose FN3 fold has been experimentally confirmed (PDBID: 2W1N). The SMART database predicted PHOFN3 with 108 (P1873-A1980) amino acid residues, but the C-terminal 26 residues (E1955-A1980) did not have detectable homology to the sequence of the bacterial homologue (FIG. 9).

[0236] In order to confirm whether the C-terminal extended region is a part of the core structure of PHOFN3, the inventors constructed expression vectors for different segments of PHOFN3 with the same N-terminus but with different truncations from the C-terminus, PHOFN3 (P1873-A1980), PHOFN3ΔC21 (P1873-E1959), PHOFN3ΔC25 (P1873-E1955) and PHOFN3ΔC35 (P1873-E1945), and analyzed their soluble expressions in *E. coli* and thermal stabilities. It was found that PHOFN3, PHOFN3ΔC21 and PHOFN3ΔC25 were robustly produced in *E. coli* as soluble proteins that were predominantly monomeric. Melting temperatures obtained by differential scanning fluorimetry (DSF) were >95° C. for PHOFN3, PHOFN3ΔC21 and PHOFN3ΔC25 at pH 7.4, confirming that they were highly thermostable proteins (FIG. 10). These results suggested that the C-terminal extended region, which was not aligned with the bacterial homologue, was not a part of the core structure of PHOFN3. In contrast, PHOFN3ΔC35 was expressed mostly as insoluble proteins and the melting temperature was 32.2° C. at pH 7.4 (FIG. 10), suggesting that the region of K1946-E1955 was important for folding a proper structure and for thermal stability as the region was aligned with the last β-strand (G strand) of the bacterial homologue.

[0237] To confirm that PHOFN3 had the FN3 fold, the inventors determined the crystal structure of PHOFN3ΔC25 at 1.7 Å resolution with Se-Met labeled at M1882 and I1905M. The PHOFN3ΔC25 segment indeed adopts the FN3 fold consisting of seven anti-parallel β-strands and six loops (FIG. 11A). Superposition with the FN3fn10 (PDBID: 1FNA), the most well characterized FN3 domain, demonstrated that the structure of the segment is highly similar to FN3fn10 with an average RMSD value of 2.2 Å for aligned backbone Ca atoms excluding residues in loops (FIG. 11B). Figure E shows the schematic drawing of the amino acid sequence of PHOFN3ΔC25 in its secondary structure context.

[0238] Phage display of PHOFN3. A vector for phage display of PHOFN3 (without C-terminal deletion, P1873-A1980) was constructed based on the previously reported DsbA-based vector containing the V5 tag sequence for display detection (Wojcik et al., NSMB). Phage particles were produced using helper phage, M13KO7 or hyperphage, and the display of PHOFN3 on phage particles was examined by phage ELISA using an anti-V5 tag antibody, a HRP conjugated anti-M13 phage antibody and 1-step Ultra TMB ELISA (Thermo Scientific). The absorbance changes at 405 nm after the HRP reaction for 10 min for M13KO7-produced phages or 1 min for hyperphage-produced phages are shown in FIG. 12. FIG. 13 shows the significant levels of display of PHOFN3 on phage particles. The phage ELISA signals for PHOFN3 with M13KO7 and hyperphage were lower than those for FN3fn10 only by 2.9 and 1.3-folds, respectively, strongly suggesting that a combinatorial library similar to those for FN3fn10 can be constructed using phage display from which novel binding proteins can be identified.

[0239] Although certain embodiments have been described above with a certain degree of particularity, or with reference to one or more individual embodiments, those skilled in the art could make numerous alterations to the disclosed embodiments without departing from the scope of this invention. Further, where appropriate, aspects of any of the examples described above may be combined with aspects of any of the other examples described to form further examples having comparable or different properties and addressing the same or different problems. Similarly, it will be understood that the benefits and advantages described above may relate to one embodiment or may relate to several embodiments. Any reference to a patent publication or other publication is a herein a specific incorporation by reference of the disclosure of that publication. The claims are not to be interpreted as including means-plus- or step-plus-function limitations, unless such a limitation is explicitly recited in a given claim using the phrase(s) “means for” or “step for,” respectively.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 7

<210> SEQ ID NO 1
<211> LENGTH: 84
<212> TYPE: PRT
<213> ORGANISM: Sulfolobus tokodaii

<400> SEQUENCE: 1

Pro Pro Pro Lys Pro Gln Ile Ala Ser Ile Ala Ser Gly Asn Glu Thr
1 5 10 15

-continued

```

Ile Thr Val Lys Trp Tyr Asp Thr Asn Ala Ser Gly Tyr Tyr Ile Thr
      20                      25                      30
Tyr Trp Ser Asn Phe Ser Gln Lys Val Thr Ile Asn Val Gly Asn Val
      35                      40                      45
Thr Ser Tyr Thr Ile Lys His Leu Lys Asp Gly Val Thr Tyr Tyr Ile
      50                      55                      60
Gln Ile Val Pro Tyr Asn Ser Leu Gly Asn Gly Thr Pro Ser Asp Ile
      65                      70                      75                      80
Ile Ser Ala Thr

```

```

<210> SEQ ID NO 2
<211> LENGTH: 86
<212> TYPE: PRT
<213> ORGANISM: Sulfolobus tokodaii

```

```

<400> SEQUENCE: 2

```

```

Pro Ser Ser Val Pro Asn Pro Pro Ile Ile Lys Val Lys Ile Gly Asn
 1      5                      10                      15
Leu Asn Ala Thr Leu Thr Trp Tyr Asp Thr Phe Asn Gly Gly Tyr Pro
      20                      25                      30
Ile Glu Gly Tyr Tyr Leu Tyr Val Asn Gly Lys Gly Ile Asn Val Gly
      35                      40                      45
Asn Ile Thr Ser Tyr Val Leu Thr Asn Leu Thr Ala Gly Glu Leu Tyr
      50                      55                      60
Thr Ile Glu Leu Ile Ala Tyr Asn Lys Ile Gly Asn Ser Ser Ile Ser
      65                      70                      75                      80
Ser Val Ser Phe Ile Ala
      85

```

```

<210> SEQ ID NO 3
<211> LENGTH: 79
<212> TYPE: PRT
<213> ORGANISM: Sulfolobus tokodaii

```

```

<400> SEQUENCE: 3

```

```

Ala Asn Leu Thr Val Thr Val Tyr Lys Lys Ile Asn Gly Phe Leu Val
 1      5                      10                      15
Ser Trp Asn Ser Thr Ser Lys Ala Lys Tyr Ile Leu Thr Val Ser Lys
      20                      25                      30
Glu Asn Val Val Leu Leu Asn Val Ser Thr Thr Asn Thr Ser Tyr Phe
      35                      40                      45
Val Lys Val Pro Phe Gly Val Tyr Asn Ile Ser Leu Glu Ala Val Asn
      50                      55                      60
Ile Val Gly Ile Thr Lys Tyr Ala Phe Ile Leu Ile Tyr Tyr Ile
      65                      70                      75

```

```

<210> SEQ ID NO 4
<211> LENGTH: 76
<212> TYPE: PRT
<213> ORGANISM: Sulfolobus tokodaii

```

```

<400> SEQUENCE: 4

```

```

Pro Ala Ser Pro Thr Val Asn Trp Ser Ile Thr Leu Asn Thr Val Ser
 1      5                      10                      15
Leu Asn Trp Ser Lys Val Ser Gly Ala Glu Tyr Tyr Leu Ile Tyr Asp

```


-continued

	20	25	30	
Asn Gly Lys Leu Ile Thr Asn Thr Thr Asn Thr Ala Phe Thr Phe Asn	35	40	45	
Leu Thr Ile Gly Gln Asn Glu Ile Glu Val Tyr Ala Ala Asn Ala Tyr	50	55	60	
Tyr Lys Ser Ala Pro Tyr Ile Ile Asn Asp Val Arg	65	70	75	
 <210> SEQ ID NO 5				
<211> LENGTH: 709				
<212> TYPE: PRT				
<213> ORGANISM: Sulfolobus tokodaii				
 <400> SEQUENCE: 5				
Met Lys Arg Asn Thr Leu Leu Ala Leu Val Leu Val Ile Leu Ile Phe	1	5	10	15
Pro Thr Leu Ser Thr Ala Tyr Ile Glu Phe Thr Thr Ser Ile Asn Gln	20	25	30	
Ala Ile Pro Asp Ser Leu Val Tyr Ala Thr Ser Ala Tyr Tyr Asp Gly	35	40	45	
Lys Ile Phe Leu Ile Gly Gly Glu Asn Leu Tyr Ser Thr Pro Val Asn	50	55	60	
Ser Val Tyr Val Tyr Glu Asn Gly Ser Trp Tyr Leu Gly Pro Ser Leu	65	70	75	80
Pro Phe Ser Leu Ser Ser Ala Gly Ala Thr Val Cys Asn Asn Thr Leu	85	90	95	
Tyr Val Val Gly Gly Ala Asn Ser Thr Ser Ile Phe Gly Gly Ile Leu	100	105	110	
Glu Phe Ile Gly Asn Gly Trp Lys Val Ile Thr Asn Ser Met Pro Ile	115	120	125	
Pro Val Tyr Gly Ala Ile Val Phe Ser Tyr Asp Tyr Lys Ile Tyr Val	130	135	140	
Ile Gly Gly Met Asn Tyr Ser Gly Asn Ser Leu Val Pro Pro Val Asn	145	150	155	160
Tyr Ile Gln Val Tyr Asn Leu Lys Thr Asn Ser Trp Gln Ile Ile Gly	165	170	175	
Asn Ala Pro Leu Arg Leu Ala Tyr Ser Ala Tyr Tyr Phe Asn Gly Ser	180	185	190	
Ala Leu Phe Val Val Gly Gly Phe Thr Gln Ser Ala Thr Leu Thr Ser	195	200	205	
Ser Val Phe Val Tyr Tyr Pro Glu Asn Asn Thr Trp Ile Ser Leu Pro	210	215	220	
Ser Leu Pro Gly Val Glu Ala Gly Gly Val Leu Gly Tyr Tyr Asn Gly	225	230	235	240
Tyr Met Tyr Leu Val Gly Gly Leu Tyr Tyr Val Ser Gly Ala Tyr Gln	245	250	255	
Leu Gly Glu Ile Leu Tyr Tyr Tyr Asn Gly Thr Trp Arg Asn Thr Asn	260	265	270	
Ile Gln Glu Gln Ile Pro Thr Gln Phe Ser Thr Ser Val Gln Ile Gly	275	280	285	
Asn Lys Leu Ile Ile Leu Gly Gly Phe Gly Pro Gly Asn Ile Pro Ser	290	295	300	

-continued

Asn Ala Met Gln Thr Val Ser Ile Tyr Leu Pro Pro Pro Lys Pro Gln
 305 310 315 320
 Ile Ala Ser Ile Ala Ser Gly Asn Glu Thr Ile Thr Val Lys Trp Tyr
 325 330 335
 Asp Thr Asn Ala Ser Gly Tyr Tyr Ile Thr Tyr Trp Ser Asn Phe Ser
 340 345 350
 Gln Lys Val Thr Ile Asn Val Gly Asn Val Thr Ser Tyr Thr Ile Lys
 355 360 365
 His Leu Lys Asp Gly Val Thr Tyr Tyr Ile Gln Ile Val Pro Tyr Asn
 370 375 380
 Ser Leu Gly Asn Gly Thr Pro Ser Asp Ile Ile Ser Ala Thr Pro Ser
 385 390 395 400
 Ser Val Pro Asn Pro Pro Ile Ile Lys Val Lys Ile Gly Asn Leu Asn
 405 410 415
 Ala Thr Leu Thr Trp Tyr Asp Thr Phe Asn Gly Gly Tyr Pro Ile Glu
 420 425 430
 Gly Tyr Tyr Leu Tyr Val Asn Gly Lys Gly Ile Asn Val Gly Asn Ile
 435 440 445
 Thr Ser Tyr Val Leu Thr Asn Leu Thr Ala Gly Glu Leu Tyr Thr Ile
 450 455 460
 Glu Leu Ile Ala Tyr Asn Lys Ile Gly Asn Ser Ser Ile Ser Ser Val
 465 470 475 480
 Ser Phe Ile Ala Ala Ser Lys Ala Asn Leu Thr Val Thr Val Tyr Lys
 485 490 495
 Lys Ile Asn Gly Phe Leu Val Ser Trp Asn Ser Thr Ser Lys Ala Lys
 500 505 510
 Tyr Ile Leu Thr Val Ser Lys Glu Asn Val Val Leu Leu Asn Val Ser
 515 520 525
 Thr Thr Asn Thr Ser Tyr Phe Val Lys Val Pro Phe Gly Val Tyr Asn
 530 535 540
 Ile Ser Leu Glu Ala Val Asn Ile Val Gly Ile Thr Lys Tyr Ala Phe
 545 550 555 560
 Ile Leu Ile Tyr Tyr Ile Gln Pro Ala Ser Pro Thr Val Asn Trp Ser
 565 570 575
 Ile Thr Leu Asn Thr Val Ser Leu Asn Trp Ser Lys Val Ser Gly Ala
 580 585 590
 Glu Tyr Tyr Leu Ile Tyr Asp Asn Gly Lys Leu Ile Thr Asn Thr Thr
 595 600 605
 Asn Thr Ala Phe Thr Phe Asn Leu Thr Ile Gly Gln Asn Glu Ile Glu
 610 615 620
 Val Tyr Ala Ala Asn Ala Tyr Tyr Lys Ser Ala Pro Tyr Ile Ile Asn
 625 630 635 640
 Asp Val Arg Asn Tyr Ile Val Val Val Asn Ser Thr Ala Ile Ser Ile
 645 650 655
 Ser Val Pro Gln Ile Lys Val Val Ser Gly Glu Asn Thr Asp Ala Pro
 660 665 670
 Leu Gln Thr Asn Asn Ile Asp Leu Lys Ser Ala Ile Ile Val Ile Thr
 675 680 685
 Val Phe Val Ile Ala Leu Leu Met Ile Leu Val Ile Leu Arg Glu Arg
 690 695 700
 Ser Asp Asn Tyr Trp

-continued

705

<210> SEQ ID NO 6
 <211> LENGTH: 108
 <212> TYPE: PRT
 <213> ORGANISM: Pyrococcus horikoshii

<400> SEQUENCE: 6

```

Pro Ser Pro Pro Ser Gly Val Thr Leu Met Leu Asn Gly Ser Tyr Val
1          5          10          15
Glu Leu Ser Trp Leu Pro Ser Pro Asp Ser Asp Val Ala Gly Tyr Phe
20          25          30
Ile Tyr Lys Asp Gly Lys Arg Leu Asn Glu Val Pro Ile Glu Lys Pro
35          40          45
Asn Phe Arg Asp Ile Tyr Ser Gly Thr Leu Asn Tyr Ser Ile Ser Ala
50          55          60
Ile Asp Phe Ser Gly Phe Glu Ser Glu Lys Thr Glu Val Phe Pro Val
65          70          75          80
Lys Leu Glu Val Asp Glu Glu Asn Leu Thr Ala Gly Tyr Pro Gly Ala
85          90          95
Val Lys Val Lys Val Glu Asn Leu Asp Gly Glu Ala
100          105

```

<210> SEQ ID NO 7
 <211> LENGTH: 4436
 <212> TYPE: PRT
 <213> ORGANISM: Pyrococcus horikoshii

<400> SEQUENCE: 7

```

Met Ile Asn Ile Lys Gly Leu Ile Leu Thr Leu Ile Leu Phe Ile Ser
1          5          10          15
Leu Ile Pro Pro Trp Ala Leu Gly Glu Gly Ser Lys Asp Thr Lys Val
20          25          30
Phe Ala Asp Tyr Tyr Leu Ala Gly Asp Ser Val Val Ile Asn Ala Thr
35          40          45
Leu Tyr Asp Ala Gly Ser Cys Asn Leu Thr Phe Ser Val Phe Ser Pro
50          55          60
Ile Glu Ala Pro Asn Val Ser Glu Ile Ser Phe Thr Trp Met Asn Leu
65          70          75          80
Ser Glu Tyr Ile Glu Ser Ala Thr Glu Ala Thr Tyr Gly Glu Tyr Leu
85          90          95
Arg Asp Gly Asn Val Ile Met Arg Glu Asp Asp Gly Tyr Phe Ile Tyr
100          105          110
Glu Leu Pro Phe Ser Leu Asn Tyr Phe Gly Arg Glu Ile Lys Lys Ile
115          120          125
Ala Val Asn Thr Asn Gly Leu Ile Glu Leu Leu Glu Glu Tyr Glu Glu
130          135          140
Pro Arg Ile Glu Asp Tyr Tyr Gly Ile His Glu Glu Gly Glu Phe Tyr
145          150          155          160
Glu Ser Asp Val Ile Phe Gly Leu Asp Glu Asp Leu Val Thr Tyr Asp
165          170          175
Gly Tyr Leu Leu Leu Val Asn Leu Gln Asp Lys Ile Val Ile Glu Trp
180          185          190
Leu Ala Ser Thr Tyr Glu Asp Tyr Glu Ser Glu Ile Val Asp Asn Ile

```

-continued

195					200					205					
Asn	Phe	Gln	Val	Ile	Ile	Asn	Ser	Asn	Gly	Thr	Ile	Thr	Trp	Ser	Tyr
210						215					220				
Lys	Ser	Leu	Glu	Tyr	Ser	Tyr	His	Asp	Tyr	Asp	Leu	Phe	Ser	Gly	Tyr
225					230					235					240
Tyr	Ser	Lys	Val	Ser	Gly	Asp	Val	Lys	Gly	Phe	Thr	Lys	Gly	Glu	Gly
				245					250					255	
Lys	Ser	Phe	Ala	Ile	Gln	Val	Pro	Leu	Gly	Thr	Pro	Lys	Leu	Tyr	Thr
			260					265					270		
Tyr	Gln	Val	Arg	Glu	Ser	Gly	Ser	Tyr	Leu	Leu	Thr	Leu	Pro	Leu	Ser
			275					280					285		
Asn	Tyr	His	Val	Glu	Val	Phe	Ala	Asn	Cys	Met	Asp	Asp	Pro	Asp	Leu
290						295					300				
Ser	Asn	Asn	Leu	Ala	Glu	Val	Gly	Val	Trp	Pro	Gly	Asp	Tyr	Trp	Val
305					310					315					320
Glu	Asn	Ala	Ser	Ile	Asn	Asn	Leu	Ile	Pro	Gly	Glu	Phe	Ala	Ser	Ile
				325					330						335
Asn	Phe	Lys	Val	Arg	Thr	Thr	Ser	Lys	Ile	Pro	Ser	Ala	Lys	Val	Lys
			340					345						350	
Leu	Leu	Arg	Asn	Gly	Val	Glu	Glu	Lys	Ile	Glu	Tyr	Leu	Ser	Phe	Tyr
		355					360						365		
Asn	Gly	Ile	Ala	Glu	Gly	Glu	Ile	Ser	Trp	Leu	Val	Gln	Gly	Gly	Asn
370						375					380				
Tyr	Thr	Leu	Ala	Leu	Leu	Val	Glu	Gly	Lys	Gly	Asp	Ile	Asn	Ser	Ser
385					390					395					400
Asn	Asn	Ile	Tyr	Leu	Leu	Gly	Asn	Tyr	Asn	Phe	Pro	Leu	Pro	Asn	Phe
				405					410						415
Glu	Val	Gly	Asn	Tyr	Ser	Ile	Asp	Leu	Pro	Thr	Cys	Val	Asp	Ser	Thr
			420					425					430		
Gly	Glu	Val	Arg	Val	Asn	Val	Thr	Ser	Thr	Ala	Asn	Trp	Ser	Ile	Pro
		435					440						445		
Val	Arg	Leu	Thr	Leu	Val	Tyr	Glu	Glu	Gly	Asn	Arg	Ser	Tyr	Thr	Arg
450						455					460				
Tyr	Ile	Ser	Thr	Lys	Gly	Glu	Glu	Glu	Ser	Glu	Val	Ile	Phe	Thr	Pro
465					470					475					480
Met	Ile	Lys	Ala	Gly	Thr	Leu	Glu	Lys	Val	Val	Ile	Glu	Ile	Asp	Pro
				485					490						495
Trp	Asn	Glu	Val	Glu	Glu	Ser	Asn	Glu	Ser	Asp	Asn	Lys	Val	Glu	Val
		500						505						510	
Pro	Tyr	His	Ile	Ile	Ile	Glu	Lys	Pro	Asp	Phe	Thr	Val	Lys	Ser	Leu
		515						520						525	
Asn	Ile	Pro	Gly	Asn	Val	Ser	Ile	Gly	Asn	Leu	Tyr	Glu	Val	Asn	Val
530						535								540	
Thr	Leu	Asp	Asn	Leu	Gly	Gly	Cys	Tyr	Gly	Arg	Asn	Val	Leu	Val	Lys
545					550					555					560
Leu	Tyr	Glu	Asn	Gly	Thr	Ser	Lys	Asp	Trp	Arg	Arg	Val	Arg	Ile	Asn
				565					570					575	
Asn	Glu	Thr	Asn	Val	Thr	Leu	Thr	Trp	Lys	Pro	Gly	Asn	Ala	Gly	Leu
				580					585					590	
Val	Asn	Leu	Thr	Val	Val	Val	Asp	Pro	Tyr	Ser	Tyr	Val	Asp	Glu	Ile
		595						600						605	

-continued

Asn Glu Gly Asn Asn Arg Leu Ser Arg Leu Ile Phe Val Asn Ala Pro
 610 615 620
 Asp Phe Lys Ile Ser Lys Val Glu Leu Leu Ser Phe Asp Gly Ile Ala
 625 630 635 640
 Gly Ser Lys Ala Lys Phe Asn Val Thr Val Lys Asn Glu Gly Glu Asp
 645 650 655
 Tyr Ser Gly Tyr Phe Ser Ile Ala Val Tyr Gly Gly Leu Arg Ser Ser
 660 665 670
 Ile Ala Tyr Leu Arg Gly Ile Lys Ser Gly Glu Glu Lys Trp Thr Ile
 675 680 685
 Ile Ser Leu Pro Ile Asn Gly Gly Asn Ser Thr Leu Ile Phe Val Val
 690 695 700
 Asp Pro His Asn Val Ile Ser Glu Thr Asn Glu Gly Asn Asn Val Ile
 705 710 715 720
 Phe Tyr Asn Met Gly Tyr Ile Pro Lys Pro Asn Phe Val Val Lys Glu
 725 730 735
 Ile Ser Leu Pro Asn Asn Thr Val Gly Tyr Ile Pro Leu Asn Ile Thr
 740 745 750
 Ile Gly Asn Val Gly Ala Pro Tyr Asn Ala Thr Ser Tyr Gln Val Pro
 755 760 765
 Val Lys Ile Lys Thr Glu Tyr Gly Trp Lys Val Ser Tyr Leu Arg Gly
 770 775 780
 Ile Ile Arg Asp Asn Tyr Thr Ile Ser Ile Asp Ser Leu Ala Met Leu
 785 790 795 800
 Pro Pro Gly Ser Thr Ile Asn Val Thr Val Asn Tyr Asn Met Lys Val
 805 810 815
 Asn Glu Thr Ser Tyr Ser Asp Asn Ser Leu Ile Ile Asn Tyr Thr Thr
 820 825 830
 Gly Tyr Pro Asp Leu Glu Leu Gly Ile Ile Pro Pro Ser Gly Glu Leu
 835 840 845
 Ser Ala Gly Lys Asp Val Lys Ile Thr Phe Leu Val Lys Asn Val Gly
 850 855 860
 Asn Ala Thr Leu Arg Ile Asp Arg Ser Ser Trp Tyr Ser Pro Tyr Leu
 865 870 875 880
 Gly Leu Tyr Val Thr Leu Glu Asp Glu Asn Gly Lys Thr His Thr Leu
 885 890 895
 Gly Arg Tyr Glu Leu Ala Pro Ala Thr Leu Ser Pro Gly Ala Asn Ile
 900 905 910
 Ser Gln Val Val Trp Ile Thr Leu Asn Gly Gly Thr Asn Lys Ile Met
 915 920 925
 Gly Arg Ile Val Asp Glu Tyr Glu Asn Ile Tyr Glu Asn Asn Asn Asp
 930 935 940
 Thr Leu Ile Leu Thr Leu Glu Lys Pro Asp Phe Ala Ile Leu Asn Tyr
 945 950 955 960
 Ser Ile Pro Asp Glu Ile Leu Asn Gly Thr Ala Tyr Leu Tyr Lys Ala
 965 970 975
 Tyr Pro Ile Val Leu Asn Ile Ser Asn Leu Gly Gly Asn Phe Ser Asp
 980 985 990
 Gly Ile Arg Val Asp Leu Phe Asp Asn Gly Ile Ile Lys Thr Ser Thr
 995 1000 1005

-continued

Ser	Val	Tyr	Gly	Leu	Glu	Ser	Gly	Ala	Ser	Arg	Asp	Val	Thr	Leu
1010						1015					1020			
Arg	Tyr	Leu	Pro	Ser	Ser	Gly	Lys	His	Asn	Leu	Ser	Ile	Val	Leu
1025						1030					1035			
Asp	Pro	Tyr	Asn	Arg	Trp	Ile	Glu	Glu	Asn	Glu	Glu	Asn	Asn	Asn
1040						1045					1050			
Leu	Thr	Phe	Ser	Leu	Ser	Phe	Gly	Lys	Pro	Asp	Leu	Lys	Val	Glu
1055						1060					1065			
Gly	Ile	Thr	Trp	Ala	Pro	Tyr	Asn	Phe	Thr	Ser	Gly	Glu	Asn	Val
1070						1075					1080			
Leu	Phe	Thr	Ile	Tyr	Val	Lys	Asn	Leu	Gly	Gln	Pro	Phe	Leu	Lys
1085						1090					1095			
Ser	Phe	Thr	Val	Arg	Ala	Glu	Ile	Trp	Asn	Gly	Thr	Arg	Lys	Ile
1100						1105					1110			
Tyr	Ser	Thr	Asn	Ala	Tyr	Pro	Arg	Asn	Trp	Ser	Phe	Gly	Lys	Gly
1115						1120					1125			
Glu	Thr	Lys	Glu	Phe	Asn	Trp	Arg	Trp	Tyr	Asn	Ala	Lys	Pro	Gly
1130						1135					1140			
Asn	Leu	Thr	Val	Lys	Ile	Val	Val	Asp	Tyr	Tyr	Asn	Ser	Ile	Pro
1145						1150					1155			
Glu	Gly	Asn	Glu	Ser	Asn	Asn	Glu	Phe	Ser	Ala	Phe	Leu	Gly	Asn
1160						1165					1170			
Val	Gly	Thr	Pro	Asp	Phe	Lys	Leu	Glu	Asn	Leu	Ser	Val	Glu	Asp
1175						1180					1185			
Leu	Ala	Tyr	Gly	Lys	Phe	Val	Arg	Ile	Asn	Ala	Thr	Val	Lys	Asn
1190						1195					1200			
Leu	Gly	Asp	Ser	Ile	Tyr	Arg	Pro	Ile	Thr	Val	Leu	Phe	Asn	Val
1205						1210					1215			
Ser	Gly	Glu	Arg	Tyr	Tyr	Arg	Thr	Val	Tyr	Gly	Ile	Lys	Glu	Asn
1220						1225					1230			
Glu	Ser	Lys	Ser	Val	Thr	Leu	Pro	Trp	Tyr	Val	Asp	Arg	Val	Gly
1235						1240					1245			
Glu	Val	Arg	Val	Lys	Val	Glu	Val	Asp	Pro	Gly	Asn	Arg	Ile	Val
1250						1255					1260			
Glu	Gly	Asn	Glu	Ser	Asn	Asn	Ile	Ile	Glu	Arg	Thr	Tyr	Tyr	Val
1265						1270					1275			
Glu	Ser	Pro	Glu	Leu	Met	Leu	Ser	Gly	Tyr	Glu	Trp	Leu	Glu	Glu
1280						1285					1290			
Glu	Val	Arg	Arg	Gly	Tyr	Leu	Ala	Tyr	Lys	Val	Asn	Val	Thr	Asn
1295						1300					1305			
Thr	Gly	Gly	Asp	Val	Tyr	Arg	Gly	Phe	Tyr	Val	Gln	Met	Phe	Val
1310						1315					1320			
Asp	Gly	Glu	Pro	Lys	Ser	Ser	Val	Trp	Ile	Asn	Lys	Leu	Leu	His
1325						1330					1335			
Gly	Glu	Thr	Ala	Glu	Arg	Thr	Leu	Arg	Trp	Arg	Phe	Ser	Ser	Gly
1340						1345					1350			
Gly	Arg	Lys	Glu	Val	Arg	Ile	Val	Val	Asp	Pro	Gln	Asp	Tyr	Ile
1355						1360					1365			
Pro	Glu	Ser	Asn	Glu	Asp	Asn	Asn	Ala	Ile	Val	Glu	Asn	Val	Thr
1370						1375					1380			
Ile	Val	Leu	Pro	Asp	Ile	Glu	Val	Leu	Ser	Leu	Asn	Ile	Pro	Ser

-continued

Thr	Val	Lys	Ala	Val	Asp	Arg	Ala	Gly	Asn	Glu	Ile	Thr	Tyr	Ser
1775						1780					1785			
Trp	Asn	Phe	Thr	Leu	Asp	Arg	Glu	Pro	Pro	Arg	Ile	Val	Cys	Asn
1790						1795					1800			
Leu	Thr	Asp	Gly	Thr	Leu	Tyr	Asn	Gly	Thr	Val	Val	Pro	Gly	Val
1805						1810					1815			
Gln	Val	Ile	Asp	Asp	Asn	Leu	Asp	Trp	Tyr	Lys	Val	Lys	Val	Asn
1820						1825					1830			
Gly	Arg	Glu	Phe	Ser	Gly	Pro	Ile	Lys	Leu	Asp	Gly	Thr	Tyr	Thr
1835						1840					1845			
Leu	Asn	Val	Thr	Ala	Lys	Asp	Lys	Ala	Gly	Asn	Leu	Ala	Glu	Lys
1850						1855					1860			
Ile	Ile	Arg	Phe	Thr	Val	Asn	Gly	Val	Pro	Ser	Pro	Pro	Ser	Gly
1865						1870					1875			
Val	Thr	Leu	Met	Leu	Asn	Gly	Ser	Tyr	Val	Glu	Leu	Ser	Trp	Leu
1880						1885					1890			
Pro	Ser	Pro	Asp	Ser	Asp	Val	Ala	Gly	Tyr	Phe	Ile	Tyr	Lys	Asp
1895						1900					1905			
Gly	Lys	Arg	Leu	Asn	Glu	Val	Pro	Ile	Glu	Lys	Pro	Asn	Phe	Arg
1910						1915					1920			
Asp	Ile	Tyr	Ser	Gly	Thr	Leu	Asn	Tyr	Ser	Ile	Ser	Ala	Ile	Asp
1925						1930					1935			
Phe	Ser	Gly	Phe	Glu	Ser	Glu	Lys	Thr	Glu	Val	Phe	Pro	Val	Lys
1940						1945					1950			
Leu	Glu	Val	Asp	Glu	Glu	Asn	Leu	Thr	Ala	Gly	Tyr	Pro	Gly	Ala
1955						1960					1965			
Val	Lys	Val	Lys	Val	Glu	Asn	Leu	Asp	Gly	Glu	Ala	Asn	Gly	Thr
1970						1975					1980			
Leu	Ser	Ile	Ile	Leu	Ile	Asp	Glu	Phe	Gly	Asn	Glu	Ile	Glu	Lys
1985						1990					1995			
Leu	Ser	Arg	Lys	Val	Glu	Val	Pro	Arg	Gly	Arg	Ser	Ser	His	Glu
2000						2005					2010			
Phe	Val	Phe	Met	Val	Pro	Arg	Gly	Leu	Thr	Leu	Ile	Arg	Gly	Glu
2015						2020					2025			
Leu	Lys	Val	Gly	Asn	Ser	Thr	Ala	Arg	Ile	Ile	His	Arg	Ala	Lys
2030						2035					2040			
Val	Arg	Glu	Gly	Glu	Asn	Pro	Glu	Ile	Arg	Val	Gly	Lys	Leu	Leu
2045						2050					2055			
Ala	Gly	Phe	Pro	Gly	Leu	Val	Glu	Val	Glu	Ile	Arg	Asn	Cys	Gly
2060						2065					2070			
Ile	Val	Glu	Leu	Asn	Thr	Ser	Glu	Thr	Leu	Met	Lys	Leu	Asp	Asn
2075						2080					2085			
Ser	Ser	Gly	Glu	Leu	Ile	Glu	Ala	Pro	Leu	Thr	Ile	Pro	Pro	Gly
2090						2095					2100			
Lys	Lys	Thr	Val	Leu	Arg	Tyr	Lys	Ile	Val	Pro	Pro	Lys	Lys	Gly
2105						2110					2115			
Ser	Tyr	Asn	Leu	Thr	Phe	Arg	Ile	Ala	Asp	Val	Glu	Val	Arg	Lys
2120						2125					2130			
Ile	Val	Asn	Val	Ser	Glu	Ser	Val	Leu	Asn	Pro	Ile	Thr	Ile	Ser
2135						2140					2145			

-continued

Thr 2150	Glu	Asn	Phe	Ile	Lys	Gly 2155	Gly	Lys	Ala	Lys	Ile 2160	Tyr	Val	Ser
Phe 2165	Arg	Asn	Ile	Gly	Ser	Ala 2170	Pro	Ile	Phe	Val	Lys 2175	Ser	Ile	Glu
Leu 2180	Asn	Gly	Met	Ser	Lys	Arg 2185	Leu	Ser	Ile	Glu	Leu 2190	Pro	Pro	Asn
Leu 2195	Ser	Val	Glu	Glu	Ser	Phe 2200	Glu	Tyr	Leu	Ile	Ser 2205	Glu	Glu	Asn
Val 2210	Glu	Ile	Asn	Ala	Thr	Val 2215	Asn	Thr	Asp	Val	Gly 2220	Lys	Phe	Arg
Lys 2225	Ser	Leu	Thr	Leu	Thr	Ala 2230	Glu	Gln	Pro	Glu	Tyr 2235	Asn	Ala	Asp
Val 2240	Ser	Val	Ser	Ser	Val	Tyr 2245	Glu	Val	Gly	Lys	Glu 2250	Ile	Leu	Ile
Thr 2255	Gly	Val	Ala	Tyr	Asn	Glu 2260	Ser	Gly	Met	Leu	Ser 2265	Asn	Val	Pro
Val 2270	Lys	Val	Ser	Ile	Ala	Arg 2275	Gly	Gly	Phe	Val	Arg 2280	Glu	Tyr	Ile
Val 2285	Thr	Thr	Asn	Glu	Asn	Gly 2290	Tyr	Phe	Asn	Leu	Thr 2295	Phe	Arg	Pro
Phe 2300	Lys	Gly	Glu	Ser	Gly	His 2305	Phe	Ile	Val	Ser	Ala 2310	Thr	His	Pro
Lys 2315	Ile	Glu	Leu	Leu	Glu	Arg 2320	Asp	Ala	Glu	Phe	Asp 2325	Val	Val	Gly
Ile 2330	Glu	Val	Ile	Pro	Ser	Leu 2335	Tyr	Leu	Leu	Thr	Val 2340	Pro	Val	Glu
Phe 2345	Asn	Gly	Thr	Val	Arg	Val 2350	Arg	Leu	Ile	Asn	Tyr 2355	Trp	Arg	Ala
Ser 2360	Asp	Val	Ser	Val	Ser	Val 2365	Lys	Ala	Pro	Pro	Glu 2370	Tyr	Glu	Val
Ser 2375	Ile	Pro	Lys	Val	Leu	His 2380	Leu	Lys	Pro	Gly	Ser 2385	Asn	Ile	Ile
Asn 2390	Ile	Gly	Leu	Ser	Ser	Lys 2395	Asn	Ala	Val	Asn	Gly 2400	Ser	Ile	Met
Ile 2405	Thr	Phe	Lys	Ala	Arg	Gln 2410	Leu	Gly	Leu	Asn	Ile 2415	Thr	Arg	Ser
Leu 2420	Thr	Leu	Lys	Leu	Lys	Val 2425	Leu	Pro	Pro	Ala	Pro 2430	Ala	Ile	Val
Thr 2435	Ser	Pro	Asn	Phe	Leu	Asp 2440	Val	Gly	Val	Leu	Thr 2445	Asn	Glu	Thr
Ala 2450	Ser	Ala	Glu	Val	Val	Val 2455	Arg	Asn	Leu	Gly	Phe 2460	Thr	Ala	Leu
Arg 2465	Asn	Val	Ser	Ile	Arg	Ser 2470	Ser	Ile	Pro	Trp	Val 2475	Lys	Val	Val
Ser 2480	Asn	Phe	Thr	Glu	Val	Asp 2485	Pro	Lys	Asp	Asn	Glu 2490	Ser	Ile	Ser
Leu 2495	Tyr	Ile	Glu	Pro	Pro	Arg 2500	Asn	Val	Thr	Gly	Thr 2505	Phe	Lys	Gly
Glu 2510	Ile	Thr	Ile	Ser	Ser	Ser 2515	Asn	Tyr	Asn	Pro	Ile 2520	Lys	Val	Pro
Met	Arg	Ile	Arg	Val	Thr	Pro	Asn	Ala	Thr	Gly	Gly	Val	Lys	Val

-continued

2525	2530	2535
Thr Val Met Asp Pro Asn Ala	Thr Arg Leu Glu Asn Val Lys Leu	
2540	2545	2550
Thr Leu Tyr Asn Gly Tyr Phe	His Phe Glu Gly Tyr Thr Asn Lys	
2555	2560	2565
Asn Gly Thr Leu Glu Val Glu	Asn Val Pro Ile Gly Glu Tyr Lys	
2570	2575	2580
Leu Phe Ala Ser Leu Glu Gly	Tyr Tyr Gly Tyr Ser Thr Ser Ile	
2585	2590	2595
Thr Ile Glu Val Gly Val Glu	Lys Asn Val Ser Ile Ile Leu Thr	
2600	2605	2610
Pro Ser Ile Leu Glu Val Glu	Trp Glu Val Val Pro Val Thr Ile	
2615	2620	2625
Gln Asp Val Tyr Ile Ile Lys	His Glu Met Trp Tyr Ser Thr His	
2630	2635	2640
Val Pro Ala Pro Glu Ile Arg	Met Glu Gly Gly Asp Leu Glu Val	
2645	2650	2655
Tyr Val Asp Tyr Glu Lys Leu	Ala Glu Glu Gly Met Leu Glu Phe	
2660	2665	2670
Arg Gly Gln Val Ile Val Arg	Asn Thr His Gln Tyr Ile Ser Val	
2675	2680	2685
Tyr Asn Val Thr Phe Glu Ser	Gly Gly Ser His Tyr Ile Asp Val	
2690	2695	2700
Glu Phe Gly Ile Asn Arg Ile	Asp Glu Leu Lys Pro Gly Glu Ala	
2705	2710	2715
Val Ile Val Pro Tyr Val Val	Arg Ile Tyr Tyr Ser Arg Ser Pro	
2720	2725	2730
Pro Ile Asn Pro Cys Leu His	Glu Thr Lys Val Phe Lys Leu Lys	
2735	2740	2745
Ala Gly Val Val Cys Val Glu	Glu Ala Gly Lys Ile Thr Leu Lys	
2750	2755	2760
Ala Gln Arg Ile His Gln Ile	Val Val Lys Pro Thr Cys Lys Gly	
2765	2770	2775
Cys Trp Glu Ser Val Phe Pro	Val Ala Gly Lys Leu Ala Phe Met	
2780	2785	2790
Ala Ile Ala Gln Lys Val Gly	Gln Ala Leu Gly Asn Ile Asp Asp	
2795	2800	2805
Thr Gly Val Leu Ser Thr Leu	Ala Gly Glu Ala Leu Asn Asn Leu	
2810	2815	2820
Glu Ser Leu Phe Asp Ala Tyr	Asn Ala Tyr Lys Ala Asn Pro Thr	
2825	2830	2835
Lys Glu Asn Met Glu Asn Tyr	Val Lys Thr Phe Asn Ser Val Lys	
2840	2845	2850
Ala Asn Leu Ala Ser Leu Phe	Met Phe Asp Pro Val Ala Tyr Gln	
2855	2860	2865
Glu Ile Asn Ser Leu Gln Leu	Thr Leu Ile Lys Thr Pro Lys Gly	
2870	2875	2880
Asp Ile Ala Gly Phe Ala Val	Ser Arg Thr Ala Thr Pro Val Tyr	
2885	2890	2895
Ala Leu Gly Met Gly Val Gly	Lys Ile Glu Asn Gly Gln Leu Lys	
2900	2905	2910

-continued

Val	Asp	Tyr	Lys	Lys	Ala	Val	Asn	Ile	Ala	Asn	Gly	Ile	Val	Leu
2915						2920					2925			
Asn	Val	Met	Ser	Lys	Met	Gly	Gly	Ala	Leu	Gly	Gly	Ile	Ala	Ser
2930						2935					2940			
Gly	Val	Gly	Leu	Leu	Gln	Leu	Leu	Asp	Lys	Ala	Ala	Glu	Asp	Leu
2945						2950					2955			
Pro	Pro	Tyr	Ile	Ala	Gln	Leu	Phe	Leu	Asn	Cys	Ala	Ile	Cys	Leu
2960						2965					2970			
Met	Arg	Asn	Asp	Cys	Thr	Leu	Pro	Glu	Gly	Glu	Glu	Ile	Arg	Pro
2975						2980					2985			
Ile	Gln	Ile	Ile	Ala	Ser	Gly	Ser	Leu	Gly	Gly	Tyr	Pro	Ser	Met
2990						2995					3000			
Ile	Pro	Ser	Gly	Leu	Ala	Gly	Gly	Gly	Asp	Gly	Gly	Thr	Ala	Val
3005						3010					3015			
Gly	Arg	Phe	Thr	Cys	Gly	Gly	Leu	Pro	Thr	Val	Lys	Lys	Ser	Ser
3020						3025					3030			
Thr	Ser	Met	Ser	Cys	Ser	Thr	Cys	Ser	Ser	Glu	Asp	Val	Val	Lys
3035						3040					3045			
Glu	Arg	Val	Cys	Arg	Leu	Phe	Arg	Glu	Ser	Glu	Thr	His	Asp	Glu
3050						3055					3060			
Glu	Glu	Pro	Ser	Asn	Thr	Leu	His	Met	Cys	Val	Asp	Leu	Val	Leu
3065						3070					3075			
Thr	Ile	Glu	Gln	Arg	Leu	Thr	Phe	Glu	Arg	Gln	Ala	Phe	Arg	Ala
3080						3085					3090			
Ser	Leu	Lys	Phe	Thr	Asn	Thr	Asn	Arg	Asn	Tyr	Ser	Leu	Glu	Asn
3095						3100					3105			
Val	Ser	Val	Arg	Val	Ile	Phe	Phe	Asp	Glu	Glu	Gly	Asn	Arg	Val
3110						3115					3120			
Asp	Asp	Lys	Phe	Phe	Val	Arg	Leu	Asp	Glu	Lys	Ala	Gly	Leu	Ser
3125						3130					3135			
Gly	Ser	Ser	Leu	Glu	Pro	Glu	Lys	Thr	Ala	Glu	Met	Lys	Trp	Leu
3140						3145					3150			
Ile	Ile	Pro	Lys	Val	Gly	Ala	Ala	Glu	Lys	Phe	Arg	Ala	Arg	Tyr
3155						3160					3165			
Tyr	Val	Met	Ala	Asn	Ile	Thr	Ala	Arg	Val	Gly	Ser	Thr	Lys	Leu
3170						3175					3180			
Val	Tyr	Glu	Thr	Trp	Pro	Ala	Met	Ile	Glu	Val	Glu	Pro	Val	Pro
3185						3190					3195			
Gln	Leu	Val	Leu	Asp	Tyr	Val	Leu	Pro	Ser	Tyr	Val	Phe	Gly	Asp
3200						3205					3210			
Asp	Pro	Tyr	Thr	Pro	Glu	Lys	Glu	Leu	Pro	Ile	Pro	Phe	Ile	Phe
3215						3220					3225			
Gly	Val	Arg	Val	Lys	Asn	Val	Gly	Tyr	Gly	Thr	Ala	Arg	Lys	Leu
3230						3235					3240			
Arg	Ile	Ala	Ser	Ala	Gln	Pro	Lys	Ile	Glu	Arg	Ser	Asn	Tyr	Pro
3245						3250					3255			
Gly	Val	Tyr	Ile	Asp	Phe	Lys	Ile	Ile	Gly	Thr	Leu	Val	Asn	Gly
3260						3265					3270			
Lys	Lys	Val	Pro	Asn	Ser	Leu	Thr	Ile	Asp	Phe	Gly	Asp	Leu	Asn
3275						3280					3285			

-continued

Pro Gly 3290	Glu Ser Ser Thr 3295	Ala Trp Leu Met 3300	Ile Ala Glu Val 3300
Ser Gly 3305	Lys Phe Leu Gln Tyr 3310	Asn Ala Thr Phe 3315	Lys His Ser Asp 3315
Glu Leu 3320	Gly Gly Asn Glu Thr 3325	Ser Leu Ile Lys 3330	Glu Val Arg Thr 3330
His Phe 3335	Leu Ile Arg Ala Phe 3340	Asn Asn Thr Glu 3345	Asn Asp Asp Gly 3345
Met Leu 3350	Asp Phe Leu Val Asp 3355	Asp Asp Gly Asp 3360	Gly Lys Pro Glu 3360
Lys Ile 3365	Ile Asp Ser Arg Gly 3370	Phe Asp Tyr Asn 3375	Val Leu Leu Leu 3375
Asn Phe 3380	Thr Glu Val Glu Glu 3385	Gly Ser Met Arg 3390	Lys Ile Ile Pro 3390
Glu Met 3395	Lys Thr Pro Phe Trp 3400	Val Tyr Phe Thr 3405	Val Pro Phe Lys 3405
Gly Ser 3410	Val Val Arg Ser Asp 3415	Gly Lys Asn Pro 3420	Met Asp Gln Trp 3420
Met Glu 3425	Asn Gly Thr Leu His 3430	Val Leu Asp Leu 3435	Gly Thr Pro Glu 3435
Phe Tyr 3440	Ile Leu Lys Ser Asn 3445	Gln Pro Pro Ile 3450	Pro Arg Ile Tyr 3450
Val Lys 3455	Glu Pro Val Ile Ala 3460	Asn Glu Thr Val 3465	Val Leu Asp Gly 3465
Ser Leu 3470	Ser Tyr Asp Pro Asp 3475	Gly Ser Ile Ile 3480	Ala Tyr Thr Trp 3480
Lys Ile 3485	Gly Asn Glu Ser Phe 3490	Val Gly Asp Lys 3495	Val Ser Tyr Val 3495
Phe Arg 3500	Glu Pro Gly Thr Tyr 3505	Asn Val Thr Leu 3510	Thr Val Arg Asp 3510
Asp Lys 3515	Gly Thr Glu Ser Ser 3520	Lys Thr Met Glu 3525	Ile Lys Val Tyr 3525
Leu Gly 3530	Pro Lys Phe Asn Glu 3535	Ser Leu Lys Val 3540	Glu Pro Gln Trp 3540
Gly Ile 3545	Val Pro Phe Asn Leu 3550	Ser Ile Thr Phe 3555	Asn Val Thr Asn 3555
Val Gly 3560	Asp Val Ser Gly Glu 3565	Tyr Ser Tyr Ile 3570	Ile Lys Leu Gly 3570
Asn Ser 3575	Thr Ile Ala Glu Gly 3580	Ser Glu Ile Ile 3585	Glu Ser Gly Arg 3585
Trp Lys 3590	Val Ile Asn Ser Thr 3595	Val Glu Ile Arg 3600	Lys Glu Gly Asn 3600
Tyr Thr 3605	Val Thr Ala Asn Asn 3610	Leu Ser Lys Thr 3615	Val Thr Ala Tyr 3615
Arg Lys 3620	Val Tyr Gly Asn Leu 3625	Thr Glu Asn Tyr 3630	Ile Lys Glu Lys 3630
Asp Phe 3635	Gly His Tyr Lys Ser 3640	Phe Tyr Trp Asn 3645	Glu Phe Lys Arg 3645
Asp Phe 3650	Glu Gly Trp Val Glu 3655	Glu Ala Leu Ser 3660	Thr Ile Glu Leu 3660
Pro Lys 3665	Val Asn Phe Lys Val 3670	Leu Asn Tyr Phe 3675	Pro Gly Asn Trp 3675

-continued

3665	3670	3675
Ser Leu Leu Asn Tyr Ser Glu Met Leu Asn Ile Thr Lys Gly Trp 3680 3685 3690		
Gly Trp Ile Asn Ala Thr Tyr Ala Arg Arg Val Arg Val Glu Gly 3695 3700 3705		
Leu Glu Glu Phe Lys Tyr Leu Ile Val Asn Val Thr Gln Leu Val 3710 3715 3720		
Val Leu Leu Gly Asn Ala Thr His Glu Leu Asp Glu Ser Pro Pro 3725 3730 3735		
Thr Leu Asn Val Thr Pro Ser Ser Gly Ile Tyr Ser Glu Ile Pro 3740 3745 3750		
Lys Ile Gln Val Arg Thr Cys Asp Glu Thr Gly Ile Thr Leu Val 3755 3760 3765		
Trp Gly Ala Val Gly Asn Tyr Thr Lys Glu Phe Thr Glu Val Glu 3770 3775 3780		
Ser Asn Gly Thr Cys Ser Thr Trp Glu Gly Ile Val Pro Leu Asn 3785 3790 3795		
Ile Gly Asn Asn Thr Val Ala Ile Tyr Ala Glu Asp Glu Phe Gly 3800 3805 3810		
Asn Arg Gly Asn Val Ser Leu Trp Ile Tyr Leu Asn Pro Glu Ala 3815 3820 3825		
Pro Val Ile Tyr Ile Glu Ser Pro Glu Glu Lys Val Tyr Asn Ser 3830 3835 3840		
Arg Glu Val Met Ile Asn Tyr Thr Val Val Asn His Asp Leu Val 3845 3850 3855		
Gly Val Val Ala Tyr Leu Asn Gly Glu Leu Ile Ser Ser Asn Ala 3860 3865 3870		
Ser Tyr Ser Gly Phe Ile Lys Leu Asp Tyr Gly Trp His Asn Phe 3875 3880 3885		
Thr Ile Tyr Ala Trp Asp Val Ser Tyr Asn Val Ser Lys Ser Val 3890 3895 3900		
Ile Phe Arg Val Asn Glu Pro Pro Ser Val Asp Phe Ser Trp Glu 3905 3910 3915		
Val Asp Asn Leu Thr Val Lys Phe Glu Ala Asn Ala Ser Asp Glu 3920 3925 3930		
Asp Gly Ile Ser Lys Tyr Leu Trp Asp Phe Gly Asp Asn Glu Ser 3935 3940 3945		
Ser Leu Leu Val Asn Pro Thr His Thr Tyr Arg Lys Gly Gly Arg 3950 3955 3960		
Tyr Asn Val Thr Leu Thr Val Trp Asp Ser Tyr Asn Leu Ser Ser 3965 3970 3975		
Ser Ile Ser Lys Glu Val Val Val Phe Gly Ser Ser Thr Leu Thr 3980 3985 3990		
Met Val Lys Glu Tyr Ser Tyr Thr Lys Asp Phe Gly Phe Tyr Asn 3995 4000 4005		
Thr Thr Ser Trp Lys Asp Phe Leu Lys Asp Phe Glu Val Trp Val 4010 4015 4020		
Asn Leu Thr Leu Arg Asn Val Thr Leu Pro Leu Glu Tyr Phe Glu 4025 4030 4035		
Glu Ile Ile Glu Val Asn Val Glu Asn Trp Ser Leu Ile Ser Val 4040 4045 4050		

-continued

Glu	Lys	Asn	Leu	Lys	Asn	Asp	Ile	Gly	Glu	Met	Ser	Ala	Glu	Tyr
4055						4060					4065			
Glu	Arg	Asn	Ala	Thr	Ile	Val	Gly	Ile	Met	Asn	Tyr	Thr	Arg	Val
4070						4075					4080			
Thr	Leu	Lys	Leu	Thr	Gln	Glu	Val	Ile	Leu	Ser	Gly	Arg	Ala	Arg
4085						4090					4095			
Lys	Val	Glu	Asp	Lys	Ile	Pro	Pro	Leu	Val	Glu	Ile	Leu	Phe	Pro
4100						4105					4110			
Arg	Asn	Met	Thr	Tyr	Asn	Glu	Thr	Ile	Arg	Glu	Ile	Lys	Val	Arg
4115						4120					4125			
Ala	Thr	Asp	Glu	Ser	Gly	Ile	Ala	Asn	Val	Thr	Ala	Thr	Ile	Asn
4130						4135					4140			
Gly	Glu	Ser	Leu	Ser	Leu	Glu	Lys	Val	Asn	Glu	Thr	Trp	Ile	Gly
4145						4150					4155			
Arg	Val	Glu	Leu	Asp	Asp	Gly	Lys	Tyr	Glu	Leu	Asn	Val	Phe	Ala
4160						4165					4170			
Ser	Asp	Lys	Trp	Gly	Asn	Val	Gly	Cys	Ser	Thr	Val	Asn	Phe	Thr
4175						4180					4185			
Ile	Asn	Arg	Ser	Val	Lys	Val	Arg	Ile	Ile	Asn	Gly	Thr	Glu	Ile
4190						4195					4200			
Val	Thr	Ile	Pro	Gly	Asp	Ile	Lys	Thr	Arg	Val	Tyr	Phe	Glu	Gly
4205						4210					4215			
Asp	Ile	Ile	Val	Glu	Ile	Val	Lys	Glu	Ser	Leu	Arg	Phe	Lys	Ile
4220						4225					4230			
Pro	Ser	Gly	Gly	Thr	Leu	Val	Ile	Asp	Glu	Arg	Gly	Arg	Lys	Asp
4235						4240					4245			
Pro	Trp	Leu	Leu	Ala	Arg	Ile	Asn	Ser	Thr	Ile	Glu	Asn	Ile	Ser
4250						4255					4260			
Lys	Thr	Ser	Arg	Ile	Phe	Glu	Glu	Asn	Gly	Lys	Lys	Val	His	Glu
4265						4270					4275			
Ile	Arg	Tyr	Arg	Ile	Ser	Ile	Ser	Arg	Gly	Tyr	Ala	Ile	Leu	Val
4280						4285					4290			
Val	Pro	Leu	Glu	Gly	Met	Lys	Val	Ser	Ser	Ile	Arg	Ile	Ile	Lys
4295						4300					4305			
Asn	Gly	Thr	Val	Thr	Arg	Asp	Glu	Lys	His	Gly	Asn	Tyr	Tyr	Lys
4310						4315					4320			
Leu	Ser	Lys	Gly	Tyr	Leu	Phe	Ile	Phe	Leu	Ser	Glu	Asp	Pro	Ile
4325						4330					4335			
Val	Glu	Val	Thr	Leu	Ser	Lys	Ile	Glu	Lys	Lys	Asp	Ile	Phe	Arg
4340						4345					4350			
Val	Leu	Tyr	Tyr	Ala	Gly	Ile	Ile	Trp	Glu	Arg	Asn	Tyr	Leu	Arg
4355						4360					4365			
Leu	Lys	Glu	Glu	Phe	Ile	Met	Lys	Met	Ser	Asn	Glu	Thr	Ser	Gln
4370						4375					4380			
Glu	Ala	Ile	Arg	Leu	His	Glu	Glu	Ala	Glu	Lys	Tyr	Tyr	Leu	Lys
4385						4390					4395			
Gly	Arg	Glu	Tyr	Tyr	Pro	Arg	Ile	Pro	Ser	Pro	Ser	Ala	Ile	Tyr
4400						4405					4410			
Trp	Tyr	Ala	Val	Tyr	Met	Arg	Lys	Ala	Tyr	Leu	Thr	Glu	Arg	Lys
4415						4420					4425			

-continued

Ala	Leu	Glu	Leu	Leu	Ser	Ile	Ser
4430						4435	

1. A polypeptide comprising a variant fibronectin type III (FN3) domain from *Sulfolobus tokodaii* comprising one or more amino acid substitutions or insertions in a loop region of FN3, in a non-loop region of FN3, or in both.

2. The polypeptide of claim 1, wherein the non-variant FN3 domain comprises at least 70% identity to SEQ ID NO:1.

3. The polypeptide of claim 2, wherein the non-variant FN3 domain comprises SEQ ID NO:1.

4. The polypeptide of claim 1, wherein the non-variant FN3 domain comprises a polypeptide with at least 70% identity to SEQ ID NO:2 (STOFN3-2), 3 (STOFN3-3), or 4 (STOFN3-4).

5. The polypeptide of claim 1, wherein the non-variant FN3 domain comprises SEQ ID NO:2, 3, or 4.

6. The polypeptide of any one of claims 1-4, wherein the amino acid substitution comprises one or more substitutions of the FN3 domain corresponding to amino acid positions 1, 2, and 3 of SEQ ID NO:1.

7. The polypeptide of claim 6, wherein the amino acid substitutions at one or more positions corresponding to amino acid positions 1, 2, and/or 3 of SEQ ID NO:1 are a substitution with a hydrophilic amino acid or combinations thereof.

8. The polypeptide of any one of claims 1-7, wherein the amino acid insertion comprises one or more amino acid insertions of the FN3 domain after the amino acid corresponding to amino acid position 84 of SEQ ID NO:1.

9. The polypeptide of claim 8, wherein the amino acid insertion comprises 1-3 amino acids.

10. The polypeptide of claim 9, wherein the inserted amino acids are proline and serine or combinations thereof.

11. The polypeptide of any one of claims 1-10, wherein the substitution comprises one or more substitutions of the FN3 domain corresponding to amino acid positions 4, 14, 28, and 58 of SEQ ID NO:1.

12. The polypeptide of claim 11, wherein the substitution corresponding to amino acid position 4 of SEQ ID NO:1 is with a valine.

13. The polypeptide of claim 11 or 12, wherein the substitution corresponding to amino acid position 14 of SEQ ID NO:1 is with a proline.

14. The polypeptide of any one of claims 11-13, wherein the substitution corresponding to amino acid position 28 of SEQ ID NO:1 is with a tyrosine.

15. The polypeptide of any one of claims 11-14, wherein the substitution corresponding to amino acid position 58 of SEQ ID NO:1 is with a proline.

16. The polypeptide of any one of claims 1-15, wherein the one or more amino acid substitutions or insertions in the non-loop segment is in one or more of beta strand C or beta strand D.

17. The polypeptide of claim 16, wherein the one or more amino acid substitutions or insertions correspond to or after positions 28, 29, 30, 31, 32, 33, 34, or 35 of SEQ ID NO:1.

18. The polypeptide of claim 16, wherein the one or more amino acid substitutions or insertions correspond to or after positions 38, 39, 40, 41, 42, 43, 44, 45, or 46 of SEQ ID NO:1.

19. The polypeptide of any one of claims 1-18, wherein the one or more amino acid substitutions or insertions in the loop region of FN3 are in the BC loop, DE loop, FG loop and/or CD loop.

20. The polypeptide of any one of claims 1-19, wherein the DE loop does not comprise any inserted amino acids.

21. The polypeptide of any one of claims 1-20, wherein the EF loop comprises less than 3 substitutions or insertions.

22. The polypeptide of any one of claims 1-21, wherein the FN3 domain does not have substitutions at amino acid positions corresponding to positions 69 and 77 of SEQ ID NO:1.

23. The polypeptide of any one of claims 1-22, further comprising a non-FN3 polypeptide that enhances the FN3 polypeptide binding affinity for a target molecule.

24. The polypeptide of any one of claims 1-23, comprising or further comprising an insertion or deletion of at least 1 amino acids in at least one loop region of FN3.

25. The polypeptide of claim 24, wherein at least one loop region of FN3 comprises an insertion of at least 2 amino acids.

26. The polypeptide of claim 25, wherein at least one region of FN3 comprises an insertion of 2 to 25 amino acids in at least one loop region.

27. The polypeptide of claim 24, wherein at least two loop regions comprise an insertion.

28. The polypeptide of claim 24, wherein at least one loop region of FN3 comprises a deletion of at least 1 amino acid.

29. The polypeptide of claim 28, wherein at least one loop region of FN3 comprises a deletion of 2 to 10 amino acids.

30. The polypeptide of claim 28, wherein at least two loop regions comprise a deletion of at least 1 amino acid.

31. The polypeptide of claim 24, wherein the polypeptide comprises at least 1 amino acid insertion and 1 amino acid deletion in at least one loop region.

32. The polypeptide of claim 31, wherein the polypeptide comprises an insertion and deletion of at least 1 amino acid in the same loop region.

33. The polypeptide of any one of claims 1-32, wherein the polypeptide further comprises one or more serine residues immediately before amino acid 1 or immediately after amino acid 84 of SEQ ID NO:1.

34. A polypeptide comprising a variant fibronectin type III (FN3) domain from *Pyrococcus horikoshii* comprising one or more amino acid substitutions or insertions in a loop region of FN3, in a non-loop region of FN3, or in both.

35. The polypeptide of claim 34, wherein the non-variant FN3 domain comprises at least 70% identity to SEQ ID NO:6

36. The polypeptide of claim 35, wherein the non-variant FN3 domain comprises SEQ ID NO:6.

37. The polypeptide of any one of claims **34-36**, wherein the one or more amino acid substitutions or insertions in the non-loop segment is in one or more of beta strand C or beta strand D.

38. The polypeptide of claim **37**, wherein the one or more amino acid substitutions or insertions correspond to or after positions 13 and/or 14 of SEQ ID NO:6.

39. The polypeptide of claim **37** or **38**, wherein the one or more amino acid substitutions or insertions correspond to or after positions 20, 21, 22, 23, 24, 25, 26, and/or 27 of SEQ ID NO:6.

40. The polypeptide of any one of claims **37-39**, wherein the one or more amino acid substitutions or insertions correspond to or after positions 36 and/or 37 of SEQ ID NO:6.

41. The polypeptide of any one of claims **37-40**, wherein the one or more amino acid substitutions or insertions correspond to or after positions 40, 41, 42, 43, 44, 45, 46, 47, and/or 48 of SEQ ID NO:6.

42. The polypeptide of any one of claims **37-41**, wherein the one or more amino acid substitutions or insertions correspond to or after positions 54, 55, 56, 57, 58, and/or 59 of SEQ ID NO:6.

43. The polypeptide of any one of claims **37-42**, wherein the one or more amino acid substitutions or insertions correspond to or after positions 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, and/or 74 of SEQ ID NO:6.

44. The polypeptide of any one of claims **34-43**, wherein the one or more amino acid substitutions or insertions in the loop region of FN3 are in the AB loop, BC loop, CD loop, DE loop, EF loop, and/or FG loop.

45. The polypeptide of any one of claims **34-44**, further comprising a non-FN3 polypeptide that enhances the FN3 polypeptide binding affinity for a target molecule.

46. The polypeptide of any one of claims **34-45**, comprising or further comprising an insertion or deletion of at least 1 amino acids in at least one loop region of FN3.

47. The polypeptide of claim **46**, wherein at least one loop region of FN3 comprises an insertion of at least 2 amino acids.

48. The polypeptide of claim **47**, wherein at least one region of FN3 comprises an insertion of 2 to 25 amino acids in at least one loop region.

49. The polypeptide of claim **46**, wherein at least two loop regions comprise an insertion.

50. The polypeptide of claim **46**, wherein at least one loop region of FN3 comprises a deletion of at least 1 amino acid.

51. The polypeptide of claim **50**, wherein at least one loop region of FN3 comprises a deletion of 2 to 10 amino acids.

52. The polypeptide of claim **50**, wherein at least two loop regions comprise a deletion of at least 1 amino acid.

53. The polypeptide of claim **46**, wherein the polypeptide comprises at least 1 amino acid insertion and 1 amino acid deletion in at least one loop region.

54. The polypeptide of claim **53**, wherein the polypeptide comprises an insertion and deletion of at least 1 amino acid in the same loop region.

55. The polypeptide of any one of claims **1-54**, wherein the polypeptide is isolated.

56. The polypeptide of any one of claims **1-55**, wherein the polypeptide is recombinant.

57. The polypeptide of any one of claims **1-56**, wherein the polypeptide is synthetic.

58. The polypeptide of any one of claims **1-57**, wherein the polypeptide binds specifically to a target motif.

59. The polypeptide of claim **58**, wherein the polypeptide further comprises a polypeptide comprising a biorecognition module including a molecular recognition domain; wherein the polypeptide comprising the biorecognition molecule and the polypeptide comprising the variant FN3 domain are operatively linked together either directly or indirectly via the linker, and are spatially oriented to bind on the same, overlapping, or distinct sites within the target motif.

60. The polypeptide of claim **59**, wherein the K_d for the polypeptide being equal to or lower than one μM .

61. The polypeptide of claim **59** or **60**, wherein the target motif is a peptide, a phosphorylated peptide or a methylated peptide.

62. The polypeptide of any of claims **59-61**, wherein the target motif is present within a protein.

63. The polypeptide of any of claims **59-62**, wherein the molecular recognition domain comprises an interaction domain or mutants of interaction domains.

64. The polypeptide of claim **63**, wherein the interaction domain is selected from the group consisting of PDZ, WW, SH2, PTB, SH3, Bromo, Chromo, PHD, Polo-box and FHA domains.

65. The polypeptide of any of claims **59-64**, wherein the polypeptide comprising the variant FN3 domain further comprises a first signaling moiety and the polypeptide comprising the biorecognition molecule further comprises a second signaling moiety and wherein the first and second signaling moieties are capable of interacting to produce a detectable signal.

66. The polypeptide of claim **65**, wherein the signaling moiety is a dye, a quencher, a reporter protein, or a quantum dot.

67. The polypeptide of claim **65**, wherein the first and second signaling moieties comprise a fluorescent resonance energy transfer (FRET) donor group and a FRET acceptor group, respectively, and binding of the first and second molecular recognition domains to the target motif results in a change in the FRET efficiency between the FRET donor and FRET acceptor groups.

68. A polypeptide library comprising a plurality of modified FN3 domain polypeptides comprising one or more amino acid substitutions, insertions, or deletions a loop region of FN3, in a non-loop region of FN3, or in both; wherein the unmodified FN3 domain comprises a polypeptide comprising the amino acid sequence of SEQ ID NOS: 1-4 or 6.

69. The polypeptide library of claim **68**, wherein the FN3 domain polypeptides comprises one or more amino acid substitutions corresponding to amino acid positions corresponding to positions 1, 2, 3, 4, 14, 28, and/or 58 of SEQ ID NO:1.

70. The polypeptide library of claim **68** or **69**, wherein the FN3 domain polypeptides further comprise one or more amino acid substitutions corresponding to amino acid positions 13, 14, 15, 16, 22, 23, 24, 25, 26, 27, 36, 37, 47, 48, 49, 54, 55, 56, 57, 58, 59, 60, 71, 72, 73, 74, 75, 76, 77, 78, 79 and/or 80 of SEQ ID NO:1.

71. The polypeptide library of claim **68**, wherein the FN3 domain polypeptides further comprise one or more amino acid substitutions corresponding to amino acid positions of SEQ ID NO:6.

72. The polypeptide library of any one of claims **68-71**, wherein the FN3 domain polypeptides are at least 50%, 60%, 70%, 80%, or 90% identical to SEQ ID NO:1, 2, 3, 4, or 6.

73. The polypeptide library of any one of claims **68-72**, further comprising a non-FN3 polypeptide that enhances the FN3 polypeptide binding affinity for a target molecule.

74. The polypeptide library of any one of claims **68-73**, comprising or further comprising an insertion or deletion of at least 1 amino acids in at least one loop region of the FN3 domain polypeptides.

75. The polypeptide library of claim **74**, wherein at least one loop region of FN3 comprises an insertion of at least 2 amino acids.

76. The polypeptide library of claim **75**, wherein at least one region of FN3 comprises an insertion of 2 to 25 amino acids in at least one loop region.

77. The polypeptide library of any one of claims **68-76**, wherein at least two loop regions comprise an insertion.

78. The polypeptide library of any one of claims **68-77**, wherein at least one loop region of FN3 comprises a deletion of at least 1 amino acid.

79. The polypeptide library of any one of claims **68-78**, wherein at least one loop region of FN3 comprises a deletion of 2 to 10 amino acids.

80. The polypeptide library of any one of claims **68-79**, wherein at least two loop regions comprise a deletion of at least 1 amino acid.

81. The polypeptide library of claim **74**, wherein the polypeptide comprises at least 1 amino acid insertion and 1 amino acid deletion in at least one loop region.

82. The polypeptide library of any one of claims **68-81**, wherein the FN3 domain polypeptides comprise an insertion and deletion of at least 1 amino acid in the same loop region.

83. The polypeptide library of any one of claims **68-82**, wherein the library is pre-selected to bind a target.

84. A polynucleotide encoding the polypeptide of claims **1-67**.

85. The polynucleotide of claim **84**, wherein the polynucleotide is an expression construct.

86. The polynucleotide of claim **85**, wherein the expression construct is capable of expressing the encoded polypeptide in a bacteria.

87. A polynucleotide composition encoding the polypeptide library of any one of claims **68-83**.

88. A monobody library comprising a plurality of polypeptides having a variant fibronectin type III (FN3) domain, compared to wildtype (SEQ ID NOS:1-4 or 6), comprising one or more alterations in a beta strand or a loop region.

89. The monobody of claim **88**, wherein the alterations are in one or more of beta strand C, beta strand D, BC loop, CD loop, DE loop, or and/or FG loop.

90. The monobody of claim **88** or **89**, wherein the monobody comprises one or more of the variant amino acids corresponding to position 1, 2, 3, 4, 14, 28, and/or 58 of SEQ ID NO:1.

91. The monobody library of claim **88** or **90**, wherein the variant FN3 domains further comprise an insertion, substitution, or deletion of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 amino acids in at least one loop region of FN3.

92. A method for selecting a protein binding domain specific for a target comprising:

(a) detecting target specific binding of one or more members of a polypeptide library of any one of claims **68-83**; and

(b) selecting the protein binding domain that specifically binds the target.

93. The method of claim **92**, further comprising first preparing the polypeptide library.

94. The method of claim **92**, further comprising isolating the selected protein binding domain that specifically binds the target.

95. The method of claim **94**, further comprising determining the amino acid of sequence of the selected protein binding domain.

96. The method of claim **95**, further comprising generating the selected protein binding domain after the amino acid sequence has been determined.

97. A method of identifying a polypeptide that specifically binds a target comprising detecting specific binding of one or more polypeptides of a polypeptide library of any one of claims **68-83**.

98. A method of detecting a target molecule comprising contacting a sample containing the target with a fibronectin type III (FN3) binding domain that specifically binds the target.

99. A method of producing a fibronectin type III (FN3) variant comprising:

(a) expressing a polypeptide comprising a amino acid sequence selected in claim **97**; and

(b) isolating and/or purifying the expressed variant FN3 domain from a host cell expressing the variant FN3.

100. A kit comprising a plurality of polypeptides as described in any one of claims **1-67**.

101. A kit comprising a polypeptide library as described in any one of claims **68-83**.

102. A kit comprising a plurality of polynucleotides encoding FN3 domain variants described in claims **1-57**. non-variant FN3 domain comprises SEQ ID NO:1.

103. A method of making a polypeptide or polypeptide library according to any one of claims **1-83** comprising expressing the polynucleotide of any of claims **84-87** in a host cell.

104. A method of testing a polypeptide according to any one of claims **1-67**, the method comprising contacting the polypeptide with a target molecule and testing for binding activity between the polypeptide and the target molecule.

* * * * *