

advances.sciencemag.org/cgi/content/full/6/50/eabc9959/DC1

Supplementary Materials for

Titanium isotopes constrain a magmatic transition at the Hadean-Archean boundary in the Acasta Gneiss Complex

Sarah M. Aarons*, Jesse R. Reimink, Nicolas D. Greber, Andy W. Heard, Zhe Zhang, Nicolas Dauphas

*Corresponding author. Email: smaarons@ucsd.edu

Published 9 December 2020, *Sci. Adv.* **6**, eabc9959 (2020)
DOI: [10.1126/sciadv.abc9959](https://doi.org/10.1126/sciadv.abc9959)

The PDF file includes:

Supplementary Text
Figs. S1 to S3
Legends for datasets S1 and S2
References

Other Supplementary Material for this manuscript includes the following:

(available at advances.sciencemag.org/cgi/content/full/6/50/eabc9959/DC1)

Datasets S1 and S2

Supplementary Materials

Iron isotopic fractionation

Interpretation of Fe isotope variations in igneous rocks is complicated by the fact that two Fe oxidation states often coexist in magmas. The decrease in $\delta^{56}\text{Fe}$ values with increasing SiO₂ measured in the 4.02 Ga ITG is reminiscent of what was observed by Sossi et al. (69) in the tholeiitic differentiation series found in the Red Hill intrusion, Tasmania (Fig. S1). We interpret the Fe isotopic evolution of the ITG samples to possibly represent the post-magnetite crystallization limb of the magmatic differentiation trend analogous to that documented in the Red Hill suite, however small variations within measurement error make this interpretation difficult.

Following the interpretative framework of Sossi et al. (69) and Dauphas et al. (70), the heavy Fe isotope enrichments with increasing SiO₂ documented in the <3.9 Ga AGC TTGs could reflect crystallization in a system that was open to exchange with the surrounding medium such that buffering of oxygen fugacity could have mitigated the increase in Fe³⁺/Fe_{tot} expected for closed-system crystallization. In that context, the oxides that crystallized could have been characterized by lower Fe³⁺/Fe_{tot} than those that crystallized in the 4.02 Ga ITG. The study of Roskosz et al. (71) on spinels shows that the redox state of Fe in oxides has a strong influence on the strength of Fe bonds and Fe isotopic fractionation. It is thus conceivable that crystallization of low-Fe³⁺ oxides, in concert with the crystallization of other Fe²⁺-bearing phases, drove the residual magma towards heavy isotope enrichments.

Figure S1b illustrates how the ITG and AGC samples may be defined by different trends in $\delta^{56}\text{Fe}$ vs. $\delta^{49}\text{Ti}$, however small $\delta^{56}\text{Fe}$ variations make this interpretation ambiguous. As discussed above, these trends could reflect the different paths of Fe³⁺/Fe²⁺ evolution experienced by the parental magmas during magmatic differentiation. Data is missing at the present time to

quantitatively model those trends but the systematics revealed by the present study illustrate the potential of combining Fe and Ti isotopic analyses to tease out the oxide crystallization history of igneous rocks.

Age and geochemical history of Acasta Gneiss Complex

The AGC has been extensively geochemically characterized (21), and dated using U-Pb isotope methods, with ages ranging from 4.03-3.4 Ga (27, 72-74). Reimink et al. (22) argued that the AGC saw a transition from shallow level melting of hydrated mafic crust at 3.96-3.75 Ga to greater melting depths at ~3.6 Ga (23). The ~3.6 Ga granitoids were believed to be the result of thrusting beneath a Hadean mafic plateau and subsequent melting at depths >50 km (26). These constraints on the formation depth are difficult to reconcile without invoking a horizontal tectonic regime (22).

Recently Reimink et al. (22) utilized whole-rock $^{142}\text{Nd}/^{144}\text{Nd}$ ($\mu^{142}\text{Nd}$) ratios to infer that the 4.02 Ga crust was extracted from a bulk-Earth source whereas the Eoarchean rocks from the AGC were interpreted to be the result of partial melting of a hydrated, Hadean age mafic crust at relatively shallow depths. By 3.6 Ga, granitic-rocks formed from partial melting of hydrated Archean mafic crust at increased depths within the garnet stability field. The $\mu^{142}\text{Nd}$ signatures (22, 75) and zircon Hf isotope work (24, 25, 76-78) ruled out the hypothesis of gradual thickening and melting of an ancient mafic plateau previously hypothesized in Reimink et al. (23), and supports the idea that the felsic crust observed here at 3.75 Ga was produced via plate subduction rather than melting of hydrated basaltic material at the base of a thickened crust (79).

Based on geochemical data and U-Pb geochronology, the AGC is identified to have undergone six major episodes of magmatism at 4.02, 3.96, 3.75, 3.55-3.6, 3.37, and 2.94 Ga (26). The oldest evolved silicic rocks are the Idì whaà tonalitic gneiss (4.02 Ga), which formed by shallow fractional crystallization of a basaltic magma. Hafnium isotope data from the 4.02 samples in the AGC show no indication of derivation or interaction with an older Hadean continental crust, and instead suggest that the oldest evolved crust from the AGC was generated from an older ultramafic or mafic reservoir that was present on the surface of the early Earth (10).

The later granitoids (3.96-2.94 Ga) are thought to have formed *via* partial melting of mafic rocks that underwent alteration from surface waters at low temperatures (10). The dehydration melting process is thought to have occurred at progressively deeper levels reaching the stability field of garnet at ~3.6 Ga at a depth of at least 50 km, and may have occurred in a subduction-like setting with tectonic under-thrusting or by burial or gravitational sinking of a thick oceanic plateau comprised of hydrated mafic rocks to depths of >50 km (10).




Fig. S1. $\delta^{56}\text{Fe}$ compositions as a function of SiO_2 content and the relationship of $\delta^{49}\text{Ti}$ with respect to $\delta^{56}\text{Fe}$. Samples from the AGC are plotted as gray symbols in **a** and **b**. **a)** $\delta^{56}\text{Fe}$ compositions vs. SiO_2 of rocks from Idi whaa tonalite gneiss (this study) and Red Hill, Tasmania (69) and I-, S- and A-type granites, pegmatites, and migmatites (80). Plume-derived rocks show a rapid depletion in $\delta^{56}\text{Fe}$ composition, whereas rocks from convergent settings slightly increase in $\delta^{56}\text{Fe}$ composition. **b)** $\delta^{56}\text{Fe}$ and $\delta^{49}\text{Ti}$ isotopic compositions of rocks from AGC.




Fig. S2. Calculated TIMI for plutonic rocks and igneous rocks. Calc-alkaline rocks (blue circles) from Agung and Kos (29, 36), tholeiitic rocks (red circles) from Iceland, Afar and Hawaii (30, 31), A-type (this study) and I-and S-type granites (36).




Fig. S3. Map of Acasta Gneiss Complex region modified from (26). a) Map of Slave Province with location of AGC noted. b) A close up view of the ~1300 km² exposure that encompasses the AGC. c) Simplified geologic map of the area highlighted in b).

Dataset S1 (separate file). Titanium isotope compositions and sample information for rocks analyzed in this study and plotted in Figs. 1, 2, and 3 in main text.

Dataset S2 (separate file). Iron isotope compositions and sample information for rocks analyzed in this study and plotted in Fig. S1.

REFERENCES AND NOTES

1. F. J. Vine, D. H. Matthews, Magnetic anomalies over oceanic ridges. *Nature* **199**, 947–949 (1963).
2. A. Miyashiro, Volcanic rock series in island arcs and active continental margins. *Am. J. Sci.* **274**, 321–355 (1974).
3. M. M. Zimmer, T. Plank, E. H. Hauri, G. M. Yogodzinski, P. Stelling, J. Larsen, B. Singer, B. Jicha, C. Mandeville, C. J. Nye, The role of water in generating the calc-alkaline trend: New volatile data for Aleutian magmas and a new tholeiitic index. *J. Petrol.* **51**, 2411–2444 (2010).
4. S. R. Taylor, in *Island Arcs, Deep Sea Trenches and Back-Arc Basins* (AGU, 1977), pp. 325–335.
5. M. Brown, Duality of thermal regimes is the distinctive characteristic of plate tectonics since the Neoarchean. *Geology* **34**, 961–964 (2006).
6. P. A. Cawood, A. Kröner, S. Pisarevsky, Precambrian plate tectonics: Criteria and evidence. *Geol. Soc. Am. Today* **16**, 4–11 (2006).
7. W. B. Hamilton, Plate tectonics began in Neoproterozoic time, and plumes from deep mantle have never operated. *Lithos* **123**, 1–20 (2011).
8. M. Hopkins, T. M. Harrison, C. E. Manning, Low heat flow inferred from >4 Gyr zircons suggests Hadean plate boundary interactions. *Nature* **456**, 493–496 (2008).
9. T. Komiya, S. Maruyama, T. Masuda, S. Nohda, M. Hayashi, K. Okamoto, Plate tectonics at 3.8–3.7 Ga: Field evidence from the Isua accretionary complex, southern West Greenland. *J. Geol.* **107**, 515–554 (1999).
10. A. P. Nutman, C. R. L. Friend, V. C. Bennett, Evidence for 3650–3600 Ma assembly of the northern end of the Itsaq Gneiss Complex, Greenland: Implication for early Archean tectonics. *Tectonics* **21**, 1005 (2002).

11. S. B. Shirey, B. S. Kamber, M. J. Whitehouse, P. A. Mueller, A. R. Basu, in *When Did Plate Tectonics Begin on Planet Earth?*, K. C. Condie, V. Pease, Eds. (Geological Society of America, 2008), pp. 1–29.
12. R. J. Stern, Evidence from ophiolites, blueschists, and ultrahigh-pressure metamorphic terranes that the modern episode of subduction tectonics began in Neoproterozoic time. *Geology* **33**, 557–560 (2005).
13. M. J. Van Kranendonk, R. H. Smithies, A. H. Hickman, D. C. Champion, Review: Secular tectonic evolution of Archean continental crust: Interplay between horizontal and vertical processes in the formation of the Pilbara Craton, Australia. *Terra Nova* **19**, 1–38 (2007).
14. H. Kuno, in *Basalts: The Poldervaart Treatise on Rocks of Basaltic Composition*, H. H. Hess, A. Poldervaart, Eds. (Interscience Publishers, 1968), pp. 623–688.
15. T. L. Grove, M. B. Baker, Phase equilibrium controls on the tholeiitic versus calc-alkaline differentiation trends. *J. Geophys. Res. Solid Earth* **89**, 3253–3274 (1984).
16. T. W. Sisson, T. L. Grove, Experimental investigations of the role of H₂O in calc-alkaline differentiation and subduction zone magmatism. *Contrib. Mineral. Petrol.* **113**, 143–166 (1993).
17. K. A. Kelley, E. Cottrell, Water and the oxidation state of subduction zone magmas. *Science* **325**, 605–607 (2009).
18. I. J. Parkinson, R. J. Arculus, The redox state of subduction zones: Insights from arc-peridotites. *Chem. Geol.* **160**, 409–423 (1999).
19. T. L. Grove, R. J. Kinzler, Petrogenesis of andesites. *Annu. Rev. Earth Planet. Sci.* **14**, 417–454 (1986).
20. R. W. Carlson, M. Garçon, J. O’Neil, J. Reimink, H. Rizo, The nature of Earth’s first crust. *Chem. Geol.* **530**, 119321 (2019).

21. J. R. Reimink, T. Chacko, R. A. Stern, L. M. Heaman, Earth's earliest evolved crust generated in an Iceland-like setting. *Nat. Geosci.* **7**, 529–533 (2014).
22. J. R. Reimink, T. Chacko, R. W. Carlson, S. B. Shirey, J. Liu, R. A. Stern, A. M. Bauer, D. G. Pearson, L. M. Heaman, Petrogenesis and tectonics of the Acasta Gneiss Complex derived from integrated petrology and ^{142}Nd and ^{182}W extinct nuclide-geochemistry. *Earth Planet. Sci. Lett.* **494**, 12–22 (2018).
23. J. R. Reimink, J. H. F. L. Davies, T. Chacko, R. A. Stern, L. M. Heaman, C. Sarkar, U. Schaltegger, R. A. Creaser, D. G. Pearson, No evidence for Hadean continental crust within Earth's oldest evolved rock unit. *Nat. Geosci.* **9**, 777–780 (2016).
24. Y. Amelin, D.-C. Lee, A. N. Halliday, Early-middle archaean crustal evolution deduced from Lu-Hf and U-Pb isotopic studies of single zircon grains. *Geochim. Cosmochim. Acta* **64**, 4205–4225 (2000).
25. T. Iizuka, T. Komiya, S. P. Johnson, Y. Kon, S. Maruyama, T. Hirata, Reworking of Hadean crust in the Acasta gneisses, northwestern Canada: Evidence from in-situ Lu-Hf isotope analysis of zircon. *Chem. Geol.* **259**, 230–239 (2009).
26. J. R. Reimink, T. Chacko, R. A. Stern, L. M. Heaman, The birth of a cratonic nucleus: Lithogeochemical evolution of the 4.02–2.94 Ga Acasta Gneiss Complex. *Precambrian Res.* **281**, 453–472 (2016).
27. A. M. Bauer, C. M. Fisher, J. D. Vervoort, S. A. Bowring, Coupled zircon Lu-Hf and U-Pb isotopic analyses of the oldest terrestrial crust, the >4.03 Ga Acasta Gneiss Complex. *Earth Planet. Sci. Lett.* **458**, 37–48 (2017).
28. B. R. Frost, C. D. Frost, A geochemical classification for feldspathic igneous rocks. *J. Petrol.* **49**, 1955–1969 (2008).
29. M.-A. Millet, N. Dauphas, N. D. Greber, K. W. Burton, C. W. Dale, B. Debret, C. G. Macpherson, G. M. Nowell, H. M. Williams, Titanium stable isotope investigation of magmatic processes on the Earth and Moon. *Earth Planet. Sci. Lett.* **449**, 197–205 (2016).

30. Z. Deng, M. Chaussidon, P. Savage, F. Robert, R. Pik, F. Moynier, Titanium isotopes as a tracer for the plume or island arc affinity of felsic rocks. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 1132–1135 (2019).
31. A. C. Johnson, S. M. Aarons, N. Dauphas, N. X. Nie, H. Zeng, R. T. Helz, S. J. Romanelli, A. D. Anbar, Titanium isotopic fractionation in Kilauea Iki lava lake driven by oxide crystallization. *Geochim. Cosmochim. Acta* **264**, 180–190 (2019).
32. L. Hoare, M. Klaver, N. S. Saji, J. Gillies, I. J. Parkinson, C. J. Lissenberg, M.-A. Millet, Melt chemistry and redox conditions control titanium isotope fractionation during magmatic differentiation. *Geochim. Cosmochim. Acta* **282**, 38–54 (2020).
33. K. J. Orians, E. A. Boyle, K. W. Bruland, Dissolved titanium in the open ocean. *Nature* **348**, 322–325 (1990).
34. G. E. Bebout, J. G. Ryan, W. P. Leeman, A. E. Bebout, Fractionation of trace elements by subduction-zone metamorphism—Effect of convergent-margin thermal evolution. *Earth Planet. Sci. Lett.* **171**, 63–81 (1999).
35. J. N. Rubin, C. D. Henry, J. G. Price, The mobility of zirconium and other “immobile” elements during hydrothermal alteration. *Chem. Geol.* **110**, 29–47 (1993).
36. N. D. Greber, N. Dauphas, A. Bekker, M. P. Ptáček, I. N. Bindeman, A. Hofmann, Titanium isotopic evidence for felsic crust and plate tectonics 3.5 billion years ago. *Science* **357**, 1271–1274 (2017).
37. T. H. Druitt, D. M. Pyle, T. A. Mather, Santorini volcano and its plumbing system. *Elements* **15**, 177–184 (2019).
38. X. Zhao, S. Tang, J. Li, H. Wang, R. Helz, B. Marsh, X. Zhu, H. Zhang, Titanium isotopic fractionation during magmatic differentiation. *Contrib. Mineral. Petrol.* **175**, 67 (2020).

39. S.-Q. Li, E. Hegner, Y.-Z. Yang, J.-D. Wu, F. Chen, Age constraints on late Mesozoic lithospheric extension and origin of bimodal volcanic rocks from the Hailar basin, NE China. *Lithos* **190-191**, 204–219 (2014).
40. E. J. Chin, K. Shimizu, G. M. Bybee, M. E. Erdman, On the development of the calc-alkaline and tholeiitic magma series: A deep crustal cumulate perspective. *Earth Planet. Sci. Lett.* **482**, 277–287 (2018).
41. J. R. Reimink, D. G. Pearson, S. B. Shirey, R. W. Carlson, J. W. F. Ketchum, Onset of new, progressive crustal growth in the central Slave craton at 3.55 Ga. *Geochem. Perspect. Lett.* **10**, 8–13 (2019).
42. J.-F. Moyen, H. Martin, Forty years of TTG research. *Lithos* **148**, 312–336 (2012).
43. C. O'Neill, A. Lenardic, L. Moresi, T. H. Torsvik, C.-T. A. Lee, Episodic Precambrian subduction. *Earth Planet. Sci. Lett.* **262**, 552–562 (2007).
44. E. Sizova, T. Gerya, K. Stüwe, M. Brown, Generation of felsic crust in the Archean: A geodynamic modeling perspective. *Precambrian Res.* **271**, 198–224 (2015).
45. E. Sizova, T. Gerya, M. Brown, L. L. Perchuk, Subduction styles in the Precambrian: Insight from numerical experiments. *Lithos* **116**, 209–229 (2010).
46. P. G. Silver, M. D. Behn, Intermittent plate tectonics? *Science* **319**, 85–88 (2008).
47. K. C. Condie, C. O'Neill, R. C. Aster, Evidence and implications for a widespread magmatic shutdown for 250 My on Earth. *Earth Planet. Sci. Lett.* **282**, 294–298 (2009).
48. T. E. Johnson, N. J. Gardiner, K. Miljković, C. J. Spencer, C. L. Kirkland, P. A. Bland, H. Smithies, An impact melt origin for Earth's oldest known evolved rocks. *Nat. Geosci.* **11**, 795–799 (2018).
49. Z. Deng, F. Moynier, P. A. Sossi, M. Chaussidon, Bridging the depleted MORB mantle and the continental crust using titanium isotopes. *Geochem. Perspect. Lett.* **9**, 11–15 (2018).

50. T. V. Gerya, R. J. Stern, M. Baes, S. V. Sobolev, S. A. Whattam, Plate tectonics on the Earth triggered by plume-induced subduction initiation. *Nature* **527**, 221–225 (2015).
51. N. D. Greber, N. Dauphas, I. S. Puchtel, B. A. Hofmann, N. T. Arndt, Titanium stable isotopic variations in chondrites, achondrites and lunar rocks. *Geochim. Cosmochim. Acta* **213**, 534–552 (2017).
52. A. Roman, N. Arndt, Differentiated Archean oceanic crust: Its thermal structure, mechanical stability and a test of the sagduction hypothesis. *Geochim. Cosmochim. Acta* **278**, 65–77 (2020).
53. A. B. Rozel, G. J. Golabek, C. Jain, P. J. Tackley, T. Gerya, Continental crust formation on early Earth controlled by intrusive magmatism. *Nature* **545**, 332–335 (2017).
54. A. M. Bauer, J. R. Reimink, T. Chacko, B. J. Foley, S. B. Shirey, D. G. Pearson, Hafnium isotopes in zircons document the gradual onset of mobile-lid tectonics. *Geochem. Perspect. Lett.* **14**, 1–6 (2020).
55. W. J. Collins, S. D. Beams, A. J. R. White, B. W. Chappell, Nature and origin of A-type granites with particular reference to Southeastern Australia. *Contrib. Mineral. Petrol.* **80**, 189–200 (1982).
56. J. D. Clemens, J. R. Holloway, A. J. R. White, Origin of an A-type granite: Experimental constraints. *Am. Mineral.* **71**, 317–324 (1986).
57. J. B. Whalen, K. L. Currie, B. W. Chappell, A-type granites: Geochemical characteristics, discrimination and petrogenesis. *Contrib. Mineral. Petrol.* **95**, 407–419 (1987).
58. R. F. Martin, A-type granites of crustal origin ultimately result from open-system fenitization-type reactions in an extensional environment. *Lithos* **91**, 125–136 (2006).
59. M. C. Loiselle, D. R. Wones, Characteristics of anorogenic granites. *Geol. Soc. Am. Abstr. Prog.* **11**, 468 (1979).

60. S. P. Turner, J. D. Foden, R. S. Morrison, Derivation of some A-type magmas by fractionation of basaltic magma: An example from the Padthaway Ridge, South Australia. *Lithos* **28**, 151–179 (1992).
61. J. L. Anderson, in *Proterozoic Geology: Selected Papers from an International Proterozoic Symposium*, J. G. Medaris Jr., C. W. Byers, D. M. Mickelson, W. C. Shanks, Eds. (Geological Society of America Memoir, 1983), vol. 161.
62. R. A. Creaser, R. C. Price, R. J. Wormald, A-type granites revisited: Assessment of a residual-srouce model. *Geology* **19**, 163–166 (1991).
63. M.-A. Millet, N. Dauphas, Ultra-precise titanium stable isotope measurements by double-spike high resolution MC-ICP-MS. *J. Anal. At. Spectrom* **29**, 1444–1458 (2014).
64. J. Zhang, N. Dauphas, A. M. Davis, A. Pourmand, A new method for MC-ICPMS measurement of titanium isotopic composition: Identification of correlated isotope anomalies in meteorites. *J. Anal. At. Spectrom* **26**, 2197–2205 (2011).
65. N. Dauphas, A. Pourmand, F.-Z. Teng, Routine isotopic analysis of iron by HR-MC-ICPMS: How precise and how accurate? *Chem. Geol.* **267**, 175–184 (2009).
66. N. Dauphas, P. E. Janney, R. A. Mendybaev, M. Wadhwa, F. M. Richter, A. M. Davis, M. van Zuilen, R. Hines, C. N. Foley, Chromatographic separation and multicollection-ICPMS analysis of iron. Investigating mass-dependent and independent isotope effects. *Anal. Chem.* **76**, 5855–5563 (2004).
67. P. J. Wallace, I. S. E. Carmichael, Sulfur in basaltic magmas. *Geochim. Cosmochim. Acta* **56**, 1863–1874 (1992).
68. N. Métrich, M. Bonnin-Mosbah, B. Menez, L. Galoisy, Presence of sulfite (SIV) in arc magmas: Implications for volcanic sulfur emissions. *Geophys. Res. Lett.* **29**, 1–33 (2002).

69. P. A. Sossi, J. D. Foden, G. P. Halverson, Redox-controlled iron isotope fractionation during magmatic differentiation: An example from the Red Hill intrusion, S. Tasmania. *Contrib. Mineral. Petrol.* **164**, 757–772 (2012).
70. N. Dauphas, M. Roskosz, E. E. Alp, D. R. Neuville, M. Y. Hu, C. K. Sio, F. L. H. Tissot, J. Zhao, L. Tissandier, E. Médard, C. Cordier, Magma redox and structural controls on iron isotope variations in Earth's mantle and crust. *Earth Planet. Sci. Lett.* **398**, 127–140 (2014).
71. M. Roskosz, C. K. I. Sio, N. Dauphas, W. Bi, F. L. H. Tissot, M. Y. Hu, J. Zhao, E. E. Alp, Spinel–olivine–pyroxene equilibrium iron isotopic fractionation and applications to natural peridotites. *Geochim. Cosmochim. Acta* **169**, 184–199 (2015).
72. S. A. Bowring, I. S. Williams, W. Compston, 3.96 Ga gneisses from the Slave province, Northwest Territories, Canada. *Geology* **17**, 971–975 (1989).
73. S. A. Bowring, I. S. Williams, Priscoan (4.00–4.03 Ga) orthogneisses from northwestern Canada. *Contrib. Mineral. Petrol.* **134**, 3–16 (1999).
74. R. Stern, W. Bleeker, Age of the world's oldest rocks refined using Canada's SHRIMP: The Acasta Gneiss Complex, Northwest Territories, Canada. *Geoscience Canada* **25**, 27–31 (1998).
75. A. S. G. Roth, B. Bourdon, S. J. Mojzsis, J. F. Rudge, M. Guitreau, J. Blichert-Toft, Combined $^{147,146}\text{Sm}$ - $^{143,142}\text{Nd}$ constraints on the longevity and residence time of early terrestrial crust. *Geochem. Geophys. Geosyst.* **15** (2014).
76. M. Guitreau, J. Blichert-Toft, S. J. Mojzsis, A. S. G. Roth, B. Bourdon, N. L. Cates, W. Bleeker, Lu–Hf isotope systematics of the Hadean–Eoarchean Acasta Gneiss Complex (Northwest Territories, Canada). *Geochim. Cosmochim. Acta* **135**, 251–269 (2014).
77. Y. Amelin, D.-C. Lee, A. N. Halliday, R. T. Pidgeon, Nature of the Earth's earliest crust from hafnium isotopes in single detrital zircons. *Nature* **399**, 252–255 (1999).

78. T. Iizuka, T. Hirata, Improvements of precision and accuracy in in situ Hf isotope microanalysis of zircon using the laser ablation-MC-ICPMS technique. *Chem. Geol.* **220**, 121–137 (2005).
79. R. H. Smithies, The Archaean tonalite-trondhjemite-granodiorite (TTG) series is not an analogue of Cenozoic adakite. *Earth Planet. Sci. Lett.* **182**, 115–125 (2000).
80. M. Telus, N. Dauphas, F. Moynier, F. L. H. Tissot, F.-Z. Teng, P. I. Nabelek, P. R. Craddock, L. A. Groat, Iron, zinc, magnesium and uranium isotopic fractionation during continental crust differentiation: The tale from migmatites, granitoids, and pegmatites. *Geochim. Cosmochim. Acta* **97**, 247–265 (2012).
81. K. Kreissig, T. F. Nägler, J. D. Kramers, D. D. Reenen, C. A. Smit, An isotopic and geochemical study of the northern Kaapvaal Craton and the Southern Marginal Zone of the Limpopo Belt: Are they juxtaposed terranes? *Lithos* **50**, 1–25 (2000).
82. R. Batiza, Y. Niu, Petrology and magma chamber processes at the East Pacific Rise ~ 9°30'N. *J. Geophys. Res. Solid Earth* **97**, 6779–6797 (1992).
83. H. Puchelt, R. Emmermann, Petrogenetic implications of tholeiitic basalt glasses from the East Pacific Rise and the Galápagos Spreading Center. *Chem. Geol.* **38**, 39–56 (1983).
84. S. Escrig, F. Capmas, B. Dupré, C.J. Allègre, Osmium isotopic constraints on the nature of the DUPAL anomaly from Indian mid-ocean-ridge basalts. *Nature* **431**, 59–63 (2004).
85. W. Yi, A. N. Halliday, J. C. Alt, D.-C. Lee, M. Rehkämper, M. O. Garcia, C. H. Langmuir, Y. Su, Cadmium, indium, tin, tellurium, and sulfur in oceanic basalts: Implications for chalcophile element fractionation in the Earth. *J. Geophys. Res. Solid Earth* **105**, 18927–18948 (2000).
86. P. E. Baker, F. Buckley, J. G. Holland, Petrology and geochemistry of easter island. *Contrib. Mineral. Petrol.* **44**, 85–100 (1974).

87. M.-A. Millet, R. Doucelance, P. Schiano, K. David, C. Bosq, Mantle plume heterogeneity versus shallow-level interactions: A case study, the São Nicolau Island, Cape Verde archipelago. *J. Volcanol. Geotherm. Res.* **176**, 265–276 (2008).
88. S. Turner, C. Hawkesworth, N. Rogers, P. King, U-Th isotope disequilibria and ocean island basalt generation in the Azores. *Chem. Geol.* **139**, 145–164 (1997).
89. M.-A. Millet, R. Doucelance, J. A. Baker, P. Schiano, Reconsidering the origins of isotopic variations in Ocean Island Basalts: Insights from fine-scale study of São Jorge Island, Azores archipelago. *Chem. Geol.* **265**, 289–302 (2009).
90. R. T. Helz, H. K. Kirschenbaum, J. W. Marinenko, R. Qian, Whole-rock analyses of core samples from the 1967, 1975, 1979 and 1981 drillings of Kilauea Iki lava lake, Hawaii (U.S. Geological Survey Professional Paper, Open-File Report 94-684, 1994).
91. K. J. Murata, D. H. Richter, Chemistry of the lavas of the 1959–60 eruption of Kilauea Volcano, Hawaii (U.S. Geological Survey Professional Paper, Report no. 537-A, 1966).
92. F. A. Frey, M. O. Garcia, M. F. Roden, Geochemical characteristics of Koolau Volcano: Implications of intershield geochemical differences among Hawaiian volcanoes. *Geochim. Cosmochim. Acta* **58**, 1441–1462 (1994)
93. P. S. Savage, R. B. Georg, H. M. Williams, K. W. Burton, A. N. Halliday, Silicon isotope fractionation during magmatic differentiation. *Geochim. Cosmochim. Acta* **75**, 6124–6139 (2011).
94. R. Pik, B. Marty, D. R. Hilton, How many mantle plumes in Africa? The geochemical point of view. *Chem. Geol.*, **226**, 100–114 (2006).
95. L. Field, A. Calvert, G. Yirgu, Magmatic history of Dabbahu, a composite volcano in the Afar Rift, Ethiopia. *Geol. Soc. Am. Bull.* **125**, 128–147 (2013)
96. K. J. Chamberlin, J. Barclay, K. J. Preece, R. J. Brown, J. P. Davidson, Lower crustal heterogeneity and fractional crystallization control evolution of small-volume magma batches

- at ocean island volcanoes (Ascension Island, South Atlantic), *J. Petrol.*, **60**, 1489–1522 (2019).
97. J. Barling, “The petrogenesis of the Newer Lavas of Heard Island, southern Indian Ocean,” thesis, Monash University (1990).
98. J. Barling, S. L. Golstein, I. A. Nicholls, Geochemistry of Heard Island (southern Indian Ocean): Characterization of an enriched mantle component and implications for enrichment of the sub-Indian Ocean mantle, *J. Petrol.* **35**, 1017–1053 (1994).
99. D. A. Clague, D. W. Caress, B. M. Dreyer, L. Lundsten, J. B. Paduan, R. A. Portner, R. Spelz-Madero, J. A. Bowles, P. R. Castillo, R. Guardado-France, M. L. Saout, J. F. Martin, M. A. Santa Rosa-del Río, R. A. Zierenberg, Geology of the Alarcon rise, Southern Gulf of California, *Geochem. Geophys. Geosyst.* **19**, 807–837 (2018).
100. C. Timm, I. J. Graham, C. E. J. de Ronde, M. I. Leybourne, J. Woodhead, Geochemical evolution of Monowai volcanic center: New insights into the northern Kermadec arc subduction system, SW Pacific. *Geochem. Geophys. Geosyst.* **12**, Q0AF01 (2011).
101. S. Dempsey, “Geochemistry of volcanic rocks from the Sunda Arc,” thesis, Durham University (2013).
102. D. Peters, T. Pettke, Evaluation of major to ultra trace element bulk rock chemical analysis of nanoparticulate pressed powder pellets by LA-ICP-MS. *Geostand. Geoanal. Res.*, **41**, 5–28 (2017).
103. M. Klaver, G. R. Davies, P. Z. Vroon, Subslab mantle of African provenance infiltrating the Aegean mantle wedge. *Geology*, **44**, 367–370 (2016).
104. M. Klaver, S. Carey, P. Nomikou, I. Smet, A. Godelitsas, P. Vroon, A distinct source and differentiation history for Kolumbo submarine volcano, Santorini volcanic field, Aegean arc. *Geochem. Geophys. Geosyst.*, **17**, 3254–3273 (2016).

105. M. Klaver, “Dynamics of magma generation and differentiation in the central-eastern Aegean arc: A geochemical and petrological study of Quaternary arc volcanism in Greece,” thesis, Vrije Universiteit (2016).
106. L.-K. Sha, B. W. Chappell, Apatite chemical composition, determined by electron microprobe and laser-ablation inductively coupled plasma mass spectrometry, as a probe into granite petrogenesis. *Geochim. Cosmochim. Acta* **63**, 3861–3881 (1999).
107. B. W. Chappell, Source rocks of I-type and S-type granites in the Lachlan Fold Belt, Southeastern Australia. *Philos. Trans. A Math. Phys. Eng. Sci.* **310**, 693–707 (1984).
108. B. W. Chappell, A. J. R. White, *The Cowra Granodiorite and Its Enclaves. IAVCEI Canberra Excursion Guide* (Australian Geological Survey Organisation, 1993).
109. R. Hine, I. S. Williams, B. W. Chappell, A. J. R. White, Contrasts between I- and S-type granitoids of the Kosciusko Batholith. *J. Geol. Soc. Aus.* **25**, 219–234 (1978).
110. B. W. Chappell, A. J. R. White, I- and S-type granites in the Lachlan Fold Belt. *Phil. Trans. R. Soc. Edinb. Earth Sci.* **83**, 1–26 (1992).
111. P. I. King, A. J. R. White, B. W. Chappell, C. M. Allen, Characterization and origin of aluminous A type granites of the Lachlan Fold Belt, Southeastern Australia. *J. Petrol.* **38**, 371–391 (1997).
112. I. S. Puchtel, R. J. Walker, M. Touboul, E. G. Nisbet, G. R. Byerly, Insights into early Earth from the Pt–Re–Os isotope and highly siderophile element abundance systematics of Barberton komatiites. *Geochim. Cosmochim. Acta* **125** 394–413 (2014).
113. I. S. Puchtel, J. Blachert-Toft, M. Touboul, R. J. Walker, G. R. Byerly, E. G. Nisbet, C. R. Anhaeusser, Insights into early Earth from Barberton komatiites: Evidence from lithophile isotope and trace element systematics. *Geochim. Cosmochim. Acta* **108**, 63–90 (2013).

114. I. S. Puchtel, J. Blichert-Toft, M. Touboul, M. F. Horan, R. J. Walker, The coupled ^{182}W - ^{142}Nd record of early terrestrial mantle differentiation. *Geochem. Geophys. Geosyst.* **17**, 2168–2193 (2016).
115. Y. Lahaye, N. Arndt, Alteration of a komatiite flow from Alexo, Ontario, Canada. *J. Petrol.* **37**, 1261–1284 (1996).
116. N. Dauphas, F.-Z. Teng, N. T. Arndt, Magnesium and iron isotopes in 2.7 Ga Alexo komatiites: Mantle signatures, no evidence for Soret diffusion, and identification of diffusive transport in zoned olivine. *Geochim. Cosmochim. Acta* **74**, 3274–3291 (2014).
117. I. S. Puchtel, R. J. Walker, A. D. Brandon, E. G. Nisbet, Pt–Re–Os and Sm–Nd isotope and HSE and REE systematics of the 2.7 Ga Belingwe and Abitibi komatiites. *Geochim. Cosmochim. Acta* **73**, 6367–6389 (2009).
118. P. A. Sossi, S. M. Eggins, R. W. Nesbitt, O. Nebel, J. M. Hergt, I. H. Campbell, H. S. C. O’Neill, M. Van Kranendonk, D. R. Davies, Petrogenesis and geochemistry of Archean komatiites. *J. Petrol.* **57**, 147–184 (2016).
119. P. R. Craddock, N. Dauphas, Iron isotopic compositions of geological reference materials and chondrites. *Geostand. Geoanal. Res.*, **35**, 101–123 (2011).
120. P. I. Nabelek, C. Russ-Nabelek, J. R. Denison, The generation and crystallization conditions of the Proterozoic Harney Peak Leucogranite, Black Hills, South Dakota, USA: Petrologic and geochemical constraints. *Contrib. Mineral. Petrol.* **110**, 173–191 (1992).
121. R. J. Walker, “Origin of the Tin Mountain Pegmatites, Black Hills, South Dakota,” thesis, State University of New York at Stony Brook (1984).
122. R. J. Walker, G. N. Hanson, J. J. Papike Trace element constraints on pegmatite genesis: Tin Mountain Pegmatite, Black Hills, South Dakota. *Contrib. Mineral. Petrol.* **101**, 290–300 (1989).

123. L. A. Groat, T. Mulja, M. H. F. Mauthner, T. S. Ercit, M. Raudsepp, R. A. Gault, H. A. Rollo, Geology and mineralogy of the little nahanni rare-element granitic pegmatites, northwest territories. *Can. Mineral.* **41**, 139–160 (2003).
124. P. I. Nabelek, Trace element distribution among rock- forming minerals in Black Hills migmatites, South Dakota; a case for solid-state equilibrium. *Am. Mineral.* **84**, 1256–1269 (1999).
125. I. McDougall, Differentiation of the Tasmanian dolerites: Red Hill dolerite-granophyre association. *Geol. Soc. Am. Bull.* **73**, 279–316 (1962).