



US011059866B2

(12) **United States Patent**  
**Schneewind et al.**

(10) **Patent No.:** **US 11,059,866 B2**

(45) **Date of Patent:** **\*Jul. 13, 2021**

(54) **COMPOSITIONS AND METHODS RELATED TO PROTEIN A (SPA) VARIANTS**

(71) Applicant: **The University of Chicago**, Chicago, IL (US)

(72) Inventors: **Olaf Schneewind**, Chicago, IL (US); **Alice G. Cheng**, Boston, MA (US); **Dominique M. Missiakas**, Chicago, IL (US); **Hwan Keun Kim**, Chicago, IL (US)

(73) Assignee: **The University of Chicago**, Chicago, IL (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **16/661,155**

(22) Filed: **Oct. 23, 2019**

(65) **Prior Publication Data**

US 2020/0140494 A1 May 7, 2020

**Related U.S. Application Data**

(63) Continuation of application No. 15/702,037, filed on Sep. 12, 2017, now Pat. No. 10,464,971, which is a continuation of application No. 15/060,861, filed on Mar. 4, 2016, now abandoned, which is a continuation of application No. 14/466,514, filed on Aug. 22, 2014, now Pat. No. 9,315,554, which is a continuation of application No. 13/807,598, filed as application No. PCT/US2011/042845 on Jul. 1, 2011, now Pat. No. 8,821,894.

(60) Provisional application No. 61/370,725, filed on Aug. 4, 2010, provisional application No. 61/361,218, filed on Jul. 2, 2010.

(51) **Int. Cl.**  
**A61K 39/085** (2006.01)  
**C07K 16/12** (2006.01)  
**C07K 14/31** (2006.01)  
**A61K 39/00** (2006.01)

(52) **U.S. Cl.**  
CPC ..... **C07K 14/31** (2013.01); **A61K 39/085** (2013.01); **C07K 16/1271** (2013.01); **A61K 2039/57** (2013.01)

(58) **Field of Classification Search**  
None  
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,027,010 A 5/1977 Kiselev et al.  
4,327,082 A 4/1982 Armitage  
4,690,915 A 9/1987 Rosenberg  
4,902,616 A 2/1990 Fournier et al.

5,151,350 A 9/1992 Colbert et al.  
5,189,015 A 2/1993 Hook et al.  
5,199,942 A 4/1993 Gillis  
5,294,177 A 3/1994 Rasnick et al.  
5,320,951 A 6/1994 Hook et al.  
5,648,240 A 7/1997 Hook  
5,801,234 A 9/1998 Hodgson  
5,840,846 A 11/1998 Hook  
6,008,341 A 12/1999 Foster  
6,013,763 A 1/2000 Braisted et al.  
6,197,927 B1 3/2001 Braisted et al.  
6,288,214 B1 9/2001 Hook  
6,294,177 B1 9/2001 Fattom  
6,299,879 B1 10/2001 Wastfalt et al.  
6,340,571 B1 1/2002 Merlin et al.  
6,403,337 B1 6/2002 Bailey et al.  
6,593,114 B1 7/2003 Kunsch et al.  
6,635,473 B1 10/2003 Foster et al.  
6,680,195 B1 1/2004 Patti et al.  
6,692,739 B1 2/2004 Patti et al.  
6,703,025 B1 3/2004 Patti et al.  
6,703,492 B1 3/2004 Kimmerly  
6,737,248 B2 5/2004 Kunsch et al.  
6,753,149 B2 6/2004 Bailey et al.  
6,833,253 B2 12/2004 Choi  
6,841,154 B2 1/2005 Foster et al.  
6,984,381 B2 1/2006 Guidry et al.

(Continued)

FOREIGN PATENT DOCUMENTS

EP 0 786 519 7/1997  
EP 0 594 610 9/1998

(Continued)

OTHER PUBLICATIONS

“Policy Responses to the Growing Threat of Antibiotic Resistance: A Shot Against MRSA?” Extending the Cure (<http://www.extendingthecure.org>), Policy Brief 7, available online at [http://www.extendingthecure.org/sites/default/files/PolicyBrief7\\_1.pdf](http://www.extendingthecure.org/sites/default/files/PolicyBrief7_1.pdf), Mar. 2009.

Abdallah et al., “A specific secretion system mediates PPE41 transport in pathogenic mycobacteria”, *Mol. Microbiol.*, 62, 667-679, 2006.

Abdallah et al., “Type VII secretion—mycobacteria show the way”, *Nat. Rev. Microbiol.*, 5:883-891, 2007.

Adlam et al., “Effect of immunization with highly purified alpha- and beta-toxins on staphylococcal mastitis in rabbits,” *Infect. Immun.*, 17(2):250-6, 1977.

(Continued)

Primary Examiner — Padmavathi Baskar

(74) Attorney, Agent, or Firm — Norton Rose Fulbright US LLP

(57) **ABSTRACT**

The present invention concerns methods and compositions for treating or preventing a bacterial infection, particularly infection by a *Staphylococcus* bacterium. The invention provides methods and compositions for stimulating an immune response against the bacteria. In certain embodiments, the methods and compositions involve a non-toxicogenic Protein A (SpA) variant.

**15 Claims, 15 Drawing Sheets**

**Specification includes a Sequence Listing.**

(56)

## References Cited

## U.S. PATENT DOCUMENTS

7,045,131	B2	5/2006	Patti et al.
7,115,264	B2	10/2006	Patti et al.
7,195,763	B2	3/2007	Xu et al.
7,488,807	B2	2/2009	Mach et al.
8,821,894	B2	9/2014	Schneewind et al.
9,315,554	B2	4/2016	Schneewind et al.
10,464,971	B2*	11/2019	Schneewind ..... A61P 43/00
2002/0169288	A1	11/2002	Hook
2003/0087864	A1	5/2003	Talbot et al.
2003/0113350	A1	6/2003	Fattom et al.
2004/0006209	A1	1/2004	Patti et al.
2004/0014178	A1	1/2004	Guss et al.
2004/0043037	A1	3/2004	Kunsch et al.
2004/0101919	A1	5/2004	Hook et al.
2004/0265962	A1	12/2004	Bailey et al.
2005/0026170	A1	2/2005	Patti et al.
2005/0031625	A1	2/2005	Mohamed et al.
2005/0106597	A1	5/2005	Choi
2005/0106648	A1	5/2005	Foster et al.
2005/0220788	A1	10/2005	Nagy et al.
2005/0255478	A1	11/2005	Kimmerly
2006/0002939	A1	1/2006	Fischer et al.
2006/0051820	A1	3/2006	Horii et al.
2006/0115490	A1	6/2006	Masigani et al.
2006/0134141	A1	6/2006	Fattom et al.
2006/0117462	A1	8/2006	Anderson et al.
2006/0188515	A1	8/2006	Anderson et al.
2006/0205016	A1	9/2006	Silverman
2006/0222651	A1	10/2006	Patti et al.
2006/0228368	A1	10/2006	Fattom et al.
2006/0263792	A1	11/2006	Mohamed et al.
2007/0020746	A1	12/2007	Kunsch et al.
2008/0095792	A1	4/2008	Anderson et al.
2008/0131457	A1	6/2008	Taylor et al.
2008/0160089	A1	7/2008	Vitiello et al.
2008/0311146	A1	12/2008	Castado
2009/0053235	A1	2/2009	Taylor et al.
2009/0162902	A1	6/2009	Mach
2009/0317421	A1	12/2009	Missiakas
2010/0272743	A1	10/2010	Kimmerly
2011/0027265	A1	2/2011	Bubeck-Wardenburg
2011/0059085	A1	3/2011	Kim
2011/0206676	A1	8/2011	Missiakas
2011/0262477	A1	10/2011	Cheng
2012/0009182	A1	1/2012	Yeung et al.
2012/0107327	A1	5/2012	Anderson et al.
2012/0114686	A1	5/2012	Schneewind
2012/0282247	A1	11/2012	Schneewind
2013/0136746	A1	5/2013	Schneewind
2013/0189249	A1	7/2013	Bubeck-Wardenburg
2013/0230550	A1	9/2013	Schneewind
2013/0236419	A1	9/2013	Schneewind
2016/0194363	A1	7/2016	Schneewind et al.

## FOREIGN PATENT DOCUMENTS

EP	0 786 519	10/1998
EP	1 829 892	9/2007
EP	2 157 099	2/2010
JP	2005520853	7/2005
JP	2008518947	6/2008
WO	WO 98/57994	12/1998
WO	WO 1999/027109	6/1999
WO	WO 00/02523	1/2000
WO	WO 2000/002523	1/2000
WO	WO 00/12132	3/2000
WO	WO 00/12689	3/2000
WO	WO 00/15238	3/2000
WO	WO 2000/012131	3/2000
WO	WO 2000/012678	3/2000
WO	WO 00/69457	11/2000
WO	WO 2000/064925	11/2000
WO	WO 2000/071585	11/2000
WO	WO 01/34809	5/2001
WO	WO 01/60852	8/2001

WO	WO 2001/070267	9/2001
WO	WO 2001/070955	9/2001
WO	WO 01/98499	12/2001
WO	WO 02/59148	8/2002
WO	WO 02/94868	11/2002
WO	WO 2002/102829	12/2002
WO	WO 2003/011899	2/2003
WO	WO 2003/041726	5/2003
WO	WO 2003/076470	9/2003
WO	WO 03080106	10/2003
WO	WO 2004/025416	3/2004
WO	WO 2004/030699	4/2004
WO	WO 2004/094600	11/2004
WO	WO 2005/009378	2/2005
WO	WO 2005/009379	2/2005
WO	WO 2005/079315	9/2005
WO	WO 2006/032472	3/2006
WO	WO 2006/032475	3/2006
WO	WO 2006/032500	3/2006
WO	WO 2006/050291	5/2006
WO	WO 2006/059247	6/2006
WO	WO 2006/078213	7/2006
WO	WO 2007/001361	1/2007
WO	WO 2007/010413	1/2007
WO	WO 2007/071692	6/2007
WO	WO 2007/089470	8/2007
WO	WO 2007/095057	8/2007
WO	WO 2007/100580	9/2007
WO	WO 2007/113222	10/2007
WO	WO 2007/113223	10/2007
WO	WO 2007/145689	12/2007
WO	WO 2008/019162	2/2008
WO	WO 2008/081014	7/2008
WO	WO 2008/140487	11/2008
WO	WO 2008/140570	11/2008
WO	WO 2008/143697	11/2008
WO	WO 2008/152447	12/2008
WO	WO 2009/029132	3/2009
WO	WO 2009/140236	11/2009
WO	WO 2011/005341	1/2011
WO	WO 2011/127032	10/2011
WO	WO 2012/122533	9/2012

## OTHER PUBLICATIONS

Albus et al., "Virulence of *Staphylococcus aureus* mutants altered in type 5 capsule production," *Infect. Immun.*, 59: 1008-1014, 1991.

Allen et al., "HtaA is an iron-regulated hemin binding protein involved in the utilization of heme iron in *Corynebacterium diphtheriae*," *J. Bacteriol.*, 191:2638-2648, 2009.

Andersen et al., "Recall of Long-Lived Immunity to Mycobacterium tuberculosis Infection in Mice", *J. Immunol.*, 154:3359-3372, 1995.

Archer, "*Staphylococcus aureus*: A Well-Armed Pathogen", *Clin. Infect. Dis.*, 26:1179-1181, 1998.

Athanasopoulos et al., "The extracellular adherence protein (Eap) of *Staphylococcus aureus* inhibits wound healing by interfering with host defense and repair mechanisms," *Blood.*, 107(7):2720-2727, 2006.

Baba et al., "Genome Sequence of *Staphylococcus aureus* Strain Newman and comparative analysis of Staphylococcal Genomes: Polymorphism and Evolution of Two Major Pathogenicity Islands", *J. Bacteriol.* 190(1):300-310, 2007.

Bae and Schneewind, "Allelic replacement in *Staphylococcus aureus* with inducible counter-selection," *Plasmid.* 55:58-63, 2006.

Bae et al., "*Staphylococcus aureus* virulence genes identified by bursa aurealis mutagenesis and nematode killing", *Proc. Natl. Acad. Sci. USA*, 101(33):12312-12317, 2004.

Bhakdi and Tranum-Jensen, "Alpha-toxin of *Staphylococcus aureus*," *Microbiol. Rev.*, 55 (4): 733-751, 1991.

Bhakdi et al. "Functionally inactive *S. aureus* alpha-toxin containing a single amino acid substitution: potential usefulness as a vaccine," *Behring Inst. Mitt.*, (5):80-4, 1994. (English abstract).

Bjerketorp et al., "A novel von Willebrand factor binding protein expressed by *Staphylococcus aureus*," *Microbiology*, 148:2037-2044, 2002.

Bjerketorp et al., "The von Willebrand factor-binding protein (vWbp) of *Staphylococcus aureus* is a coagulase", *FEMS Microbiol. Lett.*, 234:309-314, 2004.

(56)

## References Cited

## OTHER PUBLICATIONS

- Boucher and Corey, "Epidemiology of Methicillin-Resistant *Staphylococcus aureus*," *Clin. Infect. Dis.*, 46:S344-S349, 2008.
- Brady et al., "Osteomyelitis and the role of biofilms in chronic infection," *FEMS Immunol. Med. Microbiol.*, 52:13-22, 2008.
- Brodin et al., "ESAT-6 proteins: protective antigens and virulence factors?" *Trends in Microbiology*, 12 (11): 500-508, 2004.
- Brodin et al., "Functional analysis of early secreted antigenic target-6, the dominant T-cell antigen of Mycobacterium tuberculosis, reveals key residues involved in secretion, complex formation, virulence, and immunogenicity," *J. of Biol. Chem.*, 280 (40): 33953-3959, 2005.
- Brown et al., "A study of the interactions between an IgG-binding domain based on the B domain of staphylococcal protein A and rabbit IgG," *Molecular Biotechnology*, 10:9-16, 1998.
- Brown et al., "Determining Protein-Protein Interactions by Oxidative Cross-Linking of a Glycine-Glycine-Histidine Fusion Protein", *Biochemistry*, 37:4397-4406, 1998.
- Bubeck Wardenburg et al., "Surface proteins and exotoxins are required for the pathogenesis of *Staphylococcus aureus* pneumonia," *Infection and Immunity*, 75(2):1040-1044, 2007.
- Bubeck Wardenburg et al., "Vaccine protection against *Staphylococcus aureus* pneumonia," *Journal of Experimental Medicine*, 205(2):287-294, 2008.
- Burts et al., "EsaC: A new secretion substrate of the staphylococcal ESAT-6 secretion pathway," *Abstracts of the General Meeting of the American Society for Microbiology*, 107:102-103, 2007.
- Burts et al., "EsaC substrate for the ESAT-6 Secretion Pathway and its role in persistent infections of *S. aureus*", *Mol. Microbiol.*, 69(3):736-746, 2008.
- Burts et al., "EsaX and EsaB are secreted by an ESAT-6-like system that is required for the pathogenesis of *Staphylococcus aureus* infections", *Proc. Natl. Acad. Sci. USA*, 102(4):1169-1174, 2005.
- Campo et al., "Subcellular sites for bacterial protein export," *Mol. Microbiol.*, 53 (6): 1583-1599, 2004.
- Cedergren et al., "Mutational analysis of the interaction between staphylococcal protein A and human IgG1", *Protein Eng.*, 6(4):441-448, 1993.
- Cespedes et al., "The Clonality of *Staphylococcus aureus* Nasal Carriage", *J. Infect. Dis.*, 191(3):444-452, 2005.
- Chavakis et al., "*Staphylococcus aureus* interactions with the endothelium," *Thromb. Haemost.*, 94:278-85, 2005.
- Cheng et al., "Contribution of Coagulases towards *Staphylococcus aureus* disease and protective immunity," *PLoS Pathogens*, 6(8):e1001036, 18 pages, 2010.
- Cheng et al., "Genetic requirements for *Staphylococcus aureus* abscess formation and persistence in host tissues", *FASEB J.*, 23:3393-3404, 2009.
- Cheung et al., "Cloning, expression, and nucleotide sequence of a *Staphylococcus aureus* gene (fbpA) encoding a fibrinogen-binding protein," *Infection and Immunity*, 63(5):1914-1920, 1995.
- Cheung et al., "Diminished virulence of a sar-/agr- mutant of *Staphylococcus aureus* in the rabbit model of endocarditis," *J. Clin. Invest.*, 94 (5): 1815-1822, 1994.
- Chhatwal, "Anchorless adhesins and invasins of Gram-positive bacteria: a new class of virulence factors," *Trends Microbiol.*, 10 (5): 205-208, 2002.
- Clarke et al., "Surface adhesins of *Staphylococcus aureus*," *Adv. Microb. Physiol.*, 51:187-224, 2006.
- Colman et al., "Effects of amino acid sequence changes on antibody-antigen interactions," *Research in Immunology*, 145:33-36, 1994.
- Craven et al., "*Staphylococcus aureus* alpha-hemolysin activates the NLRP3-inflammasome in human and mouse monocytic cells," *PLoS ONE*, 4(10):e7746, 11 pages, 2009.
- Dalbey and Wickner, "Leader Peptidase Catalyzes the Release of Exported Proteins from the Outer Surface of the *Escherichia coli* Plasma Membrane", *J. Biol. Chem.*, 260(29):15925-15931, 1985.
- DeBord et al., "Immunogenicity and protective immunity against bubonic plague and pneumonic plague by immunization of mice with the recombinant V10 antigen, a variant of LcrV," *Infect. Immun.*, 74(8):4910-4914, 2006.
- DeDent et al., "Distribution of Protein A on the Surface of *Staphylococcus aureus*", *J. Bacteriol.* 189:4473-4484, 2007.
- DeDent et al., "Signal peptides direct surface proteins to two distinct envelope locations of *Staphylococcus aureus*", *EMBO J.* 27:2656-2668, 2008.
- Deisenhofer et al., "Crystallization, Crystal Structure Analysis and Atomic Model of the Complex Formed by a Human Fc Fragment and Fragment B of Protein A from *Staphylococcus aureus*", *Hoppe-Seyh Zeitsch. Physiol. Chem.* 359:975-985, 1978.
- Deisenhofer, "Crystallographic Refinement and Atomic Models of a Human Fc Fragment and Its Complex with Fragment B of Protein A from *Staphylococcus aureus* at 2.9- and 2.8-Å Resolution", *Biochemistry* 20(9):2361-2370, 1981.
- Diep et al., "Complete genome sequence of USA300, an epidemic clone of community-acquired methicillin-resistant *Staphylococcus aureus*," *Lancet.*, 367(9512):731-739, 2006.
- Diep et al., "Roles of 34 virulence genes in the evolution of hospital- and community-associated strains of methicillin-resistant *Staphylococcus aureus*," *J. Infect. Dis.*, 193(11):1495-1503, 2006.
- Dinges et al., "Exotoxins of *Staphylococcus aureus*", *Clin. Microbiol. Rev.*, 13(1):16-34, 2000.
- Dryla et al., "High-affinity binding of the staphylococcal HarA protein to haptoglobin and hemoglobin involves a domain with an antiparallel eight-stranded beta-barrel fold," *J. Bacteriol.*, 189:254-264, 2007.
- Dryla et al., "Identification of a novel iron regulated staphylococcal surface protein with haptoglobin-haemoglobin binding activity," *Mol. Microbiol.*, 49:37-53, 2003.
- Duthie and Lorenz, "Staphylococcal coagulase: Mode of action and Antigenicity", *J. Gen. Microbiol.*, 6:95-107, 1952.
- Ekstedt and Yotis, "Effect of Coagulase on the Virulence of Coagulase Negative Strains", *iAnn. N.Y. Acad. Sci.*, 80:496-500, 1960.
- Emori and Gaynes, "An overview of nosocomial infections, including the role of the microbiology laboratory," *Clin. Microbiol. Rev.*, 6(4):428-442, 1993.
- Etz et al., Identification of in vivo expressed vaccine candidate antigens from *Staphylococcus aureus*, *PNAS*, 99(10):6573-6578, 2002.
- Fattom et al., "Development of StaphVAX, a polysaccharide conjugate vaccine against *S. aureus* infection: from the lab bench to phase III clinical trials," *Vaccine*, 880-887, 2004.
- Field and Smith, "The Coagulase Test for Staphylococci", *J. Comp. Pathol.*, 55:63-69, 1945.
- Fortune et al., "Mutually dependent secretion of proteins required for mycobacterial virulence", *Proc. Natl. Acad. Sci. USA*, 102(30):10676-10681, 2005.
- Foster, "Immune Evasion by Staphylococci", *Nat. Rev. Microbiol.*, 3:948-958, 2005.
- Fournier et al., "Purification and Characterization of *Staphylococcus aureus* Type 8 Capsular Polysaccharide", *Infect. Immun.*, 45(1):87-93, 1984.
- Friedrich et al., "Staphylocoagulase is a prototype for the mechanism of cofactor-induced zymogen activation," *Nature*, 425:535-539, 2003.
- Galan and Collmer, "Type III Secretion Machines: Bacterial Devices for Protein Delivery into Host Cells", *Science*, 284:1322-1328, 1999.
- Garcia-Lara et al., "*Staphylococcus aureus*: the search for novel targets," *Drug Discovery Today*, 10:643-651, 2005.
- GenBank Accession No. AY032850, "*Staphylococcus aureus* secreted von Willebrand factor-binding protein (vwb) gene, complete cds," 2002.
- GenBank Accession No. CAC80837, "*Staphylococcus aureus*," 2003.
- GenBank Accession No. AAA26498 (gi52953), "EryG [Saccharopolyspora erythraea NRRL 2338]," 1991.
- GenBank Accession No. AAK52333, "Secreted von Willebrand factor-binding protein [*Staphylococcus aureus* subsp. *Aureus* str. Newman]," 2001.

(56) **References Cited**

## OTHER PUBLICATIONS

- GenBank Accession No. AJ249487, "*Staphylococcus epidermidis* aap gene for accumulation-associated protein, strain RP62A," 1999.
- GenBank Accession No. CAC80837, "Autolysin [*Staphylococcus aureus*]," 2003.
- GenBank Accession No. COL (YP\_186036.1) (gi57650272), "Alpha-hemolysin precursor [*Staphylococcus aureus* subsp. *Aureus* COL]," 2005.
- GenBank Accession No. DD088871 "A Method for identification, isolation and production of antigens to a specific pathogen," Oct. 14, 2004.
- GenBank Accession No. DD120800, "*Staphylococcus aureus* Proteins and Nucleic Acids," Jan. 27, 2005.
- GenBank Accession No. DD120801, "*Staphylococcus aureus* Proteins and Nucleic Acids," Jan. 27, 2005.
- GenBank Accession No. JH1 (YP\_001316387.1) (gi50393712), "beta-channel forming cytolysin [*Staphylococcus aureus* subsp. *Aureus* JH1]," 2007.
- GenBank Accession No. JH9 (YP\_001246598.1) (gi148267655), "beta-channel forming cytolysin [*Staphylococcus aureus* subsp. *aureus* JH9]," 2007.
- GenBank Accession No. MSSA476 (YP\_043222.1) (gi49486001), "alpha-hemolysin precursor [*Staphylococcus aureus* subsp. *aureus* MSSA476]," 2004.
- GenBank Accession No. Mu50 (NP\_371687.1) (gi5924153), "alpha-hemolysin precursor [*Staphylococcus aureus* subsp. *aureus* Mu50]," 2001.
- GenBank Accession No. MW2 (NP\_645861.1) (gi21282773), "alpha-hemolysin [*Staphylococcus aureus* subsp. *aureus* MW2]," 2002.
- GenBank Accession No. N315 (NP\_374279.1) (gi150393712), "alpha-hemolysin [*Staphylococcus aureus* subsp. *aureus* N315]," 2001.
- GenBank Accession No. NCTC8325 (YP\_499665.1) (gi88194865), "alpha-hemolysin precursor [*Staphylococcus aureus* subsp. *aureus* NCTC 8325]," 2006.
- GenBank Accession No. Newman (YP\_001332107.1) (gi151221285), "alpha-hemolysin precursor [*Staphylococcus aureus* subsp. *aureus* str. Newman]," 2007.
- Genbank Accession No. NP\_371653, "Iron-regulated cell wall-anchored protein SirH [*Staphylococcus aureus* subsp. *Aureus* Mu50]," 2001.
- Genbank Accession No. NP\_371654, "Cell surface protein [*Staphylococcus aureus* subsp. *Aureus* Mu50]," 2001.
- GenBank Accession No. NP\_372518, "Anti repressor [*Staphylococcus aureus* subsp. *aureus* Mu50]," 2001.
- Genbank Accession No. NP\_373773, "Ser-Asp rich fibrinogen binding, bone sialoprotein-binding protein [*Staphylococcus aureus* subsp. *Aureus* N315]," 2001.
- Genbank Accession No. NP\_373774, "Ser-Asp rich fibrinogen binding, bone sialoprotein-binding protein [*Staphylococcus aureus* subsp. *Aureus* N315]," 2001.
- Genbank Accession No. Q99WT7, "RecName:Full= Virulence Factor esxB," 2001.
- Genbank Accession No. Q99WU4, "RecName:Full= Virulence Factor esxA," 2001.
- GenBank Accession No. USA300 (YP\_493756.1) (gi151221285), "alpha-hemolysin precursor [*Staphylococcus aureus* subsp. *aureus* USA300\_FPR3757]," 2006.
- Gomez et al., "*Staphylococcus aureus* protein A activates TNFR1 signaling through conserved IgG binding domains," *J. Biol. Chem.*, 281:20190-20196, 2006.
- Gomez et al., "Mechanisms of Signal Transduction: *Staphylococcus aureus* Protein A Activates TNFR1 Signaling through Conserved IgG Binding Domains", *J. Biol. Chem.* 281(29):20190-20196, 2006.
- Gomez et al., "*Staphylococcus aureus* protein A activates TACE through EGFR-dependent signaling", *EMBO J.* 26:701-709, 2007.
- Gomez et al., "*Staphylococcus aureus* protein A induces airway epithelial inflammatory responses by activating TNFR1", *Nature Med.* 10(8):842-848, 2004.
- Goodyear and Silverman, "Death by a B Cell Superantigen: In Vivo VH-targeted Apoptotic Supraclonal B Cell Deletion by a Staphylococcal Toxin", *J. Exp. Med.*, 197(9):1125-1139, 2003.
- Goodyear and Silverman, "Staphylococcal toxin induced preferential and prolonged in vivo deletion of innate-like B lymphocytes", *Proc. Nat. Acad. Sci. USA*, 101(31):11392-11397, 2004.
- Gouaux et al., "alpha-Hemolysin, gamma-hemolysin, and leukocidin from *Staphylococcus aureus*: distant in sequence but similar in structure," *Protein Sci.*, 6:2631-2635, 1997.
- Gouaux, "alpha-Hemolysin from *Staphylococcus aureus*: An archetype of beta-barrel, channel-forming toxins," *Journal of Structural Biology*, 121:110-122, 1998.
- Gouda et al., "NMR study of the interaction between the B domain of staphylococcal protein A and the Fc portion of immunoglobulin G," *Biochemistry*, 37:129-36, 1998.
- Gouda et al., "Three-Dimensional Solution Structure of the B Domain of Staphylococcal Protein A: Comparisons of the Solution and Crystal Structures", *Biochemistry*, 31(40):9665-72, 1992.
- Graille et al., "Crystal structure of a *Staphylococcus aureus* protein A domain complexed with the Fab fragment of a human IgM antibody: Structural basis for recognition of B-cell receptors and superantigen activity", *Proc. Nat. Acad. Sci. USA* 97(10):5399-5404, 2000.
- Greenspan and Di Cera, "Defining epitopes: It's not as easy as it seems," *Nature Biotechnology*, 17:936-937, 1999.
- Grigg et al., "Haem recognition by a *Staphylococcus aureus* NEAT domain," *Mol. Microbiol.*, 63:139-149, 2007.
- Guinn et al., "Individual RD1-region genes are required for export of ESAT-6/CFP-10 and for virulence of *Mycobacterium tuberculosis*," *Mol. Microbiol.*, 51 (2): 359-370, 2004.
- Guss et al., "Region X, the cell-wall-attachment part of staphylococcal protein A", *Eur. J. Biochem.* 138:413-420, 1984.
- Harboe et al., "Evidence for occurrence of the ESAT-6 protein in *Mycobacterium tuberculosis* and virulent *Mycobacterium bovis* and for its absence in *Mycobacterium bovis* BCG," *Infect. Immun.*, 64: 16-22, 1996.
- Harlow and Lane, in: *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Inc., pp. 23-25 and 27-33, 1988.
- Hartleib et al., "Protein A is the von Willebrand factor binding protein on *Staphylococcus aureus*", *Blood* 96:2149-2156, 2000.
- Hasegawa and Clemente, "Virulence and immunity of *Staphylococcus aureus* BB and certain deficient mutants," *Infection and Immunity*, 22(2):473-479, 1978.
- Hauck et al., "Sticky connections: extracellular matrix protein recognition and integrin-mediated cellular invasion by *Staphylococcus aureus*," *Curr. Opinion Microbiol.*, 9:5-11, 2006.
- Hoffman et al., "Chicken anti-protein A prevents *Staphylococcus aureus* protein A from binding to human and rabbit IgG in immunoassays and eliminates most false positive results," *J. Immunol. Methods*, 198(1):67-77, 1996. (Abstract only).
- Holtfreter et al., "Human immune proteome in experimental colonization with *Staphylococcus aureus*," 16(11):1607-1614, 2009.
- Houghten et al., "Relative importance of position and individual amino acid residues in peptide antigen-antibody interactions: Implications in the mechanism of antigenic drift and antigenic shift," In: *New Approaches to Immunization, Vaccines 86*, Cold Spring Harbor Laboratory Press, Inc. pp. 21-25, 1986.
- Hsu et al., "Repeated neonatal handling with maternal separation permanently alters hippocampal GABAA receptors and behavioral stress responses," *PNAS*, 100:12420-12425, 2003.
- Hsu et al., "The primary mechanism of attenuation of bacillus Calmette-Guerin is a loss of secreted lytic function required for invasion of lung interstitial tissue", *Proc. Natl. Acad. Sci. USA*, 100(21):12420-12425, 2003.
- Hume et al., "Immunization with alpha-toxin toxoid protects the cornea against tissue damage during experimental *Staphylococcus aureus* keratitis," *Infect. Immun.*, 68(10):6052-6055, 2000.
- Iaschenko et al., "Changes in the peripheral blood lymphocytes after immunization and its effects on the course of experimental inflammatory process in the lung" *Zh Mikrobiol Epidemiol Immunobiol.*, 4:88-92, 1978. (English Abstract).

(56)

## References Cited

## OTHER PUBLICATIONS

- Jansson et al., "All individual domains of staphylococcal protein A show Fab binding", *FEMS Immunol. Med. Microbiol.* 20:69-78, 1998.
- Jensen, "A Normally Occurring Staphylococcus Antibody in Human Serum", *Acta Path. Microbiol. Scand.* 44:421-428, 1958.
- Johnson et al., "Iron-regulated biofilm formation in *Staphylococcus aureus* Newman requires ica and the secreted protein Emp", *Infect. Immun.*, 76(4):1756-65, 2008.
- Jonsson et al., "Virulence of *Staphylococcus aureus* in a Mouse Mastitis Model: Studies of Alpha Hemolysin, Coagulase, and Protein A as Possible Virulence Determinants with Protoplast Fusion and Gene Cloning", *Infection and Immunity*, 49(3):765-769, 1985.
- Josefsson et al., "Protection against experimental *Staphylococcus aureus* arthritis by vaccination with clumping factor A, a novel virulence determinant", *J. Infect. Dis.*, 184 (2): 1572-1580, 2001.
- Jursch et al., "Histidine residues near the N terminus of staphylococcal alpha-toxin as reporters of regions that are critical for oligomerization and pore formation," *Infection and Immunity*, 62(6):2249-2256, 1994.
- Kelly, "Immunotherapy against antibiotic-resistant bacteria: the Russian experience with an antistaphylococcal hyperimmune plasma and immunoglobulin," *Microbes and Infection*, 2:1383-1392, 2000.
- Kennedy et al., "Targeting of alpha-hemolysin by active or passive immunization decreases severity of USA300 skin infections in a mouse model," *J. Infect. Dis.*, 202(7):1050-1058, 2010.
- Kennedy et al., "Epidemic community-associated methicillin-resistant *Staphylococcus aureus*: Recent clonal expansion and diversification", *Proc. Natl. Acad. Sci. USA* 105(4):1327-1332, 2008.
- Kim and Schneewind, "Development of Non-toxogenic Protein A vaccine against MRSA" Gordon Conference Poster Presentation, Staphylococcal diseases, Aug. 31, 2009.
- Kim et al., "IsdA and IsdB antibodies protect mice against *Staphylococcus aureus* abscess formation and lethal challenge," *Vaccine*, 28(38):6382-6392, 2010.
- Kim et al., "Nontoxic protein A vaccine for methicillin-resistant *Staphylococcus aureus* infections in mice," *J. Exp. Med.*, 207(9):1863-1870, 2010.
- Klevens et al., "Invasive methicillin-resistant *Staphylococcus aureus* infections in the United States," *JAMA*, 298:1763-1771, 2007.
- Krishnasastri et al., "Surface labeling of key residues during assembly of the transmembrane pore formed by staphylococcal alpha-hemolysin," *FEBS Letters*, 356:66-71, 1994.
- Kuklin et al., "A Novel *Staphylococcus aureus* Vaccine: Iron Surface Determinant B Induces Rapid Antibody Responses in Rhesus Macaques and Specific Increased Survival in a Murine *S. aureus* Sepsis Model", *Infect. Immun.*, 74(4):2215-23, 2006.
- Kuroda et al., "Whole Genome sequencing of methicillin-resistant *Staphylococcus aureus*," *Lancet*, 357 (9264): 1225-1240, 2001.
- Lagergard et al., "Determination of Neutralizing Antibodies and Specific Immunoglobulin Isotype Levels in Infants after Vaccination against Diphtheria", *Eur. J. Clin. Microbiol. Infect. Dis.*, 11(4):341-345, 1992.
- Lam et al., "Abscess-Forming Factor(s) Produced by *Staphylococcus Aureus*", *J. Bacteriol.*, 86:87-91, 1963.
- Lancefield, "Current knowledge of type-specific M antigens of group A streptococci," *J. Immunol.*, 89:307-313, 1962.
- Lancefield, "The antigenic complex of *Streptococcus haemolyticus*. I. Demonstration of a type-specific substance in extracts of *Streptococcus haemolyticus*," *J. Exp. Med.*, 47:91-103, 1928.
- Lee et al., "Development of antistaphylococcal vaccines," *Current Infectious Disease Reports*, 3:517-524, 2001.
- Lee, "The prospects for developing a vaccine against *Staphylococcus aureus*", *Trends Microbiol.* 4(4):162-166, 1996.
- Lee, Jean C., Harvard Medical School "S. aureus vaccine development," available online at [www.ischemo.org/pdf/Lee.pdf](http://www.ischemo.org/pdf/Lee.pdf), accessed Aug. 13, 2010.
- Lindsay et al., "Microarrays reveal that each of the ten dominant lineages of *Staphylococcus aureus* has a unique combination of surface-associated and regulatory genes," *J. Bacteriol.*, 188:669-676, 2006.
- Liu et al., "Direct heme transfer from IsdA to IsdC in the iron-regulated surface determinant (Isd) heme acquisition system of *Staphylococcus aureus*," *J. Biol. Chem.*, 283:6668-6676, 2008.
- Lowy, "Staphylococcus aureus Infections", *New Engl. J. Med.*, 339(8):520-532, 1998.
- MacGurn et al., "A non-RD1 gene cluster is required for Snm secretion in *Mycobacterium tuberculosis*", *Mol. Microbiol.*, 57(6):1653-1663, 2005.
- Madden et al., "Cytolysin-mediated translocation (CMT): a functional equivalent of type III secretion in gram-positive bacteria," *Cell*, 104 (1): 143-152, 2001.
- Mahairas et al., "Molecular analysis of genetic differences between *Mycobacterium bovis* BCG and virulent *M. bovis*," *J. Bacteriol.*, 178 (5): 1274-1282, 1996.
- Maione et al., "Identification of a universal Group B streptococcus vaccine by multiple genome screen," *Science*, 309 (5731):148-150, 2005.
- Maira-Litran et al., "Comparative opsonic and protective activities of *Staphylococcus aureus* conjugate vaccines containing native or deacetylated Staphylococcal Poly-N-acetyl-beta-(1-6)-glucosamine," *Infect. Immun.* 73 (10): 6762, 2005.
- Mamo et al., "Vaccination against *Staphylococcus aureus* mastitis: immunological response of mice vaccinated with fibronectin-binding protein (FnBP-A) to challenge with *S. aureus*," *Vaccine*, 12:988-992, 1994.
- Manolova et al., "The creation of specific immunity to staphylococcal infection in newborn infants by the intranasal administration of absorbed staphylococcal anatoxin," *Zh Mikrobiol Epidemiol Immunobiol.*, 8:64-7, 1989. (Russian Publication. English Abstract).
- Marraffini and Schneewind, "Anchor structure of staphylococcal surface proteins. V. Anchor structure of the sortase B substrate IsdC," *J. Biol. Chem.*, 280:16263-16271, 2005.
- Mazmanian et al., "An iron-regulated sortase-enzyme anchors a class of surface protein during *Staphylococcus aureus* pathogenesis," *Proc. Natl. Acad. Sci. USA*, 99:2293-2298, 2002.
- Mazmanian et al., "Passage of heme-iron across the envelope of *Staphylococcus aureus*," *Science*, 299:906-909, 2003.
- Mazmanian et al., "Sortase-catalysed anchoring of surface proteins to the cell wall of *Staphylococcus aureus*", *Mol. Microbiol.*, 40(5):1049-1057, 2001.
- Mazmanian et al., "Staphylococcus aureus sortase mutants defective in the display of surface proteins and in the pathogenesis of animal infections", *Proc. Natl. Acad. Sci. USA*, 97(10):5510-5515, 2000.
- Mazmanian et al., "Staphylococcus aureus Sortase, an Enzyme that Anchors Surface Proteins to the Cell Wall", *Science*, 285(5428):760-3, 1999.
- McElroy et al., "Alpha-toxin damages the air-blood barrier of the lung in a rat model of *Staphylococcus aureus*-induced pneumonia," *Infect. Immun.*, 67(10):5541-5544, 1999.
- McLaughlin et al., "A *Mycobacterium* ESX-1-Secreted Virulence Factor with Unique Requirements for Export", *PLoS Pathog.*, 3(8):1051-1061, 2007.
- Mempel et al., *The Journal of Investigative Dermatology* 111(3): 452-456, 1998.
- Mendoza et al., "Identification of staphylococcus species by 16S-23S rDNA intergenic spacer PCR analysis," *International Journal of Systematic Bacteriology*, 48:1049-1055, 1998.
- Menestrina et al., "Mode of action of beta-barrel pore-forming toxins of the staphylococcal alpha-hemolysin family," *Toxicon*, 39:1661-1672, 2001.
- Menzies and Kernodle, "Passive immunization with antiserum to a nontoxic alpha-toxin mutant from *Staphylococcus aureus* is protective in a murine model," *Infection and Immunity*, 64(5): 1839-1841, 1996.
- Menzies and Kernodle, "Site-directed mutagenesis of the alpha-toxin gene of *Staphylococcus aureus*: Role of histidines in toxin activity in vitro and in a murine model," *Infection and Immunity*, 62(5):1843-1847, 1994.

(56)

## References Cited

## OTHER PUBLICATIONS

- Mills et al., "Yersinia enterocolitica induces apoptosis in macrophages by a process requiring functional type III secretion and translocation mechanisms and involving YopP, presumably acting as an effector protein," *PNAS*, 94 (23): 12638-12643, 1997.
- Moreau et al., "Structure of the type 5 capsular polysaccharide of *Staphylococcus aureus*", *Carbohydrate Res.*, 201(2):285-297, 1990.
- Moreillon et al., "Role of *Staphylococcus aureus* Coagulase and Clumping Factor in Pathogenesis of Experimental Endocarditis", *Infect. Immun.*, 63(12):4738-4743, 1995.
- Muryoi et al., "Demonstration of the iron-regulated surface determinant (Isd) heme transfer pathway in *Staphylococcus aureus*," *J. Biol. Chem.*, 283:28125-28136S, 2008.
- Navarre et al., "Multiple enzymatic activities of the murein hydrolase from staphylococcal phage phi11. Identification of a D-alanyl-glycine endopeptidase activity," *J. Biol. Chem.*, 274:15847-15856, 1999.
- Ni Eidhin et al., "Clumping factor B (ClfB), a new surface-located fibrinogen-binding adhesin of *Staphylococcus aureus*," *Mol. Microbiol.*, 30 (2): 245-257, 1998.
- Nitsche-Smitz et al., "Invasion mechanisms of Gram-positive pathogenic cocci," *Thrombosis and Haemostasis*, 98(3):488-496, 2007.
- Nordhaug et al., "A field trial with an experimental vaccine against *Staphylococcus aureus* mastitis in cattle. 2. Antibody response," *J. Dairy Sci.*, 77:1276-1284, 1994.
- Novick, "Autoinduction and signal transduction in the regulation of staphylococcal virulence", *Mol. Microbiol.*, 48(6):1429-1449, 2003.
- O'Brien et al., "Multiple mechanisms for the activation of human platelet aggregation by *Staphylococcus aureus*: roles for the clumping factors ClfA and ClfB, the serine-aspartate repeat protein SdrE and protein A", *Mol. Microbiol.* 44(4):1033-1044, 2002.
- O'Seaghdha et al., "Staphylococcus aureus protein A binding to von Willebrand factor A1 domain is mediated by conserved IgG binding regions", *FEBS J.* 273:4831-4841, 2006.
- Office Action in Singapore Patent Application No. 201107080-2 dated Sep. 19, 2013.
- Office Communication issued in European Patent Application No. 07 840 104.9, dated May 19, 2009.
- Office Communication issued in European Patent Application No. 88 828 277, dated Apr. 7, 2011.
- Office Communication issued in U.S. Appl. No. 12/842,811, dated Feb. 29, 2012.
- Office Communication issued in U.S. Appl. No. 12/161,315, dated Mar. 8, 2011.
- Office Communication issued in U.S. Appl. No. 12/161,315, dated Jun. 14, 2010.
- Office Communication issued in U.S. Appl. No. 12/161,315, dated Apr. 1, 2010.
- Office Communication issued in U.S. Appl. No. 12/161,315, dated Nov. 18, 2011.
- O'Reilly et al., "Cryptic alpha-toxin gene in toxic shock syndrome and septicaemia strains of *Staphylococcus aureus*," *Mol. Microbiol.*, 4:1947-1955, 1990.
- O'Reilly et al., "Inactivation of the alpha-haemolysin gene of *Staphylococcus aureus* 8325-4 by site-directed mutagenesis and studies on the expression of its haemolysins," *Microb. Pathog.*, 1:125-138, 1986.
- Overheim et al., "LcrV plague vaccine with altered immunomodulatory properties," *Infect. Immun.*, 73:5152-5159, 2005.
- Pal et al., "Design of potent, non-toxic antimicrobial agents based upon the structure of the frog skin peptide, pseudin-2," *Regul. Pept.*, 129(1-3):85-91, 2005.
- Pallen, "The ESAT-6/WXG100 superfamily—and a new Gram-positive secretion system?," *Trends Microbiol.*, 10(5):209-212, 2002.
- Palmqvist et al., "Bacterial cell wall-expressed protein A triggers supraclonal B-cell responses upon in vivo infection with *Staphylococcus aureus*", *Microbes. Infect.*, 7:1501-1511, 2005.
- Pancholi and Fischetti, "A major surface protein on group A streptococci is a glyceraldehyde-phosphate-3-dehydrogenase with multiple binding activity," *J. Exp. Med.*, 176 (2): 415-426, 1992.
- Panizzi et al., "Fibrinogen Substrate Recognition by Staphylocoagulase-(Pro) thrombin Complexes", *J. Biol. Chem.*, 281(2):1179-1187, 2006.
- Pankey et al., "Evaluation of Protein A and a Commercial Bacterin as Vaccines Against *Staphylococcus aureus* Mastitis by Experimental Challenge", *J. Dairy Sci.* 68:726-731, 1985.
- Park et al., "Immunogenicity of alpha-toxin, capsular polysaccharide (CPS) and recombinant fibronectin-binding protein (r-FnBP) of *Staphylococcus aureus* in rabbit," *J. Vet. Med. Sci.*, 61(9):995-1000, 1999.
- PCT International Preliminary Report on Patentability issued in International application No. PCT/US10/29959, dated Oct. 13, 2011.
- PCT International Search Report and Written Opinion issued in International application No. PCT/US10/29959, dated Apr. 11, 2011.
- PCT International Search Report and Written Opinion issued in International App. No. PCT/US2007/060720, dated Jun. 9, 2008.
- PCT International Search Report and Written Opinion issued in International App. No. PCT/US2009/059648, dated Feb. 16, 2010.
- PCT International Search Report issued in International App. No. PCT/US2008/074849, dated Dec. 9, 2008.
- PCT Invitation to Pay Additional Fees, issued in International App. No. PCT/US2007/060720, dated Apr. 9, 2008.
- PCT Invitation to Pay Additional Fees issued in International application No. PCT/US10/29959, dated Feb. 4, 2011.
- PCT Invitation to Pay Additional Fees issued in International application No. PCT/US11/42845, dated Nov. 10, 2011.
- Philipp et al., "Physical mapping of *Mycobacterium bovis* BCG Pasteur reveals differences from the genome map of *Mycobacterium tuberculosis* H37Rv and from *M. bovis*," *Microbiology*, 142: 3135-3145, 2003.
- Phonimdaeng et al., "The coagulase of *Staphylococcus aureus* 8325-4. Sequence analysis and virulence of site-specific coagulase-deficient mutants", *Mol. Microbiol.*, 4(3):393-404, 1990.
- Pilpa et al., "Functionally distinct NEAT (NEAr Transporter) domains within the *Staphylococcus aureus* IsdH/HarA protein extract heme from methemoglobin," *J. Biol. Chem.*, 284:1166-1176, 2009.
- Pilpa et al., "Solution structure of the NEAT (NEAr Transporter) domain from IsdH/HarA: the human hemoglobin receptor in *Staphylococcus aureus*," *J. Mol. Biol.*, 360:435-447, 2006.
- Poole-Warren et al., "Vaccination for prevention of CAPD associated staphylococcal infection: results of a prospective multicenter clinical trial," *Clin. Nephrol.*, 35(5):198-206, 1991.
- Projan et al., "Staphylococcal vaccines and immunotherapy: to dream the impossible dream?" *Curr. Opin. Pharmacol.*, 6:473-479, 2006.
- Pym et al., "Recombinant BCG exporting ESAT-6 confers enhanced protection against tuberculosis," *Nature Medicine*, 9 (5): 533-539, 2003.
- Pym et al., "Loss of RD1 contributed to the attenuation of the live tuberculosis vaccines *Mycobacterium bovis* BCG and *Mycobacterium microti*", *Molecular Microbiology*, 46(3):709-717, 2002.
- Raedler et al., "Serologic assay to quantify human immunoglobulin antibodies to *Staphylococcus aureus* iron surface determinant B antigen," *Clin. Vaccine Immunol.*, 16(5):739-48, 2009.
- Ragle and Wardenburg, "Anti-alpha-hemolysin monoclonal antibodies mediate protection against *Staphylococcus aureus* pneumonia," *Infection and Immunity*, 77(7):2712-2718, 2009.
- Ragle et al., "Prevention and treatment of *Staphylococcus aureus* pneumonia with a beta-cyclodextrin derivative," *Antimicrobial Agents and Chemotherapy*, 54(1):298-304, 2010.
- Renshaw et al., "Conclusive evidence that the major T-cell antigens of the *Mycobacterium tuberculosis* complex ESAT-6 and CFP=10 form a tight, 1:1 complex and characterization of the structural properties of ESAT-6, CFP-10, and the ESAT-6CFP-10 complex. Implications for pathogenesis and virulence," *J. of Biol. Chem.*, 277 (24): 21598-21603, 2002.
- Renshaw et al., "Structure and function of the complex formed by the tuberculosis virulence factors CFP-10 and ESAT-6," *Embo Journal*, 24 (14): 2491-2498, 2005.
- Rivera et al., "Fibrinogen-binding proteins of gram-positive bacteria," *Thromb. Haemost.*, 98:503-511, 2007.

(56)

## References Cited

## OTHER PUBLICATIONS

- Roben et al., "VH3 Family Antibodies Bind Domain D of Staphylococcal Protein A1", *J. Immunol.* 154:6437-6445, 1995.
- Rosch and Caparon, "A microdomain for protein secretion in Gram-positive bacteria," *Science*, 304: 1513-1515, 2004.
- Rose et al., "Mediator generation and signaling events in alveolar epithelial cells attacked by *S. aureus* alpha-toxin," *Am. J. Physiol. Lung Cell Mol. Physiol.*, 282:L207-L214, 2002.
- Said-Salim et al., "Community-Acquired Methicillin-Resistant *Staphylococcus aureus*: An Emerging Pathogen", *Infect. Control Hosp. Epidemiol.* 24(6):451-455, 2003.
- Schaffer et al., "Immunization with *Staphylococcus aureus* clumping factor B, a major determinant in nasal carriage, reduces nasal colonization in a murine model," *Infect. Immun.*, 74 (4): 2145-2153, 2006.
- Schneewind et al., "Cell wall sorting signals in surface proteins of gram-positive bacteria," *EMBO*, 12(12):4803-4811, 1993.
- Schneewind et al., "Structure of the cell wall anchor of surface proteins in *Staphylococcus aureus*," *Science*, 268:103-6, 1995.
- Schneewind et al., "Sorting of Protein A to the Staphylococcal Cell Wall", *Cell* 70:267-281, 1992.
- Scriba et al., "The *Staphylococcus aureus* Eap protein activates expression of proinflammatory cytokines," *Infect. Immun.*, 76(5):2164-2168, 2008.
- Search Report and Written Opinion in PCT/US11/42845 dated Feb. 10, 2012.
- Search Report and Written Opinion in Singapore Application No. 2012091104 dated Feb. 27, 2014.
- Seeger et al., "Staphylococcal alpha-toxin elicits hypertension in isolated rabbit lungs. Evidence for thromboxane formation and the role of extracellular calcium," *J. Clin. Invest.*, 74, 849-858, 1984.
- Seeger et al., "Staphylococcal alpha-toxin-induced vascular leakage in isolated perfused rabbit lungs," *Lab. Invest.*, 63:341-349, 1990.
- Sequence 2913 from Patent EP 1829892, NCBI accession No. CS710373, Sep. 5, 2007.
- Sequence 2913 from Patent WO02094868, NCBI accession No. AX619950, Nov. 28, 2002.
- Sequence 2915 from Patent EP1829892, NCBI accession No. CS710375, Sep. 5, 2007.
- Sequence 2915 from Patent WO02094868, NCBI accession No. AX619952, Nov. 28, 2002.
- Sequence 42 from Patent EP1616876, NCBI accession No. CS252757, Jan. 18, 2006.
- Sequence 42 from Patent EP1630172, NCBI accession No. CS274094, Mar. 1, 2006.
- Sequence 42 from Patent WO02059148, NCBI accession No. AX583665, Aug. 1, 2002.
- Sequence 785 from patent U.S. Pat. No. 6,593,114, NCBI accession No. AR354667, Jul. 15, 2003.
- Sequence 785 from patent U.S. Pat. No. 6,737,248, NCBI accession No. AR536223, May 18, 2004.
- Sequence 94 from patent U.S. Pat. No. 6,348,582, NCBI accession No. AR194545, Feb. 19, 2002.
- Sharp et al., "Crystal structure of the heme-IsdC complex, the central conduit of the Isd iron/heme uptake system in *Staphylococcus aureus*," *J. Biol. Chem.*, 282:10625-10631, 2007.
- Shaw et al., "The role and regulation of the extracellular proteases of *Staphylococcus aureus*", *Microbiology*, 150:217-228, 2004.
- Sheagren, "*Staphylococcus aureus*: The Persistent Pathogen", *N. Engl. J. Med.* 310(21):1368-1373, 1984.
- Shopsin et al., "Evaluation of protein A gene polymorphic region DNA sequencing for typing of *Staphylococcus aureus* strains," *J. Clin. Microbiol.*, 37(11):3556-63, 1999.
- Sibbald et al., "Mapping the Pathways to Staphylococcal Pathogenesis by Comparative Secretomics", *Microbiol. Mol Biol. Rev.*, 70(3):755-788, 2006.
- Silverman and Goodyear, "Confounding B-cell defences: lessons from a staphylococcal superantigen," *Nat. Rev. Immunol.*, 6(6):465-75, 2006.
- Sjodahl, "Repetitive Sequences in Protein A from *Staphylococcus aureus*", *Eur. J. Biochem.* 73:343-351, 1977.
- Sjoquist et al., "Protein A Isolated from *Staphylococcus aureus* after Digestion with Lysostaphin", *Eur. J. Biochem.* 29 :572-578, 1972.
- Skaar et al., "Iron-regulated surface determinants (Isd) of *Staphylococcus aureus*: stealing iron from heme," *Microbes Infect.*, 6:390-397, 2004.
- Skaar et al., "Iron-source preference of *Staphylococcus aureus* infections," *Science*, 305:1626-1628, 2004.
- Skaar et al., "IsdG and IsdI, heme degrading enzymes in the cytoplasm of *Staphylococcus aureus*," *J. Biol. Chem.*, 279:436-443, 2004.
- Smith et al., "The Role of Coagulase in Staphylococcal Infections", *Brit. J. Exp. Pathol.*, 28:57-67, 1947.
- Song et al., "Structure of staphylococcal alpha-hemolysin, a heptameric transmembrane pore," *Science*, 274:1859-1866, 1996.
- Sorenson et al., "Purification and characterization of a low-molecular-mass T-cell antigen secreted by *Mycobacterium tuberculosis*," *Infect. Immun.*, 63 (5): 1710-1717, 1995.
- Stanley et al., "Acute infection and macrophage subversion by *Mycobacterium tuberculosis* require a specialized secretion system", *Proc. Natl. Acad. Sci. USA*, 100(22):13001-13006, 2003.
- Stranger-Jones et al., "Vaccine assembly from surface proteins of *Staphylococcus aureus*", *Proc. Nat. Acad. Sci. USA*, 103(45):16942-16947, 2006.
- Studier et al., "Use of T7 RNA polymerase to direct expression of cloned genes," *Methods Enzymol.* 185:60-89 1990.
- Stugard et al., "A 101-kilodalton heme-binding protein associated with congo red binding and virulence of *Shigella flexneri* and enteroinvasive *Escherichia coli* strains," *Infect. Immun.*, 57:3534-3539, 1989.
- Suttorp and Habben, "Effect of staphylococcal alpha-toxin on intracellular Ca<sup>2+</sup> in polymorphonuclear leukocytes," *Infect. Immun.*, 56:2228-34, 1988.
- Tenover et al., "Characterization of a strain of community-associated methicillin-resistant *Staphylococcus aureus* widely disseminated in the United States," *J. Clin. Microbiol.*, 44:108-118, 2006.
- Thammavongsa et al., "*Staphylococcus aureus* synthesizes adenosine to escape host immune responses," *J. Exp. Med.*, 206(11):2417-2427, 2009.
- Tiedemann et al., "Isolation of HLA-DR1.(staphylococcal enterotoxin A)<sub>2</sub> trimers in solution," *PNAS*, 92(26):12156-9, 1995.
- Tollersrud et al., "Antibody responses in sheep vaccinated against *Staphylococcus aureus* mastitis: A comparison of two experimental vaccines containing different adjuvants," *Veterinary Research Communications*, 26:587-600, 2002.
- Ton-That et al., "Fatigue characterization of a hydroxyapatite-reinforced polyethylene composite. II. Biaxial fatigue," *J. Biomed. Mater Res.*, 51 (3): 461-468, 2000.
- Ton-That et al., "Purification and characterization of sortase, the transpeptidase that cleaves surface proteins of *Staphylococcus aureus* at the LPXTG motif", *Proc. Natl. Acad. Sci. USA*, 96(22):12424-12429, 1999.
- Tones et al., "*Staphylococcus aureus* IsdB is a hemoglobin receptor required for heme-iron utilization," *J. Bacteriol.*, 188:8421-8429, 2006.
- Uhlen et al., "Complete Sequence of the Staphylococcal Gene Encoding Protein A: A gene evolved through multiple duplications", *J. Biol. Chem.* 259(3):1695-1702, 1984.
- U.S. Appl. No. 61/103,196, entitled "Compositions and Methods Related to Bacterial Eap and/or Emp Proteins," by Alice Cheng, filed Oct. 6, 2009.
- U.S. Appl. No. 61/166,432, entitled "Compositions and Methods Related to Protein A (Spa) Variants," by Olaf Schneewind, filed Apr. 3, 2009.
- U.S. Appl. No. 61/170,779, entitled "Compositions and Methods Related to Bacterial Eap and/or Emp Proteins," by Alice Cheng, filed Apr. 20, 2009.
- Valeva et al., "Staphylococcal alpha-toxin: Formation of the heptameric pore is partially cooperative and proceeds through multiple intermediate stages," *Biochemistry*, 36:13298-13304, 1997.

(56)

**References Cited**

## OTHER PUBLICATIONS

- Van Wely et al., "Translocation of proteins across the cell envelope of Gram-positive bacteria", *FEMS Microbiol. Rev.*, 25:437-454, 2001.
- Verkaik et al., "Immunogenicity of toxins using *Staphylococcus aureus* infections," *Clinical Infectious Diseases*, 50:61-8, 2010.
- Villareal et al., "The IsdC protein from *Staphylococcus aureus* uses a flexible binding pocket to capture heme," *J. Biol. Chem.*, 283:31591-31600, 2008.
- Walker and Bayley, "Key residues for membrane binding, oligomerization, and pore forming activity of staphylococcal alpha-hemolysin identified by cysteine scanning mutagenesis and targeted chemical modification," *The Journal of Biological Chemistry*, 270(39):23065-23071, 1995.
- Walker and Bayley, "Restoration of pore-forming activity in staphylococcal alpha-hemolysin by targeted covalent modification," *Protein Engineering*, 8(5):491-495, 1995.
- Walker et al., "An intermediate in the assembly of a pore-forming protein trapped with a genetically-engineered switch," *Chemistry & Biology*, 2:99-105, 1995.
- Wardenburg et al., "Poring over pores: alpha-hemolysin and Panton-Valentine leukocidin in *Staphylococcus aureus* pneumonia," *Nature Medicine*, 13(12):1405-1406, 2007.
- Wardenburg et al., "Vaccines for *Staphylococcus aureus* infections," In: *New Generation Vaccines*, 4<sup>th</sup> edition, Dr. Myron Levine, Ed., Informa Healthcare, Chapter 67, 2009.
- Weichhart et al., "Functional selection of vaccine candidate peptides from *Staphylococcus aureus* whole-genome expression libraries in vitro," *Infect. Immun.*, 71 (8): 4633-4641, 2003.
- Weiss et al., "Effect of srtA and srtB gene expression on the virulence of *Staphylococcus aureus* in animal models of infection", *J. Antimicrob. Chemother.*, 53:480-486, 2004.
- Wilke and Wardenburg, "Role of a disintegrin and metalloprotease 10 in *Staphylococcus aureus* alpha-hemolysin-mediated cellular injury," *PNAS*, 107(30):13473-8. Epub Jul. 12, 2010.
- Wlekinski et al., "Protective effects of active immunization against alpha hemolysin of *Staphylococcus aureus*," *Zentralbl. Veterinarmed B.*, 29(8):596-603, 1982. (German Publication. English summary).
- Wu et al., "*Staphylococcus aureus* IsdG and IsdI, heme degrading enzymes with structural similarity to monooxygenases," *J. Biol. Chem.*, 2004.
- Xie et al., "Suppression of experimental autoimmune encephalomyelitis by extracellular adherence protein of *Staphylococcus aureus*," *J Exp. Med.*, 203(4):985-94, 2006.
- Xu et al., "A unique *Mycobacterium* ESX-1 protein co-secretes with CFP-10/ESAT-6 and is necessary for inhibiting phagosome maturation", *Mol. Microbiol.*, 66(3):787-800, 2007.
- Yanagisawa et al., "Neutralization of staphylococcal exotoxins in vitro by human-origin intravenous immunoglobulin," *J. Infect. Chemother.*, 13:368-372, 2007.
- Yoshida et al., "Induction of resistance with heat-killed compact-type strains of *Staphylococcus aureus* against challenge with the diffuse variant of the Smith strain of *Staphylococcus aureus*," *Infection and Immunity*, 12(5):939-942, 1975.
- Zhang et al., "Construction and expression of fused gene vaccine esat6-cfp10 of *Mycobacterium tuberculosis*," *Disi Junyi Daxue Xuebao*, 26(3):193-195, 2005. (Chinese abstract). publication. English.
- Zhou et al., "An immunogenicity study of a newly fusion protein Cna-FnBP vaccinated against *Staphylococcus aureus* infections in a mice model," *Vaccine*, 24 (22): 4830-4837, 2006.
- Zhu et al., "Pathway for heme uptake from human methemoglobin by the iron-regulated surface determinants system of *Staphylococcus aureus*," *J. Biol. Chem.*, 283:18450-18460, 2008.

\* cited by examiner




FIG. 1A

**B**




FIG. 1B




FIG. 2A





FIG. 2B-2C




FIG. 3




FIG. 4




FIG. 5




FIG. 6A-6B




FIG. 6C-6E




FIG. 7






FIG. 8A-8C




FIG. 9A-9D




FIG. 10




FIG. 11




FIG. 12A-12C

## COMPOSITIONS AND METHODS RELATED TO PROTEIN A (SPA) VARIANTS

This application is a continuation of U.S. patent application Ser. No. 15/702,037 filed Sep. 12, 2017, which is a continuation of U.S. patent application Ser. No. 15/060,861, filed Mar. 4, 2016, which is a continuation of U.S. patent application Ser. No. 14/466,514, filed Aug. 22, 2014, now U.S. Pat. No. 9,315,554, which is a continuation of U.S. patent application Ser. No. 13/807,598, filed Mar. 19, 2013, now U.S. Pat. No. 8,821,894, which is a national phase application under 35 U.S.C. § 371 of International Application No. PCT/US2011/042845, filed Jul. 1, 2011, which claims the benefit of priority to U.S. Provisional Patent Application Ser. No. 61/361,218 filed Jul. 2, 2010, and 61/370,725 filed Aug. 4, 2010. The entire contents of each of the above-referenced disclosures are specifically incorporated herein by reference without disclaimer.

This invention was made with government support under AI057153, AI052474, and GM007281 awarded by the National Institutes of Health. The government has certain rights in the invention.

### BACKGROUND OF THE INVENTION

#### I. Field of the Invention

The present invention relates generally to the fields of immunology, microbiology, and pathology. More particularly, it concerns methods and compositions involving bacterial Protein A variants, which can be used to invoke an immune response against the bacteria.

#### II. Background

The number of both community acquired and hospital acquired infections have increased over recent years with the increased use of intravascular devices. Hospital acquired (nosocomial) infections are a major cause of morbidity and mortality, more particularly in the United States, where it affects more than 2 million patients annually. The most frequent infections are urinary tract infections (33% of the infections), followed by pneumonia (15.5%), surgical site infections (14.8%) and primary bloodstream infections (13%) (Emorl and Gaynes, 1993).

The major nosocomial pathogens include *Staphylococcus aureus*, coagulase-negative Staphylococci (mostly *Staphylococcus epidermidis*), *enterococcus* spp., *Escherichia coli* and *Pseudomonas aeruginosa*. Although these pathogens cause approximately the same number of infections, the severity of the disorders they can produce combined with the frequency of antibiotic resistant isolates balance this ranking towards *S. aureus* and *S. epidermidis* as being the most significant nosocomial pathogens.

Staphylococci can cause a wide variety of diseases in humans and other animals through either toxin production or invasion. Staphylococcal toxins are also a common cause of food poisoning, as the bacteria can grow in improperly-stored food.

*Staphylococcus epidermidis* is a normal skin commensal which is also an important opportunistic pathogen responsible for infections of impaired medical devices and infections at sites of surgery. Medical devices infected by *S. epidermidis* include cardiac pacemakers, cerebrospinal fluid shunts, continuous ambulatory peritoneal dialysis catheters, orthopedic devices and prosthetic heart valves.

*Staphylococcus aureus* is the most common cause of nosocomial infections with a significant morbidity and mortality. It is the cause of some cases of osteomyelitis, endocarditis, septic arthritis, pneumonia, abscesses, and toxic shock syndrome. *S. aureus* can survive on dry surfaces, increasing the chance of transmission. Any *S. aureus* infection can cause the staphylococcal scalded skin syndrome, a cutaneous reaction to exotoxin absorbed into the bloodstream. It can also cause a type of septicemia called pyaemia that can be life-threatening. Problematically, Methicillin-resistant *Staphylococcus aureus* (MRSA) has become a major cause of hospital-acquired infections.

*S. aureus* and *S. epidermidis* infections are typically treated with antibiotics, with penicillin being the drug of choice, whereas vancomycin is used for methicillin resistant isolates. The percentage of staphylococcal strains exhibiting wide-spectrum resistance to antibiotics has become increasingly prevalent, posing a threat for effective antimicrobial therapy. In addition, the recent emergence of vancomycin resistant *S. aureus* strain has aroused fear that MRSA strains are emerging and spreading for which no effective therapy is available.

An alternative to antibiotic treatment for staphylococcal infections is under investigation that uses antibodies directed against staphylococcal antigens. This therapy involves administration of polyclonal antisera (WO00/15238, WO00/12132) or treatment with monoclonal antibodies against lipoteichoic acid (WO98/57994).

An alternative approach would be the use of active vaccination to generate an immune response against staphylococci. The *S. aureus* genome has been sequenced and many of the coding sequences have been identified (WO02/094868, EP0786519), which can lead to the identification of potential antigens. The same is true for *S. epidermidis* (WO01/34809). As a refinement of this approach, others have identified proteins that are recognized by hyperimmune sera from patients who have suffered staphylococcal infection (WO01/98499, WO02/059148).

*S. aureus* secretes a plethora of virulence factors into the extracellular milieu (Archer, 1998; Dinges et al., 2000; Foster, 2005; Shaw et al., 2004; Sibbald et al., 2006). Like most secreted proteins, these virulence factors are translocated by the Sec machinery across the plasma membrane. Proteins secreted by the Sec machinery bear an N-terminal leader peptide that is removed by leader peptidase once the pre-protein is engaged in the Sec translocon (Dalbey and Wickner, 1985; van Wely et al., 2001). Recent genome analysis suggests that Actinobacteria and members of the Firmicutes encode an additional secretion system that recognizes a subset of proteins in a Sec-independent manner (Pallen, 2002). ESAT-6 (early secreted antigen target 6 kDa) and CFP-10 (culture filtrate antigen 10 kDa) of *Mycobacterium tuberculosis* represent the first substrates of this novel secretion system termed ESX-1 or Snm in *M. tuberculosis* (Andersen et al., 1995; Hsu et al., 2003; Pym et al., 2003; Stanley et al., 2003). In *S. aureus*, two ESAT-6 like factors designated EsxA and EsxB are secreted by the Ess pathway (ESAT-6 secretion system) (Burts et al., 2005).

The first generation of vaccines targeted against *S. aureus* or against the exoproteins it produces have met with limited success (Lee, 1996). There remains a need to develop effective vaccines against staphylococcal infections. Additional compositions for treating staphylococcal infections are also needed.

### SUMMARY OF THE INVENTION

Protein A (SpA)(SEQ ID NO:33), a cell wall anchored surface protein of *Staphylococcus aureus*, provides for bac-

terial evasion from innate and adaptive immune responses. Protein A binds immunoglobulins at their Fc portion, interacts with the VH3 domain of B cell receptors inappropriately stimulating B cell proliferation and apoptosis, binds to von Willebrand factor A1 domains to activate intracellular clotting, and also binds to the TNF Receptor-1 to contribute to the pathogenesis of staphylococcal pneumonia. Due to the fact that Protein A captures immunoglobulin and displays toxic attributes, the possibility that this surface molecule may function as a vaccine in humans has not been rigorously pursued. Here the inventors demonstrate that Protein A variants no longer able to bind to immunoglobulins, which are thereby removed of their toxigenic potential, i.e., are non-toxigenic, stimulate humoral immune responses that protect against staphylococcal disease.

In certain embodiments the SpA variant is a full length SpA variant comprising a variant A, B, C, D, and/or E domain. In certain aspects, the SpA variant comprises or consists of the amino acid sequence that is 80, 90, 95, 98, 99, or 100% identical to the amino acid sequence of SEQ ID NO:34. In other embodiments the SpA variant comprises a segment of SpA. The SpA segment can comprise at least or at most 1, 2, 3, 4, 5 or more IgG binding domains. The IgG domains can be at least or at most 1, 2, 3, 4, 5 or more variant A, B, C, D, or E domains. In certain aspects the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant A domains. In a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant B domains. In still a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant C domains. In yet a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant D domains. In certain aspects the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant E domains. In a further aspect the SpA variant comprises a combination of A, B, C, D, and E domains in various combinations and permutations. The combinations can include all or part of a SpA signal peptide segment, a SpA region X segment, and/or a SpA sorting signal segment. In other aspects the SpA variant does not include a SpA signal peptide segment, a SpA region X segment, and/or a SpA sorting signal segment. In certain aspects a variant A domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:4. In another aspect a variant B domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:6. In still another aspect a variant C domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:5. In certain aspects a variant D domain comprises a substitution at position(s) 9, 10, 36, and/or 37 of SEQ ID NO:2. In a further aspect a variant E domain comprises a substitution at position(s) 6, 7, 33, and/or 34 of SEQ ID NO:3.

In certain aspects, an SpA domain D variant or its equivalent can comprise a mutation at position 9 and 36; 9 and 37; 9 and 10; 36 and 37; 10 and 36; 10 and 37; 9, 36, and 37; 10, 36, and 37, 9, 10 and 36; or 9, 10 and 37 of SEQ ID NO:2. In a further aspect, analogous mutations can be included in one or more of domains A, B, C, or E.

In further aspects, the amino acid glutamine (Q) at position 9 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an aspartic acid (D), a cysteine (C), a glutamic acid (E), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the glutamine at position 9 can be substituted with an arginine (R). In a further aspect, the glutamine at position 9

of SEQ ID NO:2, or its equivalent, can be substituted with a lysine or a glycine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

In another aspect, the amino acid glutamine (Q) at position 10 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an aspartic acid (D), a cysteine (C), a glutamic acid (E), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the glutamine at position 10 can be substituted with an arginine (R). In a further aspect, the glutamine at position 10 of SEQ ID NO:2, or its equivalent, can be substituted with a lysine or a glycine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

In certain aspects, the aspartic acid (D) at position 36 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an arginine (R), a cysteine (C), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a glutamine (Q), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the aspartic acid at position 36 can be substituted with a glutamic acid (E). In certain aspects, an aspartic acid at position 36 of SEQ ID NO:2, or its equivalent, can be substituted with an alanine or a serine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

In another aspect, the aspartic acid (D) at position 37 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an arginine (R), a cysteine (C), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a glutamine (Q), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the aspartic acid at position 37 can be substituted with a glutamic acid (E). In certain aspects, an aspartic acid at position 37 of SEQ ID NO:2, or its equivalent, can be substituted with an alanine or a serine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

In a particular embodiment the amino at position 9 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 9 of SEQ ID NO:2 is replaced by a glycine. In a further aspect the amino acid at position 9 of SEQ ID NO:2 is replaced by a lysine.

In a particular embodiment the amino at position 10 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 10 of SEQ ID NO:2 is replaced by a glycine. In a further aspect the amino acid at position 10 of SEQ ID NO:2 is replaced by a lysine.

In a particular embodiment the amino at position 36 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 36 of SEQ ID NO:2 is replaced by a serine. In a further aspect the amino acid at position 36 of SEQ ID NO:2 is replaced by an alanine.

In a particular embodiment the amino acid at position 37 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 37 of SEQ ID NO:2 is replaced by a serine. In a further aspect the amino acid at position 37 of SEQ ID NO:2 is replaced by an alanine.

In certain aspects the SpA variant includes a substitution of (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a  $V_H3$  binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to  $V_H3$ . In still further aspects the amino acid sequence of a SpA variant comprises an amino acid sequence that is at least 50%, 60%, 70%, 80%, 90%, 95%, or 100% identical, including all values and ranges there between, to the amino acid sequence of SEQ ID NOs:2-6.

In a further aspect the SpA variant includes (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain D, or at a corresponding amino acid position in other IgG domains, that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a  $V_H3$  binding sub-domain of SpA domain D, or at a corresponding amino acid position in other IgG domains, that disrupts or decreases binding to  $V_H3$ . In certain aspects amino acid residue F5, Q9, Q10, S11, F13, Y14, L17, N28, 131, and/or K35 (SEQ ID NO:2, QQNNFNKDDQSSAFYEILNMPNLNEAQRNG-FIQSLKDDPSQSTNVLGEAKKLNES) of the IgG Fc binding sub-domain of domain D are modified or substituted. In certain aspects amino acid residue Q26, G29, F30, S33, D36, D37, Q40, N43, and/or E47 (SEQ ID NO:2) of the  $V_H3$  binding sub-domain of domain D are modified or substituted such that binding to Fc or  $V_H3$  is attenuated. In further aspects corresponding modifications or substitutions can be engineered in corresponding positions of the domain A, B, C, and/or E. Corresponding positions are defined by alignment of the domain D amino acid sequence with one or more of the amino acid sequences from other IgG binding domains of SpA, for example see FIG. 2A. In certain aspects the amino acid substitution can be any of the other 20 amino acids. In a further aspect conservative amino acid substitutions can be specifically excluded from possible amino acid substitutions. In other aspects only non-conservative substitutions are included. In any event, any substitution or combination of substitutions that reduces the binding of the domain such that SpA toxicity is significantly reduced is contemplated. The significance of the reduction in binding refers to a variant that produces minimal to no toxicity when introduced into a subject and can be assessed using in vitro methods described herein.

In certain embodiments, a variant SpA comprises at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more variant SpA domain D peptides. In certain aspects 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 or more amino acid residues of the variant SpA are substituted or modified—including but not limited to amino acids F5, Q9, Q10, S11, F13, Y14, L17, N28, 131, and/or K35 (SEQ ID NO:2) of the IgG Fc binding sub-domain of domain D and amino acid residue Q26, G29, F30, S33, D36, D37, Q40, N43, and/or E47 (SEQ ID NO:2) of the  $V_H3$  binding sub-domain of domain D. In one aspect of the invention glutamine residues at position 9 and/or 10 of SEQ ID NO:2 (or corresponding positions in other domains) are mutated. In another aspect, aspartic acid residues 36 and/or 37 of SEQ ID NO:2 (or

corresponding positions in other domains) are mutated. In a further aspect, glutamine 9 and 10, and aspartic acid residues 36 and 37 are mutated. Purified non-toxicogenic SpA or SpA-D mutants/variants described herein are no longer able to significantly bind (i.e., demonstrate attenuated or disrupted binding affinity) Fcγ or F(ab)<sub>2</sub>  $V_H3$  and also do not stimulate B cell apoptosis. These non-toxicogenic Protein A variants can be used as subunit vaccines and raise humoral immune responses and confer protective immunity against *S. aureus* challenge. Compared to wild-type full-length Protein A or the wild-type SpA-domain D, immunization with SpA-D variants resulted in an increase in Protein A specific antibody. Using a mouse model of staphylococcal challenge and abscess formation, it was observed that immunization with the non-toxicogenic Protein A variants generated significant protection from staphylococcal infection and abscess formation. As virtually all *S. aureus* strains express Protein A, immunization of humans with the non-toxicogenic Protein A variants can neutralize this virulence factor and thereby establish protective immunity. In certain aspects the protective immunity protects or ameliorates infection by drug resistant strains of *Staphylococcus*, such as USA300 and other MRSA strains.

Embodiments include the use of Protein A variants in methods and compositions for the treatment of bacterial and/or staphylococcal infection. This application also provides an immunogenic composition comprising a Protein A variant or immunogenic fragment thereof. In certain aspects, the immunogenic fragment is a Protein A domain D segment. Furthermore, the present invention provides methods and compositions that can be used to treat (e.g., limiting staphylococcal abscess formation and/or persistence in a subject) or prevent bacterial infection. In some cases, methods for stimulating an immune response involve administering to the subject an effective amount of a composition including or encoding all or part of a Protein A variant polypeptide or antigen, and in certain aspects other bacterial proteins. Other bacterial proteins include, but are not limited to (i) a secreted virulence factor, and/or a cell surface protein or peptide, or (ii) a recombinant nucleic acid molecule encoding a secreted virulence factor, and/or a cell surface protein or peptide.

In other aspects, the subject can be administered all or part of a Protein A variant, such as a variant Protein A domain D segment. The polypeptide of the invention can be formulated in a pharmaceutically acceptable composition. The composition can further comprise one or more of at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 additional staphylococcal antigen or immunogenic fragment thereof (e.g., Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla (e.g., H35 mutants), IsdC, SasF, vWbp, or vWh). Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa (GenBank CAC80837), Aap (GenBank accession AJ249487), Ant (GenBank accession NP\_372518), autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg2+ transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/





TABLE 1-continued

SpA and staphylococcal antigen combinations.														
vWh														+
EsxB	+	+	+	+	+	+	+	+	+	+	+	+	+	+
SdrC		+	+	+	+	+	+	+	+	+	+	+	+	+
SdrD			+	+	+	+	+	+	+	+	+	+	+	+
SdrE				+	+	+	+	+	+	+	+	+	+	+
IsdA					+	+	+	+	+	+	+	+	+	+
IsdB						+	+	+	+	+	+	+	+	+
ClfA							+	+	+	+	+	+	+	+
ClfB								+	+	+	+	+	+	+
Coa									+	+	+	+	+	+
Hla										+	+	+	+	+
Hla <sub>II35,4</sub>											+	+	+	+
IsdC												+	+	+
SasF													+	+
vWbp														+
vWh														+
SdrC		+	+	+	+	+	+	+	+	+	+	+	+	+
SdrD			+	+	+	+	+	+	+	+	+	+	+	+
SdrE				+	+	+	+	+	+	+	+	+	+	+
IsdA					+	+	+	+	+	+	+	+	+	+
IsdB						+	+	+	+	+	+	+	+	+
ClfA							+	+	+	+	+	+	+	+
ClfB								+	+	+	+	+	+	+
Coa									+	+	+	+	+	+
Hla										+	+	+	+	+
Hla <sub>II35,4</sub>											+	+	+	+
IsdC												+	+	+
SasF													+	+
vWbp														+
vWh														+
SdrD			+	+	+	+	+	+	+	+	+	+	+	+
SdrE				+	+	+	+	+	+	+	+	+	+	+
IsdA					+	+	+	+	+	+	+	+	+	+
IsdB						+	+	+	+	+	+	+	+	+
ClfA							+	+	+	+	+	+	+	+
ClfB								+	+	+	+	+	+	+
Coa									+	+	+	+	+	+
Hla										+	+	+	+	+
Hla <sub>II35,4</sub>											+	+	+	+
IsdC												+	+	+
SasF													+	+
vWbp														+
vWh														+
SdrE				+	+	+	+	+	+	+	+	+	+	+
IsdA					+	+	+	+	+	+	+	+	+	+
IsdB						+	+	+	+	+	+	+	+	+
ClfA							+	+	+	+	+	+	+	+
ClfB								+	+	+	+	+	+	+
Coa									+	+	+	+	+	+
Hla										+	+	+	+	+
Hla <sub>II35,4</sub>											+	+	+	+
IsdC												+	+	+
SasF													+	+
vWbp														+
vWh														+
IsdA					+	+	+	+	+	+	+	+	+	+
IsdB						+	+	+	+	+	+	+	+	+
ClfA							+	+	+	+	+	+	+	+
ClfB								+	+	+	+	+	+	+
Coa									+	+	+	+	+	+
Hla										+	+	+	+	+
Hla <sub>II35,4</sub>											+	+	+	+
IsdC												+	+	+
SasF													+	+
vWbp														+
vWh														+
IsdB						+	+	+	+	+	+	+	+	+
ClfA							+	+	+	+	+	+	+	+
ClfB								+	+	+	+	+	+	+
Coa									+	+	+	+	+	+
Hla										+	+	+	+	+
Hla <sub>II35,4</sub>											+	+	+	+
IsdC												+	+	+
SasF													+	+
vWbp														+
vWh														+
ClfA							+	+	+	+	+	+	+	+
ClfB								+	+	+	+	+	+	+



thereof. A secreted protein or cell surface protein includes, but is not limited to Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, and/or vWh proteins and immunogenic fragments thereof. Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg2+ transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein.

Embodiments of the invention include compositions that include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to Protein A, or a second protein or peptide that is a secreted bacterial protein or a bacterial cell surface protein. In a further embodiment of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Protein A domain D polypeptide (SEQ ID NO:2), domain E (SEQ ID NO:3), domain A (SEQ ID NO:4), domain C (SEQ ID NO:5), domain B (SEQ ID NO:6), or a nucleic acid sequence encoding a Protein A domain D, domain E, domain A, domain C, or domain B polypeptide. In certain aspects a Protein A polypeptide segment will have an amino acid sequence of SEQ ID NO:8. Similarity or identity, with identity being preferred, is known in the art and a number of different programs can be used to identify whether a protein (or nucleic acid) has sequence identity or similarity to a known sequence. Sequence identity and/or similarity is determined using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith & Waterman (1981), by the sequence identity alignment algorithm of Needleman & Wunsch (1970), by the search for similarity method of Pearson & Lipman (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux et al. (1984), preferably using the default settings, or by inspection. Preferably, percent identity is calculated by using alignment tools known to and readily ascertainable to those of skill in the art. Percent identity is essentially the number of identical amino acids divided by the total number of amino acids compared times one hundred.

Still further embodiments include methods for stimulating in a subject a protective or therapeutic immune response against a *Staphylococcus* bacterium comprising administering to the subject an effective amount of a composition including (i) a SpA variant, e.g., a variant SpA domain D polypeptide or peptide thereof; or, (ii) a nucleic acid molecule encoding such a SpA variant polypeptide or peptide thereof, or (iii) administering a SpA variant domain D polypeptide with any combination or permutation of bacte-

rial proteins described herein. In a preferred embodiment the composition is not a *Staphylococcus* bacterium. In certain aspects the subject is a human or a cow. In a further aspect the composition is formulated in a pharmaceutically acceptable formulation. The staphylococci may be *Staphylococcus aureus*.

Yet still further embodiments include vaccines comprising a pharmaceutically acceptable composition having an isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein, wherein the composition is capable of stimulating an immune response against a *Staphylococcus* bacterium. The vaccine may comprise an isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described. In certain aspects of the invention the isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described are multimerized, e.g., dimerized or concatamerized. In a further aspect, the vaccine composition is contaminated by less than about 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, 0.5, 0.25, 0.05% (or any range derivable therein) of other Staphylococcal proteins. A composition may further comprise an isolated non-SpA polypeptide. Typically the vaccine comprises an adjuvant. In certain aspects a protein or peptide of the invention is linked (covalently or non-covalently) to the adjuvant, preferably the adjuvant is chemically conjugated to the protein.

In still yet further embodiments, a vaccine composition is a pharmaceutically acceptable composition having a recombinant nucleic acid encoding all or part of a SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein, wherein the composition is capable of stimulating an immune response against a *Staphylococcus* bacteria. The vaccine composition may comprise a recombinant nucleic acid encoding all or part of a SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein. In certain embodiments the recombinant nucleic acid contains a heterologous promoter. Preferably the recombinant nucleic acid is a vector. More preferably the vector is a plasmid or a viral vector. In some aspects the vaccine includes a recombinant, non-*Staphylococcus* bacterium containing the nucleic acid. The recombinant non-staphylococci may be *Salmonella* or another gram-positive bacteria. The vaccine may comprise a pharmaceutically acceptable excipient, more preferably an adjuvant.

Still further embodiments include methods for stimulating in a subject a protective or therapeutic immune response against a *Staphylococcus* bacterium comprising administering to the subject an effective amount of a composition of a SpA variant polypeptide or segment/fragment thereof and further comprising one or more of a Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, or vWh protein or peptide thereof. In a preferred embodiment the composition comprises a non-*Staphylococcus* bacterium. In a further aspect the composition is formulated in a pharmaceutically acceptable formulation. The staphylococci for which a subject is being treated may be *Staphylococcus aureus*. Methods of the invention also include SpA variant compositions that contain 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or more secreted virulence factors and/or cell surface proteins, such as Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, or vWh in various combinations. In certain aspects a vaccine formulation includes Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla,

IsdC, SasF, vWbp, and vWh. In certain aspects an antigen combination can include (1) a SpA variant and IsdA; (2) SpA variant and ClfB; (3) SpA variant and SdrD; (4) SpA variant and Hla or Hla variant; (5) SpA variant and ClfB, SdrD, and Hla or Hla variant; (6) SpA variant, IsdA, SdrD, and Hla or Hla variant; (7) SpA variant, IsdA, ClfB, and Hla or Hla variant; (8) SpA variant, IsdA, ClfB, and SdrD; (9) SpA variant, IsdA, ClfB, SdrD and Hla or Hla variant; (10) SpA variant, IsdA, ClfB, and SdrD; (11) SpA variant, IsdA, SdrD, and Hla or Hla variant; (12) SpA variant, IsdA, and Hla or Hla variant; (13) SpA variant, IsdA, ClfB, and Hla or Hla variant; (14) SpA variant, ClfB, and SdrD; (15) SpA variant, ClfB, and Hla or Hla variant; or (16) SpA variant, SdrD, and Hla or Hla variant.

In certain aspects, a bacterium delivering a composition of the invention will be limited or attenuated with respect to prolonged or persistent growth or abscess formation. In yet a further aspect, SpA variant(s) can be overexpressed in an attenuated bacterium to further enhance or supplement an immune response or vaccine formulation.

The term "EsxA protein" refers to a protein that includes isolated wild-type EsxA polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria EsxA proteins.

The term "EsxB protein" refers to a protein that includes isolated wild-type EsxB polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria EsxB proteins.

The term "SdrD protein" refers to a protein that includes isolated wild-type SdrD polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria SdrD proteins.

The term "SdrE protein" refers to a protein that includes isolated wild-type SdrE polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria SdrE proteins.

The term "IsdA protein" refers to a protein that includes isolated wild-type IsdA polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria IsdA proteins.

The term "IsdB protein" refers to a protein that includes isolated wild-type IsdB polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria IsdB proteins.

The term "Eap protein" refers to a protein that includes isolated wild-type Eap polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria Eap proteins.

The term "Ebh protein" refers to a protein that includes isolated wild-type Ebh polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria Ebh proteins.

The term "Emp protein" refers to a protein that includes isolated wild-type Emp polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria Emp proteins.

The term "EsaB protein" refers to a protein that includes isolated wild-type EsaB polypeptides from *Staphylococcus*

bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria EsaB proteins.

The term "EsaC protein" refers to a protein that includes isolated wild-type EsaC polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria EsaC proteins.

The term "SdrC protein" refers to a protein that includes isolated wild-type SdrC polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria SdrC proteins.

The term "ClfA protein" refers to a protein that includes isolated wild-type ClfA polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria ClfA proteins.

The term "ClfB protein" refers to a protein that includes isolated wild-type ClfB polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria ClfB proteins.

The term "Coa protein" refers to a protein that includes isolated wild-type Coa polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria Coa proteins.

The term "Hla protein" refers to a protein that includes isolated wild-type Hla polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria Hla proteins.

The term "IsdC protein" refers to a protein that includes isolated wild-type IsdC polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria IsdC proteins.

The term "SasF protein" refers to a protein that includes isolated wild-type SasF polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria SasF proteins.

The term "vWbp protein" refers to a protein that includes isolated wild-type vWbp (von Willebrand factor binding protein) polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria vWbp proteins.

The term "vWh protein" refers to a protein that includes isolated wild-type vWh (von Willebrand factor binding protein homolog) polypeptides from *Staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *Staphylococcus* bacteria vWh proteins.

An immune response refers to a humoral response, a cellular response, or both a humoral and cellular response in an organism. An immune response can be measured by assays that include, but are not limited to, assays measuring the presence or amount of antibodies that specifically recognize a protein or cell surface protein, assays measuring T-cell activation or proliferation, and/or assays that measure modulation in terms of activity or expression of one or more cytokines.

In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%,

98%, or 99% identical or similar to an EsxA protein. In certain aspects the EsxA protein will have all or part of the amino acid sequence of SEQ ID NO:11.

In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsxB protein. In certain aspects the EsxB protein will have all or part of the amino acid sequence of SEQ ID NO:12.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an SdrD protein. In certain aspects the SdrD protein will have all or part of the amino acid sequence of SEQ ID NO:13.

In further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an SdrE protein. In certain aspects the SdrE protein will have all or part of the amino acid sequence of SEQ ID NO:14.

In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdA protein. In certain aspects the IsdA protein will have all or part of the amino acid sequence of SEQ ID NO:15.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdB protein. In certain aspects the IsdB protein will have all or part of the amino acid sequence of SEQ ID NO:16.

Embodiments of the invention include compositions that include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a EsaB protein. In certain aspects the EsaB protein will have all or part of the amino acid sequence of SEQ ID NO:17.

In a further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a ClfB protein. In certain aspects the ClfB protein will have all or part of the amino acid sequence of SEQ ID NO:18.

In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdC protein. In certain aspects the IsdC protein will have all or part of the amino acid sequence of SEQ ID NO:19.

In yet further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a SasF protein. In certain aspects the SasF protein will have all or part of the amino acid sequence of SEQ ID NO:20.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a SdrC protein. In certain aspects the SdrC protein will have all or part of the amino acid sequence of SEQ ID NO:21.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a ClfA protein. In certain

aspects the ClfA protein will have all or part of the amino acid sequence of SEQ ID NO:22.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Eap protein. In certain aspects the Eap protein will have all or part of the amino acid sequence of SEQ ID NO:23.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Ebh protein. In certain aspects the Ebh protein will have all or part of the amino acid sequence of SEQ ID NO:24.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Emp protein. In certain aspects the Emp protein will have all or part of the amino acid sequence of SEQ ID NO:25.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsaC protein. In certain aspects the EsaC protein will have all or part of the amino acid sequence of SEQ ID NO:26. Sequence of EsaC polypeptides can be found in the protein databases and include, but are not limited to accession numbers ZP\_02760162 (GI:168727885), NP\_645081.1 (GI:21281993), and NP\_370813.1 (GI:15923279), each of which is incorporated herein by reference as of the priority date of this application.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Coa protein. In certain aspects the Coa protein will have all or part of the amino acid sequence of SEQ ID NO:27.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Hla protein. In certain aspects the Hla protein will have all or part of the amino acid sequence of SEQ ID NO:28.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a vWa protein. In certain aspects the vWa protein will have all or part of the amino acid sequence of SEQ ID NO:29.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a vWbp protein. In certain aspects the vWbp protein will have all or part of the amino acid sequence of SEQ ID NO:32.

In certain aspects, a polypeptide or segment/fragment can have a sequence that is at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% or more identical to the amino acid sequence of the reference polypeptide. The term "similarity" refers to a polypeptide that has a sequence that has a certain percentage of amino acids that are either identical with the reference polypeptide or constitute conservative substitutions with the reference polypeptides.

The polypeptides described herein may include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more variant amino acids within at least, or at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12,

13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:2-30, or SEQ ID NO:32-34.

A polypeptide segment as described herein may include 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:2-30, or SEQ ID NO:33-34.

The compositions may be formulated in a pharmaceutically acceptable composition. In certain aspects of the invention the *Staphylococcus* bacterium is an *S. aureus* bacterium.

In further aspects, a composition may be administered more than one time to the subject, and may be administered 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20 or more times. The administration of the compositions include, but is not limited to oral, parenteral, subcutaneous, intramuscular, intravenous, or various combinations thereof, including inhalation or aspiration.

In still further embodiments, a composition comprises a recombinant nucleic acid molecule encoding a polypeptide described herein or segments/fragments thereof. Typically a recombinant nucleic acid molecule encoding a polypeptide described herein contains a heterologous promoter. In certain aspects, a recombinant nucleic acid molecule of the invention is a vector, in still other aspects the vector is a plasmid. In certain embodiments the vector is a viral vector. In certain aspects a composition includes a recombinant, non-*Staphylococcus* bacterium containing or expressing a polypeptide described herein. In particular aspects the recombinant non-*Staphylococcus* bacteria is *Salmonella* or another gram-positive bacteria. A composition is typically

administered to mammals, such as human subjects, but administration to other animals that are capable of eliciting an immune response is contemplated. In further aspects the *Staphylococcus* bacterium containing or expressing the polypeptide is *Staphylococcus aureus*. In further embodiments the immune response is a protective immune response.

In further embodiments a composition comprises a recombinant nucleic acid molecule encoding all or part of one or more of a Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, SpA, vWbp, or vWh protein or peptide or variant thereof. Additional staphylococcal antigens that can be used in combination with the polypeptides described herein include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg2+ transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein. In particular aspects, a bacteria is a recombinant non-*Staphylococcus* bacteria, such as a *Salmonella* or other gram-positive bacteria.

Compositions of the invention are typically administered to human subjects, but administration to other animals that are capable of eliciting an immune response to a *Staphylococcus* bacterium is contemplated, particularly cattle, horses, goats, sheep and other domestic animals, i.e., mammals.

In certain aspects the *Staphylococcus* bacterium is a *Staphylococcus aureus*. In further embodiments the immune response is a protective immune response. In still further aspects, the methods and compositions of the invention can be used to prevent, ameliorate, reduce, or treat infection of tissues or glands, e.g., mammary glands, particularly mastitis and other infections. Other methods include, but are not limited to prophylactically reducing bacterial burden in a subject not exhibiting signs of infection, particularly those subjects suspected of or at risk of being colonized by a target bacteria, e.g., patients that are or will be at risk or susceptible to infection during a hospital stay, treatment, and/or recovery.

Any embodiment discussed with respect to one aspect of the invention applies to other aspects of the invention as well. In particular, any embodiment discussed in the context of a SpA variant polypeptide or peptide or nucleic acid may be implemented with respect to other antigens, such as Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD,

MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein (or nucleic acids), and vice versa. It is also understood that any one or more of Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein can be specifically excluded from a claimed composition.

Embodiments of the invention include compositions that contain or do not contain a bacterium. A composition may or may not include an attenuated or viable or intact staphylococcal bacterium. In certain aspects, the composition comprises a bacterium that is not a staphylococcal bacterium or does not contain staphylococcal bacteria. In certain embodiments a bacterial composition comprises an isolated or recombinantly expressed staphylococcal Protein A variant or a nucleotide encoding the same. The composition may be or include a recombinantly engineered *Staphylococcus* bacterium that has been altered in a way that comprises specifically altering the bacterium with respect to a secreted virulence factor or cell surface protein. For example, the bacteria may be recombinantly modified to express more of the virulence factor or cell surface protein than it would express if unmodified.

The term "isolated" can refer to a nucleic acid or polypeptide that is substantially free of cellular material, bacterial material, viral material, or culture medium (when produced by recombinant DNA techniques) of their source of origin, or chemical precursors or other chemicals (when chemically synthesized). Moreover, an isolated compound refers to one that can be administered to a subject as an isolated compound; in other words, the compound may not simply be considered "isolated" if it is adhered to a column or embedded in an agarose gel. Moreover, an "isolated nucleic acid fragment" or "isolated peptide" is a nucleic acid or protein fragment that is not naturally occurring as a fragment and/or is not typically in the functional state.

Moieties of the invention, such as polypeptides, peptides, antigens, or immunogens, may be conjugated or linked covalently or noncovalently to other moieties such as adjuvants, proteins, peptides, supports, fluorescence moieties, or labels. The term "conjugate" or "immunoconjugate" is broadly used to define the operative association of one moiety with another agent and is not intended to refer solely to any type of operative association, and is particularly not limited to chemical "conjugation." Recombinant fusion proteins are particularly contemplated. Compositions of the

invention may further comprise an adjuvant or a pharmaceutically acceptable excipient. An adjuvant may be covalently or non-covalently coupled to a polypeptide or peptide of the invention. In certain aspects, the adjuvant is chemically conjugated to a protein, polypeptide, or peptide.

The term "providing" is used according to its ordinary meaning to indicate "to supply or furnish for use." In some embodiments, the protein is provided directly by administering the protein, while in other embodiments, the protein is effectively provided by administering a nucleic acid that encodes the protein. In certain aspects the invention contemplates compositions comprising various combinations of nucleic acid, antigens, peptides, and/or epitopes.

The subject will have (e.g., are diagnosed with a staphylococcal infection), will be suspected of having, or will be at risk of developing a staphylococcal infection. Compositions of the present invention include immunogenic compositions wherein the antigen(s) or epitope(s) are contained in an amount effective to achieve the intended purpose. More specifically, an effective amount means an amount of active ingredients necessary to stimulate or elicit an immune response, or provide resistance to, amelioration of, or mitigation of infection. In more specific aspects, an effective amount prevents, alleviates or ameliorates symptoms of disease or infection, or prolongs the survival of the subject being treated. Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any preparation used in the methods of the invention, an effective amount or dose can be estimated initially from in vitro studies, cell culture, and/or animal model assays. For example, a dose can be formulated in animal models to achieve a desired immune response or circulating antibody concentration or titer. Such information can be used to more accurately determine useful doses in humans.

The embodiments in the Example section are understood to be embodiments of the invention that are applicable to all aspects of the invention.

The use of the term "or" in the claims is used to mean "and/or" unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and "and/or." It is also contemplated that anything listed using the term "or" may also be specifically excluded.

Throughout this application, the term "about" is used to indicate that a value includes the standard deviation of error for the device or method being employed to determine the value.

Following long-standing patent law, the words "a" and "an," when used in conjunction with the word "comprising" in the claims or specification, denotes one or more, unless specifically noted.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

#### DESCRIPTION OF THE DRAWINGS

So that the matter in which the above-recited features, advantages and objects of the invention as well as others which will become clear are attained and can be understood

in detail, more particular descriptions and certain embodiments of the invention briefly summarized above are illustrated in the appended drawings. These drawings form a part of the specification. It is to be noted, however, that the appended drawings illustrate certain embodiments of the invention and therefore are not to be considered limiting in their scope.

FIGS. 1A-1B. (FIG. 1A) Primary structure of the Protein A precursor with an N-terminal YSIRK motif signal peptide, five immunoglobulin binding domains as tandem repeats designated E, D, A, B, C, region X, and the LPXTG sorting signal. (FIG. 1B) Following synthesis of the Protein A precursor, staphylococci secrete this product via the Sec pathway, and sortase A cleaves the LPXTG sorting signal between the T and G residues. Nucleophilic attack of the amino group within lipid II at the sortase-Protein A thioester-linked intermediate forms the amide bond that links Protein A to the cell wall envelope and enables its display on the bacterial surface.

FIG. 2. Three dimensional model of the molecular interactions between the SpA-domain D of Protein A, the VH3 Fab domain of the B cell receptor, and of the Fc $\gamma$  domain of immunoglobulin. The model is derived from two crystal structures (Graille et al., 2000 and Gouda et al., 1992) that revealed side chain residues involved in the formation of ionic bonds that enable these complexes. Gln-9 and Gln-10 of SpA-D promote binding to Fc $\gamma$ , whereas Asp-36 and Asp-37 enable complex formation with VH3 Fab.

FIG. 3. Left panel—Coomassie Blue stained SDS-PAGE reveals the migrational position of purified His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub>, human IgG, and sortase A (SrtA), a control protein. Right panel—Coomassie Blue stained SDS-PAGE to reveal the elution of Protein A immunoglobulin complexes eluted following affinity chromatography of human IgG on Ni-NTA columns pre-charged with His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or SrtA.

FIG. 4. ELISA assays to quantify human immunoglobulin (hIgG), human F(ab)<sub>2</sub> IgG fragments and human Fc fragments of immunoglobulin (hFc). Plates were coated with equal amounts of His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or SrtA. hIgG-HRP, F(ab)<sub>2</sub>-HRP and hFc-HRP were added onto the plates and incubated for an hour. Absorbance at 450 nm was recorded and plotted to determine the half maximal titers.

FIG. 5. Purified SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or a PBS mock control were injected into the peritoneum of mice and analyzed for their ability to reduce the B cell population in the spleen of experimental BALB/c mice. Animals were killed 4 hours following injection, their spleen removed, tissue homogenized and stained with CD19 antibodies directed against B cells. The number of B cells was quantified by FACS sorting.

FIG. 6 Generation of a non-toxicogenic protein A vaccine. a, Translational protein A (SpA) product of *S. aureus* Newman and USA300 LAC with an N-terminal signal peptide (white box), five immunoglobulin binding domains (IgBDs designated E, D, A, B and C), variable region X and C-terminal sorting signal (black box). b, Amino acid sequence of the five IgBDs as well as nontoxicogenic SpA-D<sub>KKAA</sub>, with the positions of triple  $\alpha$ -helical bundles (H1, H2 and H3) as well as glutamine (Q) 9, 10 and aspartate (D) 36, 37 indicated. c, Coomassie Blue-stained SDS-PAGE of SpA, SpA-D, SpA-D<sub>KKAA</sub> or SrtA purified on Ni-NTA sepharose in the presence or absence of human immunoglobulin (hIgG). d, ELISA examining the association of immobilized SpA, SpA-D or SpA-D<sub>KKAA</sub> with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments and von Willebrand

factor (vWF). e, CD19+ B lymphocytes in splenic tissue of BALB/c mice that had been mock immunized or treated with SpA-D or SpA-D<sub>KKAA</sub> were quantified by FACS.

FIG. 7 Non-toxicogenic protein A vaccine prevents abscess formation. Histopathology of renal tissue isolated during necropsy of BALB/c mice that had been mock immunized (PBS) or vaccinated with SpA, SpA-D as well as SpA-D<sub>KKAA</sub> and challenged with *S. aureus* Newman. Thin sectioned tissues were stained with hematoxylin-eosin. White arrows identify polymorphonuclear leukocyte (PMN) infiltrates. Dark arrows identify staphylococcal abscess communities.

FIG. 8 Antibodies raised by the non-toxicogenic protein A vaccine block the B cell superantigen function of SpA. a, Rabbit antibodies raised against SpA-D<sub>KKAA</sub> were purified on a matrix with immobilized antigen and analyzed by Coomassie Blue-stained SDS-PAGE. Antibodies were cleaved with pepsin and F(ab)<sub>2</sub> fragments were purified by a second round of affinity chromatography on SpA-D<sub>KKAA</sub> matrix. b, SpA-D<sub>KKAA</sub> specific F(ab)<sub>2</sub> interfere with the binding of SpA or SpA-D to human immunoglobulin (hIgG) or, c, to von Willebrand Factor (vWF).

FIG. 9 Full-length non-toxicogenic protein A generates improved immune responses. a, Full-length SpA<sub>KKAA</sub> was purified on Ni-NTA sepharose and analyzed by Coomassie-Blue stained SDS-PAGE. b, CD19+ B lymphocytes in splenic tissue of BALB/c mice that had been mock immunized or treated with SpA or SpA<sub>KKAA</sub> were quantified by FACS. c, ELISA examining the association of immobilized SpA or SpA<sub>KKAA</sub> with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments or von Willebrand factor (vWF). d, Human or mouse serum antibody titers to diphtheria toxoid (CRM197) and non-toxicogenic SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub>. Human volunteers with a history of DTaP immunization and staphylococcal infection (n=16) as well as mice (n=20) that had been infected with *S. aureus* Newman or USA 300 LAC or immunized with SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub> were examined by quantitative dot blot.

FIG. 10 Staphylococcal infection does not generate protective immunity. BALB/c mice (n=20) were infected with *S. aureus* Newman or mock challenged (PBS) for thirty days and infection cleared with chloramphenicol treatment. Both cohorts of animals were then challenged with *S. aureus* Newman and bacterial load (CFU) in kidney tissue homogenate analyzed following necropsy on day 4.

FIG. 11 Comparison of abscess formation in mice treated with PBS, SpA, SpA-D, and SpA-D<sub>KKAA</sub>.

FIGS. 12A-12C (A) ELISA examining the association of immobilized SpA, SpA-D, SpA-D<sub>KKAA</sub> or SpA-DGGSS with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments and IgM. Statistical significance of SpA-D<sub>KKAA</sub> and SpA-DGGSS binding to each ligand was compared against SpA-D; SpA-D binding was compared against SpA (n=3); \* signifies P<0.05; \*\* signifies P<0.01. (B) ELISA examining the level of cross-reactive antibodies of hyper-immune sera samples collected from actively immunized mice (n=5) with SpA-D, SpA-D<sub>KKAA</sub> and SpA-DGGSS. (C) Abscess formation in mice treated with PBS, SpA-D, SpA-D<sub>KKAA</sub> and SpA-D<sub>GGSS</sub>.

## DETAILED DESCRIPTION

*Staphylococcus aureus* is a commensal of the human skin and nares, and the leading cause of bloodstream, skin and soft tissue infections (Klebens et al., 2007). Recent dramatic increases in the mortality of staphylococcal diseases are attributed to the spread of methicillin-resistant *S. aureus*

(MRSA) strains often not susceptible to antibiotics (Kennedy et al., 2008). In a large retrospective study, the incidence of MRSA infections was 4.6% of all hospital admissions in the United States (Klevens et al., 2007). The annual health care costs for 94,300 MRSA infected individuals in the United States exceed \$2.4 billion (Klevens et al., 2007). The current MRSA epidemic has precipitated a public health crisis that needs to be addressed by development of a preventive vaccine (Boucher and Corey, 2008). To date, an FDA licensed vaccine that prevents *S. aureus* diseases is not available.

The inventors describe here the use of Protein A, a cell wall anchored surface protein of staphylococci, for the generation of variants that can serve as subunit vaccines. The pathogenesis of staphylococcal infections is initiated as bacteria invade the skin or blood stream via trauma, surgical wounds, or medical devices (Lowy, 1998). Although the invading pathogen may be phagocytosed and killed, staphylococci can also escape innate immune defenses and seed infections in organ tissues, inducing inflammatory responses that attract macrophages, neutrophils, and other phagocytes (Lowy, 1998). The responsive invasion of immune cells to the site of infection is accompanied by liquefaction necrosis as the host seeks to prevent staphylococcal spread and allow for removal of necrotic tissue debris (Lam et al., 1963). Such lesions can be observed by microscopy as hypercellular areas containing necrotic tissue, leukocytes, and a central nidus of bacteria (Lam et al., 1963). Unless staphylococcal abscesses are surgically drained and treated with antibiotics, disseminated infection and septicemia produce a lethal outcome (Sheagren, 1984).

## II. Staphylococcal Antigens

### A. Staphylococcal Protein A (SpA)

All *Staphylococcus aureus* strains express the structural gene for Protein A (*spa*) (Jensen, 1958; Said-Salim et al., 2003), a well characterized virulence factor whose cell wall anchored surface protein product (SpA) encompasses five highly homologous immunoglobulin binding domains designated E, D, A, B, and C (Sjodahl, 1977). These domains display ~80% identity at the amino acid level, are 56 to 61 residues in length, and are organized as tandem repeats (Uhlen et al., 1984). SpA is synthesized as a precursor protein with an N-terminal YSIRK/GS signal peptide and a C-terminal LPXTG motif sorting signal (DeDent et al., 2008; Schneewind et al., 1992). Cell wall anchored Protein A is displayed in great abundance on the staphylococcal surface (DeDent et al., 2007; Sjoquist et al., 1972). Each of its immunoglobulin binding domains is composed of anti-parallel  $\alpha$ -helices that assemble into a three helix bundle and bind the Fc domain of immunoglobulin G (IgG) (Deisenhofer, 1981; Deisenhofer et al., 1978), the VH3 heavy chain (Fab) of IgM (i.e., the B cell receptor) (Graille et al., 2000), the von Willebrand factor at its A1 domain [vWF A1 is a ligand for platelets] (O'Seaghda et al., 2006) and the tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ) receptor I (TNFRI) (Gomez et al., 2006), which is displayed on surfaces of airway epithelia (Gomez et al., 2004; Gomez et al., 2007).

SpA impedes neutrophil phagocytosis of staphylococci through its attribute of binding the Fc component of IgG (Jensen, 1958; Uhlen et al., 1984). Moreover, SpA is able to activate intravascular clotting via its binding to von Willebrand factor A1 domains (Hartleib et al., 2000). Plasma proteins such as fibrinogen and fibronectin act as bridges between staphylococci (ClfA and ClfB) and the platelet integrin GPIIb/IIIa (O'Brien et al., 2002), an activity that is supplemented through Protein A association with vWF A1, which allows staphylococci to capture platelets via the

GPIIb- $\alpha$  platelet receptor (Foster, 2005; O'Seaghda et al., 2006). SpA also binds TNFRI and this interaction contributes to the pathogenesis of staphylococcal pneumonia (Gomez et al., 2004). SpA activates proinflammatory signaling through TNFR1 mediated activation of TRAF2, the p38/c-Jun kinase, mitogen activate protein kinase (MAPK) and the Rel-transcription factor NF-KB. SpA binding further induces TNFR1 shedding, an activity that appears to require the TNF-converting enzyme (TACE)(Gomez et al., 2007). All of the aforementioned SpA activities are mediated through its five IgG binding domains and can be perturbed by the same amino acid substitutions, initially defined by their requirement for the interaction between Protein A and human IgG1 (Cedergren et al., 1993).

SpA also functions as a B cell superantigen by capturing the Fab region of VH3 bearing IgM, the B cell receptor (Gomez et al., 2007; Goodyear et al., 2003; Goodyear and Silverman, 2004; Roben et al., 1995). Following intravenous challenge, staphylococcal Protein A (SpA) mutations show a reduction in staphylococcal load in organ tissues and dramatically diminished ability to form abscesses (described herein). During infection with wildtype *S. aureus*, abscesses are formed within forty-eight hours and are detectable by light microscopy of hematoxylin-eosin stained, thin-sectioned kidney tissue, initially marked by an influx of polymorphonuclear leukocytes (PMNs). On day 5 of infection, abscesses increase in size and enclosed a central population of staphylococci, surrounded by a layer of eosinophilic, amorphous material and a large cuff of PMNs. Histopathology revealed massive necrosis of PMNs in proximity to the staphylococcal nidus at the center of abscess lesions as well as a mantle of healthy phagocytes. The inventors also observed a rim of necrotic PMNs at the periphery of abscess lesions, bordering the eosinophilic pseudocapsule that separated healthy renal tissue from the infectious lesion. Staphylococcal variants lacking Protein A are unable to establish the histopathology features of abscesses and are cleared during infection.

In previous studies, Cedergren et al. (1993) engineered five individual substitutions in the Fc fragment binding sub-domain of the B domain of SpA, L17D, N28A, I31A and K35A. These authors created these proteins to test data gathered from a three dimensional structure of a complex between one domain of SpA and Fc. Cedergren et al. determined the effects of these mutations on stability and binding, but did not contemplate use of such substitutions for the production of a vaccine antigen.

Brown et al. (1998) describe studies designed to engineer new proteins based on SpA that allow the use of more favorable elution conditions when used as affinity ligands. The mutations studied included single mutations of Q13A, Q14H, N15A, N15H, F17H, Y18F, L21H, N32H, or K39H. Brown et al. report that Q13A, N15A, N15H, and N32H substitutions made little difference to the dissociation constant values and that the Y18F substitution resulted in a 2 fold decrease in binding affinity as compared to wild type SpA. Brown et al. also report that L21H and F17H substitutions decrease the binding affinity by five-fold and a hundred-fold respectively. The authors also studied analogous substitutions in two tandem domains. Thus, the Brown et al. studies were directed to generating a SpA with a more favorable elution profile, hence the use of His substitutions to provide a pH sensitive alteration in the binding affinity. Brown et al. is silent on the use of SpA as a vaccine antigen.

Graille et al. (2000) describe a crystal structure of domain D of SpA and the Fab fragment of a human IgM antibody. Graille et al. define by analysis of a crystal structure the D

domain amino acid residues that interact with the Fab fragment as residues Q26, G29, F30, Q32, S33, D36, D37, Q40, N43, E47, or L51, as well as the amino acid residues that form the interface between the domain D sub-domains. Graille et al. define the molecular interactions of these two proteins, but is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

O'Seaghda et al. (2006) describe studies directed at elucidating which sub-domain of domain D binds vWF. The authors generated single mutations in either the Fc or VH3 binding sub-domains, i.e., amino acid residues F5A, Q9A, Q10A, F13A, Y14A, L17A, N28A, I31A, K35A, G29A, F30A, S33A, D36A, D37A, Q40A, E47A, or Q32A. The authors discovered that vWF binds the same sub-domain that binds Fc. O'Seaghda et al. define the sub-domain of domain D responsible for binding vWF, but is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

Gomez et al. (2006) describe the identification of residues responsible for activation of the TNFR1 by using single mutations of F5A, F13A, Y14A, L17A, N21A, I31A, Q32A, and K35A. Gomez et al. is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

Recombinant affinity tagged Protein A, a polypeptide encompassing the five IgG domains (EDCAB) (Sjodahl, 1977) but lacking the C-terminal Region X (Guss et al., 1984), was purified from recombinant *E. coli* and used as a vaccine antigen (Stranger-Jones et al., 2006). Because of the attributes of SpA in binding the Fc portion of IgG, a specific humoral immune response to Protein A could not be measured (Stranger-Jones et al., 2006). The inventors have overcome this obstacle through the generation of SpA-DQ9, 10K;D36,37A. BALB/c mice immunized with recombinant Protein A (SpA) displayed significant protection against intravenous challenge with *S. aureus* strains: a 2.951 log reduction in staphylococcal load as compared to the wild-type ( $P > 0.005$ ; Student's t-test) (Stranger-Jones et al., 2006). SpA specific antibodies may cause phagocytic clearance prior to abscess formation and/or impact the formation of the aforementioned eosinophilic barrier in abscesses that separate staphylococcal communities from immune cells since these do not form during infection with Protein A mutant strains. Each of the five SpA domains (i.e., domains formed from three helix bundles designated E, D, A, B, and C) exerts similar binding properties (Jansson et al., 1998). The solution and crystal structure of the domain D has been solved both with and without the Fc and VH3 (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille et al., 2000). Mutations in residues known to be involved in IgG binding (FS, Q9, Q10, S11, F13, Y14, L17, N28, 131 and K35) are also required for vWF A1 and TNFR1 binding (Cedergren et al., 1993; Gomez et al., 2006; O'Seaghda et al., 2006), whereas residues important for the VH3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) appear to have no impact on the other binding activities (Graille et al., 2000; Jansson et al., 1998). SpA specifically targets a subset of B cells that express VH3 family related IgM on their surface, i.e., VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells proliferate and commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e., marginal zone B cells and follicular B2 cells) (Goodyear et al., 2003; Goodyear et al., 2004).

Molecular basis of Protein A surface display and function. Protein A is synthesized as a precursor in the bacterial cytoplasm and secreted via its YSIRK signal peptide at the

cross wall, i.e. the cell division septum of staphylococci (FIG. 1) (DeDent et al., 2007; DeDent et al., 2008). Following cleavage of the C-terminal LPXTG sorting signal, Protein A is anchored to bacterial peptidoglycan cross-bridges by sortase A (Mazmanian et al., 1999; Schneewind et al., 1995; Mazmanian et al., 2000). Protein A is the most abundant surface protein of staphylococci; the molecule is expressed by virtually all *S. aureus* strains (Cespedes et al., 2005; Kennedy et al., 2008; Said-Salim et al., 2003). Staphylococci turn over 15-20% of their cell wall per division cycle (Navarre and Schneewind, 1999). Murine hydrolases cleave the glycan strands and wall peptides of peptidoglycan, thereby releasing Protein A with its attached C-terminal cell wall disaccharide tetrapeptide into the extracellular medium (Ton-That et al., 1999). Thus, by physiological design, Protein A is both anchored to the cell wall and displayed on the bacterial surface but also released into surrounding tissues during host infection (Marraffini et al., 2006).

Protein A captures immunoglobulins on the bacterial surface and this biochemical activity enables staphylococcal escape from host innate and acquired immune responses (Jensen, 1958; Goodyear et al., 2004). Interestingly, region X of Protein A (Guss et al., 1984), a repeat domain that tethers the IgG binding domains to the LPXTG sorting signal/cell wall anchor, is perhaps the most variable portion of the staphylococcal genome (Said-Salim, 2003; Schneewind et al., 1992). Each of the five immunoglobulin binding domains of Protein A (SpA), formed from three helix bundles and designated E, D, A, B, and C, exerts similar structural and functional properties (Sjodahl, 1977; Jansson et al., 1998). The solution and crystal structure of the domain D has been solved both with and without the Fc and V<sub>H</sub>3 (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille 2000).

In the crystal structure complex, the Fab interacts with helix II and helix III of domain D via a surface composed of four VH region  $\beta$ -strands (Graille 2000). The major axis of helix II of domain D is approximately 50° to the orientation of the strands, and the interhelical portion of domain D is most proximal to the CO strand. The site of interaction on Fab is remote from the Ig light chain and the heavy chain constant region. The interaction involves the following domain D residues: Asp-36 of helix II, Asp-37 and Gln-40 in the loop between helix II and helix III and several other residues (Graille 2000). Both interacting surfaces are composed predominantly of polar side chains, with three negatively charged residues on domain D and two positively charged residues on the 2A2 Fab buried by the interaction, providing an overall electrostatic attraction between the two molecules. Of the five polar interactions identified between Fab and domain D, three are between side chains. A salt bridge is formed between Arg-H19 and Asp-36 and two hydrogen bonds are made between Tyr-H59 and Asp-37 and between Asn-H82a and Ser-33. Because of the conservation of Asp-36 and Asp-37 in all five IgG binding domains of Protein A, the inventors mutated these residues.

The SpA-D sites responsible for Fab binding are structurally separate from the domain surface that mediates Fc $\gamma$  binding. The interaction of Fc $\gamma$  with domain D primarily involves residues in helix I with lesser involvement of helix II (Gouda et al., 1992; Deisenhofer, 1981). With the exception of the Gln-32, a minor contact in both complexes, none of the residues that mediate the Fc $\gamma$  interaction are involved in Fab binding. To examine the spatial relationship between these different Ig-binding sites, the SpA domains in these complexes have been superimposed to construct a model of a complex between Fab, the SpA-domain D, and the Fc $\gamma$

molecule. In this ternary model, Fab and Fc $\gamma$  form a sandwich about opposite faces of the helix II without evidence of steric hindrance of either interaction. These findings illustrate how, despite its small size (i.e., 56-61 aa), an SpA domain can simultaneously display both activities, explaining experimental evidence that the interactions of Fab with an individual domain are noncompetitive. Residues for the interaction between SpA-D and Fc $\gamma$  are Gln-9 and Gln-10.

In contrast, occupancy of the Fc portion of IgG on the domain D blocks its interaction with vWF A1 and probably also TNFR1 (O'Seaghdha et al., 2006). Mutations in residues essential for IgG Fc binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, 131 and K35) are also required for vWF A1 and TNFR1 binding (O'Seaghdha et al., 2006; Cedergren et al., 1993; Gomez et al., 2006), whereas residues critical for the VH3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) have no impact on the binding activities of IgG Fc, vWF A1 or TNFR1 (Jansson et al., 1998; Graille et al., 2000). The Protein A immunoglobulin Fab binding activity targets a subset of B cells that express V<sub>H</sub>3 family related IgM on their surface, i.e., these molecules function as VH3type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells rapidly proliferate and then commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e., marginal zone B cells and follicular B2 cells) (Goodyear and Silverman, 2004; Goodyear and Silverman, 2003). More than 40% of circulating B cells are targeted by the Protein A interaction and the V<sub>H</sub>3 family represents the largest family of human B cell receptors to impart protective humoral responses against pathogens (Goodyear and Silverman, 2004; Goodyear and Silverman, 2003). Thus, Protein A functions analogously to staphylococcal superantigens (Roben et al., 1995), albeit that the latter class of molecules, for example SEB, TSST-1, TSST-2, form complexes with the T cell receptor to inappropriately stimulate host immune responses and thereby precipitating characteristic disease features of staphylococcal infections (Roben et al., 1995; Tiedemann et al., 1995). Together these findings document the contributions of Protein A in establishing staphylococcal infections and in modulating host immune responses.

In sum, Protein A domains can be viewed as displaying two different interfaces for binding with host molecules and any development of Protein A based vaccines must consider the generation of variants that do not perturb host cell signaling, platelet aggregation, sequestration of immunoglobulins or the induction of B cell proliferation and apoptosis. Such Protein A variants should also be useful in analyzing vaccines for the ability of raising antibodies that block the aforementioned SpA activities and occupy the five repeat domains at their dual binding interfaces. This goal is articulated and pursued here for the first time and methods are described in detail for the generation of Protein A variants that can be used as a safe vaccine for humans. To perturb IgG Fc $\gamma$ , vWF A1 and TNFR1 binding, glutamine (Q) 9 and 10 [numbering derived from the SpA domain D as described in Uhlen et al., 1984] were mutated, and generated lysine substitutions for both glutamines with the expectation that these abolish the ligand attributes at the first binding interface. To perturb IgM Fab VH3 binding, aspartate (D) 36 and 37 were mutated, each of which is required for the association with the B cell receptor. D36 and D37 were both substituted with alanine. Q9,10K and D36,37A mutations are here combined in the recombinant molecule SpA-DQ9, 10K;D36,37A and tested for the binding attributes of Protein A. Further, SpA-D and SpA-DQ9,10K;D36,37A are subjected to immunization studies in mice and rabbits and

analyzed for [1] the production of specific antibodies (SpA-D Ab); [2] the ability of SpA-D Ab to block the association between Protein A and its four different ligands; and, [3] the attributes of SpA-D Ab to generate protective immunity against staphylococcal infections. (See Examples section below).

#### B. Staphylococcal Coagulases

Coagulases are enzymes produced by *Staphylococcus* bacteria that convert fibrinogen to fibrin. Coa and vW<sub>h</sub> activate prothrombin without proteolysis (Friedrich et al., 2003). The coagulase-prothrombin complex recognizes fibrinogen as a specific substrate, converting it directly into fibrin. The crystal structure of the active complex revealed binding of the D1 and D2 domains to prothrombin and insertion of its Ile1-Val<sup>2</sup> N-terminus into the Ile<sup>16</sup> pocket, inducing a functional active site in the zymogen through conformational change (Friedrich et al., 2003). Exosite I of  $\alpha$ -thrombin, the fibrinogen recognition site, and proexosite I on prothrombin are blocked by the D2 of Coa (Friedrich et al., 2003). Nevertheless, association of the tetramer (Coa-prothrombin)<sub>2</sub> complex binds fibrinogen at a new site with high affinity (Panizzi et al., 2006). This model explains the coagulant properties and efficient fibrinogen conversion by coagulase (Panizzi et al., 2006).

Fibrinogen is a large glycoprotein (Mr~340,000), formed by three pairs of A $\alpha$ -, B $\beta$ -, and  $\gamma$ -chains covalently linked to form a "dimer of trimers," where A and B designate the fibrinopeptides released by thrombin cleavage (Panizzi et al., 2006). The elongated molecule folds into three separate domains, a central fragment E that contains the N-termini of all six chains and two flanking fragments D formed mainly by the C-termini of the B $\beta$ - and  $\gamma$ -chains. These globular domains are connected by long triple-helical structures. Coagulase-prothrombin complexes, which convert human fibrinogen to the self-polymerizing fibrin, are not targeted by circulating thrombin inhibitors (Panizzi et al., 2006). Thus, staphylococcal coagulases bypass the physiological blood coagulation pathway.

All *S. aureus* strains secrete coagulase and vWbp (Bjerketorp et al., 2004; Field and Smith, 1945). Although early work reported important contributions of coagulase to the pathogenesis of staphylococcal infections (Ekstedt and Yotis, 1960; Smith et al., 1947), more recent investigations with molecular genetics tools challenged this view by observing no virulence phenotypes with endocarditis, skin abscess and mastitis models in mice (Moreillon et al., 1995; Phonimdaeng et al., 1990). Generating isogenic variants of *S. aureus* Newman, a fully virulent clinical isolate (Duthie et al., 1952), it is described herein that coa mutants indeed display virulence defects in a lethal bacteremia and renal abscess model in mice. In the inventors' experience, *S. aureus* 8325-4 is not fully virulent and it is presumed that mutational lesions in this strain may not be able to reveal virulence defects in vivo. Moreover, antibodies raised against Coa or vWbp perturb the pathogenesis of *S. aureus* Newman infections to a degree mirroring the impact of gene deletions. Coa and vWbp contribute to staphylococcal abscess formation and lethal bacteremia and may also function as protective antigens in subunit vaccines.

Biochemical studies document the biological value of antibodies against Coa and vWbp. By binding to antigen and blocking its association with clotting factors, the antibodies prevent the formation of Coa-prothrombin and vWbp-prothrombin complexes. Passive transfer studies revealed protection of experimental animals against staphylococcal abscess formation and lethal challenge by Coa and vWbp

antibodies. Thus, Coa and vWbp neutralizing antibodies generate immune protection against staphylococcal disease.

Earlier studies revealed a requirement of coagulase for resisting phagocytosis in blood (Smith et al., 1947) and the inventors observed a similar phenotype for  $\Delta$ coa mutants in lepirudin-treated mouse blood (see Example 3 below). As vWbp displays higher affinity for human prothrombin than the mouse counterpart, it is suspected the same may be true for  $\Delta$ vWbp variants in human blood. Further, expression of Coa and vWbp in abscess lesions as well as their striking distribution in the eosinophilic pseudocapsule surrounding (staphylococcal abscess communities (SACs) or the peripheral fibrin wall, suggest that secreted coagulases contribute to the establishment of these lesions. This hypothesis was tested and, indeed,  $\Delta$ coa mutants were defective in the establishment of abscesses. A corresponding test, blocking Coa function with specific antibodies, produced the same effect. Consequently, it is proposed that the clotting of fibrin is a critical event in the establishment of staphylococcal abscesses that can be targeted for the development of protective vaccines. Due to their overlapping function on human prothrombin, both Coa and vWbp are considered excellent candidates for vaccine development.

### C. Other Staphylococcal Antigens

Research over the past several decades identified *S. aureus* exotoxins, surface proteins and regulatory molecules as important virulence factors (Foster, 2005; Mazmanian et al., 2001; Novick, 2003). Much progress has been achieved regarding the regulation of these genes. For example, staphylococci perform a bacterial census via the secretion of auto-inducing peptides that bind to a cognate receptor at threshold concentration, thereby activating phospho-relay reactions and transcriptional activation of many of the exotoxin genes (Novick, 2003). The pathogenesis of staphylococcal infections relies on these virulence factors (secreted exotoxins, exopolysaccharides, and surface adhesins). The development of staphylococcal vaccines is hindered by the multifaceted nature of staphylococcal invasion mechanisms. It is well established that live attenuated micro-organisms are highly effective vaccines; immune responses elicited by such vaccines are often of greater magnitude and of longer duration than those produced by non-replicating immunogens. One explanation for this may be that live attenuated strains establish limited infections in the host and mimic the early stages of natural infection. Embodiments of the invention are directed to compositions and methods including variant SpA polypeptides and peptides, as well as other immunogenic extracellular proteins, polypeptides, and peptides (including both secreted and cell surface proteins or peptides) of gram positive bacteria for the use in mitigating or immunizing against infection. In particular embodiments the bacteria is a *Staphylococcus* bacteria. Extracellular proteins, polypeptides, or peptides include, but are not limited to secreted and cell surface proteins of the targeted bacteria.

The human pathogen *S. aureus* secretes EsxA and EsxB, two ESAT-6 like proteins, across the bacterial envelope (Burts et al., 2005, which is incorporated herein by reference). Staphylococcal *esxA* and *esxB* are clustered with six other genes in the order of transcription: *esxA esaA essA esaB essB essC esaC esxB*. The acronyms *esa*, *ess*, and *esx* stand for ESAT-6 secretion accessory, system, and extracellular, respectively, depending whether the encoded proteins play an accessory (*esa*) or direct (*ess*) role for secretion, or are secreted (*esx*) in the extracellular milieu. The entire cluster of eight genes is herein referred to as the *Ess* cluster. *EsxA*, *esxB*, *essA*, *essB*, and *essC* are all required for synthesis or secretion of *EsxA* and *EsxB*. Mutants that fail

to produce *EsxA*, *EsxB*, and *EssC* display defects in the pathogenesis of *S. aureus* murine abscesses, suggesting that this specialized secretion system may be a general strategy of human bacterial pathogenesis. Secretion of non-WXG100 substrates by the ESX-1 pathway has been reported for several antigens including *EspA*, *EspB*, *Rv3483c*, and *Rv3615c* (Fortune et al., 2005; MacGurn et al., 2005; McLaughlin et al., 2007; Xu et al., 2007). The alternate ESX-5 pathway has also been shown to secrete both WXG100 and non-WXG100 proteins in pathogenic mycobacteria (Abdallah et al., 2007; Abdallah et al., 2006).

The *Staphylococcus aureus* *Ess* pathway can be viewed as a secretion module equipped with specialized transport components (*Ess*), accessory factors (*Esa*) and cognate secretion substrates (*Esx*). *EssA*, *EssB* and *EssC* are required for *EsxA* and *EsxB* secretion. Because *EssA*, *EssB* and *EssC* are predicted to be transmembrane proteins, it is contemplated that these proteins form a secretion apparatus. Some of the proteins in the *ess* gene cluster may actively transport secreted substrates (acting as motor) while others may regulate transport (regulator). Regulation may be achieved, but need not be limited to, transcriptional or post-translational mechanisms for secreted polypeptides, sorting of specific substrates to defined locations (e.g., extracellular medium or host cells), or timing of secretion events during infection. At this point, it is unclear whether all secreted *Esx* proteins function as toxins or contribute indirectly to pathogenesis.

Staphylococci rely on surface protein mediated-adhesion to host cells or invasion of tissues as a strategy for escape from immune defenses. Furthermore, *S. aureus* utilize surface proteins to sequester iron from the host during infection. The majority of surface proteins involved in staphylococcal pathogenesis carry C-terminal sorting signals, i.e., they are covalently linked to the cell wall envelope by sortase. Further, staphylococcal strains lacking the genes required for surface protein anchoring, i.e., sortase A and B, display a dramatic defect in the virulence in several different mouse models of disease. Thus, surface protein antigens represent a validated vaccine target as the corresponding genes are essential for the development of staphylococcal disease and can be exploited in various embodiments of the invention. The sortase enzyme superfamily are Gram-positive transpeptidases responsible for anchoring surface protein virulence factors to the peptidoglycan cell wall layer. Two sortase isoforms have been identified in *Staphylococcus aureus*, *SrtA* and *SrtB*. These enzymes have been shown to recognize a LPXTG motif in substrate proteins. The *SrtB* isoform appears to be important in heme iron acquisition and iron homeostasis, whereas the *SrtA* isoform plays a critical role in the pathogenesis of Gram-positive bacteria by modulating the ability of the bacterium to adhere to host tissue via the covalent anchoring of adhesins and other proteins to the cell wall peptidoglycan. In certain embodiments the *SpA* variants described herein can be used in combination with other staphylococcal proteins such as *Coa*, *Eap*, *Ehb*, *Emp*, *EsaC*, *EsaB*, *EsxA*, *EsxB*, *Hla*, *SdrC*, *SdrD*, *SdrE*, *IsdA*, *IsdB*, *ClfA*, *ClfB*, *IsdC*, *SasF*, *vWbp*, and/or *vWh* proteins.

Certain aspects of the invention include methods and compositions concerning proteinaceous compositions including polypeptides, peptides, or nucleic acid encoding *SpA* variant(s) and other staphylococcal antigens such as other proteins transported by the *Ess* pathway, or sortase substrates. These proteins may be modified by deletion, insertion, and/or substitution.

The *Esx* polypeptides include the amino acid sequence of *Esx* proteins from bacteria in the *Staphylococcus* genus. The

Esx sequence may be from a particular *Staphylococcus* species, such as *Staphylococcus aureus*, and may be from a particular strain, such as Newman. In certain embodiments, the EsxA sequence is SAV0282 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number Q99WU4 (gil68565539), which is hereby incorporated by reference. In other embodiments, the EsxB sequence is SAV0290 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number Q99WT7 (gil68565532), which is hereby incorporated by reference. In further embodiments, other polypeptides transported by the Ess pathway may be used, the sequences of which may be identified by one of skill in the art using databases and internet accessible resources.

The sortase substrate polypeptides include, but are not limited to the amino acid sequence of SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, IsdC or SasF proteins from bacteria in the *Staphylococcus* genus. The sortase substrate polypeptide sequence may be from a particular *Staphylococcus* species, such as *Staphylococcus aureus*, and may be from a particular strain, such as Newman. In certain embodiments, the SdrD sequence is from strain N315 and can be accessed using Genbank Accession Number NP\_373773.1 (gil15926240), which is incorporated by reference. In other embodiments, the SdrE sequence is from strain N315 and can be accessed using Genbank Accession Number NP\_373774.1 (gil15926241), which is incorporated by reference. In other embodiments, the IsdA sequence is SAV1130 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number NP\_371654.1 (gil15924120), which is incorporated by reference. In other embodiments, the IsdB sequence is SAV1129 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number NP\_371653.1 (gil15924119), which is incorporated by reference. In further embodiments, other polypeptides transported by the Ess pathway or processed by sortase may be used, the sequences of which may be identified by one of skill in the art using databases and internet accessible resources.

Examples of various proteins that can be used in the context of the present invention can be identified by analysis of database submissions of bacterial genomes, including but not limited to accession numbers NC\_002951 (GI:57650036 and GenBank CP000046), NC\_002758 (GI:57634611 and GenBank BA000017), NC\_002745 (GI:29165615 and GenBank BA000018), NC\_003923 (GI:21281729 and GenBank BA000033), NC\_002952 (GI:49482253 and GenBank BX571856), NC\_002953 (GI:49484912 and GenBank BX571857), NC\_007793 (GI:87125858 and GenBank CP000255), NC\_007795 (GI:87201381 and GenBank CP000253) each of which are incorporated by reference.

As used herein, a "protein" or "polypeptide" refers to a molecule comprising at least ten amino acid residues. In some embodiments, a wild-type version of a protein or polypeptide are employed, however, in many embodiments of the invention, a modified protein or polypeptide is employed to generate an immune response. The terms described above may be used interchangeably. A "modified protein" or "modified polypeptide" or a "variant" refers to a protein or polypeptide whose chemical structure, particularly its amino acid sequence, is altered with respect to the wild-type protein or polypeptide. In some embodiments, a modified/variant protein or polypeptide has at least one modified activity or function (recognizing that proteins or polypeptides may have multiple activities or functions). It is

specifically contemplated that a modified/variant protein or polypeptide may be altered with respect to one activity or function yet retain a wild-type activity or function in other respects, such as immunogenicity.

In certain embodiments the size of a protein or polypeptide (wild-type or modified) may comprise, but is not limited to, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 875, 900, 925, 950, 975, 1000, 1100, 1200, 1300, 1400, 1500, 1750, 2000, 2250, 2500 amino molecules or greater, and any range derivable therein, or derivative of a corresponding amino sequence described or referenced herein. It is contemplated that polypeptides may be mutated by truncation, rendering them shorter than their corresponding wild-type form, but also they might be altered by fusing or conjugating a heterologous protein sequence with a particular function (e.g., for targeting or localization, for enhanced immunogenicity, for purification purposes, etc.).

As used herein, an "amino molecule" refers to any amino acid, amino acid derivative, or amino acid mimic known in the art. In certain embodiments, the residues of the proteinaceous molecule are sequential, without any non-amino molecule interrupting the sequence of amino molecule residues. In other embodiments, the sequence may comprise one or more non-amino molecule moieties. In particular embodiments, the sequence of residues of the proteinaceous molecule may be interrupted by one or more non-amino molecule moieties.

Accordingly, the term "proteinaceous composition" encompasses amino molecule sequences comprising at least one of the 20 common amino acids in naturally synthesized proteins, or at least one modified or unusual amino acid.

Proteinaceous compositions may be made by any technique known to those of skill in the art, including (i) the expression of proteins, polypeptides, or peptides through standard molecular biological techniques, (ii) the isolation of proteinaceous compounds from natural sources, or (iii) the chemical synthesis of proteinaceous materials. The nucleotide as well as the protein, polypeptide, and peptide sequences for various genes have been previously disclosed, and may be found in the recognized computerized databases. One such database is the National Center for Biotechnology Information's Genbank and GenPept databases (on the World Wide Web at [ncbi.nlm.nih.gov/](http://ncbi.nlm.nih.gov/)). The coding regions for these genes may be amplified and/or expressed using the techniques disclosed herein or as would be known to those of ordinary skill in the art.

Amino acid sequence variants of SpA, coagulases and other polypeptides of the invention can be substitutional, insertional, or deletion variants. A variation in a polypeptide of the invention may affect 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, or more non-contiguous or contiguous amino acids of the polypeptide, as compared to wild-type. A variant can comprise an amino acid sequence that is at least 50%, 60%, 70%, 80%, or 90%, including all values and ranges there between, identical to any sequence provided or referenced herein, e.g., SEQ ID NO:2-8 or SEQ ID NO:11-30, A variant can include 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13,

14, 15, 16, 17, 18, 19, 20, or more substitute amino acids. A polypeptide processed or secreted by the Ess pathway or other surface proteins (see Table 1) or sortase substrates from any *Staphylococcus* species and strain are contemplated for use in compositions and methods described herein.

Deletion variants typically lack one or more residues of the native or wild-type protein. Individual residues can be deleted or a number of contiguous amino acids can be deleted. A stop codon may be introduced (by substitution or insertion) into an encoding nucleic acid sequence to generate a truncated protein. Insertional mutants typically involve the addition of material at a non-terminal point in the polypeptide. This may include the insertion of one or more residues. Terminal additions, called fusion proteins, may also be generated. These fusion proteins include multimers or concatamers of one or more peptide or polypeptide described or referenced herein.

Substitutional variants typically contain the exchange of one amino acid for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide, with or without the loss of other functions or properties. Substitutions may be conservative, that is, one amino acid is replaced with one of similar shape and charge. Conservative substitutions are well known in the art and include, for example, the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; and valine to isoleucine or leucine. Alternatively, substitutions may be non-conservative such that a function or activity of the polypeptide is affected. Non-conservative changes typically involve substituting a residue with one that is chemically dissimilar, such as a polar or charged amino acid for a nonpolar or uncharged amino acid, and vice versa.

TABLE 2

Exemplary surface proteins of <i>S. aureus</i> strains.								
SAV #	SA#	Surface	MW2	Mu50	N315	Newman	MRSA252*	MSSA476*
SAV0111	SA0107	Spa	492	450	450	520	516	492
SAV2503	SA2291	FnBPA	1015	1038	1038	741	—	1015
SAV2502	SA2290	FnBPB	943	961	961	677	965	957
SAV0811	SA0742	ClfA	946	935	989	933	1029	928
SAV2630	SA2423	ClfB	907	877	877	913	873	905
Np	Np	Cna	1183	—	—	—	1183	1183
SAV0561	SA0519	SdrC	955	953	953	947	906	957
SAV0562	SA0520	SdrD	1347	1385	1385	1315	—	1365
SAV0563	SA0521	SdrE	1141	1141	1141	1166	1137	1141
Np	Np	Pls	—	—	—	—	—	—
SAV2654	SA2447	SasA	2275	2271	2271	2271	1351	2275
SAV2160	SA1964	SasB	686	2481	2481	2481	2222	685
	SA1577	SasC	2186	213	2186	2186	2189	2186
SAV0134	SA0129	SasD	241	241	241	241	221	241
SAV1130	SA0977	SasE/IsdA	350	350	350	350	354	350
SAV2646	SA2439	SasF	635	635	635	635	627	635
SAV2496		SasG	1371	525	927	—	—	1371
SAV0023	SA0022	SasH	772	—	772	772	786	786
SAV1731	SA1552	SasI	895	891	891	891	534	895
SAV1129	SA0976	SasJ/IsdB	645	645	645	645	652	645
	SA2381	SasK	198	211	211	—	—	197
	Np	SasL	—	232	—	—	—	—
SAV1131	SA0978	IsdC	227	227	227	227	227	227

Proteins of the invention may be recombinant, or synthesized in vitro. Alternatively, a non-recombinant or recombinant protein may be isolated from bacteria. It is also contemplated that a bacteria containing such a variant may be implemented in compositions and methods of the invention. Consequently, a protein need not be isolated.

The term “functionally equivalent codon” is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine, and also refers to codons that encode biologically equivalent amino acids (see Table 2, below).

TABLE 3

Codon Table			
Amino Acids			Codons
Alanine	Ala	A	GCA GCC GCG GCU
Cysteine	Cys	C	UGC UGU
Aspartic acid	Asp	D	GAC GAU
Glutamic acid	Glu	E	GAA GAG
Phenylalanine	Phe	F	UUC UUU
Glycine	Gly	G	GGA GGC GGG GGU
Histidine	His	H	CAC CAU
Isoleucine	Ile	I	AUA AUC AUU
Lysine	Lys	K	AAA AAG
Leucine	Leu	L	UUA UUG CUA CUC CUG CUU
Methionine	Met	M	AUG
Asparagine	Asn	N	AAC AAU
Proline	Pro	P	CCA CCC CCG CCU
Glutamine	Gln	Q	CAA CAG
Arginine	Arg	R	AGA AGG CGA CGC CGG CGU
Serine	Ser	S	AGC AGU UCA UCC UCG UCU
Threonine	Thr	T	ACA ACC ACG ACU
Valine	Val	V	GUA GUC GUG GUU
Tryptophan	Trp	W	UGG
Tyrosine	Tyr	Y	UAC UAU

It also will be understood that amino acid and nucleic acid sequences may include additional residues, such as additional N- or C-terminal amino acids, or 5' or 3' sequences, respectively, and yet still be essentially as set forth in one of the sequences disclosed herein, so long as the sequence meets the criteria set forth above, including the maintenance

of biological protein activity (e.g., immunogenicity) where protein expression is concerned. The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region.

The following is a discussion based upon changing of the amino acids of a protein to create a variant polypeptide or peptide. For example, certain amino acids may be substituted for other amino acids in a protein structure with or without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's functional activity, certain amino acid substitutions can be made in a protein sequence, and in its underlying DNA coding sequence, and nevertheless produce a protein with a desirable property. It is thus contemplated by the inventors that various changes may be made in the DNA sequences of genes.

It is contemplated that in compositions of the invention, there is between about 0.001 mg and about 10 mg of total polypeptide, peptide, and/or protein per ml. The concentration of protein in a composition can be about, at least about or at most about 0.001, 0.010, 0.050, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5, 10.0 mg/ml or more (or any range derivable therein).

Of this, about, at least about, or at most about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100% may be an SpA variant or a coagulase, and may be used in combination with other peptides or polypeptides, such as other bacterial peptides and/or antigens.

The present invention contemplates the administration of variant SpA polypeptides or peptides to effect a preventative therapy or therapeutic effect against the development of a disease or condition associated with infection by a *Staphylococcus* pathogen.

In certain aspects, combinations of staphylococcal antigens are used in the production of an immunogenic composition that is effective at treating or preventing staphylococcal infection. Staphylococcal infections progress through several different stages. For example, the staphylococcal life cycle involves commensal colonization, initiation of infection by accessing adjoining tissues or the bloodstream, and/or anaerobic multiplication in the blood. The interplay between *S. aureus* virulence determinants and the host defense mechanisms can induce complications such as endocarditis, metastatic abscess formation, and sepsis syndrome. Different molecules on the surface of the bacterium are involved in different steps of the infection cycle. Combinations of certain antigens can elicit an immune response which protects against multiple stages of staphylococcal infection. The effectiveness of the immune response can be measured either in animal model assays and/or using an opsonophagocytic assay.

#### D. Polypeptides and Polypeptide Production

The present invention describes polypeptides, peptides, and proteins and immunogenic fragments thereof for use in various embodiments of the present invention. For example, specific polypeptides are assayed for or used to elicit an immune response. In specific embodiments, all or part of the proteins of the invention can also be synthesized in solution

or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, (1984); Tam et al., (1983); Merrifield, (1986); and Barany and Merrifield (1979), each incorporated herein by reference.

Alternatively, recombinant DNA technology may be employed wherein a nucleotide sequence which encodes a peptide of the invention is inserted into an expression vector, transformed or transfected into an appropriate host cell and cultivated under conditions suitable for expression.

One embodiment of the invention includes the use of gene transfer to cells, including microorganisms, for the production and/or presentation of polypeptides or peptides. The gene for the polypeptide or peptide of interest may be transferred into appropriate cells followed by culture of cells under the appropriate conditions. The generation of recombinant expression vectors, and the elements included therein, are well known in the art and briefly discussed herein. Alternatively, the protein to be produced may be an endogenous protein normally synthesized by the cell that is isolated and purified.

Another embodiment of the present invention uses autologous B lymphocyte cell lines, which are transfected with a viral vector that expresses an immunogen product, and more specifically, a protein having immunogenic activity. Other examples of mammalian host cell lines include, but are not limited to Vero and HeLa cells, other B- and T-cell lines, such as CEM, 721.221, H9, Jurkat, Raji, as well as cell lines of Chinese hamster ovary, W138, BHK, COS-7, 293, HepG2, 3T3, RIN and MDCK cells. In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or that modifies and processes the gene product in the manner desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed.

A number of selection systems may be used including, but not limited to HSV thymidine kinase, hypoxanthine-guanine phosphoribosyltransferase, and adenine phosphoribosyltransferase genes, in tk-, hgprrt- or aprt-cells, respectively. Also, anti-metabolite resistance can be used as the basis of selection: for dhfr, which confers resistance to trimethoprim and methotrexate; gpt, which confers resistance to mycophenolic acid; neo, which confers resistance to the aminoglycoside G418; and hygromycin, which confers resistance to hygromycin.

Animal cells can be propagated in vitro in two modes: as non-anchorage-dependent cells growing in suspension throughout the bulk of the culture or as anchorage-dependent cells requiring attachment to a solid substrate for their propagation (i.e., a monolayer type of cell growth).

Non-anchorage dependent or suspension cultures from continuous established cell lines are the most widely used means of large scale production of cells and cell products. However, suspension cultured cells have limitations, such as tumorigenic potential and lower protein production than adherent cells.

Where a protein is specifically mentioned herein, it is preferably a reference to a native or recombinant protein or optionally a protein in which any signal sequence has been removed. The protein may be isolated directly from the staphylococcal strain or produced by recombinant DNA

techniques. Immunogenic fragments of the protein may be incorporated into the immunogenic composition of the invention. These are fragments comprising at least 10 amino acids, 20 amino acids, 30 amino acids, 40 amino acids, 50 amino acids, or 100 amino acids, including all values and ranges there between, taken contiguously from the amino acid sequence of the protein. In addition, such immunogenic fragments are immunologically reactive with antibodies generated against the Staphylococcal proteins or with antibodies generated by infection of a mammalian host with Staphylococci. Immunogenic fragments also include fragments that when administered at an effective dose, (either alone or as a hapten bound to a carrier), elicit a protective or therapeutic immune response against Staphylococcal infection, in certain aspects it is protective against *S. aureus* and/or *S. epidermidis* infection. Such an immunogenic fragment may include, for example, the protein lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment according to the invention comprises substantially all of the extracellular domain of a protein which has at least 80% identity, at least 85% identity, at least 90% identity, at least 95% identity, or at least 97-99% identity, including all values and ranges there between, to a sequence selected segment of a polypeptide described or referenced herein.

Also included in immunogenic compositions of the invention are fusion proteins composed of one or more Staphylococcal proteins, or immunogenic fragments of staphylococcal proteins. Such fusion proteins may be made recombinantly and may comprise one portion of at least 1, 2, 3, 4, 5, or 6 staphylococcal proteins or segments. Alternatively, a fusion protein may comprise multiple portions of at least 1, 2, 3, 4 or 5 staphylococcal proteins. These may combine different Staphylococcal proteins and/or multiples of the same protein or protein fragment, or immunogenic fragments in the same protein (forming a multimer or a concatamer). Alternatively, the invention also includes individual fusion proteins of Staphylococcal proteins or immunogenic fragments thereof, as a fusion protein with heterologous sequences such as a provider of T-cell epitopes or purification tags, for example:  $\beta$ -galactosidase, glutathione-S-transferase, green fluorescent proteins (GFP), epitope tags such as FLAG, myc tag, poly histidine, or viral surface proteins such as influenza virus haemagglutinin, or bacterial proteins such as tetanus toxoid, diphtheria toxoid, or CRM197.

## II. Nucleic Acids

In certain embodiments, the present invention concerns recombinant polynucleotides encoding the proteins, polypeptides, peptides of the invention. The nucleic acid sequences for SpA, coagulases and other bacterial proteins are included, all of which are incorporated by reference, and can be used to prepare peptides or polypeptides.

As used in this application, the term "polynucleotide" refers to a nucleic acid molecule that either is recombinant or has been isolated free of total genomic nucleic acid. Included within the term "polynucleotide" are oligonucleotides (nucleic acids of 100 residues or less in length), recombinant vectors, including, for example, plasmids, cosmids, phage, viruses, and the like. Polynucleotides include, in certain aspects, regulatory sequences, isolated substantially away from their naturally occurring genes or protein encoding sequences. Polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be RNA, DNA (genomic, cDNA or synthetic), analogs

thereof, or a combination thereof. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide.

In this respect, the term "gene," "polynucleotide," or "nucleic acid" is used to refer to a nucleic acid that encodes a protein, polypeptide, or peptide (including any sequences required for proper transcription, post-translational modification, or localization). As will be understood by those in the art, this term encompasses genomic sequences, expression cassettes, cDNA sequences, and smaller engineered nucleic acid segments that express, or may be adapted to express, proteins, polypeptides, domains, peptides, fusion proteins, and mutants. A nucleic acid encoding all or part of a polypeptide may contain a contiguous nucleic acid sequence of: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 441, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, 1000, 1010, 1020, 1030, 1040, 1050, 1060, 1070, 1080, 1090, 1095, 1100, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 9000, 10000, or more nucleotides, nucleosides, or base pairs, including all values and ranges therebetween, of a polynucleotide encoding one or more amino acid sequence described or referenced herein. It also is contemplated that a particular polypeptide may be encoded by nucleic acids containing variations having slightly different nucleic acid sequences but, nonetheless, encode the same or substantially similar protein (see Table 3 above).

In particular embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a variant SpA or coagulase. The term "recombinant" may be used in conjunction with a polynucleotide or polypeptide and generally refers to a polypeptide or polynucleotide produced and/or manipulated in vitro or that is a replication product of such a molecule.

In other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a variant SpA or coagulase polypeptide or peptide to generate an immune response in a subject. In various embodiments the nucleic acids of the invention may be used in genetic vaccines.

The nucleic acid segments used in the present invention can be combined with other nucleic acid sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant nucleic acid protocol. In some cases, a nucleic acid sequence may encode a polypeptide sequence with additional heterologous coding sequences, for example to allow for purification of the polypeptide, transport, secretion, post-translational modification, or for therapeutic benefits such as targeting or efficacy. As discussed above, a tag or other heterologous polypeptide may be added to the modified polypeptide-encoding sequence, wherein "heterologous" refers to a polypeptide that is not the same as the modified polypeptide.

In certain other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors that

include within their sequence a contiguous nucleic acid sequence from SEQ ID NO:1 (SpA domain D) or SEQ ID NO:3 (SpA) or any other nucleic acid sequences encoding coagulases or other secreted virulence factors and/or surface proteins including proteins transported by the Ess pathway, processed by sortase, or proteins incorporated herein by reference.

In certain embodiments, the present invention provides polynucleotide variants having substantial identity to the sequences disclosed herein; those comprising at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher sequence identity, including all values and ranges there between, compared to a polynucleotide sequence of this invention using the methods described herein (e.g., BLAST analysis using standard parameters).

The invention also contemplates the use of polynucleotides which are complementary to all the above described polynucleotides.

#### A. Vectors

Polypeptides of the invention may be encoded by a nucleic acid molecule comprised in a vector. The term "vector" is used to refer to a carrier nucleic acid molecule into which a heterologous nucleic acid sequence can be inserted for introduction into a cell where it can be replicated and expressed. A nucleic acid sequence can be "heterologous," which means that it is in a context foreign to the cell in which the vector is being introduced or to the nucleic acid in which is incorporated, which includes a sequence homologous to a sequence in the cell or nucleic acid but in a position within the host cell or nucleic acid where it is ordinarily not found. Vectors include DNAs, RNAs, plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (e.g., YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (for example Sambrook et al., 2001; Ausubel et al., 1996, both incorporated herein by reference). In addition to encoding a variant SpA polypeptide the vector can encode other polypeptide sequences such as a one or more other bacterial peptide, a tag, or an immunogenicity enhancing peptide. Useful vectors encoding such fusion proteins include pIN vectors (Inouye et al., 1985), vectors encoding a stretch of histidines, and pGEX vectors, for use in generating glutathione S-transferase (GST) soluble fusion proteins for later purification and separation or cleavage.

The term "expression vector" refers to a vector containing a nucleic acid sequence coding for at least part of a gene product capable of being transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. Expression vectors can contain a variety of "control sequences," which refer to nucleic acid sequences necessary for the transcription and possibly translation of an operably linked coding sequence in a particular host organism. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described herein.

#### 1. Promoters and Enhancers

A "promoter" is a control sequence. The promoter is typically a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind such as RNA polymerase and other transcription factors. The phrases "operatively positioned," "operatively linked," "under control," and "under transcriptional control" mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid

sequence to control transcriptional initiation and expression of that sequence. A promoter may or may not be used in conjunction with an "enhancer," which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

Naturally, it may be important to employ a promoter and/or enhancer that effectively directs the expression of the DNA segment in the cell type or organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression (see Sambrook et al., 2001, incorporated herein by reference). The promoters employed may be constitutive, tissue-specific, or inducible and in certain embodiments may direct high level expression of the introduced DNA segment under specified conditions, such as large-scale production of recombinant proteins or peptides.

Various elements/promoters may be employed in the context of the present invention to regulate the expression of a gene. Examples of such inducible elements, which are regions of a nucleic acid sequence that can be activated in response to a specific stimulus, include but are not limited to Immunoglobulin Heavy Chain (Banerji et al., 1983; Gilles et al., 1983; Grosschedl et al., 1985; Atchinson et al., 1986, 1987; Imler et al., 1987; Weinberger et al., 1984; Kiledjian et al., 1988; Porton et al.; 1990), Immunoglobulin Light Chain (Queen et al., 1983; Picard et al., 1984), T Cell Receptor (Luria et al., 1987; Winoto et al., 1989; Redondo et al.; 1990), HLA DQ  $\alpha$  and/or DQ  $\beta$  (Sullivan et al., 1987),  $\beta$  Interferon (Goodbourn et al., 1986; Fujita et al., 1987; Goodbourn et al., 1988), Interleukin-2 (Greene et al., 1989), Interleukin-2 Receptor (Greene et al., 1989; Lin et al., 1990), MHC Class II 5 (Koch et al., 1989), MHC Class II HLA-DR $\alpha$  (Sherman et al., 1989),  $\beta$ -Actin (Kawamoto et al., 1988; Ng et al.; 1989), Muscle Creatine Kinase (MCK) (Jaynes et al., 1988; Horlick et al., 1989; Johnson et al., 1989), Prealbumin (Transthyretin) (Costa et al., 1988), Elastase I (Ornitz et al., 1987), Metallothionein (MTII) (Karin et al., 1987; Culotta et al., 1989), Collagenase (Pinkert et al., 1987; Angel et al., 1987), Albumin (Pinkert et al., 1987; Tronche et al., 1989, 1990),  $\alpha$ -Fetoprotein (Godbout et al., 1988; Campere et al., 1989),  $\gamma$ -Globin (Bodine et al., 1987; Perez-Stable et al., 1990), 13-Globin (Trudel et al., 1987), c-fos (Cohen et al., 1987), c-Ha-Ras (Triesman, 1986; Deschamps et al., 1985), Insulin (Edlund et al., 1985), Neural Cell Adhesion Molecule (NCAM) (Hirsh et al., 1990),  $\alpha$ 1-Antitrypsin (Latimer et al., 1990), H2B (TH2B) Histone (Hwang et al., 1990), Mouse and/or Type I Collagen (Ripe et al., 1989), Glucose-Regulated Proteins (GRP94 and GRP78) (Chang et al., 1989), Rat Growth Hormone (Larsen et al., 1986), Human Serum Amyloid A (SAA) (Edbrooke et al., 1989), Troponin I (TN I) (Yutzey et al., 1989), Platelet-Derived Growth Factor (PDGF) (Pech et al., 1989), Duchenne Muscular Dystrophy (Klamut et al., 1990), SV40 (Banerji et al., 1981; Moreau et al., 1981; Sleight et al., 1985; Firak et al., 1986; Herr et al., 1986; Imbra et al., 1986; Kadesch et al., 1986; Wang et al., 1986; Ondek et al., 1987; Kuhl et al., 1987; Schaffner et al., 1988), Polyoma (Swartzendruber et al., 1975; Vasseur et al., 1980; Katinka et al., 1980, 1981; Tyndell et al., 1981; Dandolo et al., 1983; de Villiers et al., 1984; Hen et al., 1986; Satake et al., 1988; Campbell et al., 1988), Retroviruses (Kriegler et al., 1982, 1983; Levinson et al., 1982; Kriegler et al., 1983, 1984a, b, 1988; Bosze et al., 1986; Miksicek et al., 1986; Celander et al., 1987; Thiesen et al., 1988; Celander et al., 1988; Choi et al., 1988; Reisman et al., 1989), Papilloma Virus (Campo et al., 1983; Lusky et al.,

1983; Spandidos and Wilkie, 1983; Spalholz et al., 1985; Lusky et al., 1986; Cripe et al., 1987; Gloss et al., 1987; Hirochika et al., 1987; Stephens et al., 1987), Hepatitis B Virus (Bulla et al., 1986; Jameel et al., 1986; Shaul et al., 1987; Spandau et al., 1988; Vannice et al., 1988), Human Immunodeficiency Virus (Muesing et al., 1987; Hauber et al., 1988; Jakobovits et al., 1988; Feng et al., 1988; Takebe et al., 1988; Rosen et al., 1988; Berkhout et al., 1989; Laspia et al., 1989; Sharp et al., 1989; Braddock et al., 1989), Cytomegalovirus (CMV) IE (Weber et al., 1984; Boshart et al., 1985; Foecking et al., 1986), Gibbon Ape Leukemia Virus (Holbrook et al., 1987; Quinn et al., 1989).

Inducible elements include, but are not limited to MT II-Phorbol Ester (TFA)/Heavy metals (Palmiter et al., 1982; Haslinger et al., 1985; Searle et al., 1985; Stuart et al., 1985; Imagawa et al., 1987; Karin et al., 1987; Angel et al., 1987b; McNeall et al., 1989); MMTV (mouse mammary tumor virus)-Glucocorticoids (Huang et al., 1981; Lee et al., 1981; Majors et al., 1983; Chandler et al., 1983; Lee et al., 1984; Ponta et al., 1985; Sakai et al., 1988);  $\beta$ -Interferon-poly(rI) x/poly(rc) (Tavernier et al., 1983); Adenovirus 5E2-E1A (Imperiale et al., 1984); Collagenase-Phorbol Ester (TPA) (Angel et al., 1987a); Stromelysin-Phorbol Ester (TPA) (Angel et al., 1987b); SV40-Phorbol Ester (TPA) (Angel et al., 1987b); Murine MX Gene-Interferon, Newcastle Disease Virus (Hug et al., 1988); GRP78 Gene-A23187 (Resendez et al., 1988);  $\alpha$ -2-Macroglobulin-IL-6 (Kunz et al., 1989); Vimentin-Serum (Rittling et al., 1989); MHC Class I Gene H-2kb-Interferon (Blonar et al., 1989); HSP70-E1A/SV40 Large T Antigen (Taylor et al., 1989, 1990a, 1990b); Proliferin-Phorbol Ester/TPA (Mordacq et al., 1989); Tumor Necrosis Factor-PMA (Hensel et al., 1989); and Thyroid Stimulating Hormone a Gene-Thyroid Hormone (Chatterjee et al., 1989).

The particular promoter that is employed to control the expression of peptide or protein encoding polynucleotide of the invention is not believed to be critical, so long as it is capable of expressing the polynucleotide in a targeted cell, preferably a bacterial cell. Where a human cell is targeted, it is preferable to position the polynucleotide coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a bacterial, human or viral promoter.

In embodiments in which a vector is administered to a subject for expression of the protein, it is contemplated that a desirable promoter for use with the vector is one that is not down-regulated by cytokines or one that is strong enough that even if down-regulated, it produces an effective amount of a variant SpA for eliciting an immune response. Non-limiting examples of these are CMV IE and RSV LTR. Tissue specific promoters can be used, particularly if expression is in cells in which expression of an antigen is desirable, such as dendritic cells or macrophages. The mammalian MEW I and MEW II promoters are examples of such tissue-specific promoters.

## 2. Initiation Signals and Internal Ribosome Binding Sites (IRES)

A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals.

In certain embodiments of the invention, the use of internal ribosome entry sites (IRES) elements are used to

create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, 1988; Macejak and Sarnow, 1991). IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be transcribed together, each separated by an IRES, creating polycistronic messages. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message (see U.S. Pat. Nos. 5,925,565 and 5,935,819, herein incorporated by reference).

## 3. Selectable and Screenable Markers

In certain embodiments of the invention, cells containing a nucleic acid construct of the present invention may be identified *in vitro* or *in vivo* by encoding a screenable or selectable marker in the expression vector. When transcribed and translated, a marker confers an identifiable change to the cell permitting easy identification of cells containing the expression vector. Generally, a selectable marker is one that confers a property that allows for selection. A positive selectable marker is one in which the presence of the marker allows for its selection, while a negative selectable marker is one in which its presence prevents its selection. An example of a positive selectable marker is a drug resistance marker.

## B. Host Cells

As used herein, the terms "cell," "cell line," and "cell culture" may be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. In the context of expressing a heterologous nucleic acid sequence, "host cell" refers to a prokaryotic or eukaryotic cell, and it includes any transformable organism that is capable of replicating a vector or expressing a heterologous gene encoded by a vector. A host cell can, and has been, used as a recipient for vectors or viruses. A host cell may be "transfected" or "transformed," which refers to a process by which exogenous nucleic acid, such as a recombinant protein-encoding sequence, is transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny.

Host cells may be derived from prokaryotes or eukaryotes, including bacteria, yeast cells, insect cells, and mammalian cells for replication of the vector or expression of part or all of the nucleic acid sequence(s). Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection (ATCC), which is an organization that serves as an archive for living cultures and genetic materials ([www.atcc.org](http://www.atcc.org)).

## C. Expression Systems

Numerous expression systems exist that comprise at least a part or all of the compositions discussed above. Prokaryote- and/or eukaryote-based systems can be employed for use with the present invention to produce nucleic acid sequences, or their cognate polypeptides, proteins and peptides. Many such systems are commercially and widely available.

The insect cell/baculovirus system can produce a high level of protein expression of a heterologous nucleic acid segment, such as described in U.S. Pat. Nos. 5,871,986, 4,879,236, both herein incorporated by reference, and which can be bought, for example, under the name MAXBAC® 2.0 from INVITROGEN® and BACPACK™ BACULOVIRUS EXPRESSION SYSTEM FROM CLONTECH®.

In addition to the disclosed expression systems of the invention, other examples of expression systems include STRATAGENE®'s COMPLETE CONTROL™ Inducible Mammalian Expression System, which involves a synthetic ecdysone-inducible receptor, or its pET Expression System, an *E. coli* expression system. Another example of an inducible expression system is available from INVITROGEN®, which carries the T-REX™ (tetracycline-regulated expression) System, an inducible mammalian expression system that uses the full-length CMV promoter. INVITROGEN® also provides a yeast expression system called the *Pichia methanolica* Expression System, which is designed for high-level production of recombinant proteins in the methylotrophic yeast *Pichia methanolica*. One of skill in the art would know how to express a vector, such as an expression construct, to produce a nucleic acid sequence or its cognate polypeptide, protein, or peptide.

### III. Polysaccharides

The immunogenic compositions of the invention may further comprise capsular polysaccharides including one or more of PIA (also known as PNAG) and/or *S. aureus* Type V and/or type VIII capsular polysaccharide and/or *S. epidermidis* Type I, and/or Type II and/or Type III capsular polysaccharide.

#### A. PIA (PNAG)

It is now clear that the various forms of staphylococcal surface polysaccharides identified as PS/A, PIA and SAA are the same chemical entity—PNAG (Maira-Litran et al., 2004). Therefore the term PIA or PNAG encompasses all these polysaccharides or oligosaccharides derived from them.

PIA is a polysaccharide intercellular adhesin and is composed of a polymer of  $\beta$ -(1 $\rightarrow$ 6)-linked glucosamine substituted with N-acetyl and O-succinyl constituents. This polysaccharide is present in both *S. aureus* and *S. epidermidis* and can be isolated from either source (Joyce et al., 2003; Maira-Litran et al., 2002). For example, PNAG may be isolated from *S. aureus* strain MN8m (WO04/43407). PIA isolated from *S. epidermidis* is an integral constituent of biofilm. It is responsible for mediating cell-cell adhesion and probably also functions to shield the growing colony from the host's immune response. The polysaccharide previously known as poly-N-succinyl-O-(1 $\rightarrow$ 6)-glucosamine (PNSG) was recently shown not to have the expected structure since the identification of N-succinylation was incorrect (Maira-Litran et al., 2002). Therefore the polysaccharide formally known as PNSG and now found to be PNAG is also encompassed by the term PIA.

PIA (or PNAG) may be of different sizes varying from over 400 kDa to between 75 and 400 kDa to between 10 and 75 kDa to oligosaccharides composed of up to 30 repeat units (of  $\beta$ -(1 $\rightarrow$ 6)-linked glucosamine substituted with N-acetyl and O-succinyl constituents). Any size of PIA polysaccharide or oligosaccharide may be used in an immunogenic composition of the invention, in one aspect the polysaccharide is over 40 kDa. Sizing may be achieved by any method known in the art, for instance by microfluidization, ultrasonic irradiation or by chemical cleavage (WO 03/53462, EP497524, EP497525). In certain aspects PIA (PNAG) is at least or at most 40-400 kDa, 40-300 kDa, 50-350 kDa, 60-300 kDa, 50-250 kDa and 60-200 kDa.

PIA (PNAG) can have different degree of acetylation due to substitution on the amino groups by acetate. PIA produced in vitro is almost fully substituted on amino groups (95-100%). Alternatively, a deacetylated PIA (PNAG) can be used having less than 60%, 50%, 40%, 30%, 20%, 10% acetylation. Use of a deacetylated PIA (PNAG) is preferred

since non-acetylated epitopes of PNAG are efficient at mediating opsonic killing of Gram positive bacteria, preferably *S. aureus* and/or *S. epidermidis*. In certain aspects, the PIA (PNAG) has a size between 40 kDa and 300 kDa and is deacetylated so that less than 60%, 50%, 40%, 30% or 20% of amino groups are acetylated.

The term deacetylated PNAG (dPNAG) refers to a PNAG polysaccharide or oligosaccharide in which less than 60%, 50%, 40%, 30%, 20% or 10% of the amino groups are acetylated. In certain aspects, PNAG is deacetylated to form dPNAG by chemically treating the native polysaccharide. For example, the native PNAG is treated with a basic solution such that the pH rises to above 10. For instance the PNAG is treated with 0.1-5 M, 0.2-4 M, 0.3-3 M, 0.5-2 M, 0.75-1.5 M or 1 M NaOH, KOH or NH<sub>4</sub>OH. Treatment is for at least 10 to 30 minutes, or 1, 2, 3, 4, 5, 10, 15 or 20 hours at a temperature of 20-100, 25-80, 30-60 or 30-50 or 35-45° C. dPNAG may be prepared as described in WO 04/43405.

The polysaccharide(s) can be conjugated or unconjugated to a carrier protein.

#### B. Type 5 and Type 8 Polysaccharides from *S. aureus*

Most strains of *S. aureus* that cause infection in man contain either Type 5 or Type 8 polysaccharides. Approximately 60% of human strains are Type 8 and approximately 30% are Type 5. The structures of Type 5 and Type 8 capsular polysaccharide antigens are described in Moreau et al., (1990) and Fournier et al., (1984). Both have FucNAcp in their repeat unit as well as ManNAcA which can be used to introduce a sulfhydryl group. The structures are:

Type 5  
 $\rightarrow$ 4)- $\beta$ -D-ManNAcA(3OAc)-(1 $\rightarrow$ 4)- $\alpha$ -L-FucNAc  
 (1 $\rightarrow$ 3)- $\beta$ -D-FucNAc-(1 $\rightarrow$

Type 8  
 $\rightarrow$ 3)- $\beta$ -D-ManNAcA(4OAc)-(1 $\rightarrow$ 3)- $\alpha$ -L-FucNAc  
 (1 $\rightarrow$ 3)- $\beta$ -D-FucNAc-(1 $\rightarrow$

Recently (Jones, 2005) NMR spectroscopy revised the structures to:

Type 5  
 $\rightarrow$ 4)- $\beta$ -D-ManNAcA-(1 $\rightarrow$ 4)- $\alpha$ -L-FucNAc(3OAc)-  
 (1 $\rightarrow$ 3)- $\beta$ -D-FucNAc-(1 $\rightarrow$

Type 8  
 $\rightarrow$ 3)- $\beta$ -D-ManNAcA(4OAc)-(1 $\rightarrow$ 3)- $\alpha$ -L-FucNAc  
 (1 $\rightarrow$ 3)- $\alpha$ -D-FucNAc(1 $\rightarrow$

Polysaccharides may be extracted from the appropriate strain of *S. aureus* using method well known to of skill in the art, See U.S. Pat. No. 6,294,177. For example, ATCC 12902 is a Type 5 *S. aureus* strain and ATCC 12605 is a Type 8 *S. aureus* strain.

Polysaccharides are of native size or alternatively may be sized, for instance by microfluidisation, ultrasonic irradiation, or by chemical treatment. The invention also covers oligosaccharides derived from the type 5 and 8 polysaccharides from *S. aureus*. The type 5 and 8 polysaccharides included in the immunogenic composition of the invention are preferably conjugated to a carrier protein as described below or are alternatively unconjugated. The immunogenic compositions of the invention alternatively contains either type 5 or type 8 polysaccharide.

#### C. *S. aureus* 336 Antigen

In an embodiment, the immunogenic composition of the invention comprises the *S. aureus* 336 antigen described in U.S. Pat. No. 6,294,177. The 336 antigen comprises  $\beta$ -linked hexosamine, contains no O-acetyl groups, and specifically binds to antibodies to *S. aureus* Type 336 deposited under ATCC 55804. In an embodiment, the 336 antigen is a polysaccharide which is of native size or alternatively may be sized, for instance by microfluidisation,

ultrasonic irradiation, or by chemical treatment. The invention also covers oligosaccharides derived from the 336 antigen. The 336 antigen can be unconjugated or conjugated to a carrier protein.

#### D. Type I, II and III Polysaccharides from *S. epidermidis*

Amongst the problems associated with the use of polysaccharides in vaccination, is the fact that polysaccharides per se are poor immunogens. It is preferred that the polysaccharides utilized in the invention are linked to a protein carrier which provide bystander T-cell help to improve immunogenicity. Examples of such carriers which may be conjugated to polysaccharide immunogens include the Diphtheria and Tetanus toxoids (DT, DT CRM197 and TT respectively), Keyhole Limpet Haemocyanin (KLH), and the purified protein derivative of Tuberculin (PPD), *Pseudomonas aeruginosa* exoprotein A (rEPA), protein D from *Haemophilus influenzae*, pneumolysin or fragments of any of the above. Fragments suitable for use include fragments encompassing T-helper epitopes. In particular the protein D fragment from H. influenza will preferably contain the N-terminal 1/3 of the protein. Protein D is an IgD-binding protein from *Haemophilus influenzae* (EP 0 594 610 B1) and is a potential immunogen. In addition, staphylococcal proteins may be used as a carrier protein in the polysaccharide conjugates of the invention.

A carrier protein that would be particularly advantageous to use in the context of a staphylococcal vaccine is staphylococcal alpha toxoid. The native form may be conjugated to a polysaccharide since the process of conjugation reduces toxicity. Preferably genetically detoxified alpha toxins such as the His35Leu or His35Arg variants are used as carriers since residual toxicity is lower. Alternatively the alpha toxin is chemically detoxified by treatment with a cross-linking reagent, formaldehyde or glutaraldehyde. A genetically detoxified alpha toxin is optionally chemically detoxified, preferably by treatment with a cross-linking reagent, formaldehyde or glutaraldehyde to further reduce toxicity.

The polysaccharides may be linked to the carrier protein(s) by any known method (for example those methods described in U.S. Pat. Nos. 4,372,945, 4,474,757, and 4,356,170). Preferably, CDAP conjugation chemistry is carried out (see WO95/08348). In CDAP, the cyanating reagent 1-cyano-dimethylaminopyridinium tetrafluoroborate (CDAP) is preferably used for the synthesis of polysaccharide-protein conjugates. The cyanilation reaction can be performed under relatively mild conditions, which avoids hydrolysis of the alkaline sensitive polysaccharides. This synthesis allows direct coupling to a carrier protein.

Conjugation preferably involves producing a direct linkage between the carrier protein and polysaccharide. Optionally a spacer (such as adipic dihydride (ADH)) may be introduced between the carrier protein and the polysaccharide.

#### IV. Immune Response and Assays

As discussed above, the invention concerns evoking or inducing an immune response in a subject against a variant SpA or coagulase peptide. In one embodiment, the immune response can protect against or treat a subject having, suspected of having, or at risk of developing an infection or related disease, particularly those related to staphylococci. One use of the immunogenic compositions of the invention is to prevent nosocomial infections by inoculating a subject prior to undergoing procedures in a hospital or other environment having an increased risk of infection.

##### A. Immunoassays

The present invention includes the implementation of serological assays to evaluate whether and to what extent an

immune response is induced or evoked by compositions of the invention. There are many types of immunoassays that can be implemented. Immunoassays encompassed by the present invention include, but are not limited to, those described in U.S. Pat. No. 4,367,110 (double monoclonal antibody sandwich assay) and U.S. Pat. No. 4,452,901 (western blot). Other assays include immunoprecipitation of labeled ligands and immunocytochemistry, both in vitro and in vivo.

Immunoassays generally are binding assays. Certain preferred immunoassays are the various types of enzyme linked immunosorbent assays (ELISAs) and radioimmunoassays (MA) known in the art. Immunohistochemical detection using tissue sections is also particularly useful. In one example, antibodies or antigens are immobilized on a selected surface, such as a well in a polystyrene microtiter plate, dipstick, or column support. Then, a test composition suspected of containing the desired antigen or antibody, such as a clinical sample, is added to the wells. After binding and washing to remove non specifically bound immune complexes, the bound antigen or antibody may be detected. Detection is generally achieved by the addition of another antibody, specific for the desired antigen or antibody, that is linked to a detectable label. This type of ELISA is known as a "sandwich ELISA." Detection also may be achieved by the addition of a second antibody specific for the desired antigen, followed by the addition of a third antibody that has binding affinity for the second antibody, with the third antibody being linked to a detectable label.

Competition ELISAs are also possible implementations in which test samples compete for binding with known amounts of labeled antigens or antibodies. The amount of reactive species in the unknown sample is determined by mixing the sample with the known labeled species before or during incubation with coated wells. The presence of reactive species in the sample acts to reduce the amount of labeled species available for binding to the well and thus reduces the ultimate signal. Irrespective of the format employed, ELISAs have certain features in common, such as coating, incubating or binding, washing to remove non specifically bound species, and detecting the bound immune complexes.

Antigen or antibodies may also be linked to a solid support, such as in the form of plate, beads, dipstick, membrane, or column matrix, and the sample to be analyzed is applied to the immobilized antigen or antibody. In coating a plate with either antigen or antibody, one will generally incubate the wells of the plate with a solution of the antigen or antibody, either overnight or for a specified period. The wells of the plate will then be washed to remove incompletely-adsorbed material. Any remaining available surfaces of the wells are then "coated" with a nonspecific protein that is antigenically neutral with regard to the test antisera. These include bovine serum albumin (BSA), casein, and solutions of milk powder. The coating allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific binding of antisera onto the surface.

##### B. Diagnosis of Bacterial Infection

In addition to the use of proteins, polypeptides, and/or peptides, as well as antibodies binding these polypeptides, proteins, and/or peptides, to treat or prevent infection as described above, the present invention contemplates the use of these polypeptides, proteins, peptides, and/or antibodies in a variety of ways, including the detection of the presence of Staphylococci to diagnose an infection, whether in a patient or on medical equipment which may also become

infected. In accordance with the invention, a preferred method of detecting the presence of infections involves the steps of obtaining a sample suspected of being infected by one or more staphylococcal bacteria species or strains, such as a sample taken from an individual, for example, from one's blood, saliva, tissues, bone, muscle, cartilage, or skin. Following isolation of the sample, diagnostic assays utilizing the polypeptides, proteins, peptides, and/or antibodies of the present invention may be carried out to detect the presence of staphylococci, and such assay techniques for determining such presence in a sample are well known to those skilled in the art and include methods such as radio-immunoassay, western blot analysis and ELISA assays. In general, in accordance with the invention, a method of diagnosing an infection is contemplated wherein a sample suspected of being infected with staphylococci has added to it the polypeptide, protein, peptide, antibody, or monoclonal antibody in accordance with the present invention, and staphylococci are indicated by antibody binding to the polypeptides, proteins, and/or peptides, or polypeptides, proteins, and/or peptides binding to the antibodies in the sample.

Accordingly, antibodies in accordance with the invention may be used for the prevention of infection from staphylococcal bacteria (i.e., passive immunization), for the treatment of an ongoing infection, or for use as research tools. The term "antibodies" as used herein includes monoclonal, polyclonal, chimeric, single chain, bispecific, simianized, and humanized or primate antibodies as well as Fab fragments, such as those fragments which maintain the binding specificity of the antibodies, including the products of an Fab immunoglobulin expression library. Accordingly, the invention contemplates the use of single chains such as the variable heavy and light chains of the antibodies. Generation of any of these types of antibodies or antibody fragments is well known to those skilled in the art. Specific examples of the generation of an antibody to a bacterial protein can be found in U.S. Patent Application Pub. No. 20030153022, which is incorporated herein by reference in its entirety.

Any of the above described polypeptides, proteins, peptides, and/or antibodies may be labeled directly with a detectable label for identification and quantification of staphylococcal bacteria. Labels for use in immunoassays are generally known to those skilled in the art and include enzymes, radioisotopes, and fluorescent, luminescent and chromogenic substances, including colored particles such as colloidal gold or latex beads. Suitable immunoassays include enzyme-linked immunosorbent assays (ELISA).

### C. Protective Immunity

In some embodiments of the invention, proteinaceous compositions confer protective immunity to a subject. Protective immunity refers to a body's ability to mount a specific immune response that protects the subject from developing a particular disease or condition that involves the agent against which there is an immune response. An immunogenically effective amount is capable of conferring protective immunity to the subject.

As used herein in the specification and in the claims section that follows, the term polypeptide or peptide refer to a stretch of amino acids covalently linked there amongst via peptide bonds. Different polypeptides have different functionalities according to the present invention. While according to one aspect, a polypeptide is derived from an immunogen designed to induce an active immune response in a recipient, according to another aspect of the invention, a polypeptide is derived from an antibody which results

following the elicitation of an active immune response in, for example, an animal, and which can serve to induce a passive immune response in the recipient. In both cases, however, the polypeptide is encoded by a polynucleotide according to any possible codon usage.

As used herein the phrase "immune response" or its equivalent "immunological response" refers to the development of a humoral (antibody mediated), cellular (mediated by antigen-specific T cells or their secretion products) or both humoral and cellular response directed against a protein, peptide, carbohydrate, or polypeptide of the invention in a recipient patient. Such a response can be an active response induced by administration of immunogen or a passive response induced by administration of antibody, antibody containing material, or primed T-cells. A cellular immune response is elicited by the presentation of polypeptide epitopes in association with Class I or Class II MHC molecules, to activate antigen-specific CD4 (+) T helper cells and/or CD8 (+) cytotoxic T cells. The response may also involve activation of monocytes, macrophages, NK cells, basophils, dendritic cells, astrocytes, microglia cells, eosinophils or other components of innate immunity. As used herein "active immunity" refers to any immunity conferred upon a subject by administration of an antigen.

As used herein "passive immunity" refers to any immunity conferred upon a subject without administration of an antigen to the subject. "Passive immunity" therefore includes, but is not limited to, administration of activated immune effectors including cellular mediators or protein mediators (e.g., monoclonal and/or polyclonal antibodies) of an immune response. A monoclonal or polyclonal antibody composition may be used in passive immunization for the prevention or treatment of infection by organisms that carry the antigen recognized by the antibody. An antibody composition may include antibodies that bind to a variety of antigens that may in turn be associated with various organisms. The antibody component can be a polyclonal antiserum. In certain aspects the antibody or antibodies are affinity purified from an animal or second subject that has been challenged with an antigen(s). Alternatively, an antibody mixture may be used, which is a mixture of monoclonal and/or polyclonal antibodies to antigens present in the same, related, or different microbes or organisms, such as gram-positive bacteria, gram-negative bacteria, including but not limited to *Staphylococcus* bacteria.

Passive immunity may be imparted to a patient or subject by administering to the patient immunoglobulins (Ig) and/or other immune factors obtained from a donor or other non-patient source having a known immunoreactivity. In other aspects, an antigenic composition of the present invention can be administered to a subject who then acts as a source or donor for globulin, produced in response to challenge with the antigenic composition ("hyperimmune globulin"), that contains antibodies directed against *Staphylococcus* or other organism. A subject thus treated would donate plasma from which hyperimmune globulin would then be obtained, via conventional plasma-fractionation methodology, and administered to another subject in order to impart resistance against or to treat *Staphylococcus* infection. Hyperimmune globulins according to the invention are particularly useful for immune-compromised individuals, for individuals undergoing invasive procedures or where time does not permit the individual to produce their own antibodies in response to vaccination. See U.S. Pat. Nos. 6,936,258, 6,770,278, 6,756,361, 5,548,066, 5,512,282, 4,338,298, and

4,748,018, each of which is incorporated herein by reference in its entirety, for exemplary methods and compositions related to passive immunity.

For purposes of this specification and the accompanying claims the terms “epitope” and “antigenic determinant” are used interchangeably to refer to a site on an antigen to which B and/or T cells respond or recognize. B-cell epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, e.g., Epitope Mapping Protocols (1996). Antibodies that recognize the same epitope can be identified in a simple immunoassay showing the ability of one antibody to block the binding of another antibody to a target antigen. T-cells recognize continuous epitopes of about nine amino acids for CD8 cells or about 13-15 amino acids for CD4 cells. T cells that recognize the epitope can be identified by in vitro assays that measure antigen-dependent proliferation, as determined by <sup>3</sup>H-thymidine incorporation by primed T cells in response to an epitope (Burke et al., 1994), by antigen-dependent killing (cytotoxic T lymphocyte assay, Tigges et al., 1996) or by cytokine secretion.

The presence of a cell-mediated immunological response can be determined by proliferation assays (CD4 (+) T cells) or CTL (cytotoxic T lymphocyte) assays. The relative contributions of humoral and cellular responses to the protective or therapeutic effect of an immunogen can be distinguished by separately isolating IgG and T-cells from an immunized syngeneic animal and measuring protective or therapeutic effect in a second subject.

As used herein and in the claims, the terms “antibody” or “immunoglobulin” are used interchangeably and refer to any of several classes of structurally related proteins that function as part of the immune response of an animal or recipient, which proteins include IgG, IgD, IgE, IgA, IgM and related proteins.

Under normal physiological conditions antibodies are found in plasma and other body fluids and in the membrane of certain cells and are produced by lymphocytes of the type denoted B cells or their functional equivalent. Antibodies of the IgG class are made up of four polypeptide chains linked together by disulfide bonds. The four chains of intact IgG molecules are two identical heavy chains referred to as H-chains and two identical light chains referred to as L-chains.

In order to produce polyclonal antibodies, a host, such as a rabbit or goat, is immunized with the antigen or antigen fragment, generally with an adjuvant and, if necessary, coupled to a carrier. Antibodies to the antigen are subsequently collected from the sera of the host. The polyclonal antibody can be affinity purified against the antigen rendering it monospecific.

Monoclonal antibodies can be produced by hyperimmunization of an appropriate donor with the antigen or ex-vivo by use of primary cultures of splenic cells or cell lines derived from spleen (Anavi, 1998; Huston et al., 1991; Johnson et al., 1991; Mernaugh et al., 1995).

As used herein and in the claims, the phrase “an immunological portion of an antibody” includes a Fab fragment of an antibody, a Fv fragment of an antibody, a heavy chain of

an antibody, a light chain of an antibody, a heterodimer consisting of a heavy chain and a light chain of an antibody, a variable fragment of a light chain of an antibody, a variable fragment of a heavy chain of an antibody, and a single chain variant of an antibody, which is also known as scFv. In addition, the term includes chimeric immunoglobulins which are the expression products of fused genes derived from different species, one of the species can be a human, in which case a chimeric immunoglobulin is said to be humanized. Typically, an immunological portion of an antibody competes with the intact antibody from which it was derived for specific binding to an antigen.

Optionally, an antibody or preferably an immunological portion of an antibody, can be chemically conjugated to, or expressed as, a fusion protein with other proteins. For purposes of this specification and the accompanying claims, all such fused proteins are included in the definition of antibodies or an immunological portion of an antibody.

As used herein the terms “immunogenic agent” or “immunogen” or “antigen” are used interchangeably to describe a molecule capable of inducing an immunological response against itself on administration to a recipient, either alone, in conjunction with an adjuvant, or presented on a display vehicle.

#### D. Treatment Methods

A method of the present invention includes treatment for a disease or condition caused by a *Staphylococcus* pathogen. An immunogenic polypeptide of the invention can be given to induce an immune response in a person infected with *Staphylococcus* or suspected of having been exposed to *Staphylococcus*. Methods may be employed with respect to individuals who have tested positive for exposure to *Staphylococcus* or who are deemed to be at risk for infection based on possible exposure.

In particular, the invention encompasses a method of treatment for staphylococcal infection, particularly hospital acquired nosocomial infections. The immunogenic compositions and vaccines of the invention are particularly advantageous to use in cases of elective surgery. Such patients will know the date of surgery in advance and could be inoculated in advance. The immunogenic compositions and vaccines of the invention are also advantageous to use to inoculate health care workers.

In some embodiments, the treatment is administered in the presence of adjuvants or carriers or other staphylococcal antigens. Furthermore, in some examples, treatment comprises administration of other agents commonly used against bacterial infection, such as one or more antibiotics.

The use of peptides for vaccination can require, but not necessarily, conjugation of the peptide to an immunogenic carrier protein, such as hepatitis B surface antigen, keyhole limpet hemocyanin, or bovine serum albumin. Methods for performing this conjugation are well known in the art.

#### V. Vaccine and Other Pharmaceutical Compositions and Administration

##### A. Vaccines

The present invention includes methods for preventing or ameliorating staphylococcal infections, particularly hospital acquired nosocomial infections. As such, the invention contemplates vaccines for use in both active and passive immunization embodiments. Immunogenic compositions, proposed to be suitable for use as a vaccine, may be prepared from immunogenic SpA polypeptide(s), such as a SpA domain D variant, or immunogenic coagulases. In other embodiments SpA or coagulases can be used in combination with other secreted virulence proteins, surface proteins or immunogenic fragments thereof. In certain aspects, anti-

genic material is extensively dialyzed to remove undesired small molecular weight molecules and/or lyophilized for more ready formulation into a desired vehicle.

Other options for a protein/peptide-based vaccine involve introducing nucleic acids encoding the antigen(s) as DNA vaccines. In this regard, recent reports described construction of recombinant vaccinia viruses expressing either 10 contiguous minimal CTL epitopes (Thomson, 1996) or a combination of B cell, cytotoxic T-lymphocyte (CTL), and T-helper (Th) epitopes from several microbes (An, 1997), and successful use of such constructs to immunize mice for priming protective immune responses. Thus, there is ample evidence in the literature for successful utilization of peptides, peptide-pulsed antigen presenting cells (APCs), and peptide-encoding constructs for efficient *in vivo* priming of protective immune responses. The use of nucleic acid sequences as vaccines is exemplified in U.S. Pat. Nos. 5,958,895 and 5,620,896.

The preparation of vaccines that contain polypeptide or peptide sequence(s) as active ingredients is generally well understood in the art, as exemplified by U.S. Pat. Nos. 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all of which are incorporated herein by reference. Typically, such vaccines are prepared as injectables either as liquid solutions or suspensions: solid forms suitable for solution in or suspension in liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients that are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants that enhance the effectiveness of the vaccines. In specific embodiments, vaccines are formulated with a combination of substances, as described in U.S. Pat. Nos. 6,793,923 and 6,733,754, which are incorporated herein by reference.

Vaccines may be conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkalene glycols or triglycerides: such suppositories may be formed from mixtures containing the active ingredient in the range of about 0.5% to about 10%, preferably about 1% to about 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 10% to about 95% of active ingredient, preferably about 25% to about 70%.

The polypeptides and polypeptide-encoding DNA constructs may be formulated into a vaccine as neutral or salt forms. Pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the peptide) and those that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like.

Typically, vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity

to be administered depends on the subject to be treated, including the capacity of the individual's immune system to synthesize antibodies and the degree of protection desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are of the order of several hundred micrograms of active ingredient per vaccination. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed by subsequent inoculations or other administrations.

The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These are believed to include oral application within a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection and the like. The dosage of the vaccine will depend on the route of administration and will vary according to the size and health of the subject.

In certain instances, it will be desirable to have multiple administrations of the vaccine, e.g., 2, 3, 4, 5, 6 or more administrations. The vaccinations can be at 1, 2, 3, 4, 5, 6, 7, 8, to 5, 6, 7, 8, 9, 10, 11, 12 twelve week intervals, including all ranges there between. Periodic boosters at intervals of 1-5 years will be desirable to maintain protective levels of the antibodies. The course of the immunization may be followed by assays for antibodies against the antigens, as described in U.S. Pat. Nos. 3,791,932; 4,174,384 and 3,949,064.

#### 1. Carriers

A given composition may vary in its immunogenicity. It is often necessary therefore to boost the host immune system, as may be achieved by coupling a peptide or polypeptide to a carrier. Exemplary and preferred carriers are keyhole limpet hemocyanin (KLH) and bovine serum albumin (BSA). Other albumins such as ovalbumin, mouse serum albumin, or rabbit serum albumin can also be used as carriers. Means for conjugating a polypeptide to a carrier protein are well known in the art and include glutaraldehyde, m-maleimidobencoyl-N-hydroxysuccinimide ester, carbodiimide, and bis-biazotized benzidine.

#### 2. Adjuvants

The immunogenicity of polypeptide or peptide compositions can be enhanced by the use of non-specific stimulators of the immune response, known as adjuvants. Suitable adjuvants include all acceptable immunostimulatory compounds, such as cytokines, toxins, or synthetic compositions. A number of adjuvants can be used to enhance an antibody response against a variant SpA polypeptide or coagulase, or any other bacterial protein or combination contemplated herein. Adjuvants can (1) trap the antigen in the body to cause a slow release; (2) attract cells involved in the immune response to the site of administration; (3) induce proliferation or activation of immune system cells; or (4) improve the spread of the antigen throughout the subject's body.

Adjuvants include, but are not limited to, oil-in-water emulsions, water-in-oil emulsions, mineral salts, polynucleotides, and natural substances. Specific adjuvants that may be used include IL-1, IL-2, IL-4, IL-7, IL-12,  $\gamma$ -interferon, GMCSF, BCG, aluminum salts, such as aluminum hydroxide or other aluminum compound, MDP compounds, such as thur-MDP and nor-MDP, CGP (MTP-PE), lipid A, and monophosphoryl lipid A (MPL). RIBI, which contains three components extracted from bacteria, MPL, trehalose dimycolate (TDM), and cell wall skeleton (CWS) in a 2% squalene/Tween 80 emulsion. MHC antigens may even be used. Others adjuvants or methods are exemplified in U.S.

Pat. Nos. 6,814,971, 5,084,269, 6,656,462, each of which is incorporated herein by reference).

Various methods of achieving adjuvant affect for the vaccine includes use of agents such as aluminum hydroxide or phosphate (alum), commonly used as about 0.05 to about 0.1% solution in phosphate buffered saline, admixture with synthetic polymers of sugars (Carbopol®) used as an about 0.25% solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between about 70° to about 101° C. for a 30-second to 2-minute period, respectively. Aggregation by reactivating with pepsin-treated (Fab) antibodies to albumin; mixture with bacterial cells (e.g., *C. parvum*), endotoxins or lipopolysaccharide components of Gram-negative bacteria; emulsion in physiologically acceptable oil vehicles (e.g., mannide monooleate (Aracel A)); or emulsion with a 20% solution of a perfluorocarbon (Fluosol-DA®) used as a block substitute may also be employed to produce an adjuvant effect.

Examples of and often preferred adjuvants include complete Freund's adjuvant (a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*), incomplete Freund's adjuvants, and aluminum hydroxide.

In some aspects, it is preferred that the adjuvant be selected to be a preferential inducer of either a Th1 or a Th2 type of response. High levels of Th1-type cytokines tend to favor the induction of cell mediated immune responses to a given antigen, while high levels of Th2-type cytokines tend to favor the induction of humoral immune responses to the antigen.

The distinction of Th1 and Th2-type immune response is not absolute. In reality an individual will support an immune response which is described as being predominantly Th1 or predominantly Th2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4+ T cell clones by Mosmann and Coffman (Mosmann, and Coffman, 1989). Traditionally, Th1-type responses are associated with the production of the INF- $\gamma$  and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of Th1-type immune responses are not produced by T-cells, such as IL-12. In contrast, Th2-type responses are associated with the secretion of IL-4, IL-5, IL-6, IL-10.

In addition to adjuvants, it may be desirable to co-administer biologic response modifiers (BRM) to enhance immune responses. BRMs have been shown to upregulate T cell immunity or downregulate suppresser cell activity. Such BRMs include, but are not limited to, Cimetidine (CIM; 1200 mg/d) (Smith/Kline, PA); or low-dose Cyclophosphamide (CYP; 300 mg/m<sup>2</sup>) (Johnson/Mead, NJ) and cytokines such as  $\gamma$ -interferon, IL-2, or IL-12 or genes encoding proteins involved in immune helper functions, such as B-7.

#### B. Lipid Components and Moieties

In certain embodiments, the present invention concerns compositions comprising one or more lipids associated with a nucleic acid or a polypeptide/peptide. A lipid is a substance that is insoluble in water and extractable with an organic solvent. Compounds other than those specifically described herein are understood by one of skill in the art as lipids, and are encompassed by the compositions and methods of the present invention. A lipid component and a non-lipid may be attached to one another, either covalently or non-covalently.

A lipid may be a naturally occurring lipid or a synthetic lipid. However, a lipid is usually a biological substance. Biological lipids are well known in the art, and include for example, neutral fats, phospholipids, phosphoglycerides, steroids, terpenes, lysolipids, glycosphingolipids, glucolip-

ids, sulphatides, lipids with ether and ester-linked fatty acids and polymerizable lipids, and combinations thereof.

A nucleic acid molecule or a polypeptide/peptide, associated with a lipid may be dispersed in a solution containing a lipid, dissolved with a lipid, emulsified with a lipid, mixed with a lipid, combined with a lipid, covalently bonded to a lipid, contained as a suspension in a lipid or otherwise associated with a lipid. A lipid or lipid-poxvirus-associated composition of the present invention is not limited to any particular structure. For example, they may also simply be interspersed in a solution, possibly forming aggregates which are not uniform in either size or shape. In another example, they may be present in a bilayer structure, as micelles, or with a "collapsed" structure. In another non-limiting example, a lipofectamine(Gibco BRL)-poxvirus or Superfect (Qiagen)-poxvirus complex is also contemplated.

In certain embodiments, a composition may comprise about 1%, about 2%, about 3%, about 4%, about 5%, about 6%, about 7%, about 8%, about 9%, about 10%, about 11%, about 12%, about 13%, about 14%, about 15%, about 16%, about 17%, about 18%, about 19%, about 20%, about 21%, about 22%, about 23%, about 24%, about 25%, about 26%, about 27%, about 28%, about 29%, about 30%, about 31%, about 32%, about 33%, about 34%, about 35%, about 36%, about 37%, about 38%, about 39%, about 40%, about 41%, about 42%, about 43%, about 44%, about 45%, about 46%, about 47%, about 48%, about 49%, about 50%, about 51%, about 52%, about 53%, about 54%, about 55%, about 56%, about 57%, about 58%, about 59%, about 60%, about 61%, about 62%, about 63%, about 64%, about 65%, about 66%, about 67%, about 68%, about 69%, about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or any range therebetween, of a particular lipid, lipid type, or non-lipid component such as an adjuvant, antigen, peptide, polypeptide, sugar, nucleic acid or other material disclosed herein or as would be known to one of skill in the art. In a non-limiting example, a composition may comprise about 10% to about 20% neutral lipids, and about 33% to about 34% of a cerebroside, and about 1% cholesterol. In another non-limiting example, a liposome may comprise about 4% to about 12% terpenes, wherein about 1% of the micelle is specifically lycopene, leaving about 3% to about 11% of the liposome as comprising other terpenes; and about 10% to about 35% phosphatidyl choline, and about 1% of a non-lipid component. Thus, it is contemplated that compositions of the present invention may comprise any of the lipids, lipid types or other components in any combination or percentage range.

#### C. Combination Therapy

The compositions and related methods of the present invention, particularly administration of a secreted virulence factor or surface protein, including a variant SpA polypeptide or peptide, and/or other bacterial peptides or proteins to a patient/subject, may also be used in combination with the administration of traditional therapies. These include, but are not limited to, the administration of antibiotics such as streptomycin, ciprofloxacin, doxycycline, gentamycin, chloramphenicol, trimethoprim, sulfamethoxazole, ampicillin, tetracycline or various combinations of antibiotics.

In one aspect, it is contemplated that a polypeptide vaccine and/or therapy is used in conjunction with antibacterial treatment. Alternatively, the therapy may precede or

follow the other agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agents and/or a proteins or polynucleotides are administered separately, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and antigenic composition would still be able to exert an advantageously combined effect on the subject. In such instances, it is contemplated that one may administer both modalities within about 12-24 h of each other or within about 6-12 h of each other. In some situations, it may be desirable to extend the time period for administration significantly, where several days (2, 3, 4, 5, 6 or 7) to several weeks (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

Various combinations may be employed, for example antibiotic therapy is "A" and the immunogenic molecule given as part of an immune therapy regime, such as an antigen, is "B":

---

A/B/A B/A/B B/B/A A/A/B A/B/B B/A/A A/B/B B/A/B/B  
 B/B/B/A B/B/A/B A/A/B/B A/B/A/B A/B/B/A B/B/A/A  
 B/A/B/A B/A/A/B A/A/A/B B/A/A/A A/B/A/A A/A/B/A

---

Administration of the immunogenic compositions of the present invention to a patient/subject will follow general protocols for the administration of such compounds, taking into account the toxicity, if any, of the SpA composition, or other compositions described herein. It is expected that the treatment cycles would be repeated as necessary. It also is contemplated that various standard therapies, such as hydration, may be applied in combination with the described therapy.

D. General Pharmaceutical Compositions

In some embodiments, pharmaceutical compositions are administered to a subject. Different aspects of the present invention involve administering an effective amount of a composition to a subject. In some embodiments of the present invention, staphylococcal antigens, members of the Ess pathway, including polypeptides or peptides of the Esa or Esx class, and/or members of sortase substrates may be administered to the patient to protect against infection by one or more *Staphylococcus* pathogens. Alternatively, an expression vector encoding one or more such polypeptides or peptides may be given to a patient as a preventative treatment. Additionally, such compounds can be administered in combination with an antibiotic or an antibacterial. Such compositions will generally be dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium.

In addition to the compounds formulated for parenteral administration, such as those for intravenous or intramuscular injection, other pharmaceutically acceptable forms include, e.g., tablets or other solids for oral administration; time release capsules; and any other form currently used, including creams, lotions, mouthwashes, inhalants and the like.

The active compounds of the present invention can be formulated for parenteral administration, e.g., formulated for injection via the intravenous, intramuscular, sub-cutaneous, or even intraperitoneal routes. The preparation of an aqueous composition that contains a compound or compounds that increase the expression of an WIC class I molecule will be known to those of skill in the art in light of the present disclosure. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for use to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and, the preparations can also be emulsified.

Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil, or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that it may be easily injected. It also should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

The proteinaceous compositions may be formulated into a neutral or salt form. Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

The carrier also can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion, and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques, which yield a powder of the active ingredient, plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Administration of the compositions according to the present invention will typically be via any common route. This includes, but is not limited to oral, nasal, or buccal administration. Alternatively, administration may be by orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal, intranasal, or intravenous injection. In certain embodiments, a vaccine composition may be inhaled (e.g., U.S. Pat. No. 6,651,655, which is specifically incorporated by reference). Such compositions would normally be administered as pharmaceutically acceptable compositions that include physiologically acceptable carriers, buffers or other excipients. As

used herein, the term “pharmaceutically acceptable” refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem complications commensurate with a reasonable benefit/risk ratio. The term “pharmaceutically acceptable carrier,” means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting a chemical agent.

For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous, and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in isotonic NaCl solution and either added to hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, Remington’s Pharmaceutical Sciences, 1990). Some variation in dosage will necessarily occur depending on the condition of the subject. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

An effective amount of therapeutic or prophylactic composition is determined based on the intended goal. The term “unit dose” or “dosage” refers to physically discrete units suitable for use in a subject, each unit containing a predetermined quantity of the composition calculated to produce the desired responses discussed above in association with its administration, i.e., the appropriate route and regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the protection desired.

Precise amounts of the composition also depend on the judgment of the practitioner and are peculiar to each individual. Factors affecting dose include physical and clinical state of the subject, route of administration, intended goal of treatment (alleviation of symptoms versus cure), and potency, stability, and toxicity of the particular composition.

Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically or prophylactically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above.

#### E. In Vitro, Ex Vivo, or In Vivo Administration

As used herein, the term in vitro administration refers to manipulations performed on cells removed from or outside of a subject, including, but not limited to cells in culture. The term ex vivo administration refers to cells which have been manipulated in vitro, and are subsequently administered to a subject. The term in vivo administration includes all manipulations performed within a subject.

In certain aspects of the present invention, the compositions may be administered either in vitro, ex vivo, or in vivo. In certain in vitro embodiments, autologous B-lymphocyte cell lines are incubated with a virus vector of the instant invention for 24 to 48 hours or with a variant SpA and/or coagulase and/or any other composition described herein for two hours. The transduced cells can then be used for in vitro analysis, or alternatively for ex vivo administration. U.S. Pat. Nos. 4,690,915 and 5,199,942, both incorporated herein by reference, disclose methods for ex vivo manipulation of blood mononuclear cells and bone marrow cells for use in therapeutic applications.

#### F. Antibodies and Passive Immunization

Another aspect of the invention is a method of preparing an immunoglobulin for use in prevention or treatment of staphylococcal infection comprising the steps of immunizing a recipient or donor with the vaccine of the invention and isolating immunoglobulin from the recipient or donor. An immunoglobulin prepared by this method is a further aspect of the invention. A pharmaceutical composition comprising the immunoglobulin of the invention and a pharmaceutically acceptable carrier is a further aspect of the invention which could be used in the manufacture of a medicament for the treatment or prevention of staphylococcal disease. A method for treatment or prevention of staphylococcal infection comprising a step of administering to a patient an effective amount of the pharmaceutical preparation of the invention is a further aspect of the invention.

Inocula for polyclonal antibody production are typically prepared by dispersing the antigenic composition in a physiologically tolerable diluent such as saline or other adjuvants suitable for human use to form an aqueous composition. An immunostimulatory amount of inoculum is administered to a mammal and the inoculated mammal is then maintained for a time sufficient for the antigenic composition to induce protective antibodies.

The antibodies can be isolated to the extent desired by well known techniques such as affinity chromatography (Harlow and Lane, 1988). Antibodies can include antiserum preparations from a variety of commonly used animals, e.g. goats, primates, donkeys, swine, horses, guinea pigs, rats or man.

An immunoglobulin produced in accordance with the present invention can include whole antibodies, antibody fragments or subfragments. Antibodies can be whole immunoglobulins of any class (e.g., IgG, IgM, IgA, IgD or IgE), chimeric antibodies or hybrid antibodies with dual specificity to two or more antigens of the invention. They may also be fragments (e.g., F(ab')<sub>2</sub>, Fab', Fab, Fv and the like) including hybrid fragments. An immunoglobulin also includes natural, synthetic, or genetically engineered proteins that act like an antibody by binding to specific antigens to form a complex.

A vaccine of the present invention can be administered to a recipient who then acts as a source of immunoglobulin, produced in response to challenge from the specific vaccine. A subject thus treated would donate plasma from which hyperimmune globulin would be obtained via conventional plasma fractionation methodology. The hyperimmune globulin would be administered to another subject in order to impart resistance against or treat staphylococcal infection. Hyperimmune globulins of the invention are particularly useful for treatment or prevention of staphylococcal disease in infants, immune compromised individuals, or where treatment is required and there is no time for the individual to produce antibodies in response to vaccination.

An additional aspect of the invention is a pharmaceutical composition comprising two or more monoclonal antibodies (or fragments thereof; preferably human or humanised) reactive against at least two constituents of the immunogenic composition of the invention, which could be used to treat or prevent infection by Gram positive bacteria, preferably staphylococci, more preferably *S. aureus* or *S. epidermidis*. Such pharmaceutical compositions comprise monoclonal antibodies that can be whole immunoglobulins of any class, chimeric antibodies, or hybrid antibodies with specificity to two or more antigens of the invention. They may also be fragments (e.g., F(ab')<sub>2</sub>, Fab', Fab, Fv and the like) including hybrid fragments.

Methods of making monoclonal antibodies are well known in the art and can include the fusion of splenocytes with myeloma cells (Kohler and Milstein, 1975; Harlow and

Lane, 1988). Alternatively, monoclonal Fv fragments can be obtained by screening a suitable phage display library (Vaughan et al., 1998). Monoclonal antibodies may be humanized or part humanized by known methods.

## VI. EXAMPLES

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion. One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the methods described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses which are encompassed within the spirit of the invention as defined by the scope of the claims will occur to those skilled in the art.

### Example 1

#### Non-Toxicogenic Protein a Variants as Subunit Vaccines to Prevent *Staphylococcus Aureus* Infections

##### A. Results

An animal model for *S. aureus* infection BALB/c mice were infected by intravenous injection with  $1 \times 10^7$  CFU of the human clinical isolate *S. aureus* Newman (Baba et al., 2007). Within 6 hours following infection, 99.999% of staphylococci disappeared from the blood stream and were distributed via the vasculature. Staphylococcal dissemination to peripheral tissues occurred rapidly, as the bacterial load in kidney and other peripheral organ tissues reached  $1 \times 10^5$  CFU  $g^{-1}$  within the first three hours. The staphylococcal load in kidney tissues increased by 1.5 log CFU within twenty-four hours. Forty-eight hours following infection, mice developed disseminated abscesses in multiple organs, detectable by light microscopy of hematoxylin-eosin stained, thin-sectioned kidney tissue. The initial abscess diameter was 524  $\mu$ M ( $\pm 65$   $\mu$ M); lesions were initially marked by an influx of polymorphonuclear leukocytes (PMNs) and harbored no discernable organization of staphylococci, most of which appeared to reside within PMNs. On day 5 of infection, abscesses increased in size and enclosed a central population of staphylococci, surrounded by a layer of eosinophilic, amorphous material and a large cuff of

PMNs. Histopathology revealed massive necrosis of PMNs in proximity to the staphylococcal nidus at the center of abscess lesions as well as a mantle of healthy phagocytes. A rim of necrotic PMNs were observed at the periphery of abscess lesions, bordering eosinophilic, amorphous material that separates healthy renal tissue from lesions. Abscesses eventually reached a diameter of  $\geq 1,524$   $\mu$ M on day 15 or 36. At later time intervals, the staphylococcal load was increased to  $10^4$ - $10^6$  CFU  $g^{-1}$  and growing abscess lesions migrated towards the organ capsule. Peripheral lesions were prone to rupture, thereby releasing necrotic material and staphylococci into the peritoneal cavity or the retroperitoneal space. These events resulted in bacteremia as well as a secondary wave of abscesses, eventually precipitating a lethal outcome.

To enumerate staphylococcal load in renal tissue, animals were killed, their kidneys excised and tissue homogenate spread on agar media for colony formation. On day 5 of infection, a mean of  $1 \times 10^6$  CFU  $g^{-1}$  renal tissue for *S. aureus* Newman was observed. To quantify abscess formation, kidneys were visually inspected, and each individual organ was given a score of one or zero. The final sum was divided by the total number of kidneys to calculate percent surface abscesses (Table 4). In addition, randomly chosen kidneys were fixed in formalin, embedded, thin sectioned, and stained with hematoxylin-eosin. For each kidney, four sagittal sections at 200  $\mu$ M intervals were viewed by microscopy. The numbers of lesions were counted for each section and averaged to quantify the number of abscesses within the kidneys. *S. aureus* Newman caused  $4.364 \pm 0.889$  abscesses per kidney, and surface abscesses were observed on 14 out of 20 kidneys (70%) (Table 4).

When examined by scanning electron microscopy, *S. aureus* Newman was located in tightly associated lawns at the center of abscesses. Staphylococci were contained by an amorphous pseudocapsule that separated bacteria from the cuff of abscesses leukocytes. No immune cells were observed in these central nests of staphylococci, however occasional red blood cells were located among the bacteria. Bacterial populations at the abscess center, designated staphylococcal abscess communities (SAC), appeared homogenous and coated by an electron-dense, granular material. The kinetics of the appearance of infectious lesions and the morphological attributes of abscesses formed by *S. aureus* Newman were similar to those observed following mouse infection with *S. aureus* USA300 (LAC), the current epidemic community-acquired methicillin-resistant *S. aureus* (CA-MRSA) clone in the United States (Diep et al., 2006).

TABLE 4

Genotype	Genetic requirements for <i>S. aureus</i> Newman abscess formation in mice					
	Staphylococcal load in kidney tissue			Abscess formation in kidney tissue		
	<sup>a</sup> log <sub>10</sub> CFU $g^{-1}$ tissue	<sup>b</sup> Significance (P-value)	<sup>c</sup> Reduction (log <sub>10</sub> CFU $g^{-1}$ )	<sup>d</sup> Surface abscesses (%)	<sup>e</sup> Number of abscesses per kidney	<sup>f</sup> Significance (P-value)
wild-type	6.141 $\pm$ 0.192	—	—	70	4.364 $\pm$ 0.889	—
AsrtA	4.095 $\pm$ 0.347	6.7 $\times$ 10 <sup>-6</sup>	2.046	0	0.000 $\pm$ 0.000	0.0216
spa	5.137 $\pm$ 0.374	0.0144	1.004	13	0.375 $\pm$ 0.374	0.0356

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU  $g^{-1}$  in homogenized renal tissues 5 days following infection in cohorts of fifteen BALB/c mice per challenge strain. Standard error of the means ( $\pm$ SEM) is indicated.

<sup>b</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU  $g^{-1}$ .

<sup>d</sup>Abscess formation in kidney tissues five days following infection was measured by macroscopic inspection (% positive)

<sup>e</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys from eight to ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean ( $\pm$ SEM).

<sup>f</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

*S. aureus* Protein A (spa) mutants are avirulent and cannot form abscesses. Sortase A is a transpeptidase that immobilizes nineteen surface proteins in the envelope of *S. aureus* strain Newman (Mazmanian et al., 1999; Mazmanian et al., 2000). Earlier work identified sortase A as a virulence factor in multiple animal model systems, however the contributions of this enzyme and its anchored surface proteins to abscess formation or persistence have not yet been revealed (Jonsson et al., 2002; Weiss et al., 2004). Compared to the wild-type parent (Baba et al., 2007), an isogenic *srtA* variant (*AsrtA*) failed to form abscess lesions on either macroscopic or histopathology examination on days 2, 5, or 15. In mice infected with the *srtA* mutant, only  $1 \times 10^4$  CFU  $g^{-1}$  was recovered from kidney tissue on day 5 of infection, which is a  $2.046 \log_{10}$  CFU  $g^{-1}$  reduction compared to the wild-type parent strain ( $P=6.73 \times 10^{-6}$ ). A similar defect was observed for the *srtA* mutant of MRSA strain USA300 (data not shown). Scanning electron microscopy showed that *srtA* mutants were highly dispersed and often associated with leukocytes in otherwise healthy renal tissue. On day fifteen following infection, *srtA* mutants were cleared from renal tissues, a  $\geq 3.5 \log_{10}$  CFU  $g^{-1}$  reduction compared to the wild-type (Table 3). Thus, sortase A anchored surface proteins enable the formation of abscess lesions and the persistence of bacteria in host tissues, wherein staphylococci replicate as communities embedded in an extracellular matrix and shielded from surrounding leukocytes by an amorphous pseudocapsule.

Sortase A anchors a large spectrum of proteins with LPXTG motif sorting signals to the cell wall envelope, thereby providing for the surface display of many virulence factors (Mazmanian et al., 2002). To identify surface proteins required for staphylococcal abscess formation, bursa aurealis insertions were introduced in 5' coding sequences of genes that encode polypeptides with LPXTG motif proteins (Bae et al., 2004) and these mutations were transduced into *S. aureus* Newman. Mutations in the structural gene for Protein A (*spa*) reduced the staphylococcal load in infected mouse kidney tissues by  $1.004 \log_{10}$  ( $P=0.0144$ ). When analyzed for their ability to form abscesses in kidney tissues by histopathology, we observed that the *spa* mutants were unable to form abscesses as compared with the wild-type parent strain *S. aureus* Newman (wild-type *S. aureus* Newman  $4.364 \pm 0.889$  abscesses per kidney vs. the isogenic *spa* mutant with  $0.375 \pm 0.374$  lesions;  $P=0.0356$ ).

Protein A blocks innate and adaptive immune responses. Studies identified Protein A as a critical virulence factor during the pathogenesis of *S. aureus* infections. Earlier work demonstrated that Protein A impedes phagocytosis of staphylococci by binding the Fc component of immunoglobulin (Jensen 1958; Uhlén et al., 1984), activates platelet aggregation via the von Willebrand factor (Hartleib et al., 2000), functions as a B cell superantigen by capturing the F(ab)<sub>2</sub> region of VH3 bearing IgM (Roben et al., 1995), and, through its activation of TNFR1, can initiate staphylococcal pneumonia (Gomez et al., 2004). Due to the fact that Protein A captures immunoglobulin and displays toxic attributes, the possibility that this surface molecule may function as a vaccine in humans has not been rigorously pursued. The inventors demonstrate for the first time that Protein A variants no longer able to bind to immunoglobulins, vWF and TNFR-1 are removed of their toxigenic potential and are able to stimulate humoral immune responses that protect against staphylococcal disease.

Molecular basis of Protein A surface display and function. Protein A is synthesized as a precursor in the bacterial cytoplasm and secreted via its YSIRK signal peptide at the

cross wall, i.e., the cell division septum of staphylococci (FIG. 1). (DeDent et al., 2007; DeDent et al., 2008). Following cleavage of the C-terminal LPXTG sorting signal, Protein A is anchored to bacterial peptidoglycan cross-bridges by sortase A (Schneewind et al., 1995; Mazmanian et al., 1999; Mazmanian et al., 2000). Protein A is the most abundant surface protein of staphylococci; the molecule is expressed by virtually all *S. aureus* strains (Said-Salim et al., 2003; Cespedes et al., 2005; Kennedy et al., 2008). Staphylococci turn over 15-20% of their cell wall per division cycle (Navarre and Schneewind 1999). Murine hydrolases cleave the glycan strands and wall peptides of peptidoglycan, thereby releasing Protein A with its attached C-terminal cell wall disaccharide tetrapeptide into the extracellular medium (Ton-That et al., 1999). Thus, by physiological design, Protein A is both anchored to the cell wall and displayed on the bacterial surface but also released into surrounding tissues during host infection (Maraffini et al., 2006).

Protein A captures immunoglobulins on the bacterial surface and this biochemical activity enables staphylococcal escape from host innate and acquired immune responses (Jensen 1958; Goodyear and Silverman 2004). Interestingly, region X of Protein A (Guss et al., 1984), a repeat domain that tethers the IgG binding domains to the LPXTG sorting signal/cell wall anchor, is perhaps the most variable portion of the staphylococcal genome (Schneewind et al., 1992; Said-Salim et al., 2003). Each of the five immunoglobulin binding domains of Protein A (SpA), formed from three helix bundles and designated E, D, A, B, and C, exerts similar structural and functional properties (Sjödahl 1977; Jansson et al., 1998). The solution and crystal structure of domain D has been solved both with and without the Fc and V<sub>H</sub>3 (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille et al., 2000).

In the crystal structure complex, the Fab interacts with helix II and helix III of domain D via a surface composed of four VH region  $\beta$ -strands (Graille et al., 2000). The major axis of helix II of domain D is approximately 50° to the orientation of the strands, and the interhelical portion of domain D is most proximal to the CO strand. The site of interaction on Fab is remote from the Ig light chain and the heavy chain constant region. The interaction involves the following domain D residues: Asp-36 of helix II as well as Asp-37 and Gln-40 in the loop between helix II and helix III, in addition to several other residues with SpA-D (Graille et al., 2000). Both interacting surfaces are composed predominantly of polar side chains, with three negatively charged residues on domain D and two positively charged residues on the 2A2 Fab buried by the interaction, providing an overall electrostatic attraction between the two molecules. Of the five polar interactions identified between Fab and domain D, three are between side chains. A salt bridge is formed between Arg-H19 and Asp-36 and two hydrogen bonds are made between Tyr-H59 and Asp-37 and between Asn-H82a and Ser-33. Because of the conservation of Asp-36 and Asp-37 in all five IgG binding domains of Protein A, these residues were selected for mutagenesis.

The SpA-D sites responsible for Fab binding are structurally separate from the domain surface that mediates Fc $\gamma$  binding. The interaction of Fc $\gamma$  with domain B primarily involves residues in helix I with lesser involvement of helix II (Deisenhofer 1981; Gouda et al., 1992). With the exception of the Gln-32, a minor contact in both complexes, none of the residues that mediate the Fc $\gamma$  interaction are involved in Fab binding. To examine the spatial relationship between these different Ig-binding sites, the SpA domains in these complexes have been superimposed to construct a model of

a complex between Fab, the SpA-domain D, and the Fc $\gamma$  molecule. In this ternary model, Fab and Fc $\gamma$  form a sandwich about opposite faces of the helix II without evidence of steric hindrance of either interaction. These findings illustrate how, despite its small size (i.e., 56-61 aa), a SpA domain can simultaneously display both activities, explaining experimental evidence that the interactions of Fab with an individual domain are noncompetitive. Residues for the interaction between SpA-D and Fc $\gamma$  are Gln-9 and Gln-10.

In contrast, occupancy of the Fc portion of IgG on the domain D blocks its interaction with vWF A1 and probably also TNFR1 (O'Seaghda et al., 2006). Mutations in residues essential for IgG Fc binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF A1 and TNFR1 binding (Cedergren et al., 1993; Gomez et al., 2006; O'Seaghda et al. 2006), whereas residues critical for the V<sub>H</sub>3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) have no impact on the binding activities of IgG Fc, vWF A1 or TNFR1 (Jansson et al., 1998; Graille et al., 2000). The Protein A immunoglobulin Fab binding activity targets a subset of B cells that express VH3 family related IgM on their surface, i.e. these molecules function as VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells rapidly proliferate and then commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e. marginal zone B cells and follicular B2 cells) (Goodyear and Silverman 2003; Goodyear and Silverman 2004). It is important to note that more than 40% of circulating B cells are targeted by the Protein A interaction and the VH3 family represents the largest family of human B cell receptors to impart protective humoral responses against pathogens (Goodyear and Silverman 2003; Goodyear and Silverman 2004). Thus, Protein A functions analogously to staphylococcal superantigens (Roben et al., 1995), albeit that the latter class of molecules, for example SEB, TSST-1, TSST-2, form complexes with the T cell receptor to inappropriately stimulate host immune responses and thereby precipitating characteristic disease features of staphylococcal infections (Roben et al., 1995; Tiedemann et al., 1995). Together these findings document the contributions of Protein A in establishing staphylococcal infections and in modulating host immune responses.

Non-toxicogenic variant of Protein A. The inventors have developed a non-toxicogenic variant of staphylococcal Protein A and, with this reagent in hand, aimed for the first time to measure the immune response of animals to Protein A immunization. Further, the inventors address whether immunization of animals with a non-toxicogenic variant of Protein A could generate immune responses that raise protective immunity against staphylococcal infection.

To perturb the IgG Fc, vWF A1 and TNFR1 binding activities of Protein A, glutamine (Q) residues 9 and 10 [the numbering here is derived from that established for the SpA domain D] were modified generating lysine or glycine substitutions for both glutamines with the expectation that these substitutions abolish the ion bonds formed between wild-type Protein A and its ligands. The added effect of the dual lysine substitutions may be that these positively charged residues institute a repellent charge for immunoglobulins. To perturb IgM Fab VH3 binding, the inventors selected the aspartate (D) residues 36 and 37 of SpA-D, each of which is required for the association of Protein A with the B cell receptor. D36 and D37 were both substituted with alanine. The Q9,10K and D36,37A mutations were combined in the recombinant molecule SpA-D<sub>Q9,10K;D36,37A</sub> and examined for the binding attributes of Protein A.

In brief, the Protein A (spa) genomic sequence of *Staphylococcus aureus* N315 was PCR amplified with the primers (GCTGCACATATGGCGCAACACGATGAAGCTCAAC [5' primer](SEQ ID NO:35) and AGTGGATCCT-TATGCTTTGTTAGCATCTGC [3' primer] (SEQ ID NO:36)), cloned into the pET15b vector (pYSJ1, codons 48-486) (Stranger-Jones, et al., 2006) and recombinant plasmid transformed into *E. coli* BL21(DE3) (Studier et al., 1990). The Protein A product derived from pYSJ1 harbors SpA residues 36-265 fused to the N-terminal His tag (MGSSHHHHHHSSGLVPRGS (SEQ ID NO:37)). Following IPTG inducible expression, recombinant N-terminal His6-tagged SpA was purified by affinity chromatography on Ni-NTA resin (Stranger-Jones et al., 2006). The domain D of SpA (SpA-D) was PCR amplified with a pair of specific primers (AACATATGTTCAACAAAGATCAACAAAGC [5' primer](SEQ ID NO:38) and AAGGATCCAGATTCGTTTAAATTTTTCAGC [3' primer] (SEQ ID NO:39)), sub-cloned into the pET15b vector (pHAN1, spa codons 212-261) and recombinant plasmid transformed into *E. coli* BL21(DE3) to express and purify recombinant N-terminal His6-tagged protein. To generate mutations in the SpA-D coding sequence, sets of two pairs of primers were synthesized (for D to A substitutions: CTTCATCAAAGTCT-TAAAGCCGCCCAAGCCAAAGCACTAAC [5' primer] (SEQ ID NO:40) and GTTAGTGCTTTGGCTTGGGGCGGCTTTAAGACTTTT-GAATGAAG [3' primer] (SEQ ID NO:41); for Q to K substitutions CATATGTTCAACAAAGATAAAAAAAGCGCCTTCTATGAAATC [5' primer] (SEQ ID NO:42) and GATTTCATAGAAGGCGCTTTTTT-TATCTTTGTTGAACATATG [3' primer] (SEQ ID NO:43); for Q to G substitutions CATATGTTCAACAAAGATG-GAGGAAGCGCCTTCTATGAAATC [5' primer] (SEQ ID NO:44) and GATTTCATAGAAGGCGCTTCTTC-CATCTTTGTTGAACATATG' [3' primer] (SEQ ID NO:45). Primers were used for quick-change mutagenesis protocols. Following mutagenesis, DNA sequences were confirmed for each of the recombinant proteins: SpA, SpA-D and SpA-D<sub>Q9,10G;D36,37A</sub> and SpA-D<sub>Q9,10K;D36,37A</sub>. All proteins were purified from lysates of recombinant *E. coli* using Ni-NTA chromatography and subsequently dialyzed against PBS and stored at 4° C.

To measure binding of immunoglobulin to Protein A and its variants, 200  $\mu$ g of purified protein was diluted into a 1 ml volume using column buffer (50 mM Tris-HCl, 150 mM NaCl, pH7.5) and then loaded onto a pre-equilibrated Ni-NTA column (1 ml bed volume). Columns were washed with 10 ml of column buffer. 200  $\mu$ g of purified human IgG was diluted in a total volume of 1 ml column buffer and then applied to each of the columns charged with Protein A and its variants. The columns were subsequently washed with 5 ml wash buffer (10 mM imidazole in column buffer) and 5 ml column buffer. Protein samples were eluted with 2 ml elution buffer (500 mM imidazole in column buffer), fractions collected and aliquots subjected to SDS-PAGE gel electrophoresis, followed by Coomassie-Blue staining. As shown in FIG. 3, wild-type Protein A (SpA) and its SpA-domain D both retained immunoglobulin during chromatography. In contrast, the SpA-D<sub>Q9,10K;D36,37A</sub> variant did not bind to immunoglobulin.

To quantify the binding of Protein A and its variants to the Fc portion of immunoglobulin and the VH3 domain of Fab, HRP conjugated human immunoglobulin G [hIgG], the Fc portion of human IgG [hFc] and the F(ab)<sub>2</sub> portion of human IgG [hF(ab)<sub>2</sub>] as well as ELISA assays were used to quantify the relative amount binding to Protein A and its variants. The

data in FIG. 4 demonstrate the binding of SpA and SpA-D to hIgG and hFc, whereas SpA-D<sub>Q9,10G;D36,37A</sub> and SpA-D<sub>Q9,10K;D36,37A</sub> displayed only background binding activities. SpA bound similar amounts of hFc and hF(ab)<sub>2</sub>, however the binding of SpA-D to hF(ab)<sub>2</sub> was reduced compared to full length SpA. This result suggests that the presence of multiple IgG binding domains may cooperatively increase the ability of Protein A to bind to the B cell receptor. When compared with the reduced binding power of SpA-D for hF(ab)<sub>2</sub>, of the two variants only SpA-D<sub>Q9,10K;D36,37A</sub> displayed a significant reduction in the ability to bind the VH3 domain of immunoglobulin. To examine the toxicogenic attributes of SpA-D and its variants, purified proteins were injected into mice, which were sacrificed after 4 hours to remove their spleens. Organ tissue was homogenized, capsular material removed and B cells stained with fluorescent CD19 antibodies. Following FACS analysis to quantify the abundance of B cells in splenic tissues, it was observed that SpA-D caused a 5% drop in the B cell count compared to a mock (PBS) control (FIG. 5). In contrast, SpA-D<sub>Q9,10K;D36,37A</sub> did not cause a reduction in B-cell counts, indicating that the mutant molecule had lost its toxicogenic attributes of stimulating B cell proliferation and death (FIG. 5). In summary, amino acid substitutions in the SpA-D residues Q9, Q10, D36, and D37 abolished the ability of Protein A domains to bind immunoglobulins or exert toxicogenic functions in human and animal tissues.

Non-toxicogenic Protein A variants elicit vaccine protection. To test whether or not Protein A and its variants can function as vaccine antigens, SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub>, and SpA-D<sub>Q9,10G;D36,37A</sub> were emulsified with complete or incomplete Freund's adjuvant and immunized 4 week old BALB/c mice on day 1 and day 11 with 50 µg of purified protein. Cohort of animals (n=5) were analyzed for humoral immune responses to immunization by bleeding the animals before (day 0) and after the immunization schedule (day 21). Table 5 indicates that immunized mice generated only a modest humoral immune response directed at wild-type Protein A or its SpA-D module, whereas the amount of antibody raised following immunization with SpA-D<sub>Q9,10K;D36,37A</sub> or SpA-D<sub>Q9,10G;D36,37A</sub> was increased four

to five fold. Following intravenous challenge with 1×10<sup>7</sup> CFU *S. aureus* Newman, animals were killed on day 4, their kidneys removed and either analyzed for staphylococcal load (by plating tissue homogenate on agar plates and enumerating colony forming units, CFU) or histopathology. As expected, mock (PBS) immunized mice (n=19) harbored 6.46 log<sub>10</sub> (±0.25) CFU in kidney tissue and infectious lesions were organized into 3.7 (±1.2) abscesses per organ (n=10) (Table 5). Immunization of animals with SpA led to a 2.51 log<sub>10</sub> CFU reduction on day 5 (P=0.0003) with 2.1 (±1.2) abscesses per organ. The latter data indicate that there was no significant reduction in abscess formation (P=0.35). Immunization with SpA-D generated similar results: a 2.03 log<sub>10</sub> CFU reduction on day 5 (P=0.0001) with 1.5 (±0.8) abscesses per organ (P=0.15). In contrast, immunization with SpA-D<sub>Q9,10K;D36,37A</sub> or SpA-D<sub>Q9,10G;D36,37A</sub> created increased protection, with 3.07 log<sub>10</sub> and 3.03 log<sub>10</sub> CFU reduction on day 4, respectively (statistical significance P<0.0001 for both observations). Further, immunization with both SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10G;D36,37A</sub> generated significant protection from staphylococcal abscess formation, as only 0.5 (±0.4) and 0.8 (±0.5) infectious lesions per organ (P=0.02 and P=0.04) were identified. Thus, immunization with non-toxicogenic Protein A variants generates increased humoral immune responses for Protein A and provides protective immunity against staphylococcal challenge. These data indicate that Protein A is an ideal candidate for a human vaccine that prevents *S. aureus* disease.

These exciting results have several implications for the design of a human vaccine. First, the generation of substitution mutations that affect the ability of the immunoglobulin binding domains of Protein A, either alone or in combination of two or more domains, can generate non-toxicogenic variants suitable for vaccine development. It seems likely that a combination of mutant IgG binding domains closely resembling the structure of Protein A can generate even better humoral immune responses as is reported here for the SpA-domain D alone. Further, a likely attribute of Protein A specific antibodies may be that the interaction of antigen binding sites with the microbial surface can neutralize the ability of staphylococci to capture immunoglobulins via their Fc portion or to stimulate the B cell receptor via the VH3 binding activities.

TABLE 5

Non-toxicogenic Protein A variants as vaccine antigens that prevent <i>S. aureus</i> disease									
Antigen	Bacterial load in kidney (n = number of mice)			IgG titer	Abscess formation in mice (n = number of mice)				
	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> Reduction	<sup>c</sup> P value		<sup>d</sup> Surface abscess	Reduction	<sup>e</sup> Histopathology	Reduction	<sup>f</sup> p value
Mock	6.46 ± 0.25 (n = 19)	—	—	<100	14/19 (70%)	—	3.7 ± 1.2 (n = 10)	—	—
SpA	3.95 ± 0.56 (n = 20)	2.51	0.0003	1706 ± 370	10/20 (50%)	32%	2.1 ± 1.2 (n = 10)	2.2	0.35
SpA-D	4.43 ± 0.41 (n = 18)	2.03	0.0001	381 ± 27	10/18 (55%)	25%	1.5 ± 0.8 (n = 10)	2.2	0.15
SpA-D1	3.39 ± 0.50 (n = 19)	3.07	<0.0001	5600 ± 801	6/20 (30%)	59%	0.5 ± 0.4 (n = 10)	3.2	0.02
SpA-D2	3.43 ± 0.46 (n = 19)	3.03	<0.0001	3980 ± 676	6/19 (32%)	57%	0.8 ± 0.5 (n = 10)	2.9	0.04

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of 18 to 20 BALB/c mice. Standard error of the means (±SEM) is indicated.

<sup>c</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>b</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Abscess formation in kidney tissues four days following infection was measured by macroscopic inspection (% positive)

<sup>e</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys from ten animals; the number of abscesses per kidney was recorded and averaged for the final mean (±SEM).

<sup>f</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

SpA-D1 and SpA-D2 represent SpA-D<sub>Q9, 10K; D36, 37A</sub> and SpA-D<sub>Q9, 10G; D36, 37A</sub>, respectively.

Vaccine protection in murine abscess, murine lethal infection, and murine pneumonia models. Three animal models have been established for the study of *S. aureus* infectious disease. These models are used here to examine the level of protective immunity provided via the generation of Protein A specific antibodies.

Murine abscess—BALB/c mice (24-day-old female, 8-10 mice per group, Charles River Laboratories, Wilmington, Mass.) are immunized by intramuscular injection into the hind leg with purified protein (Chang et al., 2003; Schneewind et al., 1992). Purified SpA, SpA-D or SpA-DQ9,10K; D36,37A (50 µg protein) is administered on days 0 (emulsified 1:1 with complete Freund's adjuvant) and 11 (emulsified 1:1 with incomplete Freund's adjuvant). Blood samples are drawn by retroorbital bleeding on days 0, 11, and 20. Sera are examined by ELISA for IgG titers for specific SpA-D and SpA-DQ9,10K;D36,37A binding activity. Immunized animals are challenged on day 21 by retroorbital injection of 100 µl of *S. aureus* Newman or *S. aureus* USA300 suspension ( $1 \times 10^7$  cfu). For this, overnight cultures of *S. aureus* Newman are diluted 1:100 into fresh tryptic soy broth and grown for 3 h at 37° C. Staphylococci are centrifuged, washed twice, and diluted in PBS to yield an A600 of 0.4 ( $1 \times 10^8$  cfu per ml). Dilutions are verified experimentally by agar plating and colony formation. Mice are anesthetized by intraperitoneal injection of 80-120 mg of ketamine and 3-6 mg of xylazine per kilogram of body weight and infected by retroorbital injection. On day 5 or 15 following challenge, mice are euthanized by compressed CO<sub>2</sub> inhalation. Kidneys are removed and homogenized in 1% Triton X-100. Aliquots are diluted and plated on agar medium for triplicate determination of cfu. For histology, kidney tissue is incubated at room temperature in 10% formalin for 24 h. Tissues are embedded in paraffin, thin-sectioned, stained with hematoxylin-eosin, and examined by microscopy.

Murine lethal infection—BALB/c mice (24-day-old female, 8-10 mice per group, Charles River Laboratories, Wilmington, Mass.) are immunized by intramuscular injection into the hind leg with purified SpA, SpA-D or SpA-DQ9,10K;D36,37A (50 µg protein). Vaccine is administered on days 0 (emulsified 1:1 with complete Freund's adjuvant) and 11 (emulsified 1:1 with incomplete Freund's adjuvant). Blood samples are drawn by retroorbital bleeding on days 0, 11, and 20. Sera are examined by ELISA for IgG titers with specific SpA-D and SpA-DQ9,10K;D36,37A binding activity. Immunized animals are challenged on day 21 by retroorbital injection of 100 µl of *S. aureus* Newman or *S. aureus* USA300 suspension ( $15 \times 10^7$  cfu) (34). For this, overnight cultures of *S. aureus* Newman are diluted 1:100 into fresh tryptic soy broth and grown for 3 h at 37° C. Staphylococci are centrifuged, washed twice, diluted in PBS to yield an A<sub>600</sub> of 0.4 ( $1 \times 10^8$  cfu per ml) and concentrated. Dilutions are verified experimentally by agar plating and colony formation. Mice are anesthetized by intraperitoneal injection of 80-120 mg of ketamine and 3-6 mg of xylazine per kilogram of body weight. Immunized animals are challenged on day 21 by intraperitoneal inject with  $2 \times 10^{10}$  cfu of *S. aureus* Newman or  $3 \cdot 10 \times 10^9$  cfu of clinical *S. aureus* isolates. Animals are monitored for 14 days, and lethal disease is recorded.

Murine pneumonia model—*S. aureus* strains Newman or USA300 (LAC) are grown at 37° C. in tryptic soy broth/agar to OD<sub>660</sub> 0.5. 50-ml culture aliquots are centrifuged, washed in PBS, and suspended in 750 µl PBS for mortality studies ( $3 \cdot 4 \times 10^8$  CFU per 30-µl volume), or 1,250 µl PBS ( $2 \times 10^8$  CFU per 30-µl volume) for bacterial load and histopathol-

ogy experiments (2, 3). For lung infection, 7-wk-old C57BL/6J mice (The Jackson Laboratory) are anesthetized before inoculation of 30 µl of *S. aureus* suspension into the left nare. Animals are placed into the cage in a supine position for recovery and observed for 14 days. For active immunization, 4-wk-old mice receive 20 µg SpA-D or SpA-DQ9,10K;D36,37A in CFA on day 0 via the i.m. route, followed by a boost with 20 µg SpA-D or SpA-DQ9,10K;D36,37A in incomplete Freund's adjuvant (IFA) on day 10. Animals are challenged with *S. aureus* on day 21. Sera are collected before immunization and on day 20 to assess specific antibody production. For passive immunization studies, 7-wk-old mice receive 100 µl of either NRS (normal rabbit serum) or SpA-D-specific rabbit antisera via i.p. injection 24 h before challenge. To assess the pathological correlates of pneumonia, infected animals are killed via forced CO<sub>2</sub> inhalation before removal of both lungs. The right lung is homogenized for enumeration of lung bacterial load. The left lung is placed in 1% formalin and paraffin embedded, thin sectioned, stained with hematoxylin-eosin, and analyzed by microscopy.

Rabbit antibodies—Purified 200 µg SpA-D or SpA-DQ9,10K;D36,37A is used as an immunogen for the production of rabbit antisera. 200 µg protein is emulsified with CFA for injection at day 0, followed by booster injections with 200 µg protein emulsified with IFA on days 21 and 42. Rabbit antibody titers are determined by ELISA. Purified antibodies are obtained by affinity chromatography of rabbit serum on SpA-D or SpA-DQ9,10K;D36,37A sepharose. The concentration of eluted antibodies is measured by absorbance at A280 and specific antibody titers are determined by ELISA.

Active immunization with SpA-domain D variants.—To determine vaccine efficacy, animals are actively immunized with purified SpA-D or SpADQ9,10K;D36,37A. As a control, animals are immunized with adjuvant alone. Antibody titers against Protein A preparations are determined using SpA-D or SpA-DQ9,10K;D36,37A as antigens; note that the SpA-DQ9,10K;D36,37A variant cannot bind the Fc or Fab portion of IgG. Using infectious disease models described above, any reduction in bacterial load (murine abscess and pneumonia), histopathology evidence of staphylococcal disease (murine abscess and pneumonia) and protection from lethal disease (murine lethal challenge and pneumonia) is measured.

Passive immunization with affinity purified rabbit polyclonal antibodies generated against SpA-domain D variants. To determine protective immunity of Protein A specific rabbit antibodies, mice are passively immunized with 5 mg/kg of purified SpA-D or SpA-DQ9,10K;D36,37A derived rabbit antibodies. Both of these antibody preparations are purified by affinity chromatography using immobilized SpA-D or SpA-DQ9,10K;D36,37A. As a control, animals are passively immunized with rV10 antibodies (a plague protective antigen that has no impact on the outcome of staphylococcal infections). Antibody titers against all Protein A preparations are determined using SpA-DQ9,10K;D36,37A as an antigen, as this variant cannot bind the Fc or Fab portion of IgG. Using the infectious disease models described above, the reduction in bacterial load (murine abscess and pneumonia), histopathology evidence of staphylococcal disease (murine abscess and pneumonia), and the protection from lethal disease (murine lethal challenge and pneumonia) is measured.

#### Example 2

##### Non-Toxicogenic Protein A Vaccine for Methicillin-Resistant *Staphylococcus Aureus* Infection

Clinical isolates of *S. aureus* express protein A (Shopsin et al., 1999, whose primary translational product is com-

prised of an N-terminal signal peptide (DeDent et al., 2008), five Ig-BDs (designated E, D, A, B and C)(Sjodahl, 1977), region X with variable repeats of an eight residue peptide (Guss et al., 1984), and C-terminal sorting signal for the cell wall anchoring of SpA (Schneewind et al., 1992; Schneewind et al., 1995) (FIG. 6). Guided by amino acid homology (Uhlen et al., 1984), the triple  $\alpha$ -helical bundle structure of IgBDs (Deisenhofer et al., 1978; Deisenhofer et al., 1981) and their atomic interactions with Fab V<sub>H</sub>3 (Graille et al., 2000) or Fc $\gamma$  (Gouda et al., 1998), glutamine 9 and 10 were selected as well as aspartate 36 and 37 as critical for the association of SpA with antibodies or B cell receptor, respectively. Substitutions Gln9Lys, Gln10Lys, Asp36Ala and Asp37Ala were introduced into the D domain to generate SpA-D<sub>KKAA</sub> (FIG. 6). The ability of isolated SpA-D or SpA-D<sub>KKAA</sub> to bind human IgG was analyzed by affinity chromatography (FIG. 6). Polyhistidine tagged SpA-D as well as full-length SpA retained human IgG on Ni-NTA, whereas SpA-D<sub>KKAA</sub> and a negative control (SrtA) did not (FIG. 6). A similar result was observed with von Willebrand factor (Hartleib et al., 2000), which, along with tumor necrosis factor receptor 1 (TNFR1)(Gomez et al., 2004), can also bind protein A via glutamine 9 and 10 (FIG. 6). Human immunoglobulin encompasses 60-70% V<sub>H</sub>3-type IgG. The inventors distinguish between Fc domain and B cell receptor activation of Igs and measured association of human Fc $\gamma$  and F(ab)<sub>2</sub> fragments, both of which bound to full-length SpA or SpA-D, but not to SpA-D<sub>KKAA</sub> (FIG. 6). Injection of SpA-D into the peritoneal cavity of mice resulted in B cell expansion followed by apoptotic collapse of CD19+ lymphocytes in spleen tissue of BALB/c mice (Goodyear and Silverman, 2003)(FIG. 6). B cell superantigen activity was not observed following injection with SpA-D<sub>KKAA</sub>, and TUNEL-staining of splenic tissue failed to detect the increase in apoptotic cells that follows injection of SpA or SpA-D (FIG. 6).

Naive six week old BALB/c mice were injected with 50  $\mu$ g each of purified SpA, SpA-D or SpA-D<sub>KKAA</sub> emulsified in CFA and boosted with the same antigen emulsified in IFA. In agreement with the hypothesis that SpA-D promotes the apoptotic collapse of activated clonal B cell populations, the inventors observed a ten-fold higher titer of SpA-D<sub>KKAA</sub> specific antibodies following immunization of mice with the non-toxicogenic variant as compared to the B cell superantigen (SpA-D vs. SpA-D<sub>KKAA</sub> P<0.0001, Table 6). Antibody titers raised by immunization with full-length SpA were higher than those elicited by SpA-D (P=0.0022), which is likely due to the larger size and reiterative domain structure of this antigen (Table 6). Nevertheless, even SpA elicited lower antibody titers than SpA-D<sub>KKAA</sub> (P=0.0003), which encompasses only 50 amino acids of protein A (520 residues, SEQ ID NO:33). Immunized mice were challenged by intravenous inoculation with *S. aureus* Newman and the ability of staphylococci to seed abscesses in renal tissues was examined by necropsy four days after challenge. In homogenized renal tissue of mock (PBS/adjuvant) immunized mice, an average staphylococcal load of 6.46 log<sub>10</sub> CFU g<sup>-1</sup> was enumerated (Table 6). Immunization of mice with SpA or SpA-D led to a reduction in staphylococcal load, however SpA-D<sub>KKAA</sub> vaccinated animals displayed an even greater, 3.07 log<sub>10</sub> CFU g<sup>-1</sup> reduction of *S. aureus* Newman in renal tissues (P<0.0001, Table 6). Abscess formation in kidneys was analyzed by histopathology (FIG. 7). Mock immunized animals harbored an average of 3.7 ( $\pm$ 1.2) abscesses per kidney (Table 6). Vaccination with SpA-D<sub>KKAA</sub> reduced the average number of abscesses to 0.5 ( $\pm$ 0.4)(P=0.0204), whereas immunization with SpA or SpA-D did not cause a significant reduction in the number of abscess lesions (Table

6). Lesions from SpA-D<sub>KKAA</sub> vaccinated animals were smaller in size, with fewer infiltrating PMNs and characteristically lacked staphylococcal abscess communities (Cheng et al., 2009)(FIG. 7). Abscesses in animals that had been immunized with SpA or SpA-D displayed the same overall structure of lesions in mock immunized animals (FIG. 7).

The inventors examined whether SpA-D<sub>KKAA</sub> immunization can protect mice against MRSA strains and selected the USA300 LAC isolate for animal challenge (Diep et al., 2006). This highly virulent CA-MRSA strain spread rapidly throughout the United States, causing significant human morbidity and mortality (Kennedy et al., 2008). Compared to adjuvant control mice, SpA-D<sub>KKAA</sub> immunized animals harbored a 1.07 log<sub>10</sub> CFU g<sup>-1</sup> reduction in bacterial load of infected kidney tissues. Histopathology examination of renal tissue following *S. aureus* USA300 challenge revealed that the average number of abscesses was reduced from 4.04 ( $\pm$ 0.8) to 1.6 ( $\pm$ 0.6)(P=0.02774). In contrast, SpA or SpA-D immunization did not cause a significant reduction in bacterial load or abscess formation (Table 6).

Rabbits were immunized with SpA-D<sub>KKAA</sub> and specific antibodies were purified on SpA-D<sub>KKAA</sub> affinity column followed by SDS-PAGE (FIG. 8). SpA-D<sub>KKAA</sub> specific IgG was cleaved with pepsin to generate Fc $\gamma$  and F(ab)<sub>2</sub> fragments, the latter of which were purified by chromatography on SpA-D<sub>KKAA</sub> column (FIG. 8). Binding of human IgG or vWF to SpA or SpA-D was perturbed by SpA-D<sub>KKAA</sub> specific F(ab)<sub>2</sub>, indicating that SpA-D<sub>KKAA</sub> derived antibodies neutralize the B cell superantigen function of protein A as well as its interactions with Ig (FIG. 8).

To further improve the vaccine properties for non-toxicogenic protein A, the inventors generated SpA<sub>KKAA</sub>, which includes all five IgBDs with four amino acid substitutions—substitutions corresponding to Gln9Lys, Gln10Lys, Asp36Ala and Asp37Ala of domain D—in each of its five domains (E, D, A, B and C). Polyhistidine tagged SpA<sub>KKAA</sub> was purified by affinity chromatography and analyzed by Coomassie Blue-stained SDS-PAGE (FIG. 9). Unlike full-length SpA, SpA<sub>KKAA</sub> did not bind human IgG, Fc and F(ab)<sub>2</sub> or vWF (FIG. 9). SpA<sub>KKAA</sub> failed to display B cell superantigen activity, as injection of the variant into BALB/c mice did not cause a depletion of CD19+ B cells in splenic tissue (FIG. 9). SpA<sub>KKAA</sub> vaccination generated higher specific antibody titers than SpA-D<sub>KKAA</sub> immunization and provided mice with elevated protection against *S. aureus* USA300 challenge (Table 6). Four days following challenge, SpA<sub>KKAA</sub> vaccinated animals harbored 3.54 log<sub>10</sub> CFU g<sup>-1</sup> fewer staphylococci in renal tissues (P=0.0001) and also caused a greater reduction in the number of abscess lesions (P=0.0109) (Table 6).

SpA<sub>KKAA</sub> was used to immunize rabbits. Rabbit antibodies specific for SpA-D<sub>KKAA</sub> or SpA<sub>KKAA</sub> were affinity purified on matrices with immobilized cognate antigen and injected at a concentration of 5 mg kg<sup>-1</sup> body weight into the peritoneal cavity of BALB/c mice (Table 7). Twenty-four hours later, specific antibody titers were determined in serum and animals challenged by intravenous inoculation with *S. aureus* Newman. Passive transfer reduced the staphylococcal load in kidney tissues for SpA-D<sub>KKAA</sub> (P=0.0016) or SpA<sub>KKAA</sub> (P=0.0005) specific antibodies. On histopathology examination, both antibodies reduced the abundance of abscess lesions in the kidneys of mice challenged with *S. aureus* Newman (Table 7). Together these data reveal that vaccine protection following immunization with SpA-D<sub>KKAA</sub> or SpA<sub>KKAA</sub> is conferred by antibodies that neutralize protein A.

TABLE 6

Immunization of mice with protein A vaccines.						
Staphylococcal load and abscess formation in renal tissue						
Antigen	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> P-value	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>d</sup> IgG Titer	<sup>e</sup> Number of abscesses	<sup>f</sup> P-value
<i>S. aureus</i> Newman challenge						
Mock	6.46 ± 0.25	—	—	<100	3.7 ± 1.2	—
SpA	3.95 ± 0.56	0.0003	2.51	1706 ± 370	2.1 ± 1.2	0.3581
SpA-D	4.43 ± 0.41	0.0001	2.03	381 ± 27	1.5 ± 0.8	0.1480
SpA-D <sub>KKAA</sub>	3.39 ± 0.50	<0.0001	3.07	5600 ± 801	0.5 ± 0.4	0.0204
<i>S. aureus</i> USA300 (LAC) challenge						
Mock	7.20 ± 0.24	—	—	<100	4.0 ± 0.8	—
SpA	6.81 ± 0.26	0.2819	0.39	476 ± 60	3.3 ± 1.0	0.5969
SpA-D	6.34 ± 0.52	0.1249	0.86	358 ± 19	2.2 ± 0.6	0.0912
SpA-D <sub>KKAA</sub>	6.00 ± 0.42	0.0189	1.20	3710 ± 1147	1.6 ± 0.6	0.0277
SpA <sub>KKAA</sub>	3.66 ± 0.76	0.0001	3.54	10200 ± 2476	1.2 ± 0.5	0.0109

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of fifteen to twenty BALB/c mice per immunization. Representative of two independent and reproducible animal experiments is shown. Standard error of the means (±SEM) is indicated.

<sup>b</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Means of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA.

<sup>e</sup>Histopathology of hematoxyline-eosin stained, thin sectioned kidneys from ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

TABLE 7

Passive immunization of mice with antibodies against protein A.						
Staphylococcal load and abscess formation in renal tissue						
<sup>a</sup> Antibody	<sup>b</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>c</sup> P-value	<sup>d</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>e</sup> IgG Titer	<sup>f</sup> Number of abscesses	<sup>g</sup> P-value
Mock	7.10 ± 0.14	—	—	<100	4.5 ± 0.8	—
α-SpA-D <sub>KKAA</sub>	5.53 ± 0.43	0.0016	1.57	466 ± 114	1.9 ± 0.7	0.0235
α-SpA <sub>KKAA</sub>	5.69 ± 0.34	0.0005	1.41	1575 ± 152	1.6 ± 0.5	0.0062

<sup>a</sup>Affinity purified antibodies were injected into the peritoneal cavity of BALB/c mice at a concentration of 5 mg · kg<sup>-1</sup> twenty-four hours prior to intravenous challenge with 1 × 10<sup>7</sup> CFU *S. aureus* Newman.

<sup>b</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of fifteen BALB/c mice per immunization. Representative of two independent and reproducible animal experiments is shown. Standard error of the means (±SEM) is indicated.

<sup>c</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>d</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>e</sup>Means of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA.

<sup>f</sup>Histopathology of hematoxyline-eosin stained, thin sectioned kidneys from ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

Following infection with virulent *S. aureus*, mice do not develop protective immunity against subsequent infection with the same strain (Burts et al., 2008)(FIG. 10). The average abundance of SpA-D<sub>KKAA</sub> specific IgG in these animals was determined by dot blot as 0.20 μg ml<sup>-1</sup> (±0.04) and 0.14 μg ml<sup>-1</sup> (±0.01) for strains Newman and USA300 LAC, respectively (FIG. 9). The minimal concentration of protein A-specific IgG required for disease protection in SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub> vaccinated animals (P 0.05 log<sub>10</sub> reduction in staphylococcal CFU g<sup>-1</sup> renal tissue) was calculated as 4.05 μg ml<sup>-1</sup> (±0.88). Average serum concentration of SpA-specific IgG in adult healthy human volunteers (n=16) was 0.21 μg ml<sup>-1</sup> (±0.02). Thus, *S. aureus* infections in mice or humans are not associated with immune responses that raise significant levels of neutralizing antibodies directed against protein A, which is likely due to the B cell superantigen attributes of this molecule. In contrast, the average serum concentration of IgG specific for diphtheria toxin in human volunteers, 0.068 μg ml<sup>-1</sup> (±0.20), was within range for protective immunity against diphtheria (Behring, 1890; Lagergard et al., 1992).

Clinical *S. aureus* isolates express protein A, an essential virulence factor whose B cell superantigen activity and evasive attributes towards opsono-phagocytic clearance are absolutely required for staphylococcal abscess formation (Palmqvist et al., 2005; Cheng et al., 2009; Silverman and Goodyear, 2006). Protein A can thus be thought of as a toxin, essential for pathogenesis, whose molecular attributes must be neutralized in order to achieve protective immunity. By generating non-toxic variants unable to bind Igs via Fcγ or VH<sub>3</sub>-Fab domains, the inventors measure here for the first time protein A neutralizing immune responses as a correlate for protective immunity against *S. aureus* infection. In contrast to many methicillin-sensitive strains, CA-MRSA isolate USA300 LAC is significantly more virulent (Cheng et al., 2009). For example, immunization of experimental animals with the surface protein IsdB (Kuklin et al., 2006; Stranger-Jones et al., 2006) raises antibodies that confer protection against *S. aureus* Newman (Stranger-Jones et al., 2009) but not against USA300 challenge.

The methods utilized include:

Bacterial strains and growth. *Staphylococcus aureus* strains Newman and USA300 were grown in tryptic soy broth (TSB) at 37°C. *Escherichia coli* strains DH5a and BL21 (DE3) were grown in Luria-Bertani (LB) broth with 100 µg ml<sup>-1</sup> ampicillin at 37°C.

Rabbit Antibodies. The coding sequence for SpA was PCR-amplified with two primers, gctgcatatggcgcaacacgatgaagctcaac (SEQ ID NO:35) and agtggatccttatgcttgagctttgttagcatctgc (SEQ ID NO:36) using *S. aureus* Newman template DNA. SpA-D was PCR-amplified with two primers, aacatattcaacaagaatcaacaagc (SEQ ID NO:38) and aaggatccagatctgtaatttttagc (SEQ ID NO:39). The sequence for SpA-D<sub>KKAA</sub> was mutagenized with two sets of primers catatgttcaacaagataaaaaagcgcttctatgaaate (SEQ ID NO:42) and gattcatagaaggcgctttttatcttggtaacatag (SEQ ID NO:43) for Q9K, Q10K as well as ctctcatcaaaagcttaagccgcccccaagcaagcactaac (SEQ ID NO:40) and gtagtgctttggcttggcgcgctttaagacttgaatgaag (SEQ ID NO:41) for D36A, D37A. The sequence of SpA<sub>KKAA</sub> was synthesized by Integrated DNA Technologies, Inc. PCR products were cloned into pET-15b generating N-terminal His6 tagged recombinant protein. Plasmids were transformed into BL21(DE3). Overnight cultures of transformants were diluted 1:100 into fresh media and grown at 37°C to an OD<sub>600</sub> of 0.5, at which point cultures were induced with 1 mM isopropyl 3-D-1-thiogalactopyranoside (IPTG) and grown for an additional three hours. Bacterial cells were sedimented by centrifugation, suspended in column buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl) and disrupted with a French pressure cell at 14,000 psi. Lysates were cleared of membrane and insoluble components by ultracentrifugation at 40,000×g. Proteins in the soluble lysate were subjected to nickel-nitrilotriacetic acid (Ni-NTA, Qiagen) affinity chromatography. Proteins were eluted in column buffer containing successively higher concentrations of imidazole (100-500 mM). Protein concentrations were determined by bicinchoninic acid (BCA) assay (Thermo Scientific). For antibody generation, rabbits (6 month old New-Zealand white, female, Charles River Laboratories) were immunized with 500 µg protein emulsified in Complete Freund's Adjuvant (Difco) by subscapular injection. For booster immunizations, proteins emulsified in Incomplete Freund's Adjuvant and injected 24 or 48 days following the initial immunization. On day 60, rabbits were bled and serum recovered.

Purified antigen (5 mg protein) was covalently linked to HiTrap NETS-activated HP columns (GE Healthcare). Antigen-matrix was used for affinity chromatography of 10-20 ml of rabbit serum at 4°C. Charged matrix was washed with 50 column volumes of PBS, antibodies eluted with elution buffer (1 M glycine, pH 2.5, 0.5 M NaCl) and immediately neutralized with 1M Tris-HCl, pH 8.5. Purified antibodies were dialyzed overnight against PBS at 4°C.

F(ab)<sub>2</sub> fragments. Affinity purified antibodies were mixed with 3 mg of pepsin at 37°C for 30 minutes. The reaction was quenched with 1 M Tris-HCl, pH 8.5 and F(ab)<sub>2</sub> fragments were affinity purified with specific antigen-conjugated HiTrap NETS-activated HP columns. Purified antibodies were dialyzed overnight against PBS at 4°C., loaded onto SDS-PAGE gel and visualized with Coomassie Blue staining.

Active and passive immunization. BALB/c mice (3 week old, female, Charles River Laboratories) were immunized with 50 µg protein emulsified in Complete Freund's Adjuvant (Difco) by intramuscular injection. For booster immunizations, proteins were emulsified in Incomplete Freund's

Adjuvant and injected 11 days following the initial immunization. On day 20 following immunization, 5 mice were bled to obtain sera for specific antibody titers by enzyme-linked immunosorbent assay (ELISA).

Affinity purified antibodies in PBS were injected at a concentration 5 mg kg<sup>-1</sup> of experimental animal weight into the peritoneal cavity of BALB/c mice (6 week old, female, Charles River Laboratories) 24 hours prior to challenge with *S. aureus*. Animal blood was collected via periorbital vein puncture. Blood cells were removed with heparinized microhematocrit capillary tubes (Fisher) and Z-gel serum separation micro tubes (Sarstedt) were used to collect and measure antigen specific antibody titers by ELISA.

Mouse renal abscess. Overnight cultures of *S. aureus* Newman or USA300 (LAC) were diluted 1:100 into fresh TSB and grown for 2 hours at 37°C. *Staphylococci* were sedimented, washed and suspended PBS at OD<sub>600</sub> of 0.4 (~1×10<sup>8</sup> CFU ml<sup>-1</sup>). Inocula were quantified by spreading sample aliquots on TSA and enumerating colonies formed. BALB/c mice (6 week old, female, Charles River Laboratories) were anesthetized via intraperitoneal injection with 100 mg ml<sup>-1</sup> ketamine and 20 mg ml<sup>-1</sup> xylazine per kilogram of body weight. Mice were infected by retro-orbital injection with 1×10<sup>7</sup> CFU of *S. aureus* Newman or 5×10<sup>6</sup> CFU of *S. aureus* USA300. On day 4 following challenge, mice were killed by CO<sub>2</sub> inhalation. Both kidneys were removed, and the staphylococcal load in one organ was analyzed by homogenizing renal tissue with PBS, 1% Triton X-100. Serial dilutions of homogenate were spread on TSA and incubated for colony formation. The remaining organ was examined by histopathology. Briefly, kidneys were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with hematoxylin-eosin, and inspected by light microscopy to enumerate abscess lesions. All mouse experiments were performed in accordance with the institutional guidelines following experimental protocol review and approval by the Institutional Biosafety Committee (IBC) and the Institutional Animal Care and Use Committee (IACUC) at the University of Chicago.

Protein A binding. For human IgG binding, Ni-NTA affinity columns were pre-charged with 200 µg of purified proteins (SpA, SpA-D, SpA-D<sub>KKAA</sub> and SrtA) in column buffer. After washing, 200 µg of human IgG (Sigma) was loaded onto the column. Protein samples were collected from washes and elutions and subjected to SDS-PAGE gel electrophoresis, followed by Coomassie Blue staining. Purified proteins (SpA, SpA<sub>KKAA</sub>, SpA-D and SpA-D<sub>KKAA</sub>) were coated onto MaxiSorp ELISA plates (NUNC) in 0.1M carbonate buffer (pH 9.5) at 1 µg ml<sup>-1</sup> concentration overnight at 4°C. Plates were next blocked with 5% whole milk followed by incubation with serial dilutions of peroxidase-conjugated human IgG, Fc or F(ab)<sub>2</sub> fragments for one hour. Plates were washed and developed using OptEIA ELISA reagents (BD). Reactions were quenched with 1 M phosphoric acid and A<sub>450</sub> readings were used to calculate half maximal titer and percent binding.

von Willebrand Factor (vWF) binding assays. Purified proteins (SpA, SpA<sub>KKAA</sub>, SpA D and SpA-D<sub>KKAA</sub>) were coated and blocked as described above. Plates were incubated with human vWF at 1 µg ml<sup>-1</sup> concentration for two hours, then washed and blocked with human IgG for another hour. After washing, plates were incubated with serial dilution of peroxidase-conjugated antibody directed against human vWF for one hour. Plates were washed and developed using OptEIA ELISA reagents (BD). Reactions were quenched with 1 M phosphoric acid and A<sub>450</sub> readings were

used to calculate half maximal titer and percent binding. For inhibition assays, plates were incubated with affinity purified F(ab)<sub>2</sub> fragments specific for SpA-D<sub>KKAA</sub> at 10 µg ml<sup>-1</sup> concentration for one hour prior to ligand binding assays.

Splenocyte apoptosis. Affinity purified proteins (150 µg of SpA, SpA-D, SpA<sub>KKAA</sub>, and SpA-D<sub>KKAA</sub>) were injected into the peritoneal cavity of BALB/c mice (6 week old, female, Charles River Laboratories). Four hours following injection, animals were killed by CO<sub>2</sub> inhalation. Their spleens were removed and homogenized. Cell debris were removed using cell strainer and suspended cells were transferred to ACK lysis buffer (0.15 M NH<sub>4</sub>Cl, 10 mM KHCO<sub>3</sub>, 0.1 mM EDTA) to lyse red blood cells. White blood cells were sedimented by centrifugation, suspended in PBS and stained with 1:250 diluted R-PE conjugated anti-CD19 monoclonal antibody (Invitrogen) on ice and in the dark for one hour. Cells were washed with 1% FBS and fixed with 4% formalin overnight at 4° C. The following day, cells were diluted in PBS and analyzed by flow cytometry. The remaining organ was examined for histopathology. Briefly, spleens were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with the Apoptosis detection kit (Millipore), and inspected by light microscopy.

Antibody quantification. Sera were collected from healthy human volunteers or BALB/c mice that had been either infected with *S. aureus* Newman or USA300 for 30 days or that had been immunized with SpA-D<sub>KKAA</sub>/SpA<sub>KKAA</sub> as described above. Human/mouse IgG (Jackson Immunology Laboratory), SpA<sub>KKAA</sub>, and CRM197 were blotted onto nitrocellulose membrane. Membranes were blocked with 5% whole milk, followed by incubation with either human or mouse sera. IRDye 700DX conjugated affinity purified anti-human/mouse IgG (Rockland) was used to quantify signal intensities using the Odyssey™ infrared imaging system (Li-cor). Experiments with blood from human volunteers involved protocols that were reviewed, approved and performed under regulatory supervision of The University of Chicago's Institutional Review Board (IRB).

Statistical Analysis. Two tailed Student's t tests were performed to analyze the statistical significance of renal abscess, ELISA, and B cell superantigen data.

#### REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

U.S. Pat. No. 3,791,932  
 U.S. Pat. No. 3,949,064  
 U.S. Pat. No. 4,174,384  
 U.S. Pat. No. 4,338,298  
 U.S. Pat. No. 4,356,170  
 U.S. Pat. No. 4,367,110  
 U.S. Pat. No. 4,372,945  
 U.S. Pat. No. 4,452,901  
 U.S. Pat. No. 4,474,757  
 U.S. Pat. No. 4,554,101  
 U.S. Pat. No. 4,578,770  
 U.S. Pat. No. 4,596,792  
 U.S. Pat. No. 4,599,230  
 U.S. Pat. No. 4,599,231  
 U.S. Pat. No. 4,601,903  
 U.S. Pat. No. 4,608,251  
 U.S. Pat. No. 4,683,195  
 U.S. Pat. No. 4,683,202

U.S. Pat. No. 4,684,611  
 U.S. Pat. No. 4,690,915  
 U.S. Pat. No. 4,748,018  
 U.S. Pat. No. 4,800,159  
 U.S. Pat. No. 4,879,236  
 U.S. Pat. No. 4,952,500  
 U.S. Pat. No. 5,084,269  
 U.S. Pat. No. 5,199,942  
 U.S. Pat. No. 5,221,605  
 U.S. Pat. No. 5,238,808  
 U.S. Pat. No. 5,302,523  
 U.S. Pat. No. 5,310,687  
 U.S. Pat. No. 5,322,783  
 U.S. Pat. No. 5,384,253  
 U.S. Pat. No. 5,464,765  
 U.S. Pat. No. 5,512,282  
 U.S. Pat. No. 5,538,877  
 U.S. Pat. No. 5,538,880  
 U.S. Pat. No. 5,548,066  
 U.S. Pat. No. 5,550,318  
 U.S. Pat. No. 5,563,055  
 U.S. Pat. No. 5,580,859  
 U.S. Pat. No. 5,589,466  
 U.S. Pat. No. 5,591,616  
 U.S. Pat. No. 5,610,042  
 U.S. Pat. No. 5,620,896  
 U.S. Pat. No. 5,648,240  
 U.S. Pat. No. 5,656,610  
 U.S. Pat. No. 5,702,932  
 U.S. Pat. No. 5,736,524  
 U.S. Pat. No. 5,780,448  
 U.S. Pat. No. 5,789,215  
 U.S. Pat. No. 5,801,234  
 U.S. Pat. No. 5,840,846  
 U.S. Pat. No. 5,843,650  
 U.S. Pat. No. 5,846,709  
 U.S. Pat. No. 5,846,783  
 U.S. Pat. No. 5,849,497  
 U.S. Pat. No. 5,849,546  
 U.S. Pat. No. 5,849,547  
 U.S. Pat. No. 5,858,652  
 U.S. Pat. No. 5,866,366  
 U.S. Pat. No. 5,871,986  
 U.S. Pat. No. 5,916,776  
 U.S. Pat. No. 5,922,574  
 U.S. Pat. No. 5,925,565  
 U.S. Pat. No. 5,928,905  
 U.S. Pat. No. 5,928,906  
 U.S. Pat. No. 5,932,451  
 U.S. Pat. No. 5,935,819  
 U.S. Pat. No. 5,935,825  
 U.S. Pat. No. 5,939,291  
 U.S. Pat. No. 5,942,391  
 U.S. Pat. No. 5,945,100  
 U.S. Pat. No. 5,958,895  
 U.S. Pat. No. 5,981,274  
 U.S. Pat. No. 5,994,624  
 U.S. Pat. No. 6,008,341  
 U.S. Pat. No. 6,288,214  
 U.S. Pat. No. 6,294,177  
 U.S. Pat. No. 6,651,655  
 U.S. Pat. No. 6,656,462  
 U.S. Pat. No. 6,733,754  
 U.S. Pat. No. 6,756,361  
 U.S. Pat. No. 6,770,278  
 U.S. Pat. No. 6,793,923  
 U.S. Pat. No. 6,814,971

U.S. Pat. No. 6,936,258  
 U.S. Patent Appln. 2002/0169288  
 U.S. Patent Appln. 2003/0153022  
 Abdallah et al., *Mol. Microbiol.*, 62, 667-679, 2006.  
 Abdallah et al., *Nat. Rev. Microbiol.*, 5, 883-891, 2007.  
 Albus et al., *Infect. Immun.*, 59:1008-1014, 1991.  
 An, *J. Virol.*, 71(3):2292-302, 1997.  
 Anavi, Sc. thesis from the department of Molecular Microbiology and Biotechnology of the Tel-Aviv University, Israel, 1998.  
 Andersén et al., *J. Immunol.*, 154, 3359-3372, 1995.  
 Angel et al., *Cell*, 49:729, 1987b.  
 Angel et al., *Mol. Cell. Biol.*, 7:2256, 1987a.  
 Archer, *Clin. Infect. Dis.*, 26, 1179-1181, 1998.  
 Atchison and Perry, *Cell*, 46:253, 1986.  
 Atchison and Perry, *Cell*, 48:121, 1987.  
 Ausubel et al., In: *Current Protocols in Molecular Biology*, John, Wiley & Sons, Inc, New York, 1996.  
 Baba et al., *J. Bacteriol.* 190:300-310, 2007.  
 Bae and Schneewind, *Plasmid*, 55:58-63, 2006.  
 Bae et al., *Proc. Natl. Acad. Sci. USA*, 101, 12312-12317, 2004.  
 Banerji et al., *Cell*, 27(2 Pt 1):299-308, 1981.  
 Banerji et al., *Cell*, 33(3):729-740, 1983.  
 Barany and Merrifield, In: *The Peptides*, Gross and Meienhofer (Eds.), Academic Press, NY, 1-284, 1979.  
 Behring E A. Über das Zustandekommen der Diphtherie—Immunität bei Thieren. *Deutsche Medizinische Wochenschrift*, 16:1145-8, 1890.  
 Bellus, *J. Macromol. Sci. Pure Appl. Chem.*, A31(1): 1355-1376, 1994.  
 Berkhout et al., *Cell*, 59:273-282, 1989.  
 Birch-Hirschfeld, L. 1934. Über die Agglutination von Staphylokokken durch Bestandteile des Säugetierblutplasmas. *Klinische Wochenschrift* 13:331.  
 Bjerketorp et al., *FEMS Microbiol. Lett.*, 234:309-314, 2004.  
 Blanar et al., *EMBO J.*, 8:1139, 1989.  
 Bodine and Ley, *EMBO J.*, 6:2997, 1987.  
 Borrebaeck, In: *Antibody Engineering—A Practical Guide*, W. H. Freeman and Co., 1992.  
 Boshart et al., *Cell*, 41:521, 1985.  
 Bosze et al., *EMBO J.*, 5(7):1615-1623, 1986.  
 Boucher and Corey. *Clin. Infect. Dis.* 46:S334-S349, 2008.  
 Braddock et al., *Cell*, 58:269, 1989.  
 Brown et al., *Biochemistry*, 37:4397-4406, 1998.  
 Bubeck Wardenburg and Schneewind. *J. Exp. Med.* 205: 287-294, 2008.  
 Bubeck-Wardenburg et al., *Infect. Immun.* 74:1040-1044, 2007.  
 Bubeck-Wardenburg et al., *Proc. Natl. Acad. Sci. USA*, 103:13831-13836, 2006.  
 Bulla and Siddiqui, *J. Virol.*, 62:1437, 1986.  
 Burke et al., *J. Inf. Dis.*, 170:1110-1119, 1994.  
 Burlak et al., *Cell Microbiol.*, 9:1172-1190, 2007.  
 Burts and Missiakas, *Mol. Microbiol.*, 69:736-46, 2008.  
 Burts et al., *Proc. Natl. Acad. Sci. USA*, 102:1169-1174, 2005.  
 Campbell and Villarreal, *Mol. Cell. Biol.*, 8:1993, 1988.  
 Campere and Tilghman, *Genes and Dev.*, 3:537, 1989.  
 Campo et al., *Nature*, 303:77, 1983.  
 Carbonelli et al., *FEMS Microbiol. Lett.*, 177(1):75-82, 1999.  
 Cedergren et al., *Protein Eng.*, 6:441-448, 1993.  
 Celander and Haseltine, *J. Virology*, 61:269, 1987.  
 Celander et al., *J. Virology*, 62:1314, 1988.  
 Cespedes et al., *J. Infect. Dis.*, 191(3):444-52, 2005.

Champion et al., *Science*, 313:1632-1636, 2006.  
 Chandler et al., *Cell*, 33:489, 1983.  
 Chandler et al., *Proc. Natl. Acad. Sci. USA*, 94(8):3596-601, 1997.  
 5 Chang et al., *Lancet.*, 362(9381):362-369, 2003.  
 Chang et al., *Mol. Cell. Biol.*, 9:2153, 1989.  
 Chatterjee et al., *Proc. Natl. Acad. Sci. USA*, 86:9114, 1989.  
 Chen and Okayama, *Mol. Cell Biol.*, 7(8):2745-2752, 1987.  
 Cheng et al., *FASEB J.*, 23:1-12, 2009.  
 10 Choi et al., *Cell*, 53:519, 1988.  
 Cocea, *Biotechniques*, 23(5):814-816, 1997.  
 Cohen et al., *J. Cell. Physiol.*, 5:75, 1987.  
 Cosgrove et al., *Infect. Control Hosp. Epidemiol.* 26:166-174, 2005.  
 15 Costa et al., *Mol. Cell. Biol.*, 8:81, 1988.  
 Cripe et al., *EMBO J.*, 6:3745, 1987.  
 Culotta and Hamer, *Mol. Cell. Biol.*, 9:1376, 1989.  
 Dalbey and Wickner, *J. Biol. Chem.*, 260:15925-15931, 1985.  
 20 Dandolo et al., *J. Virology*, 47:55-64, 1983.  
 De Villiers et al., *Nature*, 312(5991):242-246, 1984.  
 DeBord et al., *Infect. Immun.*, 74:4910-4914, 2006.  
 DeDent et al., *EMBO J.* 27:2656-2668, 2008.  
 25 DeDent et al., *J. Bacteriol.* 189:4473-4484, 2007.  
 Deisenhofer et al., *Hoppe-Seyh Zeitsch. Physiol. Chem.* 359:975-985, 1978.  
 Deisenhofer, *Biochemistry* 20:2361-2370, 1981.  
 Deschamps et al., *Science*, 230:1174-1177, 1985.  
 30 Devereux et al., *Nucl. Acid Res.*, 12:387-395, 1984.  
 Diep et al., *J. Infect. Dis.*, 193:1495-1503, 2006a.  
 Diep et al., *Lancet.*, 367:731-739, 2006b.  
 Dinges et al., *Clin. Microbiol. Rev.* 13:16-34, 2000.  
 Duthie and Lorenz, *J. Gen. Microbiol.*, 6:95-107, 1952.  
 35 Edbrooke et al., *Mol. Cell. Biol.*, 9:1908, 1989.  
 Edlund et al., *Science*, 230:912-916, 1985.  
 Ekstedt and Yotis, *Ann. N.Y. Acad. Sci.*, 80:496-500, 1960.  
 Emorl and Gaynes, *Clin. Microbiol. Rev.*, 6:428-442, 1993.  
 EP 0786519  
 EP 497524  
 EP 497525  
 Epitope Mapping Protocols In: *Methods in Molecular Biology*, Vol. 66, Morris (Ed.), 1996.  
 40 Fechtmeier, et al., *Proc Natl. Acad. Sci. USA*, 84:8463-8467, 1987.  
 45 Feng and Holland, *Nature*, 334:6178, 1988.  
 Field and Smith, *J. Comp. Pathol.*, 55:63, 1945.  
 Firak and Subramanian, *Mol. Cell. Biol.*, 6:3667, 1986.  
 Foeking and Hofstetter, *Gene*, 45(1):101-105, 1986.  
 50 Fortune et al., *Proc Natl. Acad. Sci. USA*, 102:10676-10681, 2005.  
 Foster, *Nat. Rev. Microbiol.*, 3:948-958, 2005.  
 Fournier et al., *Infect. Immun.*, 45:87-93, 1984.  
 Fraley et al., *Proc. Natl. Acad. Sci. USA*, 76:3348-3352, 1979.  
 55 Friedrich et al., *Nature*, 425:535-539, 2003.  
 Fujita et al., *Cell*, 49:357, 1987.  
 GB Appln. 2 202 328  
 Gilles et al., *Cell*, 33:717, 1983.  
 60 Gloss et al., *EMBO J.*, 6:3735, 1987.  
 Godbout et al., *Mol. Cell. Biol.*, 8:1169, 1988.  
 Gomez et al., *EMBO J.* 26:701-709, 2007.  
 Gomez et al., *J. Biol. Chem.* 281:20190-20196, 2006.  
 Gomez et al., *Nature Med.* 10:842-8, 2004.  
 65 Goodbourn and Maniatis, *Proc. Natl. Acad. Sci. USA*, 85:1447, 1988.  
 Goodbourn et al., *Cell*, 45:601, 1986.

Goodyear and Silverman, *J. Exp. Med.*, 197:1125-1139, 2003.

Goodyear and Silverman, *Proc. Nat. Acad. Sci. USA*, 101: 11392-11397, 2004.

Gopal, *Mol. Cell Biol.*, 5:1188-1190, 1985.

Gouda et al., *Biochemistry*, 31(40):9665-72, 1992.

Gouda et al., *Biochemistry*, 37:129-36, 1998.

Graham and Van Der Eb, *Virology*, 52:456-467, 1973.

Graille et al., *Proc. Nat. Acad. Sci. USA* 97:5399-5404, 2000.

Greene et al., *Immunology Today*, 10:272, 1989

Grosschedl and Baltimore, *Cell*, 41:885, 1985.

Guinn et al., *Mol. Microbiol.*, 51:359-370, 2004.

Guss et al., *Eur. J. Biochem.* 138:413-420, 1984.

Harland and Weintraub, *J. Cell Biol.*, 101(3):1094-1099, 1985.

Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., Chapter 8, 1988.

Hartleib et al., *Blood* 96:2149-2156, 2000.

Harvey et al., *Proc. Natl. Acad. Sci. USA*, 83:1084-1088, 1986.

Haslinger and Karin, *Proc. Natl. Acad. Sci. USA*, 82:8572, 1985.

Hauber and Cullen, *J. Virology*, 62:673, 1988.

Hen et al., *Nature*, 321:249, 1986.

Hensel et al., *Lymphokine Res.*, 8:347, 1989.

Herr and Clarke, *Cell*, 45:461, 1986.

Hirochika et al., *J. Virol.*, 61:2599, 1987.

Hirsch et al., *Mol. Cell Biol.*, 10:1959, 1990.

Holbrook et al., *Virology*, 157:211, 1987.

Horlick and Benfield, *Mol. Cell Biol.*, 9:2396, 1989.

Hsu et al., *Proc. Natl. Acad. Sci. USA*, 100:12420-12425, 2003.

Huang et al., *Cell*, 27:245, 1981.

Hug et al., *Mol. Cell Biol.*, 8:3065, 1988.

Huston et al., In: *Methods in Enzymology*, Langone (Ed.), Academic Press, NY, 203:46-88, 1991.

Hwang et al., *Mol. Cell Biol.*, 10:585, 1990.

Imagawa et al., *Cell*, 51:251, 1987.

Imbra and Karin, *Nature*, 323:555, 1986.

Imler et al., *Mol. Cell Biol.*, 7:2558, 1987.

Imperiale and Nevins, *Mol. Cell Biol.*, 4:875, 1984.

Innis et al., *Proc Natl Acad Sci USA*, 85(24):9436-9440, 1988.

Inouye and Inouye, *Nucleic Acids Res.*, 13: 3101-3109, 1985.

Jakovovits et al., *Mol. Cell Biol.*, 8:2555, 1988.

Jameel and Siddiqui, *Mol. Cell Biol.*, 6:710, 1986.

Jansson et al., *FEMS Immunol. Med. Microbiol.* 20:69-78 1998.

Jaynes et al., *Mol. Cell Biol.*, 8:62, 1988.

Jensen, *Acta Path. Microbiol. Scandim.* 44:421-428, 1958.

Johnson et al., *Methods in Enzymol.*, 203:88-99, 1991.

Johnson et al., *Mol. Cell Biol.*, 9:3393, 1989.

Jones, *Carb. Research*, 340:1097-1106, 2005.

Jonsson et al., *Oral Dis.*, 8(3):130-140, 2002.

Joyce et al., *Carbohydrate Research* 338:903-922 (2003

Kadesch and Berg, *Mol. Cell Biol.*, 6:2593, 1986.

Kaeppeler et al., *Plant Cell Rep.*, 8:415-418, 1990.

Kaneda et al., *Science*, 243:375-378, 1989.

Karin et al., *Mol. Cell Biol.*, 7:606, 1987.

Katinka et al., *Cell*, 20:393, 1980.

Kato et al., *J. Biol. Chem.*, 266:3361-3364, 1991.

Kawamoto et al., *Mol. Cell Biol.*, 8:267, 1988.

Kennedy et al., *Proc. Natl. Acad. Sci. USA* 105:1327-1332, 2008.

Kiledjian et al., *Mol. Cell Biol.*, 8:145, 1988.

Kinoshita, M., N. Kobayashi, S. Nagashima, M. Ishino, S. Otokozaawa, K. Mise, A. Sumi, H. Tsutsumi, N. Uehara, N. Watanabe, and M. Endo. 2008. Diversity of staphylocoagulase and identification of novel variants of staphylocoagulase gene in *Staphylococcus aureus*. *Microbiol. Immunol.s* 52:334-348.

Klamut et al., *Mol. Cell Biol.*, 10:193, 1990.

Klevens et al., *Clin. Infect. Dis.*, 2008; 47:927-30, 2008.

Klevens et al., *JAMA*, 298:1763-1771, 2007.

Koch et al., *Mol. Cell Biol.*, 9:303, 1989.

Kohler and Milstein, *Nature* 256:495-497 (1975

Kriegler and Botchan, In: *Eukaryotic Viral Vectors*, Gluzman (Ed.), Cold Spring Harbor: Cold Spring Harbor Laboratory, N Y, 1982.

Kriegler and Botchan, *Mol. Cell Biol.*, 3:325, 1983.

Kriegler et al., *Cell*, 38:483, 1984a.

Kriegler et al., *Cell*, 53:45, 1988.

Kriegler et al., In: *Cancer Cells 2/Oncogenes and Viral Genes*, Van de Woude et al. eds, Cold Spring Harbor, Cold Spring Harbor Laboratory, 1984b.

Kroh et al., *Proc. Natl. Acad. Sci. USA*, 106:7786-7791, 2009.

Kuhl et al., *Cell*, 50:1057, 1987.

Kuklin et al., *Infect. Immun.*, 74:2215-23, 2006.

Kunz et al., *Nucl. Acids Res.*, 17:1121, 1989.

Kyoda et al., *Lancet.*, 357:1225-1240, 2001.

Kyte and Doolittle, *J Mol. Biol.*, 157(1):105-132, 1982.

Lagergard et al., *Eur. J Clin. Microbiol. Infect. Dis.*, 11:341-5, 1992.

Lam et al., *J. Bacteriol.*, 86:87-91, 1963.

Larsen et al., *Proc Natl. Acad. Sci. USA.*, 83:8283, 1986, 1963.

Laspia et al., *Cell*, 59:283, 1989.

Latimer et al., *Mol. Cell Biol.*, 10:760, 1990.

Lee et al., *Nature*, 294:228, 1981.

Lee et al., *Nucleic Acids Res.*, 12:4191-206, 1984.

Lee, *Trends Microbiol.* 4(4):162-166, 1996.

Levenson et al., *Hum. Gene Ther.*, 9(8):1233-1236, 1998.

Levinson et al., *Nature*, 295:79, 1982.

Lin et al., *Mol. Cell Biol.*, 10:850, 1990.

Lowy, *New Engl. J. Med.*, 339:520-532, 1998.

Luria et al., *EMBO J.*, 6:3307, 1987.

Lusky and Botchan, *Proc. Natl. Acad. Sci. USA*, 83:3609, 1986.

Lusky et al., *Mol. Cell Biol.*, 3:1108, 1983.

Macejak and Sarnow, *Nature*, 353:90-94, 1991.

MacGurn et al., *Mol. Microbiol.*, 57:1653-1663, 2005.

Maira-Litran et al., *Infect. Immun.*, 70:4433-4440, 2002.

Maira-Litran et al., *Vaccine*, 22:872-879, 2004.

Majors and Varmus, *Proc. Natl. Acad. Sci. USA*, 80:5866, 1983.

Markwardt, *Untersuchungen über Hirudin. Naturwissenschaften*, 41:537-538, 1955.

Mazmanian et al., *Mol. Microbiol.* 40, 1049-1057, 2001.

Mazmanian et al., *Mol. Microbiol.*, 40(5):1049-1057, 2001.

Mazmanian et al., *Proc. Natl. Acad. Sci. USA*, 97:5510-5515, 2000.

Mazmanian et al., *Science*, 285(5428):760-3, 1999.

McLaughlin et al., *PLoS Pathog.*, 3:e105, 2007.

McNeall et al., *Gene*, 76:81, 1989.

Mernaugh et al., In: *Molecular Methods in Plant Pathology*, Singh et al. (Eds.), CRC Press Inc., Boca Raton, Fla., 359-365, 1995.

Merrifield, *Science*, 232(4748):341-347, 1986.

Miksicek et al., *Cell*, 46:203, 1986.

Mordacq and Linzer, *Genes and Dev.*, 3:760, 1989.

Moreau et al., *Carbohydrate Res.*, 201:285-297, 1990.  
 Moreau et al., *Nucl. Acids Res.*, 9:6047, 1981.  
 Moreillon et al., *Infect. Immun.*, 63:4738-4743, 1995.  
 Mosmann and Coffman, *Ann. Rev. Immunol.*, 7:145-173, 1989.  
 Muesing et al., *Cell*, 48:691, 1987.  
 Musher et al., *Medicine (Baltimore)*, 73:186-208, 1994.  
 Navarre and Schneewind, *J Biol. Chem.*, 274:15847-15856, 1999.  
 Needleman & Wunsch, *J. Mol. Biol.*, 48:443, 1970.  
 Ng et al., *Nuc. Acids Res.*, 17:601, 1989.  
 Nicolau and Sene, *Biochim. Biophys. Acta*, 721:185-190, 1982.  
 Nicolau et al., *Methods Enzymol.*, 149:157-176, 1987.  
 Novick, *Mol. Microbiol.*, 48:1429-1449, 2003.  
 O'Brien et al., *Mol. Microbiol.* 44:1033-1044, 2002.  
 O'Seaghda et al., *FEBS J.* 273:4831-4841, 2006.  
 Omirulleh et al., *Plant Mol. Biol.*, 21(3):415-28, 1993.  
 Ondek et al., *EMBO J.*, 6:1017, 1987.  
 Ornitz et al., *Mol. Cell. Biol.*, 7:3466, 1987.  
 Pallen, *Trends Microbiol.*, 10:209-212, 2002.  
 Palmiter et al., *Nature*, 300:611, 1982.  
 Palmqvist et al., *Microbes. Infect.*, 7:1501-11, 2005.  
 Panizzi et al., *J. Biol. Chem.*, 281:1179-1187, 2006.  
 PCT Appln. PCT/US89/01025  
 PCT Appln. WO 00/02523  
 PCT Appln. WO 00/12132  
 PCT Appln. WO 00/12689  
 PCT Appln. WO 00/15238  
 PCT Appln. WO 01/34809  
 PCT Appln. WO 01/60852  
 PCT Appln. WO 01/98499  
 PCT Appln. WO 02/059148  
 PCT Appln. WO 02/094868  
 PCT Appln. WO 03/53462  
 PCT Appln. WO 04/43407  
 PCT Appln. WO 06/032472  
 PCT Appln. WO 06/032475  
 PCT Appln. WO 06/032500  
 PCT Appln. WO 07/113222  
 PCT Appln. WO 07/113223  
 PCT Appln. WO 94/09699  
 PCT Appln. WO 95/06128  
 PCT Appln. WO 95/08348  
 PCT Appln. WO 98/57994  
 Pearson & Lipman, *Proc. Natl. Acad. Sci. USA*, 85:2444, 1988.  
 Pech et al., *Mol. Cell. Biol.*, 9:396, 1989.  
 Pelletier and Sonenberg, *Nature*, 334(6180):320-325, 1988.  
 Perez-Stable and Constantini, *Mol. Cell. Biol.*, 10:1116, 1990.  
 Phonimdaeng et al., *Mol. Microbiol.*, 4:393-404, 1990.  
 Picard and Schaffner, *Nature*, 307:83, 1984.  
 Pinkert et al., *Genes and Dev.*, 1:268, 1987.  
 Ponta et al., *Proc. Natl. Acad. Sci. USA*, 82:1020, 1985.  
 Porton et al., *Mol. Cell. Biol.*, 10:1076, 1990.  
 Potrykus et al., *Mol. Gen. Genet.*, 199(2):169-177, 1985.  
 Pugsley, *Microbiol. Rev.*, 57:50-108, 1993.  
 Pym et al., *Mol. Microbiol.*, 46: 709-717, 2002.  
 Pym et al., *Nat. Med.*, 9:533-539, 2003.  
 Queen and Baltimore, *Cell*, 35:741, 1983.  
 Quinn et al., *Mol. Cell. Biol.*, 9:4713, 1989.  
 Redondo et al., *Science*, 247:1225, 1990.  
 Reisman and Rotter, *Mol. Cell. Biol.*, 9:3571, 1989.  
 Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1289-1329, 1990.  
 Resendez Jr. et al., *Mol. Cell. Biol.*, 8:4579, 1988.

Ripe et al., *Mol. Cell. Biol.*, 9:2224, 1989.  
 Rippe, et al., *Mol. Cell Biol.*, 10:689-695, 1990.  
 Rittling et al., *Nuc. Acids Res.*, 17:1619, 1989.  
 Roben et al., *J. Immunol.* 154:6437-6445, 1995.  
 5 Rosen et al., *Cell*, 41:813, 1988.  
 Sakai et al., *Genes and Dev.*, 2:1144, 1988.  
 Salid-Salim et al., *Infect. Control Hosp. Epidemiol.* 24:451-455, 2003.  
 Sambrook et al., *In: Molecular cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001.  
 10 Schaffner et al., *J. Mol. Biol.*, 201:81, 1988.  
 Schneewind et al., *Cell* 70:267-281, 1992.  
 Schneewind et al., *EMBO*, 12:4803-4811, 1993.  
 Schneewind et al., *Science*, 268:103-6, 1995.  
 15 Searle et al., *Mol. Cell. Biol.*, 5:1480, 1985.  
 Sharp and Marciniak, *Cell*, 59:229, 1989.  
 Shaul and Ben-Levy, *EMBO J.*, 6:1913, 1987.  
 Shaw et al., *Microbiology*, 150:217-228, 2004.  
 Sheagren, *N. Engl. J. Med.* 310:1368-1373, 1984.  
 20 Sherman et al., *Mol. Cell. Biol.*, 9:50, 1989.  
 Shopsis et al., *J. Clin. Microbiol.*, 37:3556-63, 1999.  
 Sibbald et al., *Microbiol. Mol Biol. Rev.*, 70:755-788, 2006.  
 Silverman and Goodyear. *Nat. Rev. Immunol.*, 6:465-75, 2006.  
 25 Sjodahl, *Eur. J. Biochem.* 73:343-351, 1977.  
 Sjoquist et al., *Eur. J. Biochem.* 30:190-194, 1972.  
 Sleight and Lockett, *J. EMBO*, 4:3831, 1985.  
 Smith & Waterman, *Adv. Appl. Math.*, 2:482, 1981.  
 Smith et al., *Brit. J. Exp. Pathol.*, 28:57, 1947.  
 30 Sorensen et al., *Infect. Immun.*, 63:1710-1717, 1995.  
 Spalholz et al., *Cell*, 42:183, 1985.  
 Spandau and Lee, *J. Virology*, 62:427, 1988.  
 Spandidos and Wilkie, *EMBO J.*, 2:1193, 1983.  
 Stanley et al., *Proc. Natl. Acad. Sci. USA*, 100:13001-13006, 2003.  
 35 Stephens and Hentschel, *Biochem.* 1, 248:1, 1987.  
 Stewart and Young, In: *Solid Phase Peptide Synthesis*, 2d. ed., Pierce Chemical Co., 1984.  
 Stranger-Jones et al., *Proc. Nat. Acad. Sci. USA*, 103:16942-16947, 2006.  
 40 Stuart et al., *Nature*, 317:828, 1985.  
 Studier et al., *Methods Enzymol.* 185:60-89 1990.  
 Sullivan and Peterlin, *Mol. Cell. Biol.*, 7:3315, 1987.  
 Swartzendruber and Lehman, *J. Cell. Physiology*, 85:179, 1975.  
 45 Takebe et al., *Mol. Cell. Biol.*, 8:466, 1988.  
 Tam et al., *J. Am. Chem. Soc.*, 105:6442, 1983.  
 Tavernier et al., *Nature*, 301:634, 1983.  
 Taylor and Kingston, *Mol. Cell. Biol.*, 10:165, 1990a.  
 Taylor and Kingston, *Mol. Cell. Biol.*, 10:176, 1990b.  
 Taylor et al., *J. Biol. Chem.*, 264:15160, 1989.  
 Thiesen et al., *J. Virology*, 62:614, 1988.  
 Thomson et al., *J. Immunol.*, 157(2):822-826, 1996.  
 Tigges et al., *J. Immunol.*, 156(10):3901-3910, 1996.  
 55 Ton-That et al., *Proc. Natl. Acad. Sci. USA*, 96(22):12424-9, 1999.  
 Treisman, *Cell*, 42:889, 1985.  
 Tronche et al., *Mol. Biol. Med.*, 7:173, 1990.  
 Trudel and Constantini, *Genes and Dev.*, 6:954, 1987.  
 60 Tyndell et al., *Nuc. Acids. Res.*, 9:6231, 1981.  
 Uhlen et al., *J. Biol. Chem.* 259:1695-1702 and 13628 (Corr.) 1984.  
 van den Ent and Lowe, *FEBS Lett.*, 579:3837-3841, 2005.  
 van Wely et al., *FEMS Microbiol. Rev.*, 25:437-454, 2001.  
 Vannice and Levinson, *J. Virology*, 62:1305, 1988.  
 Vasseur et al., *Proc Natl. Acad. Sci. USA*, 77:1068, 1980.  
 Vaughan, et al., *Nat. Biotech.* 16; 535-539, 1998.

Wang and Calame, *Cell*, 47:241, 1986.  
 Weber et al., *Cell*, 36:983, 1984.  
 Weinberger et al. *Mol. Cell. Biol.*, 8:988, 1984.  
 Weiss et al., *J. Antimicrob. Chemother.*, 53(3):480-6, 2004.  
 Winoto and Baltimore, *Cell*, 59:649, 1989.

Wong et al., *Gene*, 10:87-94, 1980.  
 Xu et al., *J. Infect. Dis.*, 189:2323-2333, 2004.  
 Xu et al., *Mol. Microbiol.*, 66(3):787-800, 2007.  
 Yutzey et al. *Mol. Cell. Biol.*, 9:1397, 1989.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 64

<210> SEQ ID NO 1  
 <211> LENGTH: 150  
 <212> TYPE: DNA  
 <213> ORGANISM: *Staphylococcus* sp.

<400> SEQUENCE: 1

```
ttcaacaaag atcaacaaag cgccttctat gaaatcttga acatgcctaa cttaaacgaa    60
gcgcaacgta acggttcat tcaaagtctt aaagacgacc caagccaaag cactaatggt    120
ttaggtgaag ctaaaaaatt aaacgaatct                                     150
```

<210> SEQ ID NO 2  
 <211> LENGTH: 54  
 <212> TYPE: PRT  
 <213> ORGANISM: *Staphylococcus* sp.

<400> SEQUENCE: 2

```
Gln Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile
1          5          10          15
Leu Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln
20          25          30
Ser Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala
35          40          45
Lys Lys Leu Asn Glu Ser
50
```

<210> SEQ ID NO 3  
 <211> LENGTH: 51  
 <212> TYPE: PRT  
 <213> ORGANISM: *Staphylococcus* sp.

<400> SEQUENCE: 3

```
Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr Gln Val Leu Asn Met
1          5          10          15
Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys
20          25          30
Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys Leu
35          40          45
Asn Asp Ser
50
```

<210> SEQ ID NO 4  
 <211> LENGTH: 52  
 <212> TYPE: PRT  
 <213> ORGANISM: *Staphylococcus* sp.

<400> SEQUENCE: 4

```
Asn Asn Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn
1          5          10          15
Met Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
20          25          30
Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys
35          40          45
```

-continued

Leu Asn Glu Ser  
50

<210> SEQ ID NO 5  
<211> LENGTH: 52  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 5

Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His  
1 5 10 15

Leu Pro Asn Leu Thr Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu  
20 25 30

Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys  
35 40 45

Leu Asn Asp Ala  
50

<210> SEQ ID NO 6  
<211> LENGTH: 52  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 6

Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His  
1 5 10 15

Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu  
20 25 30

Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys  
35 40 45

Leu Asn Asp Ala  
50

<210> SEQ ID NO 7  
<211> LENGTH: 52  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (7)..(8)  
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (34)..(35)  
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 7

Asn Asn Phe Asn Lys Asp Xaa Xaa Ser Ala Phe Tyr Glu Ile Leu Asn  
1 5 10 15

Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu  
20 25 30

Lys Xaa Xaa Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys  
35 40 45

Leu Asn Glu Ser  
50

<210> SEQ ID NO 8  
<211> LENGTH: 52  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.  
<220> FEATURE:  
<221> NAME/KEY: MISC\_FEATURE

-continued

<222> LOCATION: (7)..(8)  
 <223> OTHER INFORMATION: where X is any amino acid other than Q  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (12)..(35)  
 <223> OTHER INFORMATION: where Y is any amion acid other than D

&lt;400&gt; SEQUENCE: 8

Asn Asn Phe Asn Lys Asp Xaa Xaa Ser Ala Phe Tyr Glu Ile Leu Asn  
 1 5 10 15  
 Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu  
 20 25 30  
 Lys Tyr Tyr Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys  
 35 40 45  
 Leu Asn Glu Ser  
 50

<210> SEQ ID NO 9  
 <211> LENGTH: 450  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 9

Met Lys Lys Lys Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile  
 1 5 10 15  
 Ala Ser Val Thr Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro  
 20 25 30  
 Ala Ala Asn Ala Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr  
 35 40 45  
 Gln Val Leu Asn Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe  
 50 55 60  
 Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly  
 65 70 75 80  
 Glu Ala Gln Lys Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln  
 85 90 95  
 Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu  
 100 105 110  
 Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser  
 115 120 125  
 Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys  
 130 135 140  
 Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys  
 145 150 155 160  
 Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn  
 165 170 175  
 Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser  
 180 185 190  
 Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln  
 195 200 205  
 Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe  
 210 215 220  
 Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly  
 225 230 235 240  
 Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu  
 245 250 255  
 Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Glu Glu Asp  
 260 265 270

-continued

Asn Lys Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp  
 275 280 285  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys Lys Pro Gly Lys Glu Asp  
 290 295 300  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp  
 305 310 315 320  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp  
 325 330 335  
 Gly Asn Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp  
 340 345 350  
 Gly Asn Gly Val His Val Val Lys Pro Gly Asp Thr Val Asn Asp Ile  
 355 360 365  
 Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys Ile Ala Ala Asp Asn Lys  
 370 375 380  
 Leu Ala Asp Lys Asn Met Ile Lys Pro Gly Gln Glu Leu Val Val Asp  
 385 390 395 400  
 Lys Lys Gln Pro Ala Asn His Ala Asp Ala Asn Lys Ala Gln Ala Leu  
 405 410 415  
 Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile Gly Thr Thr Val Phe Gly  
 420 425 430  
 Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu Leu Ala Gly Arg Arg Arg  
 435 440 445  
 Glu Leu  
 450

<210> SEQ ID NO 10  
 <211> LENGTH: 450  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 10

Met Lys Lys Lys Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile  
 1 5 10 15  
 Ala Ser Val Thr Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro  
 20 25 30  
 Ala Ala Asn Ala Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr  
 35 40 45  
 Gln Val Leu Asn Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe  
 50 55 60  
 Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly  
 65 70 75 80  
 Glu Ala Gln Lys Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln  
 85 90 95  
 Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu  
 100 105 110  
 Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser  
 115 120 125  
 Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys  
 130 135 140  
 Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys  
 145 150 155 160  
 Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn  
 165 170 175  
 Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser

-continued

180				185				190							
Gln	Ser	Ala	Asn	Leu	Leu	Ser	Glu	Ala	Lys	Lys	Leu	Asn	Glu	Ser	Gln
	195						200						205		
Ala	Pro	Lys	Ala	Asp	Asn	Lys	Phe	Asn	Lys	Glu	Gln	Gln	Asn	Ala	Phe
	210					215					220				
Tyr	Glu	Ile	Leu	His	Leu	Pro	Asn	Leu	Asn	Glu	Glu	Gln	Arg	Asn	Gly
	225				230					235					240
Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser	Val	Ser	Lys	Glu	Ile	Leu
				245					250					255	
Ala	Glu	Ala	Lys	Lys	Leu	Asn	Asp	Ala	Gln	Ala	Pro	Lys	Glu	Glu	Asp
			260					265					270		
Asn	Lys	Lys	Pro	Gly	Lys	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp
		275					280					285			
Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Lys	Lys	Pro	Gly	Lys	Glu	Asp
	290					295					300				
Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp
	305				310					315					320
Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp
			325						330					335	
Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp
			340					345					350		
Gly	Asn	Gly	Val	His	Val	Val	Lys	Pro	Gly	Asp	Thr	Val	Asn	Asp	Ile
		355					360					365			
Ala	Lys	Ala	Asn	Gly	Thr	Thr	Ala	Asp	Lys	Ile	Ala	Ala	Asp	Asn	Lys
	370					375					380				
Leu	Ala	Asp	Lys	Asn	Met	Ile	Lys	Pro	Gly	Gln	Glu	Leu	Val	Val	Asp
	385				390					395					400
Lys	Lys	Gln	Pro	Ala	Asn	His	Ala	Asp	Ala	Asn	Lys	Ala	Gln	Ala	Leu
			405						410					415	
Pro	Glu	Thr	Gly	Glu	Glu	Asn	Pro	Phe	Ile	Gly	Thr	Thr	Val	Phe	Gly
			420					425					430		
Gly	Leu	Ser	Leu	Ala	Leu	Gly	Ala	Ala	Leu	Leu	Ala	Gly	Arg	Arg	Arg
		435					440					445			
Glu	Leu														
	450														

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 97

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 11

Met	Ala	Met	Ile	Lys	Met	Ser	Pro	Glu	Glu	Ile	Arg	Ala	Lys	Ser	Gln
	1				5				10				15		
Ser	Tyr	Gly	Gln	Gly	Ser	Asp	Gln	Ile	Arg	Gln	Ile	Leu	Ser	Asp	Leu
			20						25				30		
Thr	Arg	Ala	Gln	Gly	Glu	Ile	Ala	Ala	Asn	Trp	Glu	Gly	Gln	Ala	Phe
		35					40					45			
Ser	Arg	Phe	Glu	Glu	Gln	Phe	Gln	Gln	Leu	Ser	Pro	Lys	Val	Glu	Lys
		50				55					60				
Phe	Ala	Gln	Leu	Leu	Glu	Glu	Ile	Lys	Gln	Gln	Leu	Asn	Ser	Thr	Ala
	65				70					75					80
Asp	Ala	Val	Gln	Glu	Gln	Asp	Gln	Gln	Leu	Ser	Asn	Asn	Phe	Gly	Leu
			85						90					95	

-continued

Gln

<210> SEQ ID NO 12  
 <211> LENGTH: 102  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 12

Met Gly Gly Tyr Lys Gly Ile Lys Ala Asp Gly Gly Lys Val Asn Gln  
 1 5 10 15  
 Ala Lys Gln Leu Ala Ala Lys Ile Ala Lys Asp Ile Glu Ala Cys Gln  
 20 25 30  
 Lys Gln Thr Gln Gln Leu Ala Glu Tyr Ile Glu Gly Ser Asp Trp Glu  
 35 40 45  
 Gly Gln Phe Ala Asn Lys Val Lys Asp Val Leu Leu Ile Met Ala Lys  
 50 55 60  
 Phe Gln Glu Glu Leu Val Gln Pro Met Ala Asp His Gln Lys Ala Ile  
 65 70 75 80  
 Asp Asn Leu Ser Gln Asn Leu Ala Lys Tyr Asp Thr Leu Ser Ile Lys  
 85 90 95  
 Gln Gly Leu Asp Arg Val  
 100

<210> SEQ ID NO 13  
 <211> LENGTH: 1385  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 13

Met Leu Asn Arg Glu Asn Lys Thr Ala Ile Thr Arg Lys Gly Met Val  
 1 5 10 15  
 Ser Asn Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Thr Val Gly Thr  
 20 25 30  
 Ala Ser Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Gly Asn Gln  
 35 40 45  
 Glu Ala Lys Ala Ala Glu Ser Thr Asn Lys Glu Leu Asn Glu Ala Thr  
 50 55 60  
 Thr Ser Ala Ser Asp Asn Gln Ser Ser Asp Lys Val Asp Met Gln Gln  
 65 70 75 80  
 Leu Asn Gln Glu Asp Asn Thr Lys Asn Asp Asn Gln Lys Glu Met Val  
 85 90 95  
 Ser Ser Gln Gly Asn Glu Thr Thr Ser Asn Gly Asn Lys Ser Ile Glu  
 100 105 110  
 Lys Glu Ser Val Gln Ser Thr Thr Gly Asn Lys Val Glu Val Ser Thr  
 115 120 125  
 Ala Lys Ser Asp Glu Gln Ala Ser Pro Lys Ser Thr Asn Glu Asp Leu  
 130 135 140  
 Asn Thr Lys Gln Thr Ile Ser Asn Gln Glu Gly Leu Gln Pro Asp Leu  
 145 150 155 160  
 Leu Glu Asn Lys Ser Val Val Asn Val Gln Pro Thr Asn Glu Glu Asn  
 165 170 175  
 Lys Lys Val Asp Ala Lys Thr Glu Ser Thr Thr Leu Asn Val Lys Ser  
 180 185 190  
 Asp Ala Ile Lys Ser Asn Ala Glu Thr Leu Val Asp Asn Asn Ser Asn  
 195 200 205  
 Ser Asn Asn Glu Asn Asn Ala Asp Ile Ile Leu Pro Lys Ser Thr Ala

-continued

210			215			220									
Pro	Lys	Ser	Leu	Asn	Thr	Arg	Met	Arg	Met	Ala	Ala	Ile	Gln	Pro	Asn
225					230					235					240
Ser	Thr	Asp	Ser	Lys	Asn	Val	Asn	Asp	Leu	Ile	Thr	Ser	Asn	Thr	Thr
				245					250						255
Leu	Thr	Val	Val	Asp	Ala	Asp	Asn	Ser	Lys	Thr	Ile	Val	Pro	Ala	Gln
				260					265						270
Asp	Tyr	Leu	Ser	Leu	Lys	Ser	Gln	Ile	Thr	Val	Asp	Asp	Lys	Val	Lys
		275					280						285		
Ser	Gly	Asp	Tyr	Phe	Thr	Ile	Lys	Tyr	Ser	Asp	Thr	Val	Gln	Val	Tyr
	290					295					300				
Gly	Leu	Asn	Pro	Glu	Asp	Ile	Lys	Asn	Ile	Gly	Asp	Ile	Lys	Asp	Pro
305					310					315					320
Asn	Asn	Gly	Glu	Thr	Ile	Ala	Thr	Ala	Lys	His	Asp	Thr	Ala	Asn	Asn
				325					330						335
Leu	Ile	Thr	Tyr	Thr	Phe	Thr	Asp	Tyr	Val	Asp	Arg	Phe	Asn	Ser	Val
			340					345						350	
Lys	Met	Gly	Ile	Asn	Tyr	Ser	Ile	Tyr	Met	Asp	Ala	Asp	Thr	Ile	Pro
		355					360					365			
Val	Asp	Lys	Lys	Asp	Val	Pro	Phe	Ser	Val	Thr	Ile	Gly	Asn	Gln	Ile
	370					375						380			
Thr	Thr	Thr	Thr	Ala	Asp	Ile	Thr	Tyr	Pro	Ala	Tyr	Lys	Glu	Ala	Asp
385					390					395					400
Asn	Asn	Ser	Ile	Gly	Ser	Ala	Phe	Thr	Glu	Thr	Val	Ser	His	Val	Gly
				405					410						415
Asn	Val	Glu	Asp	Pro	Gly	Tyr	Tyr	Asn	Gln	Val	Val	Tyr	Val	Asn	Pro
			420					425					430		
Met	Asp	Lys	Asp	Leu	Lys	Gly	Ala	Lys	Leu	Lys	Val	Glu	Ala	Tyr	His
		435					440					445			
Pro	Lys	Tyr	Pro	Thr	Asn	Ile	Gly	Gln	Ile	Asn	Gln	Asn	Val	Thr	Asn
	450					455					460				
Ile	Lys	Ile	Tyr	Arg	Val	Pro	Glu	Gly	Tyr	Thr	Leu	Asn	Lys	Gly	Tyr
465					470					475					480
Asp	Val	Asn	Thr	Asn	Asp	Leu	Val	Asp	Val	Thr	Asp	Glu	Phe	Lys	Asn
				485					490						495
Lys	Met	Thr	Tyr	Gly	Ser	Asn	Gln	Ser	Val	Asn	Leu	Asp	Phe	Gly	Asp
		500						505						510	
Ile	Thr	Ser	Ala	Tyr	Val	Val	Met	Val	Asn	Thr	Lys	Phe	Gln	Tyr	Thr
		515					520						525		
Asn	Ser	Glu	Ser	Pro	Thr	Leu	Val	Gln	Met	Ala	Thr	Leu	Ser	Ser	Thr
	530					535						540			
Gly	Asn	Lys	Ser	Val	Ser	Thr	Gly	Asn	Ala	Leu	Gly	Phe	Thr	Asn	Asn
545					550					555					560
Gln	Ser	Gly	Gly	Ala	Gly	Gln	Glu	Val	Tyr	Lys	Ile	Gly	Asn	Tyr	Val
				565					570						575
Trp	Glu	Asp	Thr	Asn	Lys	Asn	Gly	Val	Gln	Glu	Leu	Gly	Glu	Lys	Gly
			580						585					590	
Val	Gly	Asn	Val	Thr	Val	Thr	Val	Phe	Asp	Asn	Asn	Thr	Asn	Thr	Lys
		595						600					605		
Val	Gly	Glu	Ala	Val	Thr	Lys	Glu	Asp	Gly	Ser	Tyr	Leu	Ile	Pro	Asn
	610					615						620			
Leu	Pro	Asn	Gly	Asp	Tyr	Arg	Val	Glu	Phe	Ser	Asn	Leu	Pro	Lys	Gly
625					630					635					640



-continued

---

Thr	Thr	Lys	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	Arg	Phe	Asp	Asn	Leu
	1055					1060					1065			
Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu	Lys	Pro	Thr	Gly	Leu
	1070					1075					1080			
Thr	Gln	Thr	Gly	Thr	Asn	Thr	Thr	Glu	Asp	Asp	Lys	Asp	Ala	Asp
	1085					1090					1095			
Gly	Gly	Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp	Asp	Phe	Thr
	1100					1105					1110			
Leu	Asp	Asn	Gly	Tyr	Tyr	Glu	Glu	Glu	Thr	Ser	Asp	Ser	Asp	Ser
	1115					1120					1125			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1130					1135					1140			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1145					1150					1155			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1160					1165					1170			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1175					1180					1185			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1190					1195					1200			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1205					1210					1215			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1220					1225					1230			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1235					1240					1245			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1250					1255					1260			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1265					1270					1275			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1280					1285					1290			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1295					1300					1305			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1310					1315					1320			
Ser	Asp	Ala	Gly	Lys	His	Thr	Pro	Val	Lys	Pro	Met	Ser	Thr	Thr
	1325					1330					1335			
Lys	Asp	His	His	Asn	Lys	Ala	Lys	Ala	Leu	Pro	Glu	Thr	Gly	Asn
	1340					1345					1350			
Glu	Asn	Ser	Gly	Ser	Asn	Asn	Ala	Thr	Leu	Phe	Gly	Gly	Leu	Phe
	1355					1360					1365			
Ala	Ala	Leu	Gly	Ser	Leu	Leu	Leu	Phe	Gly	Arg	Arg	Lys	Lys	Gln
	1370					1375					1380			
Asn	Lys													
	1385													

&lt;210&gt; SEQ ID NO 14

&lt;211&gt; LENGTH: 1141

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 14

Met Ile Asn Arg Asp Asn Lys Lys Ala Ile Thr Lys Lys Gly Met Ile  
 1 5 10 15



-continued

---

Phe	Thr	Lys	Leu	Asp	Glu	Asn	Lys	Gln	Thr	Ile	Glu	Gln	Gln	Ile	Tyr
		435					440					445			
Val	Asn	Pro	Leu	Lys	Lys	Thr	Ala	Thr	Asn	Thr	Lys	Val	Asp	Ile	Ala
	450					455					460				
Gly	Ser	Gln	Val	Asp	Asp	Tyr	Gly	Asn	Ile	Lys	Leu	Gly	Asn	Gly	Ser
465				470						475					480
Thr	Ile	Ile	Asp	Gln	Asn	Thr	Glu	Ile	Lys	Val	Tyr	Lys	Val	Asn	Pro
				485					490					495	
Asn	Gln	Gln	Leu	Pro	Gln	Ser	Asn	Arg	Ile	Tyr	Asp	Phe	Ser	Gln	Tyr
			500					505					510		
Glu	Asp	Val	Thr	Ser	Gln	Phe	Asp	Asn	Lys	Lys	Ser	Phe	Ser	Asn	Asn
		515					520					525			
Val	Ala	Thr	Leu	Asp	Phe	Gly	Asp	Ile	Asn	Ser	Ala	Tyr	Ile	Ile	Lys
	530					535					540				
Val	Val	Ser	Lys	Tyr	Thr	Pro	Thr	Ser	Asp	Gly	Glu	Leu	Asp	Ile	Ala
545					550					555					560
Gln	Gly	Thr	Ser	Met	Arg	Thr	Thr	Asp	Lys	Tyr	Gly	Tyr	Tyr	Asn	Tyr
				565					570					575	
Ala	Gly	Tyr	Ser	Asn	Phe	Ile	Val	Thr	Ser	Asn	Asp	Thr	Gly	Gly	Gly
			580					585					590		
Asp	Gly	Thr	Val	Lys	Pro	Glu	Glu	Lys	Leu	Tyr	Lys	Ile	Gly	Asp	Tyr
		595					600					605			
Val	Trp	Glu	Asp	Val	Asp	Lys	Asp	Gly	Val	Gln	Gly	Thr	Asp	Ser	Lys
	610					615					620				
Glu	Lys	Pro	Met	Ala	Asn	Val	Leu	Val	Thr	Leu	Thr	Tyr	Pro	Asp	Gly
625					630					635					640
Thr	Thr	Lys	Ser	Val	Arg	Thr	Asp	Ala	Asn	Gly	His	Tyr	Glu	Phe	Gly
				645					650					655	
Gly	Leu	Lys	Asp	Gly	Glu	Thr	Tyr	Thr	Val	Lys	Phe	Glu	Thr	Pro	Ala
			660					665					670		
Gly	Tyr	Leu	Pro	Thr	Lys	Val	Asn	Gly	Thr	Thr	Asp	Gly	Glu	Lys	Asp
		675					680					685			
Ser	Asn	Gly	Ser	Ser	Ile	Thr	Val	Lys	Ile	Asn	Gly	Lys	Asp	Asp	Met
	690					695					700				
Ser	Leu	Asp	Thr	Gly	Phe	Tyr	Lys	Glu	Pro	Lys	Tyr	Asn	Leu	Gly	Asp
705					710					715					720
Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Ile	Gln	Asp	Ala	Asn	Glu
				725					730					735	
Pro	Gly	Ile	Lys	Asp	Val	Lys	Val	Thr	Leu	Lys	Asp	Ser	Thr	Gly	Lys
			740					745					750		
Val	Ile	Gly	Thr	Thr	Thr	Thr	Asp	Ala	Ser	Gly	Lys	Tyr	Lys	Phe	Thr
		755					760					765			
Asp	Leu	Asp	Asn	Gly	Asn	Tyr	Thr	Val	Glu	Phe	Glu	Thr	Pro	Ala	Gly
	770					775					780				
Tyr	Thr	Pro	Thr	Val	Lys	Asn	Thr	Thr	Ala	Glu	Asp	Lys	Asp	Ser	Asn
785					790					795					800
Gly	Leu	Thr	Thr	Thr	Gly	Val	Ile	Lys	Asp	Ala	Asp	Asn	Met	Thr	Leu
				805					810					815	
Asp	Ser	Gly	Phe	Tyr	Lys	Thr	Pro	Lys	Tyr	Ser	Leu	Gly	Asp	Tyr	Val
			820					825					830		
Trp	Tyr	Asp	Ser	Asn	Lys	Asp	Gly	Lys	Gln	Asp	Ser	Thr	Glu	Lys	Gly
		835					840					845			
Ile	Lys	Asp	Val	Lys	Val	Thr	Leu	Leu	Asn	Glu	Lys	Gly	Glu	Val	Ile

-continued

850					855					860					
Gly	Thr	Thr	Lys	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	Arg	Phe	Asp	Asn	Leu
865					870					875					880
Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu	Lys	Pro	Ala	Gly	Leu	Thr
				885					890					895	
Gln	Thr	Val	Thr	Asn	Thr	Thr	Glu	Asp	Asp	Lys	Asp	Ala	Asp	Gly	Gly
			900					905					910		
Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp	Asp	Phe	Thr	Leu	Asp	Asn
		915					920					925			
Gly	Tyr	Phe	Glu	Glu	Asp	Thr	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	930					935						940			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	945			950					955					960	
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			965					970						975	
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			980				985						990		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
		995				1000						1005			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1010					1015						1020			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Ser
	1025					1030						1035			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1040			1045					1050						
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Ser
	1055			1060					1065						
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Ala	Gly		
	1070			1075					1080						
Lys	His	Thr	Pro	Val	Lys	Pro	Met	Ser	Thr	Thr	Lys	Asp	His	His	
	1085					1090						1095			
Asn	Lys	Ala	Lys	Ala	Leu	Pro	Glu	Thr	Gly	Ser	Glu	Asn	Asn	Gly	
	1100					1105						1110			
Ser	Asn	Asn	Ala	Thr	Leu	Phe	Gly	Gly	Leu	Phe	Ala	Ala	Leu	Gly	
	1115					1120						1125			
Ser	Leu	Leu	Leu	Phe	Gly	Arg	Arg	Lys	Lys	Gln	Asn	Lys			
	1130					1135						1140			

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 350

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 15

Met	Thr	Lys	His	Tyr	Leu	Asn	Ser	Lys	Tyr	Gln	Ser	Glu	Gln	Arg	Ser
1				5					10					15	
Ser	Ala	Met	Lys	Lys	Ile	Thr	Met	Gly	Thr	Ala	Ser	Ile	Ile	Leu	Gly
		20						25					30		
Ser	Leu	Val	Tyr	Ile	Gly	Ala	Asp	Ser	Gln	Gln	Val	Asn	Ala	Ala	Thr
		35					40					45			
Glu	Ala	Thr	Asn	Ala	Thr	Asn	Asn	Gln	Ser	Thr	Gln	Val	Ser	Gln	Ala
	50					55					60				
Thr	Ser	Gln	Pro	Ile	Asn	Phe	Gln	Val	Gln	Lys	Asp	Gly	Ser	Ser	Glu
	65				70					75					80



-continued

---

Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu  
 115 120 125  
 Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser  
 130 135 140  
 Ala Pro Asn Ser Arg Pro Ile Asp Phe Glu Met Lys Lys Glu Asn Gly  
 145 150 155 160  
 Glu Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Lys Pro Ala Arg Val  
 165 170 175  
 Ile Phe Thr Asp Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly  
 180 185 190  
 Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro  
 195 200 205  
 Ile Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg  
 210 215 220  
 Phe Ser Val Ser Asn Gly Thr Lys Ala Val Lys Ile Val Ser Ser Thr  
 225 230 235 240  
 His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Glu Phe  
 245 250 255  
 Ala Gln Pro Ile Tyr Asn Ser Ala Asp Lys Phe Lys Thr Glu Glu Asp  
 260 265 270  
 Tyr Lys Ala Glu Lys Leu Leu Ala Pro Tyr Lys Lys Ala Lys Thr Leu  
 275 280 285  
 Glu Arg Gln Val Tyr Glu Leu Asn Lys Ile Gln Asp Lys Leu Pro Glu  
 290 295 300  
 Lys Leu Lys Ala Glu Tyr Lys Lys Lys Leu Glu Asp Thr Lys Lys Ala  
 305 310 315 320  
 Leu Asp Glu Gln Val Lys Ser Ala Ile Thr Glu Phe Gln Asn Val Gln  
 325 330 335  
 Pro Thr Asn Glu Lys Met Thr Asp Leu Gln Asp Thr Lys Tyr Val Val  
 340 345 350  
 Tyr Glu Ser Val Glu Asn Asn Glu Ser Met Met Asp Thr Phe Val Lys  
 355 360 365  
 His Pro Ile Lys Thr Gly Met Leu Asn Gly Lys Lys Tyr Met Val Met  
 370 375 380  
 Glu Thr Thr Asn Asp Asp Tyr Trp Lys Asp Phe Met Val Glu Gly Gln  
 385 390 395 400  
 Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile  
 405 410 415  
 Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys  
 420 425 430  
 Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile  
 435 440 445  
 Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys  
 450 455 460  
 Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr  
 465 470 475 480  
 Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln  
 485 490 495  
 Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu  
 500 505 510  
 Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys  
 515 520 525



-continued

---

Thr Ala Ala Lys Met Gln Asp Gln Thr Val Pro Gln Glu Ala Asn Ser  
 130 135 140  
 Gln Val Asp Asn Lys Thr Thr Asn Asp Ala Asn Ser Ile Ala Thr Asn  
 145 150 155 160  
 Ser Glu Leu Lys Asn Ser Gln Thr Leu Asp Leu Pro Gln Ser Ser Pro  
 165 170 175  
 Gln Thr Ile Ser Asn Ala Gln Gly Thr Ser Lys Pro Ser Val Arg Thr  
 180 185 190  
 Arg Ala Val Arg Ser Leu Ala Val Ala Glu Pro Val Val Asn Ala Ala  
 195 200 205  
 Asp Ala Lys Gly Thr Asn Val Asn Asp Lys Val Thr Ala Ser Asn Phe  
 210 215 220  
 Lys Leu Glu Lys Thr Thr Phe Asp Pro Asn Gln Ser Gly Asn Thr Phe  
 225 230 235 240  
 Met Ala Ala Asn Phe Thr Val Thr Asp Lys Val Lys Ser Gly Asp Tyr  
 245 250 255  
 Phe Thr Ala Lys Leu Pro Asp Ser Leu Thr Gly Asn Gly Asp Val Asp  
 260 265 270  
 Tyr Ser Asn Ser Asn Asn Thr Met Pro Ile Ala Asp Ile Lys Ser Thr  
 275 280 285  
 Asn Gly Asp Val Val Ala Lys Ala Thr Tyr Asp Ile Leu Thr Lys Thr  
 290 295 300  
 Tyr Thr Phe Val Phe Thr Asp Tyr Val Asn Asn Lys Glu Asn Ile Asn  
 305 310 315 320  
 Gly Gln Phe Ser Leu Pro Leu Phe Thr Asp Arg Ala Lys Ala Pro Lys  
 325 330 335  
 Ser Gly Thr Tyr Asp Ala Asn Ile Asn Ile Ala Asp Glu Met Phe Asn  
 340 345 350  
 Asn Lys Ile Thr Tyr Asn Tyr Ser Ser Pro Ile Ala Gly Ile Asp Lys  
 355 360 365  
 Pro Asn Gly Ala Asn Ile Ser Ser Gln Ile Ile Gly Val Asp Thr Ala  
 370 375 380  
 Ser Gly Gln Asn Thr Tyr Lys Gln Thr Val Phe Val Asn Pro Lys Gln  
 385 390 395 400  
 Arg Val Leu Gly Asn Thr Trp Val Tyr Ile Lys Gly Tyr Gln Asp Lys  
 405 410 415  
 Ile Glu Glu Ser Ser Gly Lys Val Ser Ala Thr Asp Thr Lys Leu Arg  
 420 425 430  
 Ile Phe Glu Val Asn Asp Thr Ser Lys Leu Ser Asp Ser Tyr Tyr Ala  
 435 440 445  
 Asp Pro Asn Asp Ser Asn Leu Lys Glu Val Thr Asp Gln Phe Lys Asn  
 450 455 460  
 Arg Ile Tyr Tyr Glu His Pro Asn Val Ala Ser Ile Lys Phe Gly Asp  
 465 470 475 480  
 Ile Thr Lys Thr Tyr Val Val Leu Val Glu Gly His Tyr Asp Asn Thr  
 485 490 495  
 Gly Lys Asn Leu Lys Thr Gln Val Ile Gln Glu Asn Val Asp Pro Val  
 500 505 510  
 Thr Asn Arg Asp Tyr Ser Ile Phe Gly Trp Asn Asn Glu Asn Val Val  
 515 520 525  
 Arg Tyr Gly Gly Gly Ser Ala Asp Gly Asp Ser Ala Val Asn Pro Lys  
 530 535 540  
 Asp Pro Thr Pro Gly Pro Pro Val Asp Pro Glu Pro Ser Pro Asp Pro



-continued

---

Ala Asn Asp Tyr Phe Asn Lys Pro Ala Lys Tyr Ile Lys Lys Asn Gly  
50 55 60

Lys Leu Tyr Val Gln Ile Thr Val Asn His Ser His Trp Ile Thr Gly  
65 70 75 80

Met Ser Ile Glu Gly His Lys Glu Asn Ile Ile Ser Lys Asn Thr Ala  
85 90 95

Lys Asp Glu Arg Thr Ser Glu Phe Glu Val Ser Lys Leu Asn Gly Lys  
100 105 110

Ile Asp Gly Lys Ile Asp Val Tyr Ile Asp Glu Lys Val Asn Gly Lys  
115 120 125

Pro Phe Lys Tyr Asp His His Tyr Asn Ile Thr Tyr Lys Phe Asn Gly  
130 135 140

Pro Thr Asp Val Ala Gly Ala Asn Ala Pro Gly Lys Asp Asp Lys Asn  
145 150 155 160

Ser Ala Ser Gly Ser Asp Lys Gly Ser Asp Gly Thr Thr Thr Gly Gln  
165 170 175

Ser Glu Ser Asn Ser Ser Asn Lys Asp Lys Val Glu Asn Pro Gln Thr  
180 185 190

Asn Ala Gly Thr Pro Ala Tyr Ile Tyr Ala Ile Pro Val Ala Ser Leu  
195 200 205

Ala Leu Leu Ile Ala Ile Thr Leu Phe Val Arg Lys Lys Ser Lys Gly  
210 215 220

Asn Val Glu  
225

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 635

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 20

Met Ala Lys Tyr Arg Gly Lys Pro Phe Gln Leu Tyr Val Lys Leu Ser  
1 5 10 15

Cys Ser Thr Met Met Ala Ser Ser Ile Ile Leu Thr Asn Ile Leu Pro  
20 25 30

Tyr Asp Ala Gln Ala Ala Ser Glu Lys Asp Thr Glu Ile Ser Lys Glu  
35 40 45

Ile Leu Ser Lys Gln Asp Leu Leu Asp Lys Val Asp Lys Ala Ile Arg  
50 55 60

Gln Ile Glu Gln Leu Lys Gln Leu Ser Ala Ser Ser Lys Ala His Tyr  
65 70 75 80

Lys Ala Gln Leu Asn Glu Ala Lys Thr Ala Ser Gln Ile Asp Glu Ile  
85 90 95

Ile Lys Arg Ala Asn Glu Leu Asp Ser Lys Glu Asn Lys Ser Ser His  
100 105 110

Thr Glu Met Asn Gly Gln Ser Asp Ile Asp Ser Lys Leu Asp Gln Leu  
115 120 125

Leu Lys Asp Leu Asn Glu Val Ser Ser Asn Val Asp Arg Gly Gln Gln  
130 135 140

Ser Gly Glu Asp Asp Leu Asn Ala Met Lys Asn Asp Met Ser Gln Thr  
145 150 155 160

Ala Thr Thr Lys Tyr Gly Glu Lys Asp Asp Lys Asn Asp Glu Ala Met  
165 170 175

Val Asn Lys Ala Leu Glu Asp Leu Asp His Leu Asn Gln Gln Ile His  
180 185 190

-continued

---

Lys Ser Lys Asp Ala Leu Lys Asp Ala Ser Lys Asp Pro Ala Val Ser  
 195 200 205  
 Thr Thr Asp Ser Asn His Glu Val Ala Lys Thr Pro Asn Asn Asp Gly  
 210 215 220  
 Ser Gly His Val Val Leu Asn Lys Phe Leu Ser Asn Glu Glu Asn Gln  
 225 230 235 240  
 Ser His Ser Asn Gln Leu Thr Asp Lys Leu Gln Gly Ser Asp Lys Ile  
 245 250 255  
 Asn His Ala Met Ile Glu Lys Leu Ala Lys Ser Asn Ala Ser Thr Gln  
 260 265 270  
 His Tyr Thr Tyr His Lys Leu Asn Thr Leu Gln Ser Leu Asp Gln Arg  
 275 280 285  
 Ile Ala Asn Thr Gln Leu Pro Lys Asn Gln Lys Ser Asp Leu Met Ser  
 290 295 300  
 Glu Val Asn Lys Thr Lys Glu Arg Ile Lys Ser Gln Arg Asn Ile Ile  
 305 310 315 320  
 Leu Glu Glu Leu Ala Arg Thr Asp Asp Lys Lys Tyr Ala Thr Gln Ser  
 325 330 335  
 Ile Leu Glu Ser Ile Phe Asn Lys Asp Glu Ala Asp Lys Ile Leu Lys  
 340 345 350  
 Asp Ile Arg Val Asp Gly Lys Thr Asp Gln Gln Ile Ala Asp Gln Ile  
 355 360 365  
 Thr Arg His Ile Asp Gln Leu Ser Leu Thr Thr Ser Asp Asp Leu Leu  
 370 375 380  
 Thr Ser Leu Ile Asp Gln Ser Gln Asp Lys Ser Leu Leu Ile Ser Gln  
 385 390 395 400  
 Ile Leu Gln Thr Lys Leu Gly Lys Ala Glu Ala Asp Lys Leu Ala Lys  
 405 410 415  
 Asp Trp Thr Asn Lys Gly Leu Ser Asn Arg Gln Ile Val Asp Gln Leu  
 420 425 430  
 Lys Lys His Phe Ala Ser Thr Gly Asp Thr Ser Ser Asp Asp Ile Leu  
 435 440 445  
 Lys Ala Ile Leu Asn Asn Ala Lys Asp Lys Lys Gln Ala Ile Glu Thr  
 450 455 460  
 Ile Leu Ala Thr Arg Ile Glu Arg Gln Lys Ala Lys Leu Leu Ala Asp  
 465 470 475 480  
 Leu Ile Thr Lys Ile Glu Thr Asp Gln Asn Lys Ile Phe Asn Leu Val  
 485 490 495  
 Lys Ser Ala Leu Asn Gly Lys Ala Asp Asp Leu Leu Asn Leu Gln Lys  
 500 505 510  
 Arg Leu Asn Gln Thr Lys Lys Asp Ile Asp Tyr Ile Leu Ser Pro Ile  
 515 520 525  
 Val Asn Arg Pro Ser Leu Leu Asp Arg Leu Asn Lys Asn Gly Lys Thr  
 530 535 540  
 Thr Asp Leu Asn Lys Leu Ala Asn Leu Met Asn Gln Gly Ser Asn Leu  
 545 550 555 560  
 Leu Asp Ser Ile Pro Asp Ile Pro Thr Pro Lys Pro Glu Lys Thr Leu  
 565 570 575  
 Thr Leu Gly Lys Gly Asn Gly Leu Leu Ser Gly Leu Leu Asn Ala Asp  
 580 585 590  
 Gly Asn Val Ser Leu Pro Lys Ala Gly Glu Thr Ile Lys Glu His Trp  
 595 600 605

-continued

---

Leu Pro Ile Ser Val Ile Val Gly Ala Met Gly Val Leu Met Ile Trp  
610 615 620

Leu Ser Arg Arg Asn Lys Leu Lys Asn Lys Ala  
625 630 635

<210> SEQ ID NO 21  
<211> LENGTH: 953  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 21

Met Asn Asn Lys Lys Thr Ala Thr Asn Arg Lys Gly Met Ile Pro Asn  
1 5 10 15  
Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Ser Val Gly Thr Ala Ser  
20 25 30  
Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Ser Gly His Glu Ala  
35 40 45  
Lys Ala Ala Glu His Thr Asn Gly Glu Leu Asn Gln Ser Lys Asn Glu  
50 55 60  
Thr Thr Ala Pro Ser Glu Asn Lys Thr Thr Glu Lys Val Asp Ser Arg  
65 70 75 80  
Gln Leu Lys Asp Asn Thr Gln Thr Ala Thr Ala Asp Gln Pro Lys Val  
85 90 95  
Thr Met Ser Asp Ser Ala Thr Val Lys Glu Thr Ser Ser Asn Met Gln  
100 105 110  
Ser Pro Gln Asn Ala Thr Ala Ser Gln Ser Thr Thr Gln Thr Ser Asn  
115 120 125  
Val Thr Thr Asn Asp Lys Ser Ser Thr Thr Tyr Ser Asn Glu Thr Asp  
130 135 140  
Lys Ser Asn Leu Thr Gln Ala Lys Asn Val Ser Thr Thr Pro Lys Thr  
145 150 155 160  
Thr Thr Ile Lys Gln Arg Ala Leu Asn Arg Met Ala Val Asn Thr Val  
165 170 175  
Ala Ala Pro Gln Gln Gly Thr Asn Val Asn Asp Lys Val His Phe Thr  
180 185 190  
Asn Ile Asp Ile Ala Ile Asp Lys Gly His Val Asn Lys Thr Thr Gly  
195 200 205  
Asn Thr Glu Phe Trp Ala Thr Ser Ser Asp Val Leu Lys Leu Lys Ala  
210 215 220  
Asn Tyr Thr Ile Asp Asp Ser Val Lys Glu Gly Asp Thr Phe Thr Phe  
225 230 235 240  
Lys Tyr Gly Gln Tyr Phe Arg Pro Gly Ser Val Arg Leu Pro Ser Gln  
245 250 255  
Thr Gln Asn Leu Tyr Asn Ala Gln Gly Asn Ile Ile Ala Lys Gly Ile  
260 265 270  
Tyr Asp Ser Lys Thr Asn Thr Thr Thr Tyr Thr Phe Thr Asn Tyr Val  
275 280 285  
Asp Gln Tyr Thr Asn Val Ser Gly Ser Phe Glu Gln Val Ala Phe Ala  
290 295 300  
Lys Arg Glu Asn Ala Thr Thr Asp Lys Thr Ala Tyr Lys Met Glu Val  
305 310 315 320  
Thr Leu Gly Asn Asp Thr Tyr Ser Lys Asp Val Ile Val Asp Tyr Gly  
325 330 335  
Asn Gln Lys Gly Gln Gln Leu Ile Ser Ser Thr Asn Tyr Ile Asn Asn  
340 345 350



-continued

---

Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 770 775 780

Asp Ser  
 785 790 795 800

Asp Ser  
 805 810 815

Asp Ser Asp Ser Asp Ser Asp Asn Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 820 825 830

Asp Ser  
 835 840 845

Asp Ser  
 850 855 860

Asp Ser  
 865 870 875 880

Asp Ser Asp Ala Gly Lys  
 885 890 895

His Thr Pro Thr Lys Pro Met Ser Thr Val Lys Asp Gln His Lys Thr  
 900 905 910

Ala Lys Ala Leu Pro Glu Thr Gly Ser Glu Asn Asn Asn Ser Asn Asn  
 915 920 925

Gly Thr Leu Phe Gly Gly Leu Phe Ala Ala Leu Gly Ser Leu Leu Leu  
 930 935 940

Phe Gly Arg Arg Lys Lys Gln Asn Lys  
 945 950

<210> SEQ ID NO 22  
 <211> LENGTH: 989  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 22

Met Asn Met Lys Lys Lys Glu Lys His Ala Ile Arg Lys Lys Ser Ile  
 1 5 10 15

Gly Val Ala Ser Val Leu Val Gly Thr Leu Ile Gly Phe Gly Leu Leu  
 20 25 30

Ser Ser Lys Glu Ala Asp Ala Ser Glu Asn Ser Val Thr Gln Ser Asp  
 35 40 45

Ser Ala Ser Asn Glu Ser Lys Ser Asn Asp Ser Ser Ser Val Ser Ala  
 50 55 60

Ala Pro Lys Thr Asp Asp Thr Asn Val Ser Asp Thr Lys Thr Ser Ser  
 65 70 75 80

Asn Thr Asn Asn Gly Glu Thr Ser Val Ala Gln Asn Pro Ala Gln Gln  
 85 90 95

Glu Thr Thr Gln Ser Ser Ser Thr Asn Ala Thr Thr Glu Glu Thr Pro  
 100 105 110

Val Thr Gly Glu Ala Thr Thr Thr Thr Thr Asn Gln Ala Asn Thr Pro  
 115 120 125

Ala Thr Thr Gln Ser Ser Asn Thr Asn Ala Glu Glu Leu Val Asn Gln  
 130 135 140

Thr Ser Asn Glu Thr Thr Ser Asn Asp Thr Asn Thr Val Ser Ser Val  
 145 150 155 160

Asn Ser Pro Gln Asn Ser Thr Asn Ala Glu Asn Val Ser Thr Thr Gln  
 165 170 175

Asp Thr Ser Thr Glu Ala Thr Pro Ser Asn Asn Glu Ser Ala Pro Gln  
 180 185 190

-continued

---

Asn Thr Asp Ala Ser Asn Lys Asp Val Val Ser Gln Ala Val Asn Pro  
 195 200 205  
 Ser Thr Pro Arg Met Arg Ala Phe Ser Leu Ala Ala Val Ala Ala Asp  
 210 215 220  
 Ala Pro Ala Ala Gly Thr Asp Ile Thr Asn Gln Leu Thr Asp Val Lys  
 225 230 235 240  
 Val Thr Ile Asp Ser Gly Thr Thr Val Tyr Pro His Gln Ala Gly Tyr  
 245 250 255  
 Val Lys Leu Asn Tyr Gly Phe Ser Val Pro Asn Ser Ala Val Lys Gly  
 260 265 270  
 Asp Thr Phe Lys Ile Thr Val Pro Lys Glu Leu Asn Leu Asn Gly Val  
 275 280 285  
 Thr Ser Thr Ala Lys Val Pro Pro Ile Met Ala Gly Asp Gln Val Leu  
 290 295 300  
 Ala Asn Gly Val Ile Asp Ser Asp Gly Asn Val Ile Tyr Thr Phe Thr  
 305 310 315 320  
 Asp Tyr Val Asp Asn Lys Glu Asn Val Thr Ala Asn Ile Thr Met Pro  
 325 330 335  
 Ala Tyr Ile Asp Pro Glu Asn Val Thr Lys Thr Gly Asn Val Thr Leu  
 340 345 350  
 Thr Thr Gly Ile Gly Thr Asn Thr Ala Ser Lys Thr Val Leu Ile Asp  
 355 360 365  
 Tyr Glu Lys Tyr Gly Gln Phe His Asn Leu Ser Ile Lys Gly Thr Ile  
 370 375 380  
 Asp Gln Ile Asp Lys Thr Asn Asn Thr Tyr Arg Gln Thr Ile Tyr Val  
 385 390 395 400  
 Asn Pro Ser Gly Asp Asn Val Val Leu Pro Ala Leu Thr Gly Asn Leu  
 405 410 415  
 Ile Pro Asn Thr Lys Ser Asn Ala Leu Ile Asp Ala Lys Asn Thr Asp  
 420 425 430  
 Ile Lys Val Tyr Arg Val Asp Asn Ala Asn Asp Leu Ser Glu Ser Tyr  
 435 440 445  
 Tyr Val Asn Pro Ser Asp Phe Glu Asp Val Thr Asn Gln Val Arg Ile  
 450 455 460  
 Ser Phe Pro Asn Ala Asn Gln Tyr Lys Val Glu Phe Pro Thr Asp Asp  
 465 470 475 480  
 Asp Gln Ile Thr Thr Pro Tyr Ile Val Val Val Asn Gly His Ile Asp  
 485 490 495  
 Pro Ala Ser Thr Gly Asp Leu Ala Leu Arg Ser Thr Phe Tyr Gly Tyr  
 500 505 510  
 Asp Ser Asn Phe Ile Trp Arg Ser Met Ser Trp Asp Asn Glu Val Ala  
 515 520 525  
 Phe Asn Asn Gly Ser Gly Ser Gly Asp Gly Ile Asp Lys Pro Val Val  
 530 535 540  
 Pro Glu Gln Pro Asp Glu Pro Gly Glu Ile Glu Pro Ile Pro Glu Asp  
 545 550 555 560  
 Ser Asp Ser Asp Pro Gly Ser Asp Ser Gly Ser Asp Ser Asn Ser Asp  
 565 570 575  
 Ser Gly Ser Asp Ser Gly Ser Asp Ser Thr Ser Asp Ser Gly Ser Asp  
 580 585 590  
 Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp  
 595 600 605

-continued

---

```

Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala
610                               615                               620

Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp
625                               630                               635                               640

Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp
645                               650                               655

Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Asp
660                               665                               670

Ser Asp Ser Asp
675                               680                               685

Ser Asp Ser Asp
690                               695                               700

Ser Asp Ser Asp
705                               710                               715                               720

Ser Asp Ser Asp
725                               730                               735

Ser Asp Ser Asp
740                               745                               750

Ser Asp Ser Asp
755                               760                               765

Ser Asp Ser Asp
770                               775                               780

Ser Asp Ser Asp
785                               790                               795                               800

Ser Asp Ser Asp
805                               810                               815

Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Asp Ser Asp Ser Glu
820                               825                               830

Ser Asp Ser Asp
835                               840                               845

Ser Asp Ser Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp
850                               855                               860

Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
865                               870                               875                               880

Ser Ala Ser Asp Ser Asp Ser Gly Ser Asp Ser Asp Ser Ser Ser Asp
885                               890                               895

Ser Asp Ser Asp Ser Thr Ser Asp Thr Gly Ser Asp Asn Asp Ser Asp
900                               905                               910

Ser Asp Ser Asn Ser Asp Ser Glu Ser Gly Ser Asn Asn Asn Val Val
915                               920                               925

Pro Pro Asn Ser Pro Lys Asn Gly Thr Asn Ala Ser Asn Lys Asn Glu
930                               935                               940

Ala Lys Asp Ser Lys Glu Pro Leu Pro Asp Thr Gly Ser Glu Asp Glu
945                               950                               955                               960

Ala Asn Thr Ser Leu Ile Trp Gly Leu Leu Ala Ser Leu Gly Ser Leu
965                               970                               975

Leu Leu Phe Arg Arg Lys Lys Glu Asn Lys Asp Lys Lys
980                               985

```

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 584

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 23

-continued

---

Met Lys Phe Lys Ser Leu Ile Thr Thr Thr Leu Ala Leu Gly Val Leu  
 1 5 10 15  
 Ala Ser Thr Gly Ala Asn Phe Asn Asn Asn Glu Ala Ser Ala Ala Ala  
 20 25 30  
 Lys Pro Leu Asp Lys Ser Ser Ser Ser Leu His His Gly Tyr Ser Lys  
 35 40 45  
 Val His Val Pro Tyr Ala Ile Thr Val Asn Gly Thr Ser Gln Asn Ile  
 50 55 60  
 Leu Ser Ser Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp  
 65 70 75 80  
 Leu Glu Asp Arg Val Lys Ser Val Leu Lys Ser Asp Arg Gly Ile Ser  
 85 90 95  
 Asp Ile Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Tyr Phe  
 100 105 110  
 Lys Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ala Gly Ile Tyr Thr  
 115 120 125  
 Ala Asp Leu Ile Asn Thr Ser Glu Ile Lys Ala Ile Asn Ile Asn Val  
 130 135 140  
 Asp Thr Lys Lys Gln Val Glu Asp Lys Lys Lys Asp Lys Ala Asn Tyr  
 145 150 155 160  
 Gln Val Pro Tyr Thr Ile Thr Val Asn Gly Thr Ser Gln Asn Ile Leu  
 165 170 175  
 Ser Asn Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp Leu  
 180 185 190  
 Glu Asp Lys Val Lys Ser Val Leu Glu Ser Asn Arg Gly Ile Thr Asp  
 195 200 205  
 Val Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Asn Phe Lys  
 210 215 220  
 Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ser Gly Ile Tyr Thr Ala  
 225 230 235 240  
 Asn Leu Ile Asn Ser Ser Asp Ile Lys Ser Ile Asn Ile Asn Val Asp  
 245 250 255  
 Thr Lys Lys His Ile Glu Asn Lys Ala Lys Arg Asn Tyr Gln Val Pro  
 260 265 270  
 Tyr Ser Ile Asn Leu Asn Gly Thr Ser Thr Asn Ile Leu Ser Asn Leu  
 275 280 285  
 Ser Phe Ser Asn Lys Pro Trp Thr Asn Tyr Lys Asn Leu Thr Ser Gln  
 290 295 300  
 Ile Lys Ser Val Leu Lys His Asp Arg Gly Ile Ser Glu Gln Asp Leu  
 305 310 315 320  
 Lys Tyr Ala Lys Lys Ala Tyr Tyr Thr Val Tyr Phe Lys Asn Gly Gly  
 325 330 335  
 Lys Arg Ile Leu Gln Leu Asn Ser Lys Asn Tyr Thr Ala Asn Leu Val  
 340 345 350  
 His Ala Lys Asp Val Lys Arg Ile Glu Ile Thr Val Lys Thr Gly Thr  
 355 360 365  
 Lys Ala Lys Ala Asp Arg Tyr Val Pro Tyr Thr Ile Ala Val Asn Gly  
 370 375 380  
 Thr Ser Thr Pro Ile Leu Ser Asp Leu Lys Phe Thr Gly Asp Pro Arg  
 385 390 395 400  
 Val Gly Tyr Lys Asp Ile Ser Lys Lys Val Lys Ser Val Leu Lys His  
 405 410 415

-continued

---

Asp Arg Gly Ile Gly Glu Arg Glu Leu Lys Tyr Ala Lys Lys Ala Thr  
                   420                                  425                                  430  
  
 Tyr Thr Val His Phe Lys Asn Gly Thr Lys Lys Val Ile Asn Ile Asn  
                   435                                  440                                  445  
  
 Ser Asn Ile Ser Gln Leu Asn Leu Leu Tyr Val Gln Asp Ile Lys Lys  
                   450                                  455                                  460  
  
 Ile Asp Ile Asp Val Lys Thr Gly Thr Lys Ala Lys Ala Asp Ser Tyr  
                   465                                  470                                  475                                  480  
  
 Val Pro Tyr Thr Ile Ala Val Asn Gly Thr Ser Thr Pro Ile Leu Ser  
                                   485                                  490                                  495  
  
 Lys Leu Lys Ile Ser Asn Lys Gln Leu Ile Ser Tyr Lys Tyr Leu Asn  
                                   500                                  505                                  510  
  
 Asp Lys Val Lys Ser Val Leu Lys Ser Glu Arg Gly Ile Ser Asp Leu  
                   515                                  520                                  525  
  
 Asp Leu Lys Phe Ala Lys Gln Ala Lys Tyr Thr Val Tyr Phe Lys Asn  
                   530                                  535                                  540  
  
 Gly Lys Lys Gln Val Val Asn Leu Lys Ser Asp Ile Phe Thr Pro Asn  
                   545                                  550                                  555                                  560  
  
 Leu Phe Ser Ala Lys Asp Ile Lys Lys Ile Asp Ile Asp Val Lys Gln  
                                   565                                  570                                  575  
  
 Tyr Thr Lys Ser Lys Lys Asn Lys  
                                   580

&lt;210&gt; SEQ ID NO 24

&lt;211&gt; LENGTH: 10419

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 24

Met Asn Tyr Arg Asp Lys Ile Gln Lys Phe Ser Ile Arg Lys Tyr Thr  
 1                  5                                  10                                  15  
  
 Val Gly Thr Phe Ser Thr Val Ile Ala Thr Leu Val Phe Leu Gly Phe  
                   20                                  25                                  30  
  
 Asn Thr Ser Gln Ala His Ala Ala Glu Thr Asn Gln Pro Ala Ser Val  
                   35                                  40                                  45  
  
 Val Lys Gln Lys Gln Gln Ser Asn Asn Glu Gln Thr Glu Asn Arg Glu  
                   50                                  55                                  60  
  
 Ser Gln Val Gln Asn Ser Gln Asn Ser Gln Asn Gly Gln Ser Leu Ser  
                   65                                  70                                  75                                  80  
  
 Ala Thr His Glu Asn Glu Gln Pro Asn Ile Ser Gln Ala Asn Leu Val  
                                   85                                  90                                  95  
  
 Asp Gln Lys Val Ala Gln Ser Ser Thr Thr Asn Asp Glu Gln Pro Ala  
                   100                                  105                                  110  
  
 Ser Gln Asn Val Asn Thr Lys Lys Asp Ser Ala Thr Ala Ala Thr Thr  
                   115                                  120                                  125  
  
 Gln Pro Asp Lys Glu Gln Ser Lys His Lys Gln Asn Glu Ser Gln Ser  
                   130                                  135                                  140  
  
 Ala Asn Lys Asn Gly Asn Asp Asn Arg Ala Ala His Val Glu Asn His  
                   145                                  150                                  155                                  160  
  
 Glu Ala Asn Val Val Thr Ala Ser Asp Ser Ser Asp Asn Gly Asn Val  
                                   165                                  170                                  175  
  
 Gln His Asp Arg Asn Glu Leu Gln Ala Phe Phe Asp Ala Asn Tyr His  
                   180                                  185                                  190  
  
 Asp Tyr Arg Phe Ile Asp Arg Glu Asn Ala Asp Ser Gly Thr Phe Asn  
                   195                                  200                                  205

-continued

---

Tyr Val Lys Gly Ile Phe Asp Lys Ile Asn Thr Leu Leu Gly Ser Asn  
 210 215 220  
 Asp Pro Ile Asn Asn Lys Asp Leu Gln Leu Ala Tyr Lys Glu Leu Glu  
 225 230 235 240  
 Gln Ala Val Ala Leu Ile Arg Thr Met Pro Gln Arg Gln Gln Thr Ser  
 245 250 255  
 Arg Arg Ser Asn Arg Ile Gln Thr Arg Ser Val Glu Ser Arg Ala Ala  
 260 265 270  
 Glu Pro Arg Ser Val Ser Asp Tyr Gln Asn Ala Asn Ser Ser Tyr Tyr  
 275 280 285  
 Val Glu Asn Ala Asn Asp Gly Ser Gly Tyr Pro Val Gly Thr Tyr Ile  
 290 295 300  
 Asn Ala Ser Ser Lys Gly Ala Pro Tyr Asn Leu Pro Thr Thr Pro Trp  
 305 310 315 320  
 Asn Thr Leu Lys Ala Ser Asp Ser Lys Glu Ile Ala Leu Met Thr Ala  
 325 330 335  
 Lys Gln Thr Gly Asp Gly Tyr Gln Trp Val Ile Lys Phe Asn Lys Gly  
 340 345 350  
 His Ala Pro His Gln Asn Met Ile Phe Trp Phe Ala Leu Pro Ala Asp  
 355 360 365  
 Gln Val Pro Val Gly Arg Thr Asp Phe Val Thr Val Asn Ser Asp Gly  
 370 375 380  
 Thr Asn Val Gln Trp Ser His Gly Ala Gly Ala Gly Ala Asn Lys Pro  
 385 390 395 400  
 Leu Gln Gln Met Trp Glu Tyr Gly Val Asn Asp Pro His Arg Ser His  
 405 410 415  
 Asp Phe Lys Ile Arg Asn Arg Ser Gly Gln Val Ile Tyr Asp Trp Pro  
 420 425 430  
 Thr Val His Ile Tyr Ser Leu Glu Asp Leu Ser Arg Ala Ser Asp Tyr  
 435 440 445  
 Phe Ser Glu Ala Gly Ala Thr Pro Ala Thr Lys Ala Phe Gly Arg Gln  
 450 455 460  
 Asn Phe Glu Tyr Ile Asn Gly Gln Lys Pro Ala Glu Ser Pro Gly Val  
 465 470 475 480  
 Pro Lys Val Tyr Thr Phe Ile Gly Gln Gly Asp Ala Ser Tyr Thr Ile  
 485 490 495  
 Ser Phe Lys Thr Gln Gly Pro Thr Val Asn Lys Leu Tyr Tyr Ala Ala  
 500 505 510  
 Gly Gly Arg Ala Leu Glu Tyr Asn Gln Leu Phe Met Tyr Ser Gln Leu  
 515 520 525  
 Tyr Val Glu Ser Thr Gln Asp His Gln Gln Arg Leu Asn Gly Leu Arg  
 530 535 540  
 Gln Val Val Asn Arg Thr Tyr Arg Ile Gly Thr Thr Lys Arg Val Glu  
 545 550 555 560  
 Val Ser Gln Gly Asn Val Gln Thr Lys Lys Val Leu Glu Ser Thr Asn  
 565 570 575  
 Leu Asn Ile Asp Asp Phe Val Asp Asp Pro Leu Ser Tyr Val Lys Thr  
 580 585 590  
 Pro Ser Asn Lys Val Leu Gly Phe Tyr Ser Asn Asn Ala Asn Thr Asn  
 595 600 605  
 Ala Phe Arg Pro Gly Gly Ala Gln Gln Leu Asn Glu Tyr Gln Leu Ser  
 610 615 620

-continued

---

Gln Leu Phe Thr Asp Gln Lys Leu Gln Glu Ala Ala Arg Thr Arg Asn  
 625 630 635 640  
 Pro Ile Arg Leu Met Ile Gly Phe Asp Tyr Pro Asp Ala Tyr Gly Asn  
 645 650 655  
 Ser Glu Thr Leu Val Pro Val Asn Leu Thr Val Leu Pro Glu Ile Gln  
 660 665 670  
 His Asn Ile Lys Phe Phe Lys Asn Asp Asp Thr Gln Asn Ile Ala Glu  
 675 680 685  
 Lys Pro Phe Ser Lys Gln Ala Gly His Pro Val Phe Tyr Val Tyr Ala  
 690 695 700  
 Gly Asn Gln Gly Asn Ala Ser Val Asn Leu Gly Gly Ser Val Thr Ser  
 705 710 715 720  
 Ile Gln Pro Leu Arg Ile Asn Leu Thr Ser Asn Glu Asn Phe Thr Asp  
 725 730 735  
 Lys Asp Trp Gln Ile Thr Gly Ile Pro Arg Thr Leu His Ile Glu Asn  
 740 745 750  
 Ser Thr Asn Arg Pro Asn Asn Ala Arg Glu Arg Asn Ile Glu Leu Val  
 755 760 765  
 Gly Asn Leu Leu Pro Gly Asp Tyr Phe Gly Thr Ile Arg Phe Gly Arg  
 770 775 780  
 Lys Glu Gln Leu Phe Glu Ile Arg Val Lys Pro His Thr Pro Thr Ile  
 785 790 795 800  
 Thr Thr Thr Ala Glu Gln Leu Arg Gly Thr Ala Leu Gln Lys Val Pro  
 805 810 815  
 Val Asn Ile Ser Gly Ile Pro Leu Asp Pro Ser Ala Leu Val Tyr Leu  
 820 825 830  
 Val Ala Pro Thr Asn Gln Thr Thr Asn Gly Gly Ser Glu Ala Asp Gln  
 835 840 845  
 Ile Pro Ser Gly Tyr Thr Ile Leu Ala Thr Gly Thr Pro Asp Gly Val  
 850 855 860  
 His Asn Thr Ile Thr Ile Arg Pro Gln Asp Tyr Val Val Phe Ile Pro  
 865 870 875 880  
 Pro Val Gly Lys Gln Ile Arg Ala Val Val Tyr Tyr Asn Lys Val Val  
 885 890 895  
 Ala Ser Asn Met Ser Asn Ala Val Thr Ile Leu Pro Asp Asp Ile Pro  
 900 905 910  
 Pro Thr Ile Asn Asn Pro Val Gly Ile Asn Ala Lys Tyr Tyr Arg Gly  
 915 920 925  
 Asp Glu Val Asn Phe Thr Met Gly Val Ser Asp Arg His Ser Gly Ile  
 930 935 940  
 Lys Asn Thr Thr Ile Thr Thr Leu Pro Asn Gly Trp Thr Ser Asn Leu  
 945 950 955 960  
 Thr Lys Ala Asp Lys Asn Asn Gly Ser Leu Ser Ile Thr Gly Arg Val  
 965 970 975  
 Ser Met Asn Gln Ala Phe Asn Ser Asp Ile Thr Phe Lys Val Ser Ala  
 980 985 990  
 Thr Asp Asn Val Asn Asn Thr Thr Asn Asp Ser Gln Ser Lys His Val  
 995 1000 1005  
 Ser Ile His Val Gly Lys Ile Ser Glu Asp Ala His Pro Ile Val  
 1010 1015 1020  
 Leu Gly Asn Thr Glu Lys Val Val Val Val Asn Pro Thr Ala Val  
 1025 1030 1035  
 Ser Asn Asp Glu Lys Gln Ser Ile Ile Thr Ala Phe Met Asn Lys

-continued

---

1040	1045	1050
Asn Gln Asn Ile Arg Gly Tyr Leu Ala Ser Thr Asp Pro Val Thr 1055 1060 1065		
Val Asp Asn Asn Gly Asn Val Thr Leu His Tyr Arg Asp Gly Ser 1070 1075 1080		
Ser Thr Thr Leu Asp Ala Thr Asn Val Met Thr Tyr Glu Pro Val 1085 1090 1095		
Val Lys Pro Glu Tyr Gln Thr Val Asn Ala Ala Lys Thr Ala Thr 1100 1105 1110		
Val Thr Ile Ala Lys Gly Gln Ser Phe Ser Ile Gly Asp Ile Lys 1115 1120 1125		
Gln Tyr Phe Thr Leu Ser Asn Gly Gln Pro Ile Pro Ser Gly Thr 1130 1135 1140		
Phe Thr Asn Ile Thr Ser Asp Arg Thr Ile Pro Thr Ala Gln Glu 1145 1150 1155		
Val Ser Gln Met Asn Ala Gly Thr Gln Leu Tyr His Ile Thr Ala 1160 1165 1170		
Thr Asn Ala Tyr His Lys Asp Ser Glu Asp Phe Tyr Ile Ser Leu 1175 1180 1185		
Lys Ile Ile Asp Val Lys Gln Pro Glu Gly Asp Gln Arg Val Tyr 1190 1195 1200		
Arg Thr Ser Thr Tyr Asp Leu Thr Thr Asp Glu Ile Ser Lys Val 1205 1210 1215		
Lys Gln Ala Phe Ile Asn Ala Asn Arg Asp Val Ile Thr Leu Ala 1220 1225 1230		
Glu Gly Asp Ile Ser Val Thr Asn Thr Pro Asn Gly Ala Asn Val 1235 1240 1245		
Ser Thr Ile Thr Val Asn Ile Asn Lys Gly Arg Leu Thr Lys Ser 1250 1255 1260		
Phe Ala Ser Asn Leu Ala Asn Met Asn Phe Leu Arg Trp Val Asn 1265 1270 1275		
Phe Pro Gln Asp Tyr Thr Val Thr Trp Thr Asn Ala Lys Ile Ala 1280 1285 1290		
Asn Arg Pro Thr Asp Gly Gly Leu Ser Trp Ser Asp Asp His Lys 1295 1300 1305		
Ser Leu Ile Tyr Arg Tyr Asp Ala Thr Leu Gly Thr Gln Ile Thr 1310 1315 1320		
Thr Asn Asp Ile Leu Thr Met Leu Lys Ala Thr Thr Thr Val Pro 1325 1330 1335		
Gly Leu Arg Asn Asn Ile Thr Gly Asn Glu Lys Ser Gln Ala Glu 1340 1345 1350		
Ala Gly Gly Arg Pro Asn Phe Arg Thr Thr Gly Tyr Ser Gln Ser 1355 1360 1365		
Asn Ala Thr Thr Asp Gly Gln Arg Gln Phe Thr Leu Asn Gly Gln 1370 1375 1380		
Val Ile Gln Val Leu Asp Ile Ile Asn Pro Ser Asn Gly Tyr Gly 1385 1390 1395		
Gly Gln Pro Val Thr Asn Ser Asn Thr Arg Ala Asn His Ser Asn 1400 1405 1410		
Ser Thr Val Val Asn Val Asn Glu Pro Ala Ala Asn Gly Ala Gly 1415 1420 1425		
Ala Phe Thr Ile Asp His Val Val Lys Ser Asn Ser Thr His Asn 1430 1435 1440		

-continued

---

Ala Ser Asp Ala Val Tyr Lys	Ala Gln Leu Tyr Leu Thr Pro Tyr
1445	1450 1455
Gly Pro Lys Gln Tyr Val Glu	His Leu Asn Gln Asn Thr Gly Asn
1460	1465 1470
Thr Thr Asp Ala Ile Asn Ile	Tyr Phe Val Pro Ser Asp Leu Val
1475	1480 1485
Asn Pro Thr Ile Ser Val Gly	Asn Tyr Thr Asn His Gln Val Phe
1490	1495 1500
Ser Gly Glu Thr Phe Thr Asn	Thr Ile Thr Ala Asn Asp Asn Phe
1505	1510 1515
Gly Val Gln Ser Val Thr Val	Pro Asn Thr Ser Gln Ile Thr Gly
1520	1525 1530
Thr Val Asp Asn Asn His Gln	His Val Ser Ala Thr Ala Pro Asn
1535	1540 1545
Val Thr Ser Ala Thr Asn Lys	Thr Ile Asn Leu Leu Ala Thr Asp
1550	1555 1560
Thr Ser Gly Asn Thr Ala Thr	Thr Ser Phe Asn Val Thr Val Lys
1565	1570 1575
Pro Leu Arg Asp Lys Tyr Arg	Val Gly Thr Ser Ser Thr Ala Ala
1580	1585 1590
Asn Pro Val Arg Ile Ala Asn	Ile Ser Asn Asn Ala Thr Val Ser
1595	1600 1605
Gln Ala Asp Gln Thr Thr Ile	Ile Asn Ser Leu Thr Phe Thr Glu
1610	1615 1620
Thr Val Pro Asn Arg Ser Tyr	Ala Arg Ala Ser Ala Asn Glu Ile
1625	1630 1635
Thr Ser Lys Thr Val Ser Asn	Val Ser Arg Thr Gly Asn Asn Ala
1640	1645 1650
Asn Val Thr Val Thr Val Thr	Tyr Gln Asp Gly Thr Thr Ser Thr
1655	1660 1665
Val Thr Val Pro Val Lys His	Val Ile Pro Glu Ile Val Ala His
1670	1675 1680
Ser His Tyr Thr Val Gln Gly	Gln Asp Phe Pro Ala Gly Asn Gly
1685	1690 1695
Ser Ser Ala Ser Asp Tyr Phe	Lys Leu Ser Asn Gly Ser Asp Ile
1700	1705 1710
Ala Asp Ala Thr Ile Thr Trp	Val Ser Gly Gln Ala Pro Asn Lys
1715	1720 1725
Asp Asn Thr Arg Ile Gly Glu	Asp Ile Thr Val Thr Ala His Ile
1730	1735 1740
Leu Ile Asp Gly Glu Thr Thr	Pro Ile Thr Lys Thr Ala Thr Tyr
1745	1750 1755
Lys Val Val Arg Thr Val Pro	Lys His Val Phe Glu Thr Ala Arg
1760	1765 1770
Gly Val Leu Tyr Pro Gly Val	Ser Asp Met Tyr Asp Ala Lys Gln
1775	1780 1785
Tyr Val Lys Pro Val Asn Asn	Ser Trp Ser Thr Asn Ala Gln His
1790	1795 1800
Met Asn Phe Gln Phe Val Gly	Thr Tyr Gly Pro Asn Lys Asp Val
1805	1810 1815
Val Gly Ile Ser Thr Arg Leu	Ile Arg Val Thr Tyr Asp Asn Arg
1820	1825 1830

-continued

---

Gln Thr 1835	Glu Asp 1835	Leu Thr 1840	Ile 1840	Leu Ser 1845	Lys Val 1845	Lys 1845	Pro Asp 1845	Pro
Pro Arg 1850	Ile Asp 1850	Ala Asn 1855	Ser 1855	Val Thr 1860	Tyr Lys 1860	Ala 1860	Gly Leu 1860	Thr
Asn Gln 1865	Glu Ile 1870	Lys Val 1870	Asn 1870	Asn Val 1875	Leu Asn 1875	Asn 1875	Ser Ser 1875	Val
Lys Leu 1880	Phe Lys 1885	Ala Asp 1885	Asn 1885	Thr Pro 1890	Leu Asn 1890	Val 1890	Thr Asn 1890	Ile
Thr His 1895	Gly Ser 1900	Gly Phe 1900	Ser 1900	Ser Val 1905	Val Thr 1905	Val 1905	Ser Asp 1905	Ala
Leu Pro 1910	Asn Gly 1915	Gly Ile 1915	Lys 1915	Ala Lys 1920	Ser Ser 1920	Ile 1920	Ser Met 1920	Asn
Asn Val 1925	Thr Tyr 1930	Thr Thr 1930	Gln 1930	Asp Glu 1935	His Gly 1935	Gln 1935	Val Val 1935	Thr
Val Thr 1940	Arg Asn 1945	Glu Ser 1945	Val 1945	Asp Ser 1950	Asn Asp 1950	Ser 1950	Ala Thr 1950	Val
Thr Val 1955	Thr Pro 1960	Gln Leu 1960	Gln 1960	Ala Thr 1965	Thr Glu 1965	Gly 1965	Ala Val 1965	Phe
Ile Lys 1970	Gly Gly 1975	Asp Gly 1975	Phe 1975	Asp Phe 1980	Gly His 1980	Val 1980	Glu Arg 1980	Phe
Ile Gln 1985	Asn Pro 1990	Pro His 1990	Gly 1990	Ala Thr 1995	Val Ala 1995	Trp 1995	His Asp 1995	Ser
Pro Asp 2000	Thr Trp 2005	Lys Asn 2005	Thr 2005	Val Gly 2010	Asn Thr 2010	His 2010	Lys Thr 2010	Ala
Val Val 2015	Thr Leu 2020	Pro Asn 2020	Gly 2020	Gln Gly 2025	Thr Arg 2025	Asn 2025	Val Glu 2025	Val
Pro Val 2030	Lys Val 2035	Tyr Pro 2035	Val 2035	Ala Asn 2040	Ala Lys 2040	Ala 2040	Pro Ser 2040	Arg
Asp Val 2045	Lys Gly 2050	Gln Asn 2050	Leu 2050	Thr Asn 2055	Gly Thr 2055	Asp 2055	Ala Met 2055	Asn
Tyr Ile 2060	Thr Phe 2065	Asp Pro 2065	Asn 2065	Thr Asn 2070	Thr Asn 2070	Gly 2070	Ile Thr 2070	Ala
Ala Trp 2075	Ala Asn 2080	Arg Gln 2080	Gln 2080	Pro Asn 2085	Asn Gln 2085	Gln 2085	Ala Gly 2085	Val
Gln His 2090	Leu Asn 2095	Val Asp 2095	Val 2095	Thr Tyr 2100	Pro Gly 2100	Ile 2100	Ser Ala 2100	Ala
Lys Arg 2105	Val Pro 2110	Val Thr 2110	Val 2110	Asn Val 2115	Tyr Gln 2115	Phe 2115	Glu Phe 2115	Pro
Gln Thr 2120	Thr Tyr 2125	Thr Thr 2125	Thr 2125	Val Gly 2130	Gly Thr 2130	Leu 2130	Ala Ser 2130	Gly
Thr Gln 2135	Ala Ser 2140	Gly Tyr 2140	Ala 2140	His Met 2145	Gln Asn 2145	Ala 2145	Thr Gly 2145	Leu
Pro Thr 2150	Asp Gly 2155	Phe Thr 2155	Tyr 2155	Lys Trp 2160	Asn Arg 2160	Asp 2160	Thr Thr 2160	Gly
Thr Asn 2165	Asp Ala 2170	Asn Trp 2170	Ser 2170	Ala Met 2175	Asn Lys 2175	Pro 2175	Asn Val 2175	Ala
Lys Val 2180	Val Asn 2185	Ala Lys 2185	Tyr 2185	Asp Val 2190	Ile Tyr 2190	Asn 2190	Gly His 2190	Thr
Phe Ala 2195	Thr Ser 2200	Leu Pro 2200	Ala 2200	Lys Phe 2205	Val Val 2205	Lys 2205	Asp Val 2205	Gln
Pro Ala 2210	Lys Pro 2215	Thr Val 2215	Thr 2215	Glu Thr 2220	Ala Ala 2220	Gly 2220	Ala Ile 2220	Thr
Ile Ala 2225	Pro Gly 2230	Ala Asn 2230	Gln 2230	Thr Val 2235	Asn Thr 2235	His 2235	Ala Gly 2235	Asn

-continued

---

2225	2230	2235
Val Thr Thr Tyr Ala Asp Lys	Leu Val Ile Lys Arg Asn Gly Asn	
2240	2245	2250
Val Val Thr Thr Phe Thr Arg	Arg Asn Asn Thr Ser Pro Trp Val	
2255	2260	2265
Lys Glu Ala Ser Ala Ala Thr	Val Ala Gly Ile Ala Gly Thr Asn	
2270	2275	2280
Asn Gly Ile Thr Val Ala Ala	Gly Thr Phe Asn Pro Ala Asp Thr	
2285	2290	2295
Ile Gln Val Val Ala Thr Gln	Gly Ser Gly Glu Thr Val Ser Asp	
2300	2305	2310
Glu Gln Arg Ser Asp Asp Phe	Thr Val Val Ala Pro Gln Pro Asn	
2315	2320	2325
Gln Ala Thr Thr Lys Ile Trp	Gln Asn Gly His Ile Asp Ile Thr	
2330	2335	2340
Pro Asn Asn Pro Ser Gly His	Leu Ile Asn Pro Thr Gln Ala Met	
2345	2350	2355
Asp Ile Ala Tyr Thr Glu Lys	Val Gly Asn Gly Ala Glu His Ser	
2360	2365	2370
Lys Thr Ile Asn Val Val Arg	Gly Gln Asn Asn Gln Trp Thr Ile	
2375	2380	2385
Ala Asn Lys Pro Asp Tyr Val	Thr Leu Asp Ala Gln Thr Gly Lys	
2390	2395	2400
Val Thr Phe Asn Ala Asn Thr	Ile Lys Pro Asn Ser Ser Ile Thr	
2405	2410	2415
Ile Thr Pro Lys Ala Gly Thr	Gly His Ser Val Ser Ser Asn Pro	
2420	2425	2430
Ser Thr Leu Thr Ala Pro Ala	Ala His Thr Val Asn Thr Thr Glu	
2435	2440	2445
Ile Val Lys Asp Tyr Gly Ser	Asn Val Thr Ala Ala Glu Ile Asn	
2450	2455	2460
Asn Ala Val Gln Val Ala Asn	Lys Arg Thr Ala Thr Ile Lys Asn	
2465	2470	2475
Gly Thr Ala Met Pro Thr Asn	Leu Ala Gly Gly Ser Thr Thr Thr	
2480	2485	2490
Ile Pro Val Thr Val Thr Tyr	Asn Asp Gly Ser Thr Glu Glu Val	
2495	2500	2505
Gln Glu Ser Ile Phe Thr Lys	Ala Asp Lys Arg Glu Leu Ile Thr	
2510	2515	2520
Ala Lys Asn His Leu Asp Asp	Pro Val Ser Thr Glu Gly Lys Lys	
2525	2530	2535
Pro Gly Thr Ile Thr Gln Tyr	Asn Asn Ala Met His Asn Ala Gln	
2540	2545	2550
Gln Gln Ile Asn Thr Ala Lys	Thr Glu Ala Gln Gln Val Ile Asn	
2555	2560	2565
Asn Glu Arg Ala Thr Pro Gln	Gln Val Ser Asp Ala Leu Thr Lys	
2570	2575	2580
Val Arg Ala Ala Gln Thr Lys	Ile Asp Gln Ala Lys Ala Leu Leu	
2585	2590	2595
Gln Asn Lys Glu Asp Asn Ser	Gln Leu Val Thr Ser Lys Asn Asn	
2600	2605	2610
Leu Gln Ser Ser Val Asn Gln	Val Pro Ser Thr Ala Gly Met Thr	
2615	2620	2625

-continued

---

Gln	Gln	Ser	Ile	Asp	Asn	Tyr	Asn	Ala	Lys	Lys	Arg	Glu	Ala	Glu
2630						2635					2640			
Thr	Glu	Ile	Thr	Ala	Ala	Gln	Arg	Val	Ile	Asp	Asn	Gly	Asp	Ala
2645						2650					2655			
Thr	Ala	Gln	Gln	Ile	Ser	Asp	Glu	Lys	His	Arg	Val	Asp	Asn	Ala
2660						2665					2670			
Leu	Thr	Ala	Leu	Asn	Gln	Ala	Lys	His	Asp	Leu	Thr	Ala	Asp	Thr
2675						2680					2685			
His	Ala	Leu	Glu	Gln	Ala	Val	Gln	Gln	Leu	Asn	Arg	Thr	Gly	Thr
2690						2695					2700			
Thr	Thr	Gly	Lys	Lys	Pro	Ala	Ser	Ile	Thr	Ala	Tyr	Asn	Asn	Ser
2705						2710					2715			
Ile	Arg	Ala	Leu	Gln	Ser	Asp	Leu	Thr	Ser	Ala	Lys	Asn	Ser	Ala
2720						2725					2730			
Asn	Ala	Ile	Ile	Gln	Lys	Pro	Ile	Arg	Thr	Val	Gln	Glu	Val	Gln
2735						2740					2745			
Ser	Ala	Leu	Thr	Asn	Val	Asn	Arg	Val	Asn	Glu	Arg	Leu	Thr	Gln
2750						2755					2760			
Ala	Ile	Asn	Gln	Leu	Val	Pro	Leu	Ala	Asp	Asn	Ser	Ala	Leu	Lys
2765						2770					2775			
Thr	Ala	Lys	Thr	Lys	Leu	Asp	Glu	Glu	Ile	Asn	Lys	Ser	Val	Thr
2780						2785					2790			
Thr	Asp	Gly	Met	Thr	Gln	Ser	Ser	Ile	Gln	Ala	Tyr	Glu	Asn	Ala
2795						2800					2805			
Lys	Arg	Ala	Gly	Gln	Thr	Glu	Ser	Thr	Asn	Ala	Gln	Asn	Val	Ile
2810						2815					2820			
Asn	Asn	Gly	Asp	Ala	Thr	Asp	Gln	Gln	Ile	Ala	Ala	Glu	Lys	Thr
2825						2830					2835			
Lys	Val	Glu	Glu	Lys	Tyr	Asn	Ser	Leu	Lys	Gln	Ala	Ile	Ala	Gly
2840						2845					2850			
Leu	Thr	Pro	Asp	Leu	Ala	Pro	Leu	Gln	Thr	Ala	Lys	Thr	Gln	Leu
2855						2860					2865			
Gln	Asn	Asp	Ile	Asp	Gln	Pro	Thr	Ser	Thr	Thr	Gly	Met	Thr	Ser
2870						2875					2880			
Ala	Ser	Ile	Ala	Ala	Phe	Asn	Glu	Lys	Leu	Ser	Ala	Ala	Arg	Thr
2885						2890					2895			
Lys	Ile	Gln	Glu	Ile	Asp	Arg	Val	Leu	Ala	Ser	His	Pro	Asp	Val
2900						2905					2910			
Ala	Thr	Ile	Arg	Gln	Asn	Val	Thr	Ala	Ala	Asn	Ala	Ala	Lys	Ser
2915						2920					2925			
Ala	Leu	Asp	Gln	Ala	Arg	Asn	Gly	Leu	Thr	Val	Asp	Lys	Ala	Pro
2930						2935					2940			
Leu	Glu	Asn	Ala	Lys	Asn	Gln	Leu	Gln	His	Ser	Ile	Asp	Thr	Gln
2945						2950					2955			
Thr	Ser	Thr	Thr	Gly	Met	Thr	Gln	Asp	Ser	Ile	Asn	Ala	Tyr	Asn
2960						2965					2970			
Ala	Lys	Leu	Thr	Ala	Ala	Arg	Asn	Lys	Ile	Gln	Gln	Ile	Asn	Gln
2975						2980					2985			
Val	Leu	Ala	Gly	Ser	Pro	Thr	Val	Glu	Gln	Ile	Asn	Thr	Asn	Thr
2990						2995					3000			
Ser	Thr	Ala	Asn	Gln	Ala	Lys	Ser	Asp	Leu	Asp	His	Ala	Arg	Gln
3005						3010					3015			

-continued

Ala	Leu	Thr	Pro	Asp	Lys	Ala	Pro	Leu	Gln	Thr	Ala	Lys	Thr	Gln
3020						3025					3030			
Leu	Glu	Gln	Ser	Ile	Asn	Gln	Pro	Thr	Asp	Thr	Thr	Gly	Met	Thr
3035						3040					3045			
Thr	Ala	Ser	Leu	Asn	Ala	Tyr	Asn	Gln	Lys	Leu	Gln	Ala	Ala	Arg
3050						3055					3060			
Gln	Lys	Leu	Thr	Glu	Ile	Asn	Gln	Val	Leu	Asn	Gly	Asn	Pro	Thr
3065						3070					3075			
Val	Gln	Asn	Ile	Asn	Asp	Lys	Val	Thr	Glu	Ala	Asn	Gln	Ala	Lys
3080						3085					3090			
Asp	Gln	Leu	Asn	Thr	Ala	Arg	Gln	Gly	Leu	Thr	Leu	Asp	Arg	Gln
3095						3100					3105			
Pro	Ala	Leu	Thr	Thr	Leu	His	Gly	Ala	Ser	Asn	Leu	Asn	Gln	Ala
3110						3115					3120			
Gln	Gln	Asn	Asn	Phe	Thr	Gln	Gln	Ile	Asn	Ala	Ala	Gln	Asn	His
3125						3130					3135			
Ala	Ala	Leu	Glu	Thr	Ile	Lys	Ser	Asn	Ile	Thr	Ala	Leu	Asn	Thr
3140						3145					3150			
Ala	Met	Thr	Lys	Leu	Lys	Asp	Ser	Val	Ala	Asp	Asn	Asn	Thr	Ile
3155						3160					3165			
Lys	Ser	Asp	Gln	Asn	Tyr	Thr	Asp	Ala	Thr	Pro	Ala	Asn	Lys	Gln
3170						3175					3180			
Ala	Tyr	Asp	Asn	Ala	Val	Asn	Ala	Ala	Lys	Gly	Val	Ile	Gly	Glu
3185						3190					3195			
Thr	Thr	Asn	Pro	Thr	Met	Asp	Val	Asn	Thr	Val	Asn	Gln	Lys	Ala
3200						3205					3210			
Ala	Ser	Val	Lys	Ser	Thr	Lys	Asp	Ala	Leu	Asp	Gly	Gln	Gln	Asn
3215						3220					3225			
Leu	Gln	Arg	Ala	Lys	Thr	Glu	Ala	Thr	Asn	Ala	Ile	Thr	His	Ala
3230						3235					3240			
Ser	Asp	Leu	Asn	Gln	Ala	Gln	Lys	Asn	Ala	Leu	Thr	Gln	Gln	Val
3245						3250					3255			
Asn	Ser	Ala	Gln	Asn	Val	Gln	Ala	Val	Asn	Asp	Ile	Lys	Gln	Thr
3260						3265					3270			
Thr	Gln	Ser	Leu	Asn	Thr	Ala	Met	Thr	Gly	Leu	Lys	Arg	Gly	Val
3275						3280					3285			
Ala	Asn	His	Asn	Gln	Val	Val	Gln	Ser	Asp	Asn	Tyr	Val	Asn	Ala
3290						3295					3300			
Asp	Thr	Asn	Lys	Lys	Asn	Asp	Tyr	Asn	Asn	Ala	Tyr	Asn	His	Ala
3305						3310					3315			
Asn	Asp	Ile	Ile	Asn	Gly	Asn	Ala	Gln	His	Pro	Val	Ile	Thr	Pro
3320						3325					3330			
Ser	Asp	Val	Asn	Asn	Ala	Leu	Ser	Asn	Val	Thr	Ser	Lys	Glu	His
3335						3340					3345			
Ala	Leu	Asn	Gly	Glu	Ala	Lys	Leu	Asn	Ala	Ala	Lys	Gln	Glu	Ala
3350						3355					3360			
Asn	Thr	Ala	Leu	Gly	His	Leu	Asn	Asn	Leu	Asn	Asn	Ala	Gln	Arg
3365						3370					3375			
Gln	Asn	Leu	Gln	Ser	Gln	Ile	Asn	Gly	Ala	His	Gln	Ile	Asp	Ala
3380						3385					3390			
Val	Asn	Thr	Ile	Lys	Gln	Asn	Ala	Thr	Asn	Leu	Asn	Ser	Ala	Met
3395						3400					3405			
Gly	Asn	Leu	Arg	Gln	Ala	Val	Ala	Asp	Lys	Asp	Gln	Val	Lys	Arg

-continued

3410	3415	3420
Thr Glu Asp Tyr Ala Asp	Ala Asp Thr Ala Lys	Gln Asn Ala Tyr
3425	3430	3435
Asn Ser Ala Val Ser Ser	Ala Glu Thr Ile Ile	Asn Gln Thr Thr
3440	3445	3450
Asn Pro Thr Met Ser Val	Asp Asp Val Asn Arg	Ala Thr Ser Ala
3455	3460	3465
Val Thr Ser Asn Lys Asn	Ala Leu Asn Gly Tyr	Glu Lys Leu Ala
3470	3475	3480
Gln Ser Lys Thr Asp Ala	Ala Arg Ala Ile Asp	Ala Leu Pro His
3485	3490	3495
Leu Asn Asn Ala Gln Lys	Ala Asp Val Lys Ser	Lys Ile Asn Ala
3500	3505	3510
Ala Ser Asn Ile Ala Gly	Val Asn Thr Val Lys	Gln Gln Gly Thr
3515	3520	3525
Asp Leu Asn Thr Ala Met	Gly Asn Leu Gln Gly	Ala Ile Asn Asp
3530	3535	3540
Glu Gln Thr Thr Leu Asn	Ser Gln Asn Tyr Gln	Asp Ala Thr Pro
3545	3550	3555
Ser Lys Lys Thr Ala Tyr	Thr Asn Ala Val Gln	Ala Ala Lys Asp
3560	3565	3570
Ile Leu Asn Lys Ser Asn	Gly Gln Asn Lys Thr	Lys Asp Gln Val
3575	3580	3585
Thr Glu Ala Met Asn Gln	Val Asn Ser Ala Lys	Asn Asn Leu Asp
3590	3595	3600
Gly Thr Arg Leu Leu Asp	Gln Ala Lys Gln Thr	Ala Lys Gln Gln
3605	3610	3615
Leu Asn Asn Met Thr His	Leu Thr Thr Ala Gln	Lys Thr Asn Leu
3620	3625	3630
Thr Asn Gln Ile Asn Ser	Gly Thr Thr Val Ala	Gly Val Gln Thr
3635	3640	3645
Val Gln Ser Asn Ala Asn	Thr Leu Asp Gln Ala	Met Asn Thr Leu
3650	3655	3660
Arg Gln Ser Ile Ala Asn	Lys Asp Ala Thr Lys	Ala Ser Glu Asp
3665	3670	3675
Tyr Val Asp Ala Asn Asn	Asp Lys Gln Thr Ala	Tyr Asn Asn Ala
3680	3685	3690
Val Ala Ala Ala Glu Thr	Ile Ile Asn Ala Asn	Ser Asn Pro Glu
3695	3700	3705
Met Asn Pro Ser Thr Ile	Thr Gln Lys Ala Glu	Gln Val Asn Ser
3710	3715	3720
Ser Lys Thr Ala Leu Asn	Gly Asp Glu Asn Leu	Ala Ala Ala Lys
3725	3730	3735
Gln Asn Ala Lys Thr Tyr	Leu Asn Thr Leu Thr	Ser Ile Thr Asp
3740	3745	3750
Ala Gln Lys Asn Asn Leu	Ile Ser Gln Ile Thr	Ser Ala Thr Arg
3755	3760	3765
Val Ser Gly Val Asp Thr	Val Lys Gln Asn Ala	Gln His Leu Asp
3770	3775	3780
Gln Ala Met Ala Ser Leu	Gln Asn Gly Ile Asn	Asn Glu Ser Gln
3785	3790	3795
Val Lys Ser Ser Glu Lys	Tyr Arg Asp Ala Asp	Thr Asn Lys Gln
3800	3805	3810

-continued

---

Gln	Glu	Tyr	Asp	Asn	Ala	Ile	Thr	Ala	Ala	Lys	Ala	Ile	Leu	Asn
3815						3820					3825			
Lys	Ser	Thr	Gly	Pro	Asn	Thr	Ala	Gln	Asn	Ala	Val	Glu	Ala	Ala
3830						3835					3840			
Leu	Gln	Arg	Val	Asn	Asn	Ala	Lys	Asp	Ala	Leu	Asn	Gly	Asp	Ala
3845						3850					3855			
Lys	Leu	Ile	Ala	Ala	Gln	Asn	Ala	Ala	Lys	Gln	His	Leu	Gly	Thr
3860						3865					3870			
Leu	Thr	His	Ile	Thr	Thr	Ala	Gln	Arg	Asn	Asp	Leu	Thr	Asn	Gln
3875						3880					3885			
Ile	Ser	Gln	Ala	Thr	Asn	Leu	Ala	Gly	Val	Glu	Ser	Val	Lys	Gln
3890						3895					3900			
Asn	Ala	Asn	Ser	Leu	Asp	Gly	Ala	Met	Gly	Asn	Leu	Gln	Thr	Ala
3905						3910					3915			
Ile	Asn	Asp	Lys	Ser	Gly	Thr	Leu	Ala	Ser	Gln	Asn	Phe	Leu	Asp
3920						3925					3930			
Ala	Asp	Glu	Gln	Lys	Arg	Asn	Ala	Tyr	Asn	Gln	Ala	Val	Ser	Ala
3935						3940					3945			
Ala	Glu	Thr	Ile	Leu	Asn	Lys	Gln	Thr	Gly	Pro	Asn	Thr	Ala	Lys
3950						3955					3960			
Thr	Ala	Val	Glu	Gln	Ala	Leu	Asn	Asn	Val	Asn	Asn	Ala	Lys	His
3965						3970					3975			
Ala	Leu	Asn	Gly	Thr	Gln	Asn	Leu	Asn	Asn	Ala	Lys	Gln	Ala	Ala
3980						3985					3990			
Ile	Thr	Ala	Ile	Asn	Gly	Ala	Ser	Asp	Leu	Asn	Gln	Lys	Gln	Lys
3995						4000					4005			
Asp	Ala	Leu	Lys	Ala	Gln	Ala	Asn	Gly	Ala	Gln	Arg	Val	Ser	Asn
4010						4015					4020			
Ala	Gln	Asp	Val	Gln	His	Asn	Ala	Thr	Glu	Leu	Asn	Thr	Ala	Met
4025						4030					4035			
Gly	Thr	Leu	Lys	His	Ala	Ile	Ala	Asp	Lys	Thr	Asn	Thr	Leu	Ala
4040						4045					4050			
Ser	Ser	Lys	Tyr	Val	Asn	Ala	Asp	Ser	Thr	Lys	Gln	Asn	Ala	Tyr
4055						4060					4065			
Thr	Thr	Lys	Val	Thr	Asn	Ala	Glu	His	Ile	Ile	Ser	Gly	Thr	Pro
4070						4075					4080			
Thr	Val	Val	Thr	Thr	Pro	Ser	Glu	Val	Thr	Ala	Ala	Ala	Asn	Gln
4085						4090					4095			
Val	Asn	Ser	Ala	Lys	Gln	Glu	Leu	Asn	Gly	Asp	Glu	Arg	Leu	Arg
4100						4105					4110			
Glu	Ala	Lys	Gln	Asn	Ala	Asn	Thr	Ala	Ile	Asp	Ala	Leu	Thr	Gln
4115						4120					4125			
Leu	Asn	Thr	Pro	Gln	Lys	Ala	Lys	Leu	Lys	Glu	Gln	Val	Gly	Gln
4130						4135					4140			
Ala	Asn	Arg	Leu	Glu	Asp	Val	Gln	Thr	Val	Gln	Thr	Asn	Gly	Gln
4145						4150					4155			
Ala	Leu	Asn	Asn	Ala	Met	Lys	Gly	Leu	Arg	Asp	Ser	Ile	Ala	Asn
4160						4165					4170			
Glu	Thr	Thr	Val	Lys	Thr	Ser	Gln	Asn	Tyr	Thr	Asp	Ala	Ser	Pro
4175						4180					4185			
Asn	Asn	Gln	Ser	Thr	Tyr	Asn	Ser	Ala	Val	Ser	Asn	Ala	Lys	Gly
4190						4195					4200			

-continued

Ile	Ile	Asn	Gln	Thr	Asn	Asn	Pro	Thr	Met	Asp	Thr	Ser	Ala	Ile
4205						4210					4215			
Thr	Gln	Ala	Thr	Thr	Gln	Val	Asn	Asn	Ala	Lys	Asn	Gly	Leu	Asn
4220						4225					4230			
Gly	Ala	Glu	Asn	Leu	Arg	Asn	Ala	Gln	Asn	Thr	Ala	Lys	Gln	Asn
4235						4240					4245			
Leu	Asn	Thr	Leu	Ser	His	Leu	Thr	Asn	Asn	Gln	Lys	Ser	Ala	Ile
4250						4255					4260			
Ser	Ser	Gln	Ile	Asp	Arg	Ala	Gly	His	Val	Ser	Glu	Val	Thr	Ala
4265						4270					4275			
Thr	Lys	Asn	Ala	Ala	Thr	Glu	Leu	Asn	Thr	Gln	Met	Gly	Asn	Leu
4280						4285					4290			
Glu	Gln	Ala	Ile	His	Asp	Gln	Asn	Thr	Val	Lys	Gln	Ser	Val	Lys
4295						4300					4305			
Phe	Thr	Asp	Ala	Asp	Lys	Ala	Lys	Arg	Asp	Ala	Tyr	Thr	Asn	Ala
4310						4315					4320			
Val	Ser	Arg	Ala	Glu	Ala	Ile	Leu	Asn	Lys	Thr	Gln	Gly	Ala	Asn
4325						4330					4335			
Thr	Ser	Lys	Gln	Asp	Val	Glu	Ala	Ala	Ile	Gln	Asn	Val	Ser	Ser
4340						4345					4350			
Ala	Lys	Asn	Ala	Leu	Asn	Gly	Asp	Gln	Asn	Val	Thr	Asn	Ala	Lys
4355						4360					4365			
Asn	Ala	Ala	Lys	Asn	Ala	Leu	Asn	Asn	Leu	Thr	Ser	Ile	Asn	Asn
4370						4375					4380			
Ala	Gln	Lys	Arg	Asp	Leu	Thr	Thr	Lys	Ile	Asp	Gln	Ala	Thr	Thr
4385						4390					4395			
Val	Ala	Gly	Val	Glu	Ala	Val	Ser	Asn	Thr	Ser	Thr	Gln	Leu	Asn
4400						4405					4410			
Thr	Ala	Met	Ala	Asn	Leu	Gln	Asn	Gly	Ile	Asn	Asp	Lys	Thr	Asn
4415						4420					4425			
Thr	Leu	Ala	Ser	Glu	Asn	Tyr	His	Asp	Ala	Asp	Ser	Asp	Lys	Lys
4430						4435					4440			
Thr	Ala	Tyr	Thr	Gln	Ala	Val	Thr	Asn	Ala	Glu	Asn	Ile	Leu	Asn
4445						4450					4455			
Lys	Asn	Ser	Gly	Ser	Asn	Leu	Asp	Lys	Thr	Ala	Val	Glu	Asn	Ala
4460						4465					4470			
Leu	Ser	Gln	Val	Ala	Asn	Ala	Lys	Gly	Ala	Leu	Asn	Gly	Asn	His
4475						4480					4485			
Asn	Leu	Glu	Gln	Ala	Lys	Ser	Asn	Ala	Asn	Thr	Thr	Ile	Asn	Gly
4490						4495					4500			
Leu	Gln	His	Leu	Thr	Thr	Ala	Gln	Lys	Asp	Lys	Leu	Lys	Gln	Gln
4505						4510					4515			
Val	Gln	Gln	Ala	Gln	Asn	Val	Ala	Gly	Val	Asp	Thr	Val	Lys	Ser
4520						4525					4530			
Ser	Ala	Asn	Thr	Leu	Asn	Gly	Ala	Met	Gly	Thr	Leu	Arg	Asn	Ser
4535						4540					4545			
Ile	Gln	Asp	Asn	Thr	Ala	Thr	Lys	Asn	Gly	Gln	Asn	Tyr	Leu	Asp
4550						4555					4560			
Ala	Thr	Glu	Arg	Asn	Lys	Thr	Asn	Tyr	Asn	Asn	Ala	Val	Asp	Ser
4565						4570					4575			
Ala	Asn	Gly	Val	Ile	Asn	Ala	Thr	Ser	Asn	Pro	Asn	Met	Asp	Ala
4580						4585					4590			
Asn	Ala	Ile	Asn	Gln	Ile	Ala	Thr	Gln	Val	Thr	Ser	Thr	Lys	Asn

-continued

---

4595	4600	4605
Ala Leu Asp Gly Thr His Asn	Leu Thr Gln Ala Lys	Gln Thr Ala
4610	4615	4620
Thr Asn Ala Ile Asp Gly Ala	Thr Asn Leu Asn Lys	Ala Gln Lys
4625	4630	4635
Asp Ala Leu Lys Ala Gln Val	Thr Ser Ala Gln Arg	Val Ala Asn
4640	4645	4650
Val Thr Ser Ile Gln Gln Thr	Ala Asn Glu Leu Asn	Thr Ala Met
4655	4660	4665
Gly Gln Leu Gln His Gly Ile	Asp Asp Glu Asn Ala	Thr Lys Gln
4670	4675	4680
Thr Gln Lys Tyr Arg Asp Ala	Glu Gln Ser Lys Lys	Thr Ala Tyr
4685	4690	4695
Asp Gln Ala Val Ala Ala Ala	Lys Ala Ile Leu Asn	Lys Gln Thr
4700	4705	4710
Gly Ser Asn Ser Asp Lys Ala	Ala Val Asp Arg Ala	Leu Gln Gln
4715	4720	4725
Val Thr Ser Thr Lys Asp Ala	Leu Asn Gly Asp Ala	Lys Leu Ala
4730	4735	4740
Glu Ala Lys Ala Ala Ala Lys	Gln Asn Leu Gly Thr	Leu Asn His
4745	4750	4755
Ile Thr Asn Ala Gln Arg Thr	Asp Leu Glu Gly Gln	Ile Asn Gln
4760	4765	4770
Ala Thr Thr Val Asp Gly Val	Asn Thr Val Lys Thr	Asn Ala Asn
4775	4780	4785
Thr Leu Asp Gly Ala Met Asn	Ser Leu Gln Gly Ser	Ile Asn Asp
4790	4795	4800
Lys Asp Ala Thr Leu Arg Asn	Gln Asn Tyr Leu Asp	Ala Asp Glu
4805	4810	4815
Ser Lys Arg Asn Ala Tyr Thr	Gln Ala Val Thr Ala	Ala Glu Gly
4820	4825	4830
Ile Leu Asn Lys Gln Thr Gly	Gly Asn Thr Ser Lys	Ala Asp Val
4835	4840	4845
Asp Asn Ala Leu Asn Ala Val	Thr Arg Ala Lys Ala	Ala Leu Asn
4850	4855	4860
Gly Ala Asp Asn Leu Arg Asn	Ala Lys Thr Ser Ala	Thr Asn Thr
4865	4870	4875
Ile Asp Gly Leu Pro Asn Leu	Thr Gln Leu Gln Lys	Asp Asn Leu
4880	4885	4890
Lys His Gln Val Glu Gln Ala	Gln Asn Val Ala Gly	Val Asn Gly
4895	4900	4905
Val Lys Asp Lys Gly Asn Thr	Leu Asn Thr Ala Met	Gly Ala Leu
4910	4915	4920
Arg Thr Ser Ile Gln Asn Asp	Asn Thr Thr Lys Thr	Ser Gln Asn
4925	4930	4935
Tyr Leu Asp Ala Ser Asp Ser	Asn Lys Asn Asn Tyr	Asn Thr Ala
4940	4945	4950
Val Asn Asn Ala Asn Gly Val	Ile Asn Ala Thr Asn	Asn Pro Asn
4955	4960	4965
Met Asp Ala Asn Ala Ile Asn	Gly Met Ala Asn Gln	Val Asn Thr
4970	4975	4980
Thr Lys Ala Ala Leu Asn Gly	Ala Gln Asn Leu Ala	Gln Ala Lys
4985	4990	4995

-continued

---

Thr	Asn	Ala	Thr	Asn	Thr	Ile	Asn	Asn	Ala	His	Asp	Leu	Asn	Gln
5000						5005					5010			
Lys	Gln	Lys	Asp	Ala	Leu	Lys	Thr	Gln	Val	Asn	Asn	Ala	Gln	Arg
5015						5020					5025			
Val	Ser	Asp	Ala	Asn	Asn	Val	Gln	His	Thr	Ala	Thr	Glu	Leu	Asn
5030						5035					5040			
Ser	Ala	Met	Thr	Ala	Leu	Lys	Ala	Ala	Ile	Ala	Asp	Lys	Glu	Arg
5045						5050					5055			
Thr	Lys	Ala	Ser	Gly	Asn	Tyr	Val	Asn	Ala	Asp	Gln	Glu	Lys	Arg
5060						5065					5070			
Gln	Ala	Tyr	Asp	Ser	Lys	Val	Thr	Asn	Ala	Glu	Asn	Ile	Ile	Ser
5075						5080					5085			
Gly	Thr	Pro	Asn	Ala	Thr	Leu	Thr	Val	Asn	Asp	Val	Asn	Ser	Ala
5090						5095					5100			
Ala	Ser	Gln	Val	Asn	Ala	Ala	Lys	Thr	Ala	Leu	Asn	Gly	Asp	Asn
5105						5110					5115			
Asn	Leu	Arg	Val	Ala	Lys	Glu	His	Ala	Asn	Asn	Thr	Ile	Asp	Gly
5120						5125					5130			
Leu	Ala	Gln	Leu	Asn	Asn	Ala	Gln	Lys	Ala	Lys	Leu	Lys	Glu	Gln
5135						5140					5145			
Val	Gln	Ser	Ala	Thr	Thr	Leu	Asp	Gly	Val	Gln	Thr	Val	Lys	Asn
5150						5155					5160			
Ser	Ser	Gln	Thr	Leu	Asn	Thr	Ala	Met	Lys	Gly	Leu	Arg	Asp	Ser
5165						5170					5175			
Ile	Ala	Asn	Glu	Ala	Thr	Ile	Lys	Ala	Gly	Gln	Asn	Tyr	Thr	Asp
5180						5185					5190			
Ala	Ser	Pro	Asn	Asn	Arg	Asn	Glu	Tyr	Asp	Ser	Ala	Val	Thr	Ala
5195						5200					5205			
Ala	Lys	Ala	Ile	Ile	Asn	Gln	Thr	Ser	Asn	Pro	Thr	Met	Glu	Pro
5210						5215					5220			
Asn	Thr	Ile	Thr	Gln	Val	Thr	Ser	Gln	Val	Thr	Thr	Lys	Glu	Gln
5225						5230					5235			
Ala	Leu	Asn	Gly	Ala	Arg	Asn	Leu	Ala	Gln	Ala	Lys	Thr	Thr	Ala
5240						5245					5250			
Lys	Asn	Asn	Leu	Asn	Asn	Leu	Thr	Ser	Ile	Asn	Asn	Ala	Gln	Lys
5255						5260					5265			
Asp	Ala	Leu	Thr	Arg	Ser	Ile	Asp	Gly	Ala	Thr	Thr	Val	Ala	Gly
5270						5275					5280			
Val	Asn	Gln	Glu	Thr	Ala	Lys	Ala	Thr	Glu	Leu	Asn	Asn	Ala	Met
5285						5290					5295			
His	Ser	Leu	Gln	Asn	Gly	Ile	Asn	Asp	Glu	Thr	Gln	Thr	Lys	Gln
5300						5305					5310			
Thr	Gln	Lys	Tyr	Leu	Asp	Ala	Glu	Pro	Ser	Lys	Lys	Ser	Ala	Tyr
5315						5320					5325			
Asp	Gln	Ala	Val	Asn	Ala	Ala	Lys	Ala	Ile	Leu	Thr	Lys	Ala	Ser
5330						5335					5340			
Gly	Gln	Asn	Val	Asp	Lys	Ala	Ala	Val	Glu	Gln	Ala	Leu	Gln	Asn
5345						5350					5355			
Val	Asn	Ser	Thr	Lys	Thr	Ala	Leu	Asn	Gly	Asp	Ala	Lys	Leu	Asn
5360						5365					5370			
Glu	Ala	Lys	Ala	Ala	Ala	Lys	Gln	Thr	Leu	Gly	Thr	Leu	Thr	His
5375						5380					5385			

-continued

Ile	Asn	Asn	Ala	Gln	Arg	Thr	Ala	Leu	Asp	Asn	Glu	Ile	Thr	Gln
5390						5395					5400			
Ala	Thr	Asn	Val	Glu	Gly	Val	Asn	Thr	Val	Lys	Ala	Lys	Ala	Gln
5405						5410					5415			
Gln	Leu	Asp	Gly	Ala	Met	Gly	Gln	Leu	Glu	Thr	Ser	Ile	Arg	Asp
5420						5425					5430			
Lys	Asp	Thr	Thr	Leu	Gln	Ser	Gln	Asn	Tyr	Gln	Asp	Ala	Asp	Asp
5435						5440					5445			
Ala	Lys	Arg	Thr	Ala	Tyr	Ser	Gln	Ala	Val	Asn	Ala	Ala	Ala	Thr
5450						5455					5460			
Ile	Leu	Asn	Lys	Thr	Ala	Gly	Gly	Asn	Thr	Pro	Lys	Ala	Asp	Val
5465						5470					5475			
Glu	Arg	Ala	Met	Gln	Ala	Val	Thr	Gln	Ala	Asn	Thr	Ala	Leu	Asn
5480						5485					5490			
Gly	Ile	Gln	Asn	Leu	Asp	Arg	Ala	Lys	Gln	Ala	Ala	Asn	Thr	Ala
5495						5500					5505			
Ile	Thr	Asn	Ala	Ser	Asp	Leu	Asn	Thr	Lys	Gln	Lys	Glu	Ala	Leu
5510						5515					5520			
Lys	Ala	Gln	Val	Thr	Ser	Ala	Gly	Arg	Val	Ser	Ala	Ala	Asn	Gly
5525						5530					5535			
Val	Glu	His	Thr	Ala	Thr	Glu	Leu	Asn	Thr	Ala	Met	Thr	Ala	Leu
5540						5545					5550			
Lys	Arg	Ala	Ile	Ala	Asp	Lys	Ala	Glu	Thr	Lys	Ala	Ser	Gly	Asn
5555						5560					5565			
Tyr	Val	Asn	Ala	Asp	Ala	Asn	Lys	Arg	Gln	Ala	Tyr	Asp	Glu	Lys
5570						5575					5580			
Val	Thr	Ala	Ala	Glu	Asn	Ile	Val	Ser	Gly	Thr	Pro	Thr	Pro	Thr
5585						5590					5595			
Leu	Thr	Pro	Ala	Asp	Val	Thr	Asn	Ala	Ala	Thr	Gln	Val	Thr	Asn
5600						5605					5610			
Ala	Lys	Thr	Gln	Leu	Asn	Gly	Asn	His	Asn	Leu	Glu	Val	Ala	Lys
5615						5620					5625			
Gln	Asn	Ala	Asn	Thr	Ala	Ile	Asp	Gly	Leu	Thr	Ser	Leu	Asn	Gly
5630						5635					5640			
Pro	Gln	Lys	Ala	Lys	Leu	Lys	Glu	Gln	Val	Gly	Gln	Ala	Thr	Thr
5645						5650					5655			
Leu	Pro	Asn	Val	Gln	Thr	Val	Arg	Asp	Asn	Ala	Gln	Thr	Leu	Asn
5660						5665					5670			
Thr	Ala	Met	Lys	Gly	Leu	Arg	Asp	Ser	Ile	Ala	Asn	Glu	Ala	Thr
5675						5680					5685			
Ile	Lys	Ala	Gly	Gln	Asn	Tyr	Thr	Asp	Ala	Ser	Gln	Asn	Lys	Gln
5690						5695					5700			
Thr	Asp	Tyr	Asn	Ser	Ala	Val	Thr	Ala	Ala	Lys	Ala	Ile	Ile	Gly
5705						5710					5715			
Gln	Thr	Thr	Ser	Pro	Ser	Met	Asn	Ala	Gln	Glu	Ile	Asn	Gln	Ala
5720						5725					5730			
Lys	Asp	Gln	Val	Thr	Ala	Lys	Gln	Gln	Ala	Leu	Asn	Gly	Gln	Glu
5735						5740					5745			
Asn	Leu	Arg	Thr	Ala	Gln	Thr	Asn	Ala	Lys	Gln	His	Leu	Asn	Gly
5750						5755					5760			
Leu	Ser	Asp	Leu	Thr	Asp	Ala	Gln	Lys	Asp	Ala	Val	Lys	Arg	Gln
5765						5770					5775			
Ile	Glu	Gly	Ala	Thr	His	Val	Asn	Glu	Val	Thr	Gln	Ala	Gln	Asn

-continued

---

5780	5785	5790
Asn Ala Asp Ala Leu Asn Thr	Ala Met Thr Asn Leu Lys Asn Gly	
5795	5800	5805
Ile Gln Asp Gln Asn Thr Ile	Lys Gln Gly Val Asn Phe Thr Asp	
5810	5815	5820
Ala Asp Glu Ala Lys Arg Asn	Ala Tyr Thr Asn Ala Val Thr Gln	
5825	5830	5835
Ala Glu Gln Ile Leu Asn Lys	Ala Gln Gly Pro Asn Thr Ser Lys	
5840	5845	5850
Asp Gly Val Glu Thr Ala Leu	Glu Asn Val Gln Arg Ala Lys Asn	
5855	5860	5865
Glu Leu Asn Gly Asn Gln Asn	Val Ala Asn Ala Lys Thr Thr Ala	
5870	5875	5880
Lys Asn Ala Leu Asn Asn Leu	Thr Ser Ile Asn Asn Ala Gln Lys	
5885	5890	5895
Glu Ala Leu Lys Ser Gln Ile	Glu Gly Ala Thr Thr Val Ala Gly	
5900	5905	5910
Val Asn Gln Val Ser Thr Thr	Ala Ser Glu Leu Asn Thr Ala Met	
5915	5920	5925
Ser Asn Leu Gln Asn Gly Ile	Asn Asp Glu Ala Ala Thr Lys Ala	
5930	5935	5940
Ala Gln Lys Tyr Thr Asp Ala	Asp Arg Glu Lys Gln Thr Ala Tyr	
5945	5950	5955
Asn Asp Ala Val Thr Ala Ala	Lys Thr Leu Leu Asp Lys Thr Ala	
5960	5965	5970
Gly Ser Asn Asp Asn Lys Ala	Ala Val Glu Gln Ala Leu Gln Arg	
5975	5980	5985
Val Asn Thr Ala Lys Thr Ala	Leu Asn Gly Asp Glu Arg Leu Asn	
5990	5995	6000
Glu Ala Lys Asn Thr Ala Lys	Gln Gln Val Ala Thr Met Ser His	
6005	6010	6015
Leu Thr Asp Ala Gln Lys Ala	Asn Leu Thr Ser Gln Ile Glu Ser	
6020	6025	6030
Gly Thr Thr Val Ala Gly Val	Gln Gly Ile Gln Ala Asn Ala Gly	
6035	6040	6045
Thr Leu Asp Gln Ala Met Asn	Gln Leu Arg Gln Ser Ile Ala Ser	
6050	6055	6060
Lys Asp Ala Thr Lys Ser Ser	Glu Asp Tyr Gln Asp Ala Asn Ala	
6065	6070	6075
Asp Leu Gln Asn Ala Tyr Asn	Asp Ala Val Thr Asn Ala Glu Gly	
6080	6085	6090
Ile Ile Ser Ala Thr Asn Asn	Pro Glu Met Asn Pro Asp Thr Ile	
6095	6100	6105
Asn Gln Lys Ala Ser Gln Val	Asn Ser Ala Lys Ser Ala Leu Asn	
6110	6115	6120
Gly Asp Glu Lys Leu Ala Ala	Ala Lys Gln Thr Ala Lys Ser Asp	
6125	6130	6135
Ile Gly Arg Leu Thr Asp Leu	Asn Asn Ala Gln Arg Thr Ala Ala	
6140	6145	6150
Asn Ala Glu Val Asp Gln Ala	Pro Asn Leu Ala Ala Val Thr Ala	
6155	6160	6165
Ala Lys Asn Lys Ala Thr Ser	Leu Asn Thr Ala Met Gly Asn Leu	
6170	6175	6180

-continued

---

Lys	His	Ala	Leu	Ala	Glu	Lys	Asp	Asn	Thr	Lys	Arg	Ser	Val	Asn
6185						6190					6195			
Tyr	Thr	Asp	Ala	Asp	Gln	Pro	Lys	Gln	Gln	Ala	Tyr	Asp	Thr	Ala
6200						6205					6210			
Val	Thr	Gln	Ala	Glu	Ala	Ile	Thr	Asn	Ala	Asn	Gly	Ser	Asn	Ala
6215						6220					6225			
Asn	Glu	Thr	Gln	Val	Gln	Ala	Ala	Leu	Asn	Gln	Leu	Asn	Gln	Ala
6230						6235					6240			
Lys	Asn	Asp	Leu	Asn	Gly	Asp	Asn	Lys	Val	Ala	Gln	Ala	Lys	Glu
6245						6250					6255			
Ser	Ala	Lys	Arg	Ala	Leu	Ala	Ser	Tyr	Ser	Asn	Leu	Asn	Asn	Ala
6260						6265					6270			
Gln	Ser	Thr	Ala	Ala	Ile	Ser	Gln	Ile	Asp	Asn	Ala	Thr	Thr	Val
6275						6280					6285			
Ala	Gly	Val	Thr	Ala	Ala	Gln	Asn	Thr	Ala	Asn	Glu	Leu	Asn	Thr
6290						6295					6300			
Ala	Met	Gly	Gln	Leu	Gln	Asn	Gly	Ile	Asn	Asp	Gln	Asn	Thr	Val
6305						6310					6315			
Lys	Gln	Gln	Val	Asn	Phe	Thr	Asp	Ala	Asp	Gln	Gly	Lys	Lys	Asp
6320						6325					6330			
Ala	Tyr	Thr	Asn	Ala	Val	Thr	Asn	Ala	Gln	Gly	Ile	Leu	Asp	Lys
6335						6340					6345			
Ala	His	Gly	Gln	Asn	Met	Thr	Lys	Ala	Gln	Val	Glu	Ala	Ala	Leu
6350						6355					6360			
Asn	Gln	Val	Thr	Thr	Ala	Lys	Asn	Ala	Leu	Asn	Gly	Asp	Ala	Asn
6365						6370					6375			
Val	Arg	Gln	Ala	Lys	Ser	Asp	Ala	Lys	Ala	Asn	Leu	Gly	Thr	Leu
6380						6385					6390			
Thr	His	Leu	Asn	Asn	Ala	Gln	Lys	Gln	Asp	Leu	Thr	Ser	Gln	Ile
6395						6400					6405			
Glu	Gly	Ala	Thr	Thr	Val	Asn	Gly	Val	Asn	Gly	Val	Lys	Thr	Lys
6410						6415					6420			
Ala	Gln	Asp	Leu	Asp	Gly	Ala	Met	Gln	Arg	Leu	Gln	Ser	Ala	Ile
6425						6430					6435			
Ala	Asn	Lys	Asp	Gln	Thr	Lys	Ala	Ser	Glu	Asn	Tyr	Ile	Asp	Ala
6440						6445					6450			
Asp	Pro	Thr	Lys	Lys	Thr	Ala	Phe	Asp	Asn	Ala	Ile	Thr	Gln	Ala
6455						6460					6465			
Glu	Ser	Tyr	Leu	Asn	Lys	Asp	His	Gly	Ala	Asn	Lys	Asp	Lys	Gln
6470						6475					6480			
Ala	Val	Glu	Gln	Ala	Ile	Gln	Ser	Val	Thr	Ser	Thr	Glu	Asn	Ala
6485						6490					6495			
Leu	Asn	Gly	Asp	Ala	Asn	Leu	Gln	Arg	Ala	Lys	Thr	Glu	Ala	Ile
6500						6505					6510			
Gln	Ala	Ile	Asp	Asn	Leu	Thr	His	Leu	Asn	Thr	Pro	Gln	Lys	Thr
6515						6520					6525			
Ala	Leu	Lys	Gln	Gln	Val	Asn	Ala	Ala	Gln	Arg	Val	Ser	Gly	Val
6530						6535					6540			
Thr	Asp	Leu	Lys	Asn	Ser	Ala	Thr	Ser	Leu	Asn	Asn	Ala	Met	Asp
6545						6550					6555			
Gln	Leu	Lys	Gln	Ala	Ile	Ala	Asp	His	Asp	Thr	Ile	Val	Ala	Ser
6560						6565					6570			

-continued

Gly	Asn	Tyr	Thr	Asn	Ala	Ser	Pro	Asp	Lys	Gln	Gly	Ala	Tyr	Thr
6575						6580					6585			
Asp	Ala	Tyr	Asn	Ala	Ala	Lys	Asn	Ile	Val	Asn	Gly	Ser	Pro	Asn
6590						6595					6600			
Val	Ile	Thr	Asn	Ala	Ala	Asp	Val	Thr	Ala	Ala	Thr	Gln	Arg	Val
6605						6610					6615			
Asn	Asn	Ala	Glu	Thr	Gly	Leu	Asn	Gly	Asp	Thr	Asn	Leu	Ala	Thr
6620						6625					6630			
Ala	Lys	Gln	Gln	Ala	Lys	Asp	Ala	Leu	Arg	Gln	Met	Thr	His	Leu
6635						6640					6645			
Ser	Asp	Ala	Gln	Lys	Gln	Ser	Ile	Thr	Gly	Gln	Ile	Asp	Ser	Ala
6650						6655					6660			
Thr	Gln	Val	Thr	Gly	Val	Gln	Ser	Val	Lys	Asp	Asn	Ala	Thr	Asn
6665						6670					6675			
Leu	Asp	Asn	Ala	Met	Asn	Gln	Leu	Arg	Asn	Ser	Ile	Ala	Asn	Lys
6680						6685					6690			
Asp	Asp	Val	Lys	Ala	Ser	Gln	Pro	Tyr	Val	Asp	Ala	Asp	Arg	Asp
6695						6700					6705			
Lys	Gln	Asn	Ala	Tyr	Asn	Thr	Ala	Val	Thr	Asn	Ala	Glu	Asn	Ile
6710						6715					6720			
Ile	Asn	Ala	Thr	Ser	Gln	Pro	Thr	Leu	Asp	Pro	Ser	Ala	Val	Thr
6725						6730					6735			
Gln	Ala	Ala	Asn	Gln	Val	Ser	Thr	Asn	Lys	Thr	Ala	Leu	Asn	Gly
6740						6745					6750			
Ala	Gln	Asn	Leu	Ala	Asn	Lys	Lys	Gln	Glu	Thr	Thr	Ala	Asn	Ile
6755						6760					6765			
Asn	Gln	Leu	Ser	His	Leu	Asn	Asn	Ala	Gln	Lys	Gln	Asp	Leu	Asn
6770						6775					6780			
Thr	Gln	Val	Thr	Asn	Ala	Pro	Asn	Ile	Ser	Thr	Val	Asn	Gln	Val
6785						6790					6795			
Lys	Thr	Lys	Ala	Glu	Gln	Leu	Asp	Gln	Ala	Met	Glu	Arg	Leu	Ile
6800						6805					6810			
Asn	Gly	Ile	Gln	Asp	Lys	Asp	Gln	Val	Lys	Gln	Ser	Val	Asn	Phe
6815						6820					6825			
Thr	Asp	Ala	Asp	Pro	Glu	Lys	Gln	Thr	Ala	Tyr	Asn	Asn	Ala	Val
6830						6835					6840			
Thr	Ala	Ala	Glu	Asn	Ile	Ile	Asn	Gln	Ala	Asn	Gly	Thr	Asn	Ala
6845						6850					6855			
Asn	Gln	Ser	Gln	Val	Glu	Ala	Ala	Leu	Ser	Thr	Val	Thr	Thr	Thr
6860						6865					6870			
Lys	Gln	Ala	Leu	Asn	Gly	Asp	Arg	Lys	Val	Thr	Asp	Ala	Lys	Asn
6875						6880					6885			
Asn	Ala	Asn	Gln	Thr	Leu	Ser	Thr	Leu	Asp	Asn	Leu	Asn	Asn	Ala
6890						6895					6900			
Gln	Lys	Gly	Ala	Val	Thr	Gly	Asn	Ile	Asn	Gln	Ala	His	Thr	Val
6905						6910					6915			
Ala	Glu	Val	Thr	Gln	Ala	Ile	Gln	Thr	Ala	Gln	Glu	Leu	Asn	Thr
6920						6925					6930			
Ala	Met	Gly	Asn	Leu	Lys	Asn	Ser	Leu	Asn	Asp	Lys	Asp	Thr	Thr
6935						6940					6945			
Leu	Gly	Ser	Gln	Asn	Phe	Ala	Asp	Ala	Asp	Pro	Glu	Lys	Lys	Asn
6950						6955					6960			
Ala	Tyr	Asn	Glu	Ala	Val	His	Asn	Ala	Glu	Asn	Ile	Leu	Asn	Lys

-continued

6965	6970	6975
Ser Thr Gly Thr Asn Val Pro Lys Asp Gln Val Glu Ala Ala Met 6980	6985	6990
Asn Gln Val Asn Ala Thr Lys Ala Ala Leu Asn Gly Thr Gln Asn 6995	7000	7005
Leu Glu Lys Ala Lys Gln His Ala Asn Thr Ala Ile Asp Gly Leu 7010	7015	7020
Ser His Leu Thr Asn Ala Gln Lys Glu Ala Leu Lys Gln Leu Val 7025	7030	7035
Gln Gln Ser Thr Thr Val Ala Glu Ala Gln Gly Asn Glu Gln Lys 7040	7045	7050
Ala Asn Asn Val Asp Ala Ala Met Asp Lys Leu Arg Gln Ser Ile 7055	7060	7065
Ala Asp Asn Ala Thr Thr Lys Gln Asn Gln Asn Tyr Thr Asp Ala 7070	7075	7080
Ser Gln Asn Lys Lys Asp Ala Tyr Asn Asn Ala Val Thr Thr Ala 7085	7090	7095
Gln Gly Ile Ile Asp Gln Thr Thr Ser Pro Thr Leu Asp Pro Thr 7100	7105	7110
Val Ile Asn Gln Ala Ala Gly Gln Val Ser Thr Thr Lys Asn Ala 7115	7120	7125
Leu Asn Gly Asn Glu Asn Leu Glu Ala Ala Lys Gln Gln Ala Ser 7130	7135	7140
Gln Ser Leu Gly Ser Leu Asp Asn Leu Asn Asn Ala Gln Lys Gln 7145	7150	7155
Thr Val Thr Asp Gln Ile Asn Gly Ala His Thr Val Asp Glu Ala 7160	7165	7170
Asn Gln Ile Lys Gln Asn Ala Gln Asn Leu Asn Thr Ala Met Gly 7175	7180	7185
Asn Leu Lys Gln Ala Ile Ala Asp Lys Asp Ala Thr Lys Ala Thr 7190	7195	7200
Val Asn Phe Thr Asp Ala Asp Gln Ala Lys Gln Gln Ala Tyr Asn 7205	7210	7215
Thr Ala Val Thr Asn Ala Glu Asn Ile Ser Lys Ala Asn Gly Asn 7220	7225	7230
Ala Thr Gln Ala Glu Val Glu Gln Ala Ile Lys Gln Val Asn Ala 7235	7240	7245
Ala Lys Gln Ala Leu Asn Gly Asn Ala Asn Val Gln His Ala Lys 7250	7255	7260
Asp Glu Ala Thr Ala Leu Ile Asn Ser Ser Asn Asp Leu Asn Gln 7265	7270	7275
Ala Gln Lys Asp Ala Leu Lys Gln Gln Val Gln Asn Ala Thr Thr 7280	7285	7290
Val Ala Gly Val Asn Asn Val Lys Gln Thr Ala Gln Glu Leu Asn 7295	7300	7305
Asn Ala Met Thr Gln Leu Lys Gln Gly Ile Ala Asp Lys Glu Gln 7310	7315	7320
Thr Lys Ala Asp Gly Asn Phe Val Asn Ala Asp Pro Asp Lys Gln 7325	7330	7335
Asn Ala Tyr Asn Gln Ala Val Ala Lys Ala Glu Ala Leu Ile Ser 7340	7345	7350
Ala Thr Pro Asp Val Val Val Thr Pro Ser Glu Ile Thr Ala Ala 7355	7360	7365



-continued

Gly	Asn	His	Asn	Leu	Gln	Val	Ala	Lys	Asp	Asn	Ala	Asn	Thr	Ala
7760						7765					7770			
Ile	Asp	Gln	Leu	Pro	Asn	Leu	Asn	Gln	Pro	Gln	Lys	Thr	Ala	Leu
7775						7780					7785			
Lys	Asp	Gln	Val	Ser	His	Ala	Glu	Leu	Val	Thr	Gly	Val	Asn	Ala
7790						7795					7800			
Ile	Lys	Gln	Asn	Ala	Asp	Ala	Leu	Asn	Asn	Ala	Met	Gly	Thr	Leu
7805						7810					7815			
Lys	Gln	Gln	Ile	Gln	Ala	Asn	Ser	Gln	Val	Pro	Gln	Ser	Val	Asp
7820						7825					7830			
Phe	Thr	Gln	Ala	Asp	Gln	Asp	Lys	Gln	Gln	Ala	Tyr	Asn	Asn	Ala
7835						7840					7845			
Ala	Asn	Gln	Ala	Gln	Gln	Ile	Ala	Asn	Gly	Ile	Pro	Thr	Pro	Val
7850						7855					7860			
Leu	Thr	Pro	Asp	Thr	Val	Thr	Gln	Ala	Val	Thr	Thr	Met	Asn	Gln
7865						7870					7875			
Ala	Lys	Asp	Ala	Leu	Asn	Gly	Asp	Glu	Lys	Leu	Ala	Gln	Ala	Lys
7880						7885					7890			
Gln	Glu	Ala	Leu	Ala	Asn	Leu	Asp	Thr	Leu	Arg	Asp	Leu	Asn	Gln
7895						7900					7905			
Pro	Gln	Arg	Asp	Ala	Leu	Arg	Asn	Gln	Ile	Asn	Gln	Ala	Gln	Ala
7910						7915					7920			
Leu	Ala	Thr	Val	Glu	Gln	Thr	Lys	Gln	Asn	Ala	Gln	Asn	Val	Asn
7925						7930					7935			
Thr	Ala	Met	Ser	Asn	Leu	Lys	Gln	Gly	Ile	Ala	Asn	Lys	Asp	Thr
7940						7945					7950			
Val	Lys	Ala	Ser	Glu	Asn	Tyr	His	Asp	Ala	Asp	Ala	Asp	Lys	Gln
7955						7960					7965			
Thr	Ala	Tyr	Thr	Asn	Ala	Val	Ser	Gln	Ala	Glu	Gly	Ile	Ile	Asn
7970						7975					7980			
Gln	Thr	Thr	Asn	Pro	Thr	Leu	Asn	Pro	Asp	Glu	Ile	Thr	Arg	Ala
7985						7990					7995			
Leu	Thr	Gln	Val	Thr	Asp	Ala	Lys	Asn	Gly	Leu	Asn	Gly	Glu	Ala
8000						8005					8010			
Lys	Leu	Ala	Thr	Glu	Lys	Gln	Asn	Ala	Lys	Asp	Ala	Val	Ser	Gly
8015						8020					8025			
Met	Thr	His	Leu	Asn	Asp	Ala	Gln	Lys	Gln	Ala	Leu	Lys	Gly	Gln
8030						8035					8040			
Ile	Asp	Gln	Ser	Pro	Glu	Ile	Ala	Thr	Val	Asn	Gln	Val	Lys	Gln
8045						8050					8055			
Thr	Ala	Thr	Ser	Leu	Asp	Gln	Ala	Met	Asp	Gln	Leu	Ser	Gln	Ala
8060						8065					8070			
Ile	Asn	Asp	Lys	Ala	Gln	Thr	Leu	Ala	Asp	Gly	Asn	Tyr	Leu	Asn
8075						8080					8085			
Ala	Asp	Pro	Asp	Lys	Gln	Asn	Ala	Tyr	Lys	Gln	Ala	Val	Ala	Lys
8090						8095					8100			
Ala	Glu	Ala	Leu	Leu	Asn	Lys	Gln	Ser	Gly	Thr	Asn	Glu	Val	Gln
8105						8110					8115			
Ala	Gln	Val	Glu	Ser	Ile	Thr	Asn	Glu	Val	Asn	Ala	Ala	Lys	Gln
8120						8125					8130			
Ala	Leu	Asn	Gly	Asn	Asp	Asn	Leu	Ala	Asn	Ala	Lys	Gln	Gln	Ala
8135						8140					8145			
Lys	Gln	Gln	Leu	Ala	Asn	Leu	Thr	His	Leu	Asn	Asp	Ala	Gln	Lys



-continued

---

Thr	Ala	Val	Lys	Gln	Asp	Leu	Thr	Glu	Ala	Gln	Ala	Leu	Asp	Gln
8555						8560					8565			
Leu	Met	Asp	Ala	Leu	Gln	Gln	Ser	Ile	Ala	Asp	Lys	Asp	Ala	Thr
8570						8575					8580			
Arg	Ala	Ser	Ser	Ala	Tyr	Val	Asn	Ala	Glu	Pro	Asn	Lys	Lys	Gln
8585						8590					8595			
Ser	Tyr	Asp	Glu	Ala	Val	Gln	Asn	Ala	Glu	Ser	Ile	Ile	Ala	Gly
8600						8605					8610			
Leu	Asn	Asn	Pro	Thr	Ile	Asn	Lys	Gly	Asn	Val	Ser	Ser	Ala	Thr
8615						8620					8625			
Gln	Ala	Val	Ile	Ser	Ser	Lys	Asn	Ala	Leu	Asp	Gly	Val	Glu	Arg
8630						8635					8640			
Leu	Ala	Gln	Asp	Lys	Gln	Thr	Ala	Gly	Asn	Ser	Leu	Asn	His	Leu
8645						8650					8655			
Asp	Gln	Leu	Thr	Pro	Ala	Gln	Gln	Gln	Ala	Leu	Glu	Asn	Gln	Ile
8660						8665					8670			
Asn	Asn	Ala	Thr	Thr	Arg	Gly	Glu	Val	Ala	Gln	Lys	Leu	Thr	Glu
8675						8680					8685			
Ala	Gln	Ala	Leu	Asn	Gln	Ala	Met	Glu	Ala	Leu	Arg	Asn	Ser	Ile
8690						8695					8700			
Gln	Asp	Gln	Gln	Gln	Thr	Glu	Ala	Gly	Ser	Lys	Phe	Ile	Asn	Glu
8705						8710					8715			
Asp	Lys	Pro	Gln	Lys	Asp	Ala	Tyr	Gln	Ala	Ala	Val	Gln	Asn	Ala
8720						8725					8730			
Lys	Asp	Leu	Ile	Asn	Gln	Thr	Asn	Asn	Pro	Thr	Leu	Asp	Lys	Ala
8735						8740					8745			
Gln	Val	Glu	Gln	Leu	Thr	Gln	Ala	Val	Asn	Gln	Ala	Lys	Asp	Asn
8750						8755					8760			
Leu	His	Gly	Asp	Gln	Lys	Leu	Ala	Asp	Asp	Lys	Gln	His	Ala	Val
8765						8770					8775			
Thr	Asp	Leu	Asn	Gln	Leu	Asn	Gly	Leu	Asn	Asn	Pro	Gln	Arg	Gln
8780						8785					8790			
Ala	Leu	Glu	Ser	Gln	Ile	Asn	Asn	Ala	Ala	Thr	Arg	Gly	Glu	Val
8795						8800					8805			
Ala	Gln	Lys	Leu	Ala	Glu	Ala	Lys	Ala	Leu	Asp	Gln	Ala	Met	Gln
8810						8815					8820			
Ala	Leu	Arg	Asn	Ser	Ile	Gln	Asp	Gln	Gln	Gln	Thr	Glu	Ser	Gly
8825						8830					8835			
Ser	Lys	Phe	Ile	Asn	Glu	Asp	Lys	Pro	Gln	Lys	Asp	Ala	Tyr	Gln
8840						8845					8850			
Ala	Ala	Val	Gln	Asn	Ala	Lys	Asp	Leu	Ile	Asn	Gln	Thr	Gly	Asn
8855						8860					8865			
Pro	Thr	Leu	Asp	Lys	Ser	Gln	Val	Glu	Gln	Leu	Thr	Gln	Ala	Val
8870						8875					8880			
Thr	Thr	Ala	Lys	Asp	Asn	Leu	His	Gly	Asp	Gln	Lys	Leu	Ala	Arg
8885						8890					8895			
Asp	Gln	Gln	Gln	Ala	Val	Thr	Thr	Val	Asn	Ala	Leu	Pro	Asn	Leu
8900						8905					8910			
Asn	His	Ala	Gln	Gln	Gln	Ala	Leu	Thr	Asp	Ala	Ile	Asn	Ala	Ala
8915						8920					8925			
Pro	Thr	Arg	Thr	Glu	Val	Ala	Gln	His	Val	Gln	Thr	Ala	Thr	Glu
8930						8935					8940			

-continued

Leu 8945	Asp	His	Ala	Met	Glu	Thr	Leu	Lys	Asn	Lys	Val	Asp	Gln	Val
							8950				8955			
Asn 8960	Thr	Asp	Lys	Ala	Gln	Pro	Asn	Tyr	Thr	Glu	Ala	Ser	Thr	Asp
							8965				8970			
Lys 8975	Lys	Glu	Ala	Val	Asp	Gln	Ala	Leu	Gln	Ala	Ala	Glu	Ser	Ile
							8980				8985			
Thr 8990	Asp	Pro	Thr	Asn	Gly	Ser	Asn	Ala	Asn	Lys	Asp	Ala	Val	Asp
							8995				9000			
Gln 9005	Val	Leu	Thr	Lys	Leu	Gln	Glu	Lys	Glu	Asn	Glu	Leu	Asn	Gly
							9010				9015			
Asn 9020	Glu	Arg	Val	Ala	Glu	Ala	Lys	Thr	Gln	Ala	Lys	Gln	Thr	Ile
							9025				9030			
Asp 9035	Gln	Leu	Thr	His	Leu	Asn	Ala	Asp	Gln	Ile	Ala	Thr	Ala	Lys
							9040				9045			
Gln 9050	Asn	Ile	Asp	Gln	Ala	Thr	Lys	Leu	Gln	Pro	Ile	Ala	Glu	Leu
							9055				9060			
Val 9065	Asp	Gln	Ala	Thr	Gln	Leu	Asn	Gln	Ser	Met	Asp	Gln	Leu	Gln
							9070				9075			
Gln 9080	Ala	Val	Asn	Glu	His	Ala	Asn	Val	Glu	Gln	Thr	Val	Asp	Tyr
							9085				9090			
Thr 9095	Gln	Ala	Asp	Ser	Asp	Lys	Gln	Asn	Ala	Tyr	Lys	Gln	Ala	Ile
							9100				9105			
Ala 9110	Asp	Ala	Glu	Asn	Val	Leu	Lys	Gln	Asn	Ala	Asn	Lys	Gln	Gln
							9115				9120			
Val 9125	Asp	Gln	Ala	Leu	Gln	Asn	Ile	Leu	Asn	Ala	Lys	Gln	Ala	Leu
							9130				9135			
Asn 9140	Gly	Asp	Glu	Arg	Val	Ala	Leu	Ala	Lys	Thr	Asn	Gly	Lys	His
							9145				9150			
Asp 9155	Ile	Asp	Gln	Leu	Asn	Ala	Leu	Asn	Asn	Ala	Gln	Gln	Asp	Gly
							9160				9165			
Phe 9170	Lys	Gly	Arg	Ile	Asp	Gln	Ser	Asn	Asp	Leu	Asn	Gln	Ile	Gln
							9175				9180			
Gln 9185	Ile	Val	Asp	Glu	Ala	Lys	Ala	Leu	Asn	Arg	Ala	Met	Asp	Gln
							9190				9195			
Leu 9200	Ser	Gln	Glu	Ile	Thr	Asp	Asn	Glu	Gly	Arg	Thr	Lys	Gly	Ser
							9205				9210			
Thr 9215	Asn	Tyr	Val	Asn	Ala	Asp	Thr	Gln	Val	Lys	Gln	Val	Tyr	Asp
							9220				9225			
Glu 9230	Thr	Val	Asp	Lys	Ala	Lys	Gln	Ala	Leu	Asp	Lys	Ser	Thr	Gly
							9235				9240			
Gln 9245	Asn	Leu	Thr	Ala	Lys	Gln	Val	Ile	Lys	Leu	Asn	Asp	Ala	Val
							9250				9255			
Thr 9260	Ala	Ala	Lys	Lys	Ala	Leu	Asn	Gly	Glu	Glu	Arg	Leu	Asn	Asn
							9265				9270			
Arg 9275	Lys	Ala	Glu	Ala	Leu	Gln	Arg	Leu	Asp	Gln	Leu	Thr	His	Leu
							9280				9285			
Asn 9290	Asn	Ala	Gln	Arg	Gln	Leu	Ala	Ile	Gln	Gln	Ile	Asn	Asn	Ala
							9295				9300			
Glu 9305	Thr	Leu	Asn	Lys	Ala	Ser	Arg	Ala	Ile	Asn	Arg	Ala	Thr	Lys
							9310				9315			
Leu 9320	Asp	Asn	Ala	Met	Gly	Ala	Val	Gln	Gln	Tyr	Ile	Asp	Glu	Gln
							9325				9330			
His 9335	Leu	Gly	Val	Ile	Ser	Ser	Thr	Asn	Tyr	Ile	Asn	Ala	Asp	Asp

-continued

---

9335	9340	9345
Asn Leu Lys Ala Asn Tyr Asp 9350	Asn Ala Ile Ala 9355	Asn Ala Ala His 9360
Glu Leu Asp Lys Val Gln Gly 9365	Asn Ala Ile Ala 9370	Lys Ala Glu Ala 9375
Glu Gln Leu Lys Gln Asn Ile 9380	Ile Asp Ala Gln 9385	Asn Ala Leu Asn 9390
Gly Asp Gln Asn Leu Ala Asn 9395	Ala Lys Asp Lys 9400	Ala Asn Ala Phe 9405
Val Asn Ser Leu Asn Gly Leu 9410	Asn Gln Gln Gln 9415	Gln Asp Leu Ala 9420
His Lys Ala Ile Asn Asn Ala 9425	Asp Thr Val Ser 9430	Asp Val Thr Asp 9435
Ile Val Asn Asn Gln Ile Asp 9440	Leu Asn Asp Ala 9445	Met Glu Thr Leu 9450
Lys His Leu Val Asp Asn Glu 9455	Ile Pro Asn Ala 9460	Glu Gln Thr Val 9465
Asn Tyr Gln Asn Ala Asp Asp 9470	Asn Ala Lys Thr 9475	Asn Phe Asp Asp 9480
Ala Lys Arg Leu Ala Asn Thr 9485	Leu Leu Asn Ser 9490	Asp Asn Thr Asn 9495
Val Asn Asp Ile Asn Gly Ala 9500	Ile Gln Ala Val 9505	Asn Asp Ala Ile 9510
His Asn Leu Asn Gly Asp Gln 9515	Arg Leu Gln Asp 9520	Ala Lys Asp Lys 9525
Ala Ile Gln Ser Ile Asn Gln 9530	Ala Leu Ala Asn 9535	Lys Leu Lys Glu 9540
Ile Glu Ala Ser Asn Ala Thr 9545	Asp Gln Asp Lys 9550	Leu Ile Ala Lys 9555
Asn Lys Ala Glu Glu Leu Ala 9560	Asn Ser Ile Ile 9565	Asn Asn Ile Asn 9570
Lys Ala Thr Ser Asn Gln Ala 9575	Val Ser Gln Val 9580	Gln Thr Ala Gly 9585
Asn His Ala Ile Glu Gln Val 9590	His Ala Asn Glu 9595	Ile Pro Lys Ala 9600
Lys Ile Asp Ala Asn Lys Asp 9605	Val Asp Lys Gln 9610	Val Gln Ala Leu 9615
Ile Asp Glu Ile Asp Arg Asn 9620	Pro Asn Leu Thr 9625	Asp Lys Glu Lys 9630
Gln Ala Leu Lys Asp Arg Ile 9635	Asn Gln Ile Leu 9640	Gln Gln Gly His 9645
Asn Gly Ile Asn Asn Ala Met 9650	Thr Lys Glu Glu 9655	Ile Glu Gln Ala 9660
Lys Ala Gln Leu Ala Gln Ala 9665	Leu Gln Asp Ile 9670	Lys Asp Leu Val 9675
Lys Ala Lys Glu Asp Ala Lys 9680	Gln Asp Val Asp 9685	Lys Gln Val Gln 9690
Ala Leu Ile Asp Glu Ile Asp 9695	Gln Asn Pro Asn 9700	Leu Thr Asp Lys 9705
Glu Lys Gln Ala Leu Lys Tyr 9710	Arg Ile Asn Gln 9715	Ile Leu Gln Gln 9720
Gly His Asn Asp Ile Asn Asn 9725	Ala Leu Thr Lys 9730	Glu Glu Ile Glu 9735

-continued

---

Gln Ala	Lys Ala	Gln Leu	Ala	Gln Ala	Leu Gln	Asp	Ile Lys	Asp	
9740			9745			9750			
Leu Val	Lys Ala	Lys Glu	Asp	Ala Lys	Asn Ala	Ile	Lys Ala	Leu	
9755			9760			9765			
Ala Asn	Ala Lys	Arg Asp	Gln	Ile Asn	Ser Asn	Pro	Asp Leu	Thr	
9770			9775			9780			
Pro Glu	Gln Lys	Ala Lys	Ala	Leu Lys	Glu Ile	Asp	Glu Ala	Glu	
9785			9790			9795			
Lys Arg	Ala Leu	Gln Asn	Val	Glu Asn	Ala Gln	Thr	Ile Asp	Gln	
9800			9805			9810			
Leu Asn	Arg Gly	Leu Asn	Leu	Gly Leu	Asp Asp	Ile	Arg Asn	Thr	
9815			9820			9825			
His Val	Trp Glu	Val Asp	Glu	Gln Pro	Ala Val	Asn	Glu Ile	Phe	
9830			9835			9840			
Glu Ala	Thr Pro	Glu Gln	Ile	Leu Val	Asn Gly	Glu	Leu Ile	Val	
9845			9850			9855			
His Arg	Asp Asp	Ile Ile	Thr	Glu Gln	Asp Ile	Leu	Ala His	Ile	
9860			9865			9870			
Asn Leu	Ile Asp	Gln Leu	Ser	Ala Glu	Val Ile	Asp	Thr Pro	Ser	
9875			9880			9885			
Thr Ala	Thr Ile	Ser Asp	Ser	Leu Thr	Ala Lys	Val	Glu Val	Thr	
9890			9895			9900			
Leu Leu	Asp Gly	Ser Lys	Val	Ile Val	Asn Val	Pro	Val Lys	Val	
9905			9910			9915			
Val Glu	Lys Glu	Leu Ser	Val	Val Lys	Gln Gln	Ala	Ile Glu	Ser	
9920			9925			9930			
Ile Glu	Asn Ala	Ala Gln	Gln	Lys Ile	Asn Glu	Ile	Asn Asn	Ser	
9935			9940			9945			
Val Thr	Leu Thr	Leu Glu	Gln	Lys Glu	Ala Ala	Ile	Ala Glu	Val	
9950			9955			9960			
Asn Lys	Leu Lys	Gln Gln	Ala	Ile Asp	His Val	Asn	Asn Ala	Pro	
9965			9970			9975			
Asp Val	His Ser	Val Glu	Glu	Ile Gln	Gln Gln	Glu	Gln Ala	His	
9980			9985			9990			
Ile Glu	Gln Phe	Asn Pro	Glu	Gln Phe	Thr Ile	Glu	Gln Ala	Lys	
9995			10000			10005			
Ser Asn	Ala Ile	Lys Ser	Ile	Glu Asp	Ala Ile	Gln	His Met	Ile	
10010			10015			10020			
Asp Glu	Ile Lys	Ala Arg	Thr	Asp Leu	Thr Asp	Lys	Glu Lys	Gln	
10025			10030			10035			
Glu Ala	Ile Ala	Lys Leu	Asn	Gln Leu	Lys Glu	Gln	Ala Ile	Gln	
10040			10045			10050			
Ala Ile	Gln Arg	Ala Gln	Ser	Ile Asp	Glu Ile	Ser	Glu Gln	Leu	
10055			10060			10065			
Glu Gln	Phe Lys	Ala Gln	Met	Lys Ala	Ala Asn	Pro	Thr Ala	Lys	
10070			10075			10080			
Glu Leu	Ala Lys	Arg Lys	Gln	Glu Ala	Ile Ser	Arg	Ile Lys	Asp	
10085			10090			10095			
Phe Ser	Asn Glu	Lys Ile	Asn	Ser Ile	Arg Asn	Ser	Glu Ile	Gly	
10100			10105			10110			
Thr Ala	Asp Glu	Lys Gln	Ala	Ala Met	Asn Gln	Ile	Asn Glu	Ile	
10115			10120			10125			

-continued

---

Val Leu Glu Thr Ile Arg Asp Ile Asn Asn Ala His Thr Leu Gln  
 10130 10135 10140

Gln Val Glu Ala Ala Leu Asn Asn Gly Ile Ala Arg Ile Ser Ala  
 10145 10150 10155

Val Gln Ile Val Thr Ser Asp Arg Ala Lys Gln Ser Ser Ser Thr  
 10160 10165 10170

Gly Asn Glu Ser Asn Ser His Leu Thr Ile Gly Tyr Gly Thr Ala  
 10175 10180 10185

Asn His Pro Phe Asn Ser Ser Thr Ile Gly His Lys Lys Lys Leu  
 10190 10195 10200

Asp Glu Asp Asp Asp Ile Asp Pro Leu His Met Arg His Phe Ser  
 10205 10210 10215

Asn Asn Phe Gly Asn Val Ile Lys Asn Ala Ile Gly Val Val Gly  
 10220 10225 10230

Ile Ser Gly Leu Leu Ala Ser Phe Trp Phe Phe Ile Ala Lys Arg  
 10235 10240 10245

Arg Arg Lys Glu Asp Glu Glu Glu Leu Glu Ile Arg Asp Asn  
 10250 10255 10260

Asn Lys Asp Ser Ile Lys Glu Thr Leu Asp Asp Thr Lys His Leu  
 10265 10270 10275

Pro Leu Leu Phe Ala Lys Arg Arg Arg Lys Glu Asp Glu Glu Asp  
 10280 10285 10290

Val Thr Val Glu Glu Lys Asp Ser Leu Asn Asn Gly Glu Ser Leu  
 10295 10300 10305

Asp Lys Val Lys His Thr Pro Phe Phe Leu Pro Lys Arg Arg Arg  
 10310 10315 10320

Lys Glu Asp Glu Glu Asp Val Glu Val Thr Asn Glu Asn Thr Asp  
 10325 10330 10335

Glu Lys Val Leu Lys Asp Asn Glu His Ser Pro Leu Leu Phe Ala  
 10340 10345 10350

Lys Arg Arg Lys Asp Lys Glu Glu Asp Val Glu Thr Thr Thr Ser  
 10355 10360 10365

Ile Glu Ser Lys Asp Glu Asp Val Pro Leu Leu Leu Ala Lys Lys  
 10370 10375 10380

Lys Asn Gln Lys Asp Asn Gln Ser Lys Asp Lys Lys Ser Ala Ser  
 10385 10390 10395

Lys Asn Thr Ser Lys Lys Val Ala Ala Lys Lys Lys Lys Lys Lys  
 10400 10405 10410

Ala Lys Lys Asn Lys Lys  
 10415

<210> SEQ ID NO 25  
 <211> LENGTH: 340  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 25

Met Lys Lys Lys Leu Leu Val Leu Thr Met Ser Thr Leu Phe Ala Thr  
 1 5 10 15

Gln Ile Met Asn Ser Asn His Ala Lys Ala Ser Val Thr Glu Ser Val  
 20 25 30

Asp Lys Lys Phe Val Val Pro Glu Ser Gly Ile Asn Lys Ile Ile Pro  
 35 40 45

Ala Tyr Asp Glu Phe Lys Asn Ser Pro Lys Val Asn Val Ser Asn Leu  
 50 55 60

-continued

Thr Asp Asn Lys Asn Phe Val Ala Ser Glu Asp Lys Leu Asn Lys Ile  
 65 70 75 80  
 Ala Asp Ser Ser Ala Ala Ser Lys Ile Val Asp Lys Asn Phe Val Val  
 85 90 95  
 Pro Glu Ser Lys Leu Gly Asn Ile Val Pro Glu Tyr Lys Glu Ile Asn  
 100 105 110  
 Asn Arg Val Asn Val Ala Thr Asn Asn Pro Ala Ser Gln Gln Val Asp  
 115 120 125  
 Lys His Phe Val Ala Lys Gly Pro Glu Val Asn Arg Phe Ile Thr Gln  
 130 135 140  
 Asn Lys Val Asn His His Phe Ile Thr Thr Gln Thr His Tyr Lys Lys  
 145 150 155 160  
 Val Ile Thr Ser Tyr Lys Ser Thr His Val His Lys His Val Asn His  
 165 170 175  
 Ala Lys Asp Ser Ile Asn Lys His Phe Ile Val Lys Pro Ser Glu Ser  
 180 185 190  
 Pro Arg Tyr Thr His Pro Ser Gln Ser Leu Ile Ile Lys His His Phe  
 195 200 205  
 Ala Val Pro Gly Tyr His Ala His Lys Phe Val Thr Pro Gly His Ala  
 210 215 220  
 Ser Ile Lys Ile Asn His Phe Cys Val Val Pro Gln Ile Asn Ser Phe  
 225 230 235 240  
 Lys Val Ile Pro Pro Tyr Gly His Asn Ser His Arg Met His Val Pro  
 245 250 255  
 Ser Phe Gln Asn Asn Thr Thr Ala Thr His Gln Asn Ala Lys Val Asn  
 260 265 270  
 Lys Ala Tyr Asp Tyr Lys Tyr Phe Tyr Ser Tyr Lys Val Val Lys Gly  
 275 280 285  
 Val Lys Lys Tyr Phe Ser Phe Ser Gln Ser Asn Gly Tyr Lys Ile Gly  
 290 295 300  
 Lys Pro Ser Leu Asn Ile Lys Asn Val Asn Tyr Gln Tyr Ala Val Pro  
 305 310 315 320  
 Ser Tyr Ser Pro Thr His Tyr Val Pro Glu Phe Lys Gly Ser Leu Pro  
 325 330 335  
 Ala Pro Arg Val  
 340

&lt;210&gt; SEQ ID NO 26

&lt;211&gt; LENGTH: 130

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 26

Met Asn Phe Asn Asp Ile Glu Thr Met Val Lys Ser Lys Phe Lys Asp  
 1 5 10 15  
 Ile Lys Lys His Ala Glu Glu Ile Ala His Glu Ile Glu Val Arg Ser  
 20 25 30  
 Gly Tyr Leu Arg Lys Ala Glu Gln Tyr Lys Arg Leu Glu Phe Asn Leu  
 35 40 45  
 Ser Phe Ala Leu Asp Asp Ile Glu Ser Thr Ala Lys Asp Val Gln Thr  
 50 55 60  
 Ala Lys Ser Ser Ala Asn Lys Asp Ser Val Thr Val Lys Gly Lys Ala  
 65 70 75 80  
 Pro Asn Thr Leu Tyr Ile Glu Lys Arg Asn Leu Met Lys Gln Lys Leu





-continued

---

Asn Lys Pro Ser Thr Val Val Ser Thr Lys Val Asn Glu Thr Arg Asp  
 65 70 75 80  
 Val Asp Thr Gln Gln Ala Ser Thr Gln Lys Pro Thr His Thr Ala Thr  
 85 90 95  
 Phe Lys Leu Ser Asn Ala Lys Thr Ala Ser Leu Ser Pro Arg Met Phe  
 100 105 110  
 Ala Ala Asn Ala Pro Gln Thr Thr Thr His Lys Ile Leu His Thr Asn  
 115 120 125  
 Asp Ile His Gly Arg Leu Ala Glu Glu Lys Gly Arg Val Ile Gly Met  
 130 135 140  
 Ala Lys Leu Lys Thr Val Lys Glu Gln Glu Lys Pro Asp Leu Met Leu  
 145 150 155 160  
 Asp Ala Gly Asp Ala Phe Gln Gly Leu Pro Leu Ser Asn Gln Ser Lys  
 165 170 175  
 Gly Glu Glu Met Ala Lys Ala Met Asn Ala Val Gly Tyr Asp Ala Met  
 180 185 190  
 Ala Val Gly Asn His Glu Phe Asp Phe Gly Tyr Asp Gln Leu Lys Lys  
 195 200 205  
 Leu Glu Gly Met Leu Asp Phe Pro Met Leu Ser Thr Asn Val Tyr Lys  
 210 215 220  
 Asp Gly Lys Arg Ala Phe Lys Pro Ser Thr Ile Val Thr Lys Asn Gly  
 225 230 235 240  
 Ile Arg Tyr Gly Ile Ile Gly Val Thr Thr Pro Glu Thr Lys Thr Lys  
 245 250 255  
 Thr Arg Pro Glu Gly Ile Lys Gly Val Glu Phe Arg Asp Pro Leu Gln  
 260 265 270  
 Ser Val Thr Ala Glu Met Met Arg Ile Tyr Lys Asp Val Asp Thr Phe  
 275 280 285  
 Val Val Ile Ser His Leu Gly Ile Asp Pro Ser Thr Gln Glu Thr Trp  
 290 295 300  
 Arg Gly Asp Tyr Leu Val Lys Gln Leu Ser Gln Asn Pro Gln Leu Lys  
 305 310 315 320  
 Lys Arg Ile Thr Val Ile Asp Gly His Ser His Thr Val Leu Gln Asn  
 325 330 335  
 Gly Gln Ile Tyr Asn Asn Asp Ala Leu Ala Gln Thr Gly Thr Ala Leu  
 340 345 350  
 Ala Asn Ile Gly Lys Ile Thr Phe Asn Tyr Arg Asn Gly Glu Val Ser  
 355 360 365  
 Asn Ile Lys Pro Ser Leu Ile Asn Val Lys Asp Val Glu Asn Val Thr  
 370 375 380  
 Pro Asn Lys Ala Leu Ala Glu Gln Ile Asn Gln Ala Asp Gln Thr Phe  
 385 390 395 400  
 Arg Ala Gln Thr Ala Glu Val Ile Ile Pro Asn Asn Thr Ile Asp Phe  
 405 410 415  
 Lys Gly Glu Arg Asp Asp Val Arg Thr Arg Glu Thr Asn Leu Gly Asn  
 420 425 430  
 Ala Ile Ala Asp Ala Met Glu Ala Tyr Gly Val Lys Asn Phe Ser Lys  
 435 440 445  
 Lys Thr Asp Phe Ala Val Thr Asn Gly Gly Gly Ile Arg Ala Ser Ile  
 450 455 460  
 Ala Lys Gly Lys Val Thr Arg Tyr Asp Leu Ile Ser Val Leu Pro Phe  
 465 470 475 480



-continued

---

Met Arg Val Leu Asn Ser Ile Lys Val Glu Arg Pro Ser Thr Ala Lys  
           115  120  125

Leu Ile Asp Leu Arg Leu Asp Thr Arg Ile Ser Tyr Thr Glu Ser Gln  
       130  135  140

Ile Lys Val Tyr Arg Thr Lys Thr Gln Tyr Thr Asp Leu Leu Phe Leu  
       145  150  155  160

Tyr Leu Glu His Ala Phe Leu Ser Gln Asp Phe Phe Asp Ile Pro Ser  
   165  170  175

Ile His Ser Asp Leu Asp Asp Ile Leu Val Asn Met Phe Leu Tyr Leu  
   180  185  190

Pro Asn Phe Phe Gln Asn Gln Asn Ser Glu Asp Asn Met Tyr Leu Ala  
   195  200  205

Gln Arg Ile Met Tyr Gln Val Asp Asp Ile Leu Lys Glu Asp Met Leu  
       210  215  220

Asn Glu Tyr Tyr Tyr Leu Pro Lys Thr Leu Tyr Asn Thr Leu Ala Ser  
       225  230  235  240

Pro Glu Phe Asp Asp Leu Lys Arg Thr Asp Ala Ser Gln Val Asp Gly  
   245  250  255

Gln Asp Asp Thr Ser Glu Asp Asp Asp Asn Glu Ser Glu Lys Ala Asp  
   260  265  270

Ser Lys Ser Ala Asp Ser Glu Ser Lys Gly Gly Ala Tyr Leu Glu Met  
   275  280  285

Glu Leu His Glu Gly Gln Asn Ser Glu Thr Leu Gly Asn Asp Glu Ala  
       290  295  300

Arg Glu Gly Asp Ala Thr Asp Asp Met Thr Asp Met Met Thr Lys Lys  
       305  310  315  320

Gly Lys Gly Ser Asn Asp Thr Leu Asn Arg Glu Glu Gly Asp Ala Val  
   325  330  335

Gly Gln Ser Gln Ala Phe Gln Leu Asp Gly Val Asn Lys Asn Val Glu  
   340  345  350

Ile Lys Trp Gln Ile Pro Glu Ile Glu Pro Gln Tyr Val Leu Glu Tyr  
   355  360  365

Gln Glu Ser Lys Gln Asp Val Gln Tyr Glu Ile Lys Asp Leu Ile Gln  
       370  375  380

Ile Ile Lys Lys Thr Ile Glu Arg Glu Gln Arg Asp Ala Arg Phe Asn  
       385  390  395  400

Leu Thr Lys Gly Arg Leu Gln Lys Asp Leu Ile Asn Trp Phe Ile Asp  
   405  410  415

Asp Gln Tyr Lys Leu Phe Tyr Lys Lys Gln Asp Leu Ser Lys Ser Phe  
   420  425  430

Asp Ala Thr Phe Thr Leu Leu Ile Asp Ala Ser Ala Ser Met His Asp  
   435  440  445

Lys Met Ala Glu Thr Lys Lys Gly Val Val Leu Phe His Glu Thr Leu  
       450  455  460

Lys Ala Leu Asn Ile Lys His Glu Ile Leu Ser Phe Ser Glu Asp Ala  
       465  470  475  480

Phe Asp Ser Asp Glu His Ala Gln Pro Asn Ile Ile Asn Glu Ile Ile  
   485  490  495

Asn Tyr Asp Tyr Ser Thr Phe Glu Lys Asp Gly Pro Arg Ile Met Ala  
   500  505  510

Leu Glu Pro Gln Asp Asp Asn Arg Asp Gly Val Ala Ile Arg Val Ala  
       515  520  525

-continued

---

Ser Glu Arg Leu Met Arg Arg Asn Gln His Gln Arg Phe Leu Ile Val  
 530 535 540

Phe Ser Asp Gly Glu Pro Ser Ala Phe Asn Tyr Ser Gln Asp Gly Ile  
 545 550 555 560

Ile Asp Thr Tyr Glu Ala Val Glu Met Ser Arg Lys Phe Gly Ile Glu  
 565 570 575

Val Phe Asn Val Phe Leu Ser Gln Asp Pro Ile Thr Glu Asp Val Glu  
 580 585 590

Gln Thr Ile His Asn Ile Tyr Gly Gln Tyr Ala Ile Phe Val Glu Gly  
 595 600 605

Val Ala His Leu Pro Gly His Leu Ser Pro Leu Leu Lys Lys Leu Leu  
 610 615 620

Leu Lys Ser Leu  
 625

<210> SEQ ID NO 30  
 <211> LENGTH: 154  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 30

Ala Glu Ile Asn Lys Gln Thr Thr Ser Gln Gly Val Thr Thr Glu Lys  
 1 5 10 15

Asn Asn Gly Ile Ala Val Leu Glu Gln Asp Val Ile Thr Pro Thr Val  
 20 25 30

Lys Pro Gln Ala Lys Gln Asp Ile Ile Gln Ala Val Thr Thr Arg Lys  
 35 40 45

Gln Gln Ile Lys Lys Ser Asn Ala Ser Leu Gln Asp Glu Lys Asp Val  
 50 55 60

Ala Asn Asp Lys Ile Gly Lys Ile Glu Thr Lys Ala Ile Lys Asp Ile  
 65 70 75 80

Asp Ala Ala Thr Thr Asn Ala Gln Val Glu Ala Ile Lys Thr Lys Ala  
 85 90 95

Ile Asn Asp Ile Asn Gln Thr Thr Pro Ala Thr Thr Ala Lys Ala Ala  
 100 105 110

Ala Leu Glu Glu Phe Asp Glu Val Val Gln Ala Gln Ile Asp Gln Ala  
 115 120 125

Pro Leu Asn Pro Asp Thr Thr Asn Glu Glu Val Ala Glu Ala Ile Glu  
 130 135 140

Arg Ile Asn Ala Ala Lys Val Ser Gly Val  
 145 150

<210> SEQ ID NO 31  
 <211> LENGTH: 584  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 31

Met Lys Phe Lys Ser Leu Ile Thr Thr Thr Leu Ala Leu Gly Val Leu  
 1 5 10 15

Ala Ser Thr Gly Ala Asn Phe Asn Asn Asn Glu Ala Ser Ala Ala Ala  
 20 25 30

Lys Pro Leu Asp Lys Ser Ser Ser Ser Leu His His Gly Tyr Ser Lys  
 35 40 45

Val His Val Pro Tyr Ala Ile Thr Val Asn Gly Thr Ser Gln Asn Ile  
 50 55 60

-continued

---

Leu Ser Ser Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp  
 65 70 80

Leu Glu Asp Arg Val Lys Ser Val Leu Lys Ser Asp Arg Gly Ile Ser  
 85 90 95

Asp Ile Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Tyr Phe  
 100 105 110

Lys Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ala Gly Ile Tyr Thr  
 115 120 125

Ala Asp Leu Ile Asn Thr Ser Glu Ile Lys Ala Ile Asn Ile Asn Val  
 130 135 140

Asp Thr Lys Lys Gln Val Glu Asp Lys Lys Lys Asp Lys Ala Asn Tyr  
 145 150 155 160

Gln Val Pro Tyr Thr Ile Thr Val Asn Gly Thr Ser Gln Asn Ile Leu  
 165 170 175

Ser Asn Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp Leu  
 180 185 190

Glu Asp Lys Val Lys Ser Val Leu Glu Ser Asn Arg Gly Ile Thr Asp  
 195 200 205

Val Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Asn Phe Lys  
 210 215 220

Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ser Gly Ile Tyr Thr Ala  
 225 230 235 240

Asn Leu Ile Asn Ser Ser Asp Ile Lys Ser Ile Asn Ile Asn Val Asp  
 245 250 255

Thr Lys Lys His Ile Glu Asn Lys Ala Lys Arg Asn Tyr Gln Val Pro  
 260 265 270

Tyr Ser Ile Asn Leu Asn Gly Thr Ser Thr Asn Ile Leu Ser Asn Leu  
 275 280 285

Ser Phe Ser Asn Lys Pro Trp Thr Asn Tyr Lys Asn Leu Thr Ser Gln  
 290 295 300

Ile Lys Ser Val Leu Lys His Asp Arg Gly Ile Ser Glu Gln Asp Leu  
 305 310 315 320

Lys Tyr Ala Lys Lys Ala Tyr Tyr Thr Val Tyr Phe Lys Asn Gly Gly  
 325 330 335

Lys Arg Ile Leu Gln Leu Asn Ser Lys Asn Tyr Thr Ala Asn Leu Val  
 340 345 350

His Ala Lys Asp Val Lys Arg Ile Glu Ile Thr Val Lys Thr Gly Thr  
 355 360 365

Lys Ala Lys Ala Asp Arg Tyr Val Pro Tyr Thr Ile Ala Val Asn Gly  
 370 375 380

Thr Ser Thr Pro Ile Leu Ser Asp Leu Lys Phe Thr Gly Asp Pro Arg  
 385 390 395 400

Val Gly Tyr Lys Asp Ile Ser Lys Lys Val Lys Ser Val Leu Lys His  
 405 410 415

Asp Arg Gly Ile Gly Glu Arg Glu Leu Lys Tyr Ala Lys Lys Ala Thr  
 420 425 430

Tyr Thr Val His Phe Lys Asn Gly Thr Lys Lys Val Ile Asn Ile Asn  
 435 440 445

Ser Asn Ile Ser Gln Leu Asn Leu Leu Tyr Val Gln Asp Ile Lys Lys  
 450 455 460

Ile Asp Ile Asp Val Lys Thr Gly Thr Lys Ala Lys Ala Asp Ser Tyr  
 465 470 475 480

Val Pro Tyr Thr Ile Ala Val Asn Gly Thr Ser Thr Pro Ile Leu Ser

-continued

---

	485		490		495										
Lys	Leu	Lys	Ile	Ser	Asn	Lys	Gln	Leu	Ile	Ser	Tyr	Lys	Tyr	Leu	Asn
			500					505						510	
Asp	Lys	Val	Lys	Ser	Val	Leu	Lys	Ser	Glu	Arg	Gly	Ile	Ser	Asp	Leu
		515						520					525		
Asp	Leu	Lys	Phe	Ala	Lys	Gln	Ala	Lys	Tyr	Thr	Val	Tyr	Phe	Lys	Asn
	530					535					540				
Gly	Lys	Lys	Gln	Val	Val	Asn	Leu	Lys	Ser	Asp	Ile	Phe	Thr	Pro	Asn
	545					550				555					560
Leu	Phe	Ser	Ala	Lys	Asp	Ile	Lys	Lys	Ile	Asp	Ile	Asp	Val	Lys	Gln
				565					570					575	
Tyr	Thr	Lys	Ser	Lys	Lys	Asn	Lys								
				580											

&lt;210&gt; SEQ ID NO 32

&lt;211&gt; LENGTH: 508

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 32

Met	Lys	Asn	Lys	Leu	Leu	Val	Leu	Ser	Leu	Gly	Ala	Leu	Cys	Val	Ser
1				5					10					15	
Gln	Ile	Trp	Glu	Ser	Asn	Arg	Ala	Ser	Ala	Val	Val	Ser	Gly	Glu	Lys
			20					25					30		
Asn	Pro	Tyr	Val	Ser	Glu	Ser	Leu	Lys	Leu	Thr	Asn	Asn	Lys	Asn	Lys
		35					40					45			
Ser	Arg	Thr	Val	Glu	Glu	Tyr	Lys	Lys	Ser	Leu	Asp	Asp	Leu	Ile	Trp
	50					55					60				
Ser	Phe	Pro	Asn	Leu	Asp	Asn	Glu	Arg	Phe	Asp	Asn	Pro	Glu	Tyr	Lys
	65			70					75					80	
Glu	Ala	Met	Lys	Lys	Tyr	Gln	Gln	Arg	Phe	Met	Ala	Glu	Asp	Glu	Ala
			85						90					95	
Leu	Lys	Lys	Phe	Phe	Ser	Glu	Glu	Lys	Lys	Ile	Lys	Asn	Gly	Asn	Thr
			100					105					110		
Asp	Asn	Leu	Asp	Tyr	Leu	Gly	Leu	Ser	His	Glu	Arg	Tyr	Glu	Ser	Val
		115					120					125			
Phe	Asn	Thr	Leu	Lys	Lys	Gln	Ser	Glu	Glu	Phe	Leu	Lys	Glu	Ile	Glu
	130					135					140				
Asp	Ile	Lys	Lys	Asp	Asn	Pro	Glu	Leu	Lys	Asp	Phe	Asn	Glu	Glu	Glu
	145				150					155					160
Gln	Leu	Lys	Cys	Asp	Leu	Glu	Leu	Asn	Lys	Leu	Glu	Asn	Gln	Ile	Leu
			165						170					175	
Met	Leu	Gly	Lys	Thr	Phe	Tyr	Gln	Asn	Tyr	Arg	Asp	Asp	Val	Glu	Ser
		180						185					190		
Leu	Tyr	Ser	Lys	Leu	Asp	Leu	Ile	Met	Gly	Tyr	Lys	Asp	Glu	Glu	Arg
		195					200					205			
Ala	Asn	Lys	Lys	Ala	Val	Asn	Lys	Arg	Met	Leu	Glu	Asn	Lys	Lys	Glu
	210					215					220				
Asp	Leu	Glu	Thr	Ile	Ile	Asp	Glu	Phe	Phe	Ser	Asp	Ile	Asp	Lys	Thr
	225				230					235					240
Arg	Pro	Asn	Asn	Ile	Pro	Val	Leu	Glu	Asp	Glu	Lys	Gln	Glu	Glu	Lys
			245						250					255	
Asn	His	Lys	Asn	Met	Ala	Gln	Leu	Lys	Ser	Asp	Thr	Glu	Ala	Ala	Lys
			260					265					270		

-continued

---

Ser Asp Glu Ser Lys Arg Ser Lys Arg Ser Lys Arg Ser Leu Asn Thr  
 275 280 285

Gln Asn His Lys Pro Ala Ser Gln Glu Val Ser Glu Gln Gln Lys Ala  
 290 295 300

Glu Tyr Asp Lys Arg Ala Glu Glu Arg Lys Ala Arg Phe Leu Asp Asn  
 305 310 315 320

Gln Lys Ile Lys Lys Thr Pro Val Val Ser Leu Glu Tyr Asp Phe Glu  
 325 330 335

His Lys Gln Arg Ile Asp Asn Glu Asn Asp Lys Lys Leu Val Val Ser  
 340 345 350

Ala Pro Thr Lys Lys Pro Thr Ser Pro Thr Thr Tyr Thr Glu Thr Thr  
 355 360 365

Thr Gln Val Pro Met Pro Thr Val Glu Arg Gln Thr Gln Gln Gln Ile  
 370 375 380

Ile Tyr Asn Ala Pro Lys Gln Leu Ala Gly Leu Asn Gly Glu Ser His  
 385 390 395 400

Asp Phe Thr Thr Thr His Gln Ser Pro Thr Thr Ser Asn His Thr His  
 405 410 415

Asn Asn Val Val Glu Phe Glu Glu Thr Ser Ala Leu Pro Gly Arg Lys  
 420 425 430

Ser Gly Ser Leu Val Gly Ile Ser Gln Ile Asp Ser Ser His Leu Thr  
 435 440 445

Glu Arg Glu Lys Arg Val Ile Lys Arg Glu His Val Arg Glu Ala Gln  
 450 455 460

Lys Leu Val Asp Asn Tyr Lys Asp Thr His Ser Tyr Lys Asp Arg Ile  
 465 470 475 480

Asn Ala Gln Gln Lys Val Asn Thr Leu Ser Glu Gly His Gln Lys Arg  
 485 490 495

Phe Asn Lys Gln Ile Asn Lys Val Tyr Asn Gly Lys  
 500 505

<210> SEQ ID NO 33  
 <211> LENGTH: 520  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 33

Met Leu Thr Leu Gln Ile His Thr Gly Gly Ile Asn Leu Lys Lys Lys  
 1 5 10 15

Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile Ala Ser Val Thr  
 20 25 30

Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro Ala Ala Asn Ala  
 35 40 45

Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr Gln Val Leu Asn  
 50 55 60

Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu  
 65 70 75 80

Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys  
 85 90 95

Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln Gln Asn Asn Phe  
 100 105 110

Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn  
 115 120 125

Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp  
 130 135 140

-continued

---

Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys Leu Asn Glu  
 145 150 155 160  
 Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys Glu Gln Gln Asn  
 165 170 175  
 Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn Glu Glu Gln Arg  
 180 185 190  
 Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn  
 195 200 205  
 Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala  
 210 215 220  
 Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu  
 225 230 235 240  
 His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser  
 245 250 255  
 Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys  
 260 265 270  
 Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys  
 275 280 285  
 Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Thr  
 290 295 300  
 Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser  
 305 310 315 320  
 Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln  
 325 330 335  
 Ala Pro Lys Glu Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Gly Asn  
 340 345 350  
 Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Asn Lys  
 355 360 365  
 Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Asn Asn  
 370 375 380  
 Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys  
 385 390 395 400  
 Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Gly Asn  
 405 410 415  
 Lys Pro Gly Lys Glu Asp Gly Asn Gly Val His Val Val Lys Pro Gly  
 420 425 430  
 Asp Thr Val Asn Asp Ile Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys  
 435 440 445  
 Ile Ala Ala Asp Asn Lys Leu Ala Asp Lys Asn Met Ile Lys Pro Gly  
 450 455 460  
 Gln Glu Leu Val Val Asp Lys Lys Gln Pro Ala Asn His Ala Asp Ala  
 465 470 475 480  
 Asn Lys Ala Gln Ala Leu Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile  
 485 490 495  
 Gly Thr Thr Val Phe Gly Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu  
 500 505 510  
 Leu Ala Gly Arg Arg Arg Glu Leu  
 515 520

&lt;210&gt; SEQ ID NO 34

&lt;211&gt; LENGTH: 291

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

-continued

<400> SEQUENCE: 34

Ala Gln His Asp Glu Ala Lys Lys Asn Ala Phe Tyr Gln Val Leu Asn  
 1 5 10 15  
 Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu  
 20 25 30  
 Lys Ala Ala Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys  
 35 40 45  
 Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln Gln Asn Asn Phe  
 50 55 60  
 Asn Lys Asp Lys Lys Ser Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn  
 65 70 75 80  
 Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Ala Ala  
 85 90 95  
 Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys Leu Asn Glu  
 100 105 110  
 Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys Glu Lys Lys Asn  
 115 120 125  
 Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn Glu Glu Gln Arg  
 130 135 140  
 Asn Gly Phe Ile Gln Ser Leu Lys Ala Ala Pro Ser Gln Ser Ala Asn  
 145 150 155 160  
 Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala  
 165 170 175  
 Asp Asn Lys Phe Asn Lys Glu Lys Lys Asn Ala Phe Tyr Glu Ile Leu  
 180 185 190  
 His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser  
 195 200 205  
 Leu Lys Ala Ala Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys  
 210 215 220  
 Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys  
 225 230 235 240  
 Glu Lys Lys Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Thr  
 245 250 255  
 Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Ala Ala Pro Ser  
 260 265 270  
 Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln  
 275 280 285  
 Ala Pro Lys  
 290

<210> SEQ ID NO 35  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 35

gctgcacata tggcgcaaca cgatgaagct caac

34

<210> SEQ ID NO 36  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 36

agtggatect tatgctttgt tagcatctgc

30

-continued

<210> SEQ ID NO 37  
 <211> LENGTH: 19  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 37

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro  
 1                   5                   10                   15

Arg Gly Ser

<210> SEQ ID NO 38  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 38

aacatattggt caacaaagat caacaaagc 29

<210> SEQ ID NO 39  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 39

aaggatccag attcggttaa ttttttagc 29

<210> SEQ ID NO 40  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 40

cttcattcaa agtcttaaag cgcaccaag ccaaagcact aac 43

<210> SEQ ID NO 41  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 41

gtagtgctt tggcttgggg cggctttaag actttgaatg aag 43

<210> SEQ ID NO 42  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 42

catatgttca acaaaagataa aaaaagcgcc ttctatgaaa tc 42

<210> SEQ ID NO 43  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 43

gatttcatag aaggcgcttt ttttatcttt gttgaacata tg 42

<210> SEQ ID NO 44  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

-continued

---

<400> SEQUENCE: 44  
 catatgttca acaaatgatgg aggaagcgcc ttctatgaaa to 42

<210> SEQ ID NO 45  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 45  
 gatttcatag aaggcgcttc ctccatcttt gttgaacata tg 42

<210> SEQ ID NO 46  
 <211> LENGTH: 52  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 46  
 ggggacaagt ttgtacaaaa aagcaggctg atgactaagt tgaaaaaaga ag 52

<210> SEQ ID NO 47  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 47  
 aaggatcccc tccaaaatgt aattgccc 28

<210> SEQ ID NO 48  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 48  
 aaggatccgt ttgtaactct atccaaagac 30

<210> SEQ ID NO 49  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 49  
 ggggaccact ttgtacaaga aagctgggtg acacctattg cacgattcg 49

<210> SEQ ID NO 50  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 50  
 ggggacaagt ttgtacaaaa aagcaggctc agatagcgat tcagattcag 50

<210> SEQ ID NO 51  
 <211> LENGTH: 31  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 51  
 aaggatccct gtattttctc cttaattttc c 31

<210> SEQ ID NO 52  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

-continued

---

<400> SEQUENCE: 52  
aaggatccca tggctgcaaa gcaaataatg 30

<210> SEQ ID NO 53  
<211> LENGTH: 51  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 53  
ggggaccact ttgtacaaga aagctgggtg ccttggtgta acaaatttat g 51

<210> SEQ ID NO 54  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 54  
gaaggatccg tttattctag ttaatatata gttaatg 37

<210> SEQ ID NO 55  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 55  
gaactgcagc tgtatgtctt tggatagagt tac 33

<210> SEQ ID NO 56  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 56  
gaaggatccg gtggcttttt tacttggatt ttc 33

<210> SEQ ID NO 57  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 57  
gaactgcagc gacaaactca ttatttgctt tgc 33

<210> SEQ ID NO 58  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 58  
gaactcgagt ctagcttatt tacatgg 27

<210> SEQ ID NO 59  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 59  
gaactcgaga tagaaggcag aatagtaaca aaggattata gtggg 45

<210> SEQ ID NO 60  
<211> LENGTH: 27  
<212> TYPE: DNA

-continued

---

 <213> ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 60

gtaggatcct gggatagagt tacaaac

27

&lt;210&gt; SEQ ID NO 61

&lt;211&gt; LENGTH: 34

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 61

gaactcgagg cattatgtgt atcacaatt tggg

34

&lt;210&gt; SEQ ID NO 62

&lt;211&gt; LENGTH: 43

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 62

gaactcgaga tagaaggcag agtggtttct ggggagaaga atc

43

&lt;210&gt; SEQ ID NO 63

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 63

gaactcgagg cagccatgca ttaattattt gcc

33

&lt;210&gt; SEQ ID NO 64

&lt;211&gt; LENGTH: 677

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 64

Met Lys Ser Asn Leu Arg Tyr Gly Ile Arg Lys His Lys Leu Gly Ala  
1 5 10 15Ala Ser Val Phe Leu Gly Thr Met Ile Val Val Gly Met Gly Gln Glu  
20 25 30Lys Glu Ala Ala Ala Ser Glu Gln Asn Asn Thr Thr Val Glu Glu Ser  
35 40 45Gly Ser Ser Ala Thr Glu Ser Lys Ala Ser Glu Thr Gln Thr Thr Thr  
50 55 60Asn Asn Val Asn Thr Ile Asp Glu Thr Gln Ser Tyr Ser Ala Thr Ser  
65 70 75 80Thr Glu Gln Pro Ser Gln Ser Thr Gln Val Thr Thr Glu Glu Ala Pro  
85 90 95Lys Thr Val Gln Ala Pro Lys Val Glu Thr Ser Arg Val Asp Leu Pro  
100 105 110Ser Glu Lys Val Ala Asp Lys Glu Thr Thr Gly Thr Gln Val Asp Ile  
115 120 125Ala Gln Pro Ser Asn Val Ser Glu Ile Lys Pro Arg Met Lys Arg Ser  
130 135 140Thr Asp Val Thr Ala Val Ala Glu Lys Glu Val Val Glu Glu Thr Lys  
145 150 155 160Ala Thr Gly Thr Asp Val Thr Asn Lys Val Glu Val Glu Glu Gly Ser  
165 170 175Glu Ile Val Gly His Lys Gln Asp Thr Asn Val Val Asn Pro His Asn  
180 185 190

-continued

---

Ala Glu Arg Val Thr Leu Lys Tyr Lys Trp Lys Phe Gly Glu Gly Ile  
 195 200 205

Lys Ala Gly Asp Tyr Phe Asp Phe Thr Leu Ser Asp Asn Val Glu Thr  
 210 215 220

His Gly Ile Ser Thr Leu Arg Lys Val Pro Glu Ile Lys Ser Thr Asp  
 225 230 235 240

Gly Gln Val Met Ala Thr Gly Glu Ile Ile Gly Glu Arg Lys Val Arg  
 245 250 255

Tyr Thr Phe Lys Glu Tyr Val Gln Glu Lys Lys Asp Leu Thr Ala Glu  
 260 265 270

Leu Ser Leu Asn Leu Phe Ile Asp Pro Thr Thr Val Thr Gln Lys Gly  
 275 280 285

Asn Gln Asn Val Glu Val Lys Leu Gly Glu Thr Thr Val Ser Lys Ile  
 290 295 300

Phe Asn Ile Gln Tyr Leu Gly Gly Val Arg Asp Asn Trp Gly Val Thr  
 305 310 315 320

Ala Asn Gly Arg Ile Asp Thr Leu Asn Lys Val Asp Gly Lys Phe Ser  
 325 330 335

His Phe Ala Tyr Met Lys Pro Asn Asn Gln Ser Leu Ser Ser Val Thr  
 340 345 350

Val Thr Gly Gln Val Thr Lys Gly Asn Lys Pro Gly Val Asn Asn Pro  
 355 360 365

Thr Val Lys Val Tyr Lys His Ile Gly Ser Asp Asp Leu Ala Glu Ser  
 370 375 380

Val Tyr Ala Lys Leu Asp Asp Val Ser Lys Phe Glu Asp Val Thr Asp  
 385 390 395 400

Asn Met Ser Leu Asp Phe Asp Thr Asn Gly Gly Tyr Ser Leu Asn Phe  
 405 410 415

Asn Asn Leu Asp Gln Ser Lys Asn Tyr Val Ile Lys Tyr Glu Gly Tyr  
 420 425 430

Tyr Asp Ser Asn Ala Ser Asn Leu Glu Phe Gln Thr His Leu Phe Gly  
 435 440 445

Tyr Tyr Asn Tyr Tyr Tyr Thr Ser Asn Leu Thr Trp Lys Asn Gly Val  
 450 455 460

Ala Phe Tyr Ser Asn Asn Ala Gln Gly Asp Gly Lys Asp Lys Leu Lys  
 465 470 475 480

Glu Pro Ile Ile Glu His Ser Thr Pro Ile Glu Leu Glu Phe Lys Ser  
 485 490 495

Glu Pro Pro Val Glu Lys His Glu Leu Thr Gly Thr Ile Glu Glu Ser  
 500 505 510

Asn Asp Ser Lys Pro Ile Asp Phe Glu Tyr His Thr Ala Val Glu Gly  
 515 520 525

Ala Glu Gly His Ala Glu Gly Thr Ile Glu Thr Glu Glu Asp Ser Ile  
 530 535 540

His Val Asp Phe Glu Glu Ser Thr His Glu Asn Ser Lys His His Ala  
 545 550 555 560

Asp Val Val Glu Tyr Glu Glu Asp Thr Asn Pro Gly Gly Gly Gln Val  
 565 570 575

Thr Thr Glu Ser Asn Leu Val Glu Phe Asp Glu Asp Ser Thr Lys Gly  
 580 585 590

Ile Val Thr Gly Ala Val Ser Asp His Thr Thr Ile Glu Asp Thr Lys  
 595 600 605

-continued

---

Glu Tyr Thr Thr Glu Ser Asn Leu Ile Glu Leu Val Asp Glu Leu Pro  
 610 615 620

Glu Glu His Gly Gln Ala Gln Gly Pro Ile Glu Glu Ile Thr Glu Asn  
 625 630 635 640

Asn His His Ile Ser His Ser Gly Leu Gly Thr Glu Asn Gly His Gly  
 645 650 655

Asn Tyr Gly Val Ile Glu Glu Ile Glu Glu Asn Ser His Val Asp Ile  
 660 665 670

Lys Ser Glu Leu Gly  
 675

---

What is claimed is:

1. A variant Protein A (SpA) comprising a domain E having alanine and/or valine residue substitutions at amino acid positions 33 and 34 of SEQ ID NO: 3.
2. The variant SpA of claim 1, wherein the domain E further comprises lysine residue substitutions at amino acid positions 6 and 7 of SEQ ID NO: 3; and/or further wherein the SpA variant further comprises a domain D having a lysine residue substitution at amino acid positions 9 and 10 of SEQ ID NO: 2; a domain A having a lysine residue substitution at amino acid positions 7 and 8 of SEQ ID NO: 4; a domain B having a lysine residue substitution at amino acid positions 7 and 8 of SEQ ID NO: 6, and/or a domain C having a lysine residue substitution at amino acid positions 7 and 8 of SEQ ID NO: 5.
3. The variant SpA of claim 1,
  - (i) wherein the domain E comprises a lysine residue substitution at amino acid positions 6 and 7 of SEQ ID NO: 3; and/or wherein the variant SpA further comprises a domain D having a lysine residue substitution at amino acid positions 9 and 10 of SEQ ID NO: 2; a domain A having a lysine residue substitution at amino acid positions 7 and 8 of SEQ ID NO: 4; a domain B having a lysine residue substitution at amino acid positions 7 and 8 of SEQ ID NO: 6, and/or a domain C having a lysine residue substitution at amino acid positions 7 and 8 of SEQ ID NO: 5; and
  - (ii) wherein the domain E comprises an alanine and/or valine residue substitution at amino acid positions 33 and 34 of SEQ ID NO: 3; and/or wherein the variant SpA further comprises a domain D having an alanine and/or valine residue substitution at amino acid positions 36 and 37 of SEQ ID NO: 2; a domain A having an alanine and/or valine residue substitution at amino acid positions 34 and 35 of SEQ ID NO: 4; a domain B having an alanine and/or valine residue substitution at amino acid positions 34 and 35 of SEQ ID NO: 6,

- and/or a domain C having an alanine and/or valine residue substitution at amino acid positions 34 and 35 of SEQ ID NO: 5.
4. The variant SpA of claim 1, wherein the variant SpA comprises a segment of SpA comprising 5 or more IgG binding domains.
5. The variant SpA of claim 1, comprising a domain E having valine residue substitutions at amino acid positions 33 and 34 of SEQ ID NO: 3.
6. An immunogenic composition comprising the variant SpA of claim 1.
7. An immunogenic composition according to claim 6, further comprising at least a second staphylococcal antigen.
8. An immunogenic composition according to claim 7, wherein the second staphylococcal antigen is selected from the group consisting of EsaB, Emp, EsxA, EsxB, EsaC, Eap, Ebh, Coa, vWh, Hla, SdrC, SdrE, IsdA, IsdB, IsdC, ClfA, ClfB, SasF peptide, a type V and/or a type VIII capsular polysaccharide or oligosaccharide from *S. aureus*.
9. A vaccine comprising a pharmaceutically acceptable composition having an isolated SpA variant polypeptide of claim 1, wherein the composition is capable of stimulating an immune response against a *Staphylococcus* bacterium.
10. A method for eliciting an immune response against a *Staphylococcus* bacterium in a subject comprising providing to the subject an effective amount of the composition of claim 6.
11. The method of claim 10, wherein the *Staphylococcus* bacterium is a *S. aureus* bacterium.
12. The method of claim 11, wherein the bacterium is methicillin resistant.
13. The method of claim 10, wherein the subject is a mammal.
14. The method of claim 10, wherein the subject is human.
15. The method of claim 10, wherein the immune response is a protective immune response.

\* \* \* \* \*