



US 20130230550A1

(19) **United States**

(12) **Patent Application Publication**  
**Schneewind et al.**

(10) **Pub. No.: US 2013/0230550 A1**

(43) **Pub. Date: Sep. 5, 2013**

(54) **METHODS AND COMPOSITIONS  
INVOLVING PROTECTIVE  
STAPHYLOCOCCAL ANTIGENS**

(75) Inventors: **Olaf Schneewind**, Chicago, IL (US);  
**Alice G. Cheng**, Boston, MA (US);  
**Dominique M. Missiakas**, Chicago, IL  
(US); **Hwan Keun Kim**, Naperville, IL  
(US); **Hye-Young Kim**, Rancho  
Cucamonga, CA (US)

(73) Assignee: **The University of Chicago**, Chicago, IL  
(US)

(21) Appl. No.: **13/821,943**

(22) PCT Filed: **Sep. 9, 2011**

(86) PCT No.: **PCT/US11/51079**

§ 371 (c)(1),  
(2), (4) Date: **May 22, 2013**

**Related U.S. Application Data**

(60) Provisional application No. 61/381,372, filed on Sep. 9, 2010, provisional application No. 61/435,617, filed on Jan. 24, 2011.

**Publication Classification**

(51) **Int. Cl.**  
**A61K 39/085** (2006.01)  
(52) **U.S. Cl.**  
CPC ..... **A61K 39/085** (2013.01)  
USPC ..... **424/190.1; 424/243.1**

(57) **ABSTRACT**

The present invention concerns methods and compositions for treating or preventing a bacterial infection, particularly infection by a *Staphylococcus* bacterium. The invention provides methods and compositions for stimulating an immune response against the bacteria. In certain embodiments, the methods and compositions involve a non-toxigenic Protein A (SpA) variant. In some embodiments, the methods and compositions involve SdrD, ClfA, and/or FnbpB polypeptides.






FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7








FIG. 8




FIG. 9




FIG. 10






FIG. 11



FIGs. 12A-12C



FIGs. 13A-13B




FIG. 14A-14H




FIG. 15



## METHODS AND COMPOSITIONS INVOLVING PROTECTIVE STAPHYLOCOCCAL ANTIGENS

**[0001]** This application claims the benefit of U.S. Provisional Patent Application Nos. 61/381,372 and 61/435,617, filed Sep. 9, 2010 and Jan. 24, 2011, respectively, the entirety of which are incorporated herein by reference.

**[0002]** This invention was made with government support under AI057153, AI052474, and GM007281 awarded by the National Institutes of Health. The government has certain rights in the invention.

### BACKGROUND OF THE INVENTION

**[0003]** I. Field of the Invention

**[0004]** The present invention relates generally to the fields of immunology, microbiology, and pathology. More particularly, it concerns methods and compositions involving bacterial Protein A variants, which can be used to invoke an immune response against the bacteria.

**[0005]** II. Background

**[0006]** The number of both community acquired and hospital acquired infections have increased over recent years with the increased use of intravascular devices. Hospital acquired (nosocomial) infections are a major cause of morbidity and mortality, more particularly in the United States, where it affects more than 2 million patients annually. The most frequent infections are urinary tract infections (33% of the infections), followed by pneumonia (15.5%), surgical site infections (14.8%) and primary bloodstream infections (13%) (Emorl and Gaynes, 1993).

**[0007]** The major nosocomial pathogens include *Staphylococcus aureus*, coagulase-negative Staphylococci (mostly *Staphylococcus epidermidis*), *enterococcus* spp., *Escherichia coli* and *Pseudomonas aeruginosa*. Although these pathogens cause approximately the same number of infections, the severity of the disorders they can produce combined with the frequency of antibiotic resistant isolates balance this ranking towards *S. aureus* and *S. epidermidis* as being the most significant nosocomial pathogens.

**[0008]** Staphylococci can cause a wide variety of diseases in humans and other animals through either toxin production or invasion. Staphylococcal toxins are also a common cause of food poisoning, as the bacteria can grow in improperly stored food.

**[0009]** *Staphylococcus epidermidis* is a normal skin commensal which is also an important opportunistic pathogen responsible for infections of impaired medical devices and infections at sites of surgery. Medical devices infected by *S. epidermidis* include cardiac pacemakers, cerebrospinal fluid shunts, continuous ambulatory peritoneal dialysis catheters, orthopedic devices and prosthetic heart valves.

**[0010]** *Staphylococcus aureus* is the most common cause of nosocomial infections with a significant morbidity and mortality. It is the cause of some cases of osteomyelitis, endocarditis, septic arthritis, pneumonia, abscesses, and toxic shock syndrome. *S. aureus* can survive on dry surfaces, increasing the chance of transmission. Any *S. aureus* infection can cause the staphylococcal scalded skin syndrome, a cutaneous reaction to exotoxin absorbed into the bloodstream. It can also cause a type of septicemia called pyaemia that can be life-threatening. Problematically, Methicillin-resistant *Staphylococcus aureus* (MRSA) has become a major cause of hospital-acquired infections.

**[0011]** *S. aureus* and *S. epidermidis* infections are typically treated with antibiotics, with penicillin being the drug of choice, whereas vancomycin is used for methicillin resistant isolates. The percentage of staphylococcal strains exhibiting wide-spectrum resistance to antibiotics has become increasingly prevalent, posing a threat for effective antimicrobial therapy. In addition, the recent emergence of vancomycin resistant *S. aureus* strain has aroused fear that MRSA strains are emerging and spreading for which no effective therapy is available.

**[0012]** An alternative to antibiotic treatment for staphylococcal infections is under investigation that uses antibodies directed against staphylococcal antigens. This therapy involves administration of polyclonal antisera (WO00/15238, WO00/12132) or treatment with monoclonal antibodies against lipoteichoic acid (WO98/57994).

**[0013]** An alternative approach would be the use of active vaccination to generate an immune response against staphylococci. The *S. aureus* genome has been sequenced and many of the coding sequences have been identified (WO02/094868, EP0786519), which can lead to the identification of potential antigens. The same is true for *S. epidermidis* (WO01/34809). As a refinement of this approach, others have identified proteins that are recognized by hyperimmune sera from patients who have suffered staphylococcal infection (WO01/98499, WO02/059148).

**[0014]** *S. aureus* secretes a plethora of virulence factors into the extracellular milieu (Archer, 1998; Dinges et al., 2000; Foster, 2005; Shaw et al., 2004; Sibbald et al., 2006). Like most secreted proteins, these virulence factors are translocated by the Sec machinery across the plasma membrane. Proteins secreted by the Sec machinery bear an N-terminal leader peptide that is removed by leader peptidase once the pre-protein is engaged in the Sec translocon (Dalbey and Wickner, 1985; van Wely et al., 2001). Recent genome analysis suggests that Actinobacteria and members of the Firmicutes encode an additional secretion system that recognizes a subset of proteins in a Sec-independent manner (Pallen, 2002). ESAT-6 (early secreted antigen target 6 kDa) and CFP-10 (culture filtrate antigen 10 kDa) of *Mycobacterium tuberculosis* represent the first substrates of this novel secretion system termed ESX-1 or 5 nm in *M. tuberculosis* (Andersen et al., 1995; Hsu et al., 2003; Pym et al., 2003; Stanley et al., 2003). In *S. aureus*, two ESAT-6 like factors designated EsxA and EsxB are secreted by the Ess pathway (ESAT-6 secretion system) (Burts et al., 2005).

**[0015]** The first generation of vaccines targeted against *S. aureus* or against the exoproteins it produces have met with limited success (Lee, 1996). There remains a need to develop effective vaccines against staphylococcal infections. Additional compositions for treating staphylococcal infections are also needed.

### SUMMARY OF THE INVENTION

**[0016]** Protein A (SpA) (SEQ ID NO:33), a cell wall anchored surface protein of *Staphylococcus aureus*, provides for bacterial evasion from innate and adaptive immune responses. Protein A binds immunoglobulins at their Fc portion, interacts with the VH3 domain of B cell receptors inappropriately stimulating B cell proliferation and apoptosis, binds to von Willebrand factor A1 domains to activate intracellular clotting, and also binds to the TNF Receptor-1 to contribute to the pathogenesis of staphylococcal pneumonia. Due to the fact that Protein A captures immunoglobulin and

displays toxic attributes, the possibility that this surface molecule may function as a vaccine in humans has not been rigorously pursued. Here the inventors demonstrate that Protein A variants no longer able to bind to immunoglobulins, which are thereby removed of their toxigenic potential, i.e., are non-toxicogenic, stimulate humoral immune responses that protect against staphylococcal disease.

**[0017]** In certain embodiments the SpA variant is a full length SpA variant comprising a variant A, B, C, D, and/or E domain. In certain aspects, the SpA variant comprises or consists of the amino acid sequence that is 80, 90, 95, 98, 99, or 100% identical to the amino acid sequence of SEQ ID NO:34. In other embodiments the SpA variant comprises a segment of SpA. The SpA segment can comprise at least or at most 1, 2, 3, 4, 5 or more IgG binding domains. The IgG domains can be at least or at most 1, 2, 3, 4, 5 or more variant A, B, C, D, or E domains. In certain aspects the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant A domains. In a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant B domains. In still a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant C domains. In yet a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant D domains. In certain aspects the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant E domains. In a further aspect the SpA variant comprises a combination of A, B, C, D, and E domains in various combinations and permutations. The combinations can include all or part of a SpA signal peptide segment, a SpA region X segment, and/or a SpA sorting signal segment. In other aspects the SpA variant does not include a SpA signal peptide segment, a SpA region X segment, and/or a SpA sorting signal segment. In certain aspects a variant A domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:4. In another aspect a variant B domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:6. In still another aspect a variant C domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:5. In certain aspects a variant D domain comprises a substitution at position(s) 9, 10, 36, and/or 37 of SEQ ID NO:2. In a further aspect a variant E domain comprises a substitution at position(s) 6, 7, 33, and/or 34 of SEQ ID NO:3.

**[0018]** In certain aspects, an SpA domain D variant or its equivalent can comprise a mutation at position 9 and 36; 9 and 37; 9 and 10; 36 and 37; 10 and 36; 10 and 37; 9, 36, and 37; 10, 36, and 37; 9, 10 and 36; or 9, 10 and 37 of SEQ ID NO:2. In a further aspect, analogous mutations can be included in one or more of domains A, B, C, or E.

**[0019]** In further aspects, the amino acid glutamine (Q) at position 9 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an aspartic acid (D), a cysteine (C), a glutamic acid (E), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the glutamine at position 9 can be substituted with an arginine (R). In a further aspect, the glutamine at position 9 of SEQ ID NO:2, or its equivalent, can be substituted with a lysine or a glycine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

**[0020]** In another aspect, the amino acid glutamine (Q) at position 10 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an

asparagine (N), an aspartic acid (D), a cysteine (C), a glutamic acid (E), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the glutamine at position 10 can be substituted with an arginine (R). In a further aspect, the glutamine at position 10 of SEQ ID NO:2, or its equivalent, can be substituted with a lysine or a glycine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

**[0021]** In certain aspects, the aspartic acid (D) at position 36 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an arginine (R), a cysteine (C), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a glutamine (Q), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the aspartic acid at position 36 can be substituted with a glutamic acid (E). In certain aspects, an aspartic acid at position 36 of SEQ ID NO:2, or its equivalent, can be substituted with an alanine or a serine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

**[0022]** In another aspect, the aspartic acid (D) at position 37 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an arginine (R), a cysteine (C), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a glutamine (Q), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the aspartic acid at position 37 can be substituted with a glutamic acid (E). In certain aspects, an aspartic acid at position 37 of SEQ ID NO:2, or its equivalent, can be substituted with an alanine or a serine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

**[0023]** In a particular embodiment the amino at position 9 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 9 of SEQ ID NO:2 is replaced by a glycine. In a further aspect the amino acid at position 9 of SEQ ID NO:2 is replaced by a lysine.

**[0024]** In a particular embodiment the amino at position 10 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 10 of SEQ ID NO:2 is replaced by a glycine. In a further aspect the amino acid at position 10 of SEQ ID NO:2 is replaced by a lysine.

**[0025]** In a particular embodiment the amino at position 36 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 36 of SEQ ID NO:2 is replaced by a serine. In a further aspect the amino acid at position 36 of SEQ ID NO:2 is replaced by an alanine.

**[0026]** In a particular embodiment the amino at position 37 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a

valine (V). In certain aspects the amino acid at position 37 of SEQ ID NO:2 is replaced by a serine. In a further aspect the amino acid at position 37 of SEQ ID NO:2 is replaced by an alanine.

**[0027]** In certain aspects the SpA variant includes (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a  $V_H3$  binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to  $V_H3$ . In still further aspects the amino acid sequence of a SpA variant comprises an amino acid sequence that is at least 50%, 60%, 70%, 80%, 90%, 95%, or 100% identical, including all values and ranges there between, to the amino acid sequence of SEQ ID NOs:2-6.

**[0028]** In a further aspect the SpA variant includes (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain D, or at a corresponding amino acid position in other IgG domains, that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a  $V_H3$  binding sub-domain of SpA domain D, or at a corresponding amino acid position in other IgG domains, that disrupts or decreases binding to  $V_H3$ . In certain aspects amino acid residue F5, Q9, Q10, S11, F13, Y14, L17, N28, I31, and/or K35 (SEQ ID NO:2, QQNNFNKDDQSSAFYEILNMPNLNEAQRNGFIQSLKDDPSQSTNVLGEAKKLNES) of the IgG Fc binding sub-domain of domain D are modified or substituted. In certain aspects amino acid residue Q26, G29, F30, S33, D36, D37, Q40, N43, and/or E47 (SEQ ID NO:2) of the  $V_H3$  binding sub-domain of domain D are modified or substituted such that binding to Fc or  $V_H3$  is attenuated. In further aspects corresponding modifications or substitutions can be engineered in corresponding positions of the domain A, B, C, and/or E. Corresponding positions are defined by alignment of the domain D amino acid sequence with one or more of the amino acid sequences from other IgG binding domains of SpA, for example see FIG. 2A. In certain aspects the amino acid substitution can be any of the other 20 amino acids. In a further aspect conservative amino acid substitutions can be specifically excluded from possible amino acid substitutions. In other aspects only non-conservative substitutions are included. In any event, any substitution or combination of substitutions that reduces the binding of the domain such that SpA toxicity is significantly reduced is contemplated. The significance of the reduction in binding refers to a variant that produces minimal to no toxicity when introduced into a subject and can be assessed using in vitro methods described herein.

**[0029]** In certain embodiments, a variant SpA comprises at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more variant SpA domain D peptides. In certain aspects 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 or more amino acid residues of the variant SpA are substituted or modified—including but not limited to amino acids F5, Q9, Q10, S11, F13, Y14, L17, N28, I31, and/or K35 (SEQ ID NO:2) of the IgG Fc binding sub-domain of domain D and amino acid residue Q26, G29, F30, S33, D36, D37, Q40, N43, and/or E47 (SEQ ID NO:2) of the  $V_H3$  binding sub-domain of domain D. In one aspect of the invention glutamine residues at position 9 and/or 10 of SEQ ID NO:2 (or corresponding positions in other domains) are mutated. In another aspect, aspartic acid residues 36 and/or 37 of SEQ ID NO:2 (or corresponding positions in other domains) are mutated. In a further aspect, glutamine 9 and 10, and aspartic acid residues 36 and 37 are

mutated. Purified non-toxicogenic SpA or SpA-D mutants/variants described herein are no longer able to significantly bind (i.e., demonstrate attenuated or disrupted binding affinity) Fc $\gamma$  or F(ab) $_2$   $V_H3$  and also do not stimulate B cell apoptosis. These non-toxicogenic Protein A variants can be used as subunit vaccines and raise humoral immune responses and confer protective immunity against *S. aureus* challenge. Compared to wild-type full-length Protein A or the wild-type SpA-domain D, immunization with SpA-D variants resulted in an increase in Protein A specific antibody. Using a mouse model of staphylococcal challenge and abscess formation, it was observed that immunization with the non-toxicogenic Protein A variants generated significant protection from staphylococcal infection and abscess formation. As virtually all *S. aureus* strains express Protein A, immunization of humans with the non-toxicogenic Protein A variants can neutralize this virulence factor and thereby establish protective immunity. In certain aspects the protective immunity protects or ameliorates infection by drug resistant strains of *Staphylococcus*, such as USA300 and other MRSA strains.

**[0030]** Embodiments include the use of Protein A variants in methods and compositions for the treatment of bacterial and/or staphylococcal infection. This application also provides an immunogenic composition comprising a Protein A variant or immunogenic fragment thereof. In certain aspects, the immunogenic fragment is a Protein A domain D segment. Furthermore, the present invention provides methods and compositions that can be used to treat (e.g., limiting staphylococcal abscess formation and/or persistence in a subject) or prevent bacterial infection. In some cases, methods for stimulating an immune response involve administering to the subject an effective amount of a composition including or encoding all or part of a Protein A variant polypeptide or antigen, and in certain aspects other bacterial proteins. Other bacterial proteins include, but are not limited to (i) a secreted virulence factor, and/or a cell surface protein or peptide, or (ii) a recombinant nucleic acid molecule encoding a secreted virulence factor, and/or a cell surface protein or peptide.

**[0031]** In other aspects, the subject can be administered all or part of a Protein A variant, such as a variant Protein A domain D segment. The polypeptide of the invention can be formulated in a pharmaceutically acceptable composition. The composition can further comprise one or more of at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 additional staphylococcal antigen or immunogenic fragment thereof (e.g., Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, an EsxA-B fusion protein (i.e., EsxAB or EsxBA), SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla (e.g., H35 mutants), IsdC, SasF, vWbp, FluD2, sta011, sta0048, sta0069 or vWh). Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa (GenBank CAC80837), Aap (GenBank accession AJ249487), Ant (GenBank accession NP\_372518), autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg $^{2+}$  transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF

(WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, Vitronectin binding protein (see PCT publications WO2007/113222, WO2007/113223, WO2006/032472, WO2006/032475, WO2006/032500, each of which is incorporated herein by reference in their entirety) and/or any of those antigens described in PCT Publ. No. WO2010119343, incorporated herein by reference.

**[0032]** In certain aspects, the SpA variant composition can further comprise SdrD, ClfA, and/or FnbpB (FnbB) staphylococcal antigens or immunogenic fragments thereof. Thus, in certain aspects, a composition of the embodiments comprises a SpA variant, SdrD, ClfA, and FnbpB (FnbB) staphylococcal antigens. Such a composition can, in some aspects be essentially free of other staphylococcal antigens, such as staphylococcal polypeptides or carbohydrates (e.g., a composition comprising staphylococcal antigens that essentially comprise the SpA variant, SdrD, ClfA, and FnbpB (FnbB) staphylococcal antigens). In a further aspect, embodiments of the invention provide for the use of a SpA variant, SdrD, ClfA, and FnbpB polypeptide in the preparation of a medicament for the treatment or prevention of a staphylococcal infection.

**[0033]** The staphylococcal antigen(s) or immunogenic fragment(s) of the embodiments can be administered concurrently with the Protein A variant. The staphylococcal antigen or immunogenic fragment and the Protein A variant can be administered in the same composition. The Protein A variant can also be a recombinant nucleic acid molecule encoding a Protein A variant. A recombinant nucleic acid molecule can encode the Protein A variant and at least one staphylococcal antigen or immunogenic fragment thereof. As used herein, the term “modulate” or “modulation” encompasses the meanings of the words “enhance,” or “inhibit.” “Modulation” of activity may be either an increase or a decrease in activity. As used herein, the term “modulator” refers to compounds that effect the function of a moiety, including up-regulation, induction, stimulation, potentiation, inhibition, down-regulation, or suppression of a protein, nucleic acid, gene, organism or the like.

**[0034]** In further aspects, an immunogenic composition comprises SdrD, ClfA, and/or FnbpB (FnbB) staphylococcal antigens or immunogenic fragments thereof. In other embodiments an immunogenic composition comprising SdrD, ClfA, and/or FnbpB (FnbB) staphylococcal antigens or immunogenic fragments thereof can be used in treating, ameliorating or inhibiting staphylococcal infection, as described herein. Thus, some embodiments of the invention concern compositions comprising SdrD, ClfA, and FnbpB (FnbB) staphylococcal antigens. Such a composition can, in some aspects, be essentially free of other staphylococcal antigens, such as staphylococcal polypeptides or carbohydrates (e.g., a composition comprising staphylococcal antigens that essentially comprise SdrD, ClfA, and FnbpB (FnbB) staphylococcal antigens). In a further aspect, embodiments of the invention provide for the use of a SdrD, ClfA, and FnbpB polypeptide in the preparation of a medicament for the treatment or prevention of a staphylococcal infection. In certain aspects, a SdrD, ClfA, and/or FnbpB (FnbB) staphylococcal antigen is from *S. aureus*.

**[0035]** In certain embodiments the methods and compositions use or include or encode all or part of the Protein A

variant or antigen. In other aspects, the Protein A variant may be used in combination with secreted factors or surface antigens including, but not limited to one or more of an isolated Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, an EsxA-B fusion protein (i.e., EsxAB or EsxBA), SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, FhuD2, sta011, sta0048, sta0069 or vWh polypeptide or immunogenic segment thereof. Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein. In certain embodiments, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more of Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein can be specifically excluded from a formulation of the invention. In further embodiments the methods and compositions use or include or encode all or part of the SdrD, ClfA and/or FnbpB (FnbB) antigens.

**[0036]** In some embodiments, the methods and compositions use, include or encode a Protein A variant in combination with the FhuD2, sta011, Hla (e.g., a H35 mutant such as HLA<sub>35L</sub> or HLA<sub>35,4</sub>) and EsxAB (i.e., an EsxA-B fusion protein) staphylococcal antigens or portions of these antigens. In further aspects, such a combination further includes SdrD, ClfA and/or FnbpB antigens

**[0037]** The following table lists (Table 1) combinations of SpA variants of the embodiments and various other Staphylococcal antigens. It will be apparent to one skilled in the art that there are, for example, 378 possible pairwise combinations selected from a set of 28 antigens, 3,276 possible three-way combinations, and 20,475 possible four-way combinations, and so on for larger subsets of antigens, all of which are contemplated herein.

**[0038]** Thus, any of the combinations of antigens of Table 1 can also be combined with one, two or more of the antigens



**[0039]** In still further aspects, the isolated Protein A variant is multimerized, e.g., dimerized or a linear fusion of two or more polypeptides or peptide segments. In certain aspects of the invention, a composition comprises multimers or concatamers of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more isolated cell surface proteins or segments thereof. Concatamers are linear polypeptides having one or more repeating peptide units. SpA polypeptides or fragments can be consecutive or separated by a spacer or other peptide sequences, e.g., one or more additional bacterial peptide. In a further aspect, the other polypeptides or peptides contained in the multimer or concatamer can include, but are not limited to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 of Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, CHB, Coa, Hla, IsdC, SasF, vWbp, vWh or immunogenic fragments thereof. Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein. In certain aspects the SpA variant is used in combination with SdrD, ClfA, and/or FnbpB (FnbB) antigens.

**[0040]** The term "Protein A variant" or "SpA variant" refers to polypeptides that include a SpA IgG domain having two or more amino acid substitutions that disrupt binding to Fc and V<sub>H</sub>3. In certain aspect, a SpA variant includes a variant domain D peptide, as well as variants of SpA polypeptides and segments thereof that are non-toxicogenic and stimulate an immune response against *staphylococcus* bacteria Protein A and/or bacteria expressing such.

**[0041]** Embodiments of the present invention include methods for eliciting an immune response against a *staphylococcus* bacterium or staphylococci in a subject comprising providing to the subject an effective amount of a Protein A variant or a segment thereof. In certain aspects, the methods for eliciting an immune response against a *staphylococcus* bacterium or staphylococci in a subject comprising providing to the subject an effective amount of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or more secreted proteins and/or cell surface proteins or segments/fragments thereof A secreted protein or cell surface protein includes, but is not limited to Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, and/or vWh proteins and immunogenic fragments thereof. Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341),

Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein. In certain aspects an SpA variant is used in combination with SdrD, ClfA, and/or FnbpB (FnbB) antigens.

**[0042]** Embodiments of the invention include compositions that include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to Protein A, or a second protein or peptide that is a secreted bacterial protein or a bacterial cell surface protein. In a further embodiment of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Protein A domain D polypeptide (SEQ ID NO:2), domain E (SEQ ID NO:3), domain A (SEQ ID NO:4), domain C (SEQ ID NO:5), domain B (SEQ ID NO:6), or a nucleic acid sequence encoding a Protein A domain D, domain E, domain A, domain C, or domain B polypeptide. In certain aspects a Protein A polypeptide segment will have an amino acid sequence of SEQ ID NO:8. Similarity or identity, with identity being preferred, is known in the art and a number of different programs can be used to identify whether a protein (or nucleic acid) has sequence identity or similarity to a known sequence. Sequence identity and/or similarity is determined using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith & Waterman (1981), by the sequence identity alignment algorithm of Needleman & Wunsch (1970), by the search for similarity method of Pearson & Lipman (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux et al. (1984), preferably using the default settings, or by inspection. Preferably, percent identity is calculated by using alignment tools known to and readily ascertainable to those of skill in the art. Percent identity is essentially the number of identical amino acids divided by the total number of amino acids compared times one hundred.

**[0043]** Still further embodiments include methods for stimulating in a subject a protective or therapeutic immune response against a *staphylococcus* bacterium comprising administering to the subject an effective amount of a composition including (i) a SpA variant, e.g., a variant SpA domain D polypeptide or peptide thereof; or, (ii) a nucleic acid molecule encoding such a SpA variant polypeptide or peptide thereof, or (iii) administering a SpA variant domain D polypeptide with any combination or permutation of bacterial proteins described herein. In a preferred embodiment the composition is not a *staphylococcus* bacterium. In certain aspects the subject is a human or a cow. In a further aspect the composition is formulated in a pharmaceutically acceptable formulation. The staphylococci may be *Staphylococcus aureus*.

**[0044]** Yet still further embodiments include vaccines comprising a pharmaceutically acceptable composition having an isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein, wherein the composition is capable of stimulating an immune response against a *staphylococcus* bacterium. The vaccine may comprise an isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described. In certain aspects of the invention the isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described are multimerized, e.g., dimerized or concatamerized. In a further aspect, the vaccine composition is contaminated by less than about 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, 0.5, 0.25, 0.05% (or any range derivable therein) of other Staphylococcal proteins. A composition may further comprise an isolated non-SpA polypeptide. Typically the vaccine comprises an adjuvant. In certain aspects a protein or peptide of the invention is linked (covalently or non-covalently) to the adjuvant, preferably the adjuvant is chemically conjugated to the protein.

**[0045]** In still yet further embodiments, a vaccine composition is a pharmaceutically acceptable composition having a recombinant nucleic acid encoding all or part of a SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein, wherein the composition is capable of stimulating an immune response against a *staphylococcus* bacteria. The vaccine composition may comprise a recombinant nucleic acid encoding all or part of a SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein. In certain embodiments the recombinant nucleic acid contains a heterologous promoter. Preferably the recombinant nucleic acid is a vector. More preferably the vector is a plasmid or a viral vector. In some aspects the vaccine includes a recombinant, non-staphylococcus bacterium containing the nucleic acid. The recombinant non-staphylococci may be *Salmonella* or another gram-positive bacteria. The vaccine may comprise a pharmaceutically acceptable excipient, more preferably an adjuvant.

**[0046]** Still further embodiments include methods for stimulating in a subject a protective or therapeutic immune response against a *staphylococcus* bacterium comprising administering to the subject an effective amount of a composition of a SpA variant polypeptide or segment/fragment thereof and further comprising one or more of a Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, or vWh protein or peptide thereof. In a preferred embodiment the composition comprises a non-staphylococcus bacterium. In a further aspect the composition is formulated in a pharmaceutically acceptable formulation. The staphylococci for which a subject is being treated may be *Staphylococcus aureus*. Methods of the invention also include SpA variant compositions that contain 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or more secreted virulence factors and/or cell surface proteins, such as Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, or vWh in various combinations. In certain aspects a vaccine formulation includes Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, and vWh. In certain aspects an antigen combination can include (1) a SpA variant and IsdA; (2) SpA variant and ClfB; (3) SpA variant and SdrD; (4) SpA variant and Hla or Hla variant; (5) SpA variant

and ClfB, SdrD, and Hla or Hla variant; (6) SpA variant, IsdA, SdrD, and Hla or Hla variant; (7) SpA variant, IsdA, ClfB, and Hla or Hla variant; (8) SpA variant, IsdA, ClfB, and SdrD; (9) SpA variant, IsdA, ClfB, SdrD and Hla or Hla variant; (10) SpA variant, IsdA, ClfB, and SdrD; (11) SpA variant, IsdA, SdrD, and Hla or Hla variant; (12) SpA variant, IsdA, and Hla or Hla variant; (13) SpA variant, IsdA, ClfB, and Hla or Hla variant; (14) SpA variant, ClfB, and SdrD; (15) SpA variant, ClfB, and Hla or Hla variant; or (16) SpA variant, SdrD, and Hla or Hla variant.

**[0047]** In certain aspects, a bacterium delivering a composition of the invention will be limited or attenuated with respect to prolonged or persistent growth or abscess formation. In yet a further aspect, SpA variant(s) can be overexpressed in an attenuated bacterium to further enhance or supplement an immune response or vaccine formulation.

**[0048]** The term "EsxA protein" refers to a protein that includes isolated wild-type EsxA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsxA proteins.

**[0049]** The term "EsxB protein" refers to a protein that includes isolated wild-type EsxB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsxB proteins.

**[0050]** The term "SdrD protein" refers to a protein that includes isolated wild-type SdrD polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrD proteins. For example, a wild type SdrD amino acid sequence is provided in NCBI accession no. CAA06651 (SEQ ID NO:65). A SrdD polypeptide for use an antigen according to the embodiments can comprise an amino acid sequence comprising SEQ ID NO:65 or a sequence at least about 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% identical to SEQ ID NO:65. In a further aspect, the SrdD polypeptide comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid segments comprising about, at least or at most 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 to 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 30, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300 or 1315 amino acids in length, including all values and ranges there between, that are at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical to amino acid segments of SEQ ID NO:65.

**[0051]** The term "SdrE protein" refers to a protein that includes isolated wild-type SdrE polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrE proteins.

**[0052]** The term "IsdA protein" refers to a protein that includes isolated wild-type IsdA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdA proteins.

**[0053]** The term "IsdB protein" refers to a protein that includes isolated wild-type IsdB polypeptides from *staphy-*

*lococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdB proteins.

**[0054]** The term “Eap protein” refers to a protein that includes isolated wild-type Eap polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Eap proteins.

**[0055]** The term “Ebh protein” refers to a protein that includes isolated wild-type Ebh polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Ebh proteins.

**[0056]** The term “Emp protein” refers to a protein that includes isolated wild-type Emp polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Emp proteins.

**[0057]** The term “EsaB protein” refers to a protein that includes isolated wild-type EsaB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsaB proteins.

**[0058]** The term “EsaC protein” refers to a protein that includes isolated wild-type EsaC polypeptides, from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsaC proteins.

**[0059]** The term “SdrC protein” refers to a protein that includes isolated wild-type SdrC polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrC proteins.

**[0060]** The term “ClfA protein” refers to a protein that includes isolated wild-type ClfA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria ClfA proteins. For example, a wild type ClfA amino acid sequence is provided in NCBI accession no. YP\_001331790 (SEQ ID NO:66). A ClfA polypeptide for use as an antigen according to the embodiments can comprise an amino acid sequence comprising SEQ ID NO:66 or a sequence at least about 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% identical to SEQ ID NO:66. In a further aspect, the ClfA polypeptide comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid segments comprising about, at least or at most 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 to 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 30, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 550, 600, 650, 700, 750, 800, 850, 900, or 933 amino acids in length, including all values and ranges there between, that are at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical to amino acid segments of SEQ ID NO:66.

**[0061]** The term “ClfB protein” refers to a protein that includes isolated wild-type ClfB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria ClfB proteins.

**[0062]** The term “Coa protein” refers to a protein that includes isolated wild-type Coa polypeptides from *staphylo-*

*coccus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Coa proteins.

**[0063]** The term “FnbpB protein” or “FnbB protein” refers to a protein that includes isolated wild-type FnbpB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria FnbpB proteins. For example, a wild type FnbpB amino acid sequence is provided in NCBI accession no. YP\_001333431 (SEQ ID NO:67). A FnbpB polypeptide for use as an antigen according to the embodiments can comprise an amino acid sequence comprising SEQ ID NO:67 or a sequence at least about 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% identical to SEQ ID NO:67. In a further aspect, the FnbpB polypeptide comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid segments comprising about, at least or at most 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 to 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 30, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 550, 600, 650 or 677 amino acids in length, including all values and ranges there between, that are at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical to amino acid segments of SEQ ID NO:67.

**[0064]** The term “Hla protein” refers to a protein that includes isolated wild-type Hla polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Hla proteins.

**[0065]** The term “IsdC protein” refers to a protein that includes isolated wild-type IsdC polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdC proteins.

**[0066]** The term “SasF protein” refers to a protein that includes isolated wild-type SasF polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SasF proteins.

**[0067]** The term “vWbp protein” refers to a protein that includes isolated wild-type vWbp (von Willebrand factor binding protein) polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria vWbp proteins.

**[0068]** The term “vWh protein” refers to a protein that includes isolated wild-type vWh (von Willebrand factor binding protein homolog) polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria vWh proteins.

**[0069]** An immune response refers to a humoral response, a cellular response, or both a humoral and cellular response in an organism. An immune response can be measured by assays that include, but are not limited to, assays measuring the presence or amount of antibodies that specifically recognize a protein or cell surface protein, assays measuring T-cell activation or proliferation, and/or assays that measure modulation in terms of activity or expression of one or more cytokines.



**[0070]** In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsxA protein. In certain aspects the EsxA protein will have all or part of the amino acid sequence of SEQ ID NO:11.

**[0071]** In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsxB protein. In certain aspects the EsxB protein will have all or part of the amino acid sequence of SEQ ID NO:12.

**[0072]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an SdrD protein. In certain aspects the SdrD protein will have all or part of the amino acid sequence of SEQ ID NO:13.

**[0073]** In further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an SdrE protein. In certain aspects the SdrE protein will have all or part of the amino acid sequence of SEQ ID NO:14.

**[0074]** In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdA protein. In certain aspects the IsdA protein will have all or part of the amino acid sequence of SEQ ID NO:15.

**[0075]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdB protein. In certain aspects the IsdB protein will have all or part of the amino acid sequence of SEQ ID NO:16.

**[0076]** Embodiments of the invention include compositions that include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a EsaB protein. In certain aspects the EsaB protein will have all or part of the amino acid sequence of SEQ ID NO:17.

**[0077]** In a further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a ClfB protein. In certain aspects the ClfB protein will have all or part of the amino acid sequence of SEQ ID NO:18.

**[0078]** In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdC protein. In certain aspects the IsdC protein will have all or part of the amino acid sequence of SEQ ID NO:19.

**[0079]** In yet further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a SasF protein. In certain aspects the SasF protein will have all or part of the amino acid sequence of SEQ ID NO:20.

**[0080]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a SdrC protein. In

certain aspects the SdrC protein will have all or part of the amino acid sequence of SEQ ID NO:21.

**[0081]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a ClfA protein. In certain aspects the ClfA protein will have all or part of the amino acid sequence of SEQ ID NO:22.

**[0082]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Eap protein. In certain aspects the Eap protein will have all or part of the amino acid sequence of SEQ ID NO:23.

**[0083]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Ebh protein. In certain aspects the Ebh protein will have all or part of the amino acid sequence of SEQ ID NO:24.

**[0084]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Emp protein. In certain aspects the Emp protein will have all or part of the amino acid sequence of SEQ ID NO:25.

**[0085]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsaC protein. In certain aspects the EsaC protein will have all or part of the amino acid sequence of SEQ ID NO:26. Sequence of EsaC polypeptides can be found in the protein databases and include, but are not limited to accession numbers ZP\_02760162 (GI:168727885), NP\_645081.1 (GI:21281993), and NP\_370813.1 (GI:15923279), each of which is incorporated herein by reference as of the priority date of this application.

**[0086]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Coa protein. In certain aspects the Coa protein will have all or part of the amino acid sequence of SEQ ID NO:27.

**[0087]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Hla protein. In certain aspects the Hla protein will have all or part of the amino acid sequence of SEQ ID NO:28.

**[0088]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a vWa protein. In certain aspects the vWa protein will have all or part of the amino acid sequence of SEQ ID NO:29.

**[0089]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a vWbp protein. In certain aspects the vWbp protein will have all or part of the amino acid sequence of SEQ ID NO:32.

**[0090]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein

that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a FnbpB protein. In certain aspects the FnbpB protein will have all or part of the amino acid sequence of SEQ ID NO:64.

**[0091]** In certain aspects, a polypeptide or segment/fragment can have a sequence that is at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% or more identical to the amino acid sequence of the reference polypeptide. The term "similarity" refers to a polypeptide that has a sequence that has a certain percentage of amino acids that are either identical with the reference polypeptide or constitute conservative substitutions with the reference polypeptides.

**[0092]** The polypeptides described herein may include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more variant amino acids within at least, or at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:2-30, or SEQ ID NO:32-34.

**[0093]** A polypeptide segment as described herein may include 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:2-30, or SEQ ID NO:33-34, or SEQ ID NO:64.

**[0094]** The compositions may be formulated in a pharmaceutically acceptable composition. In certain aspects of the invention the *staphylococcus* bacterium is an *S. aureus* bacterium.

**[0095]** In further aspects, a composition may be administered more than one time to the subject, and may be administered 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20 or more times. The

administration of the compositions include, but is not limited to oral, parenteral, subcutaneous, intramuscular, intravenous, or various combinations thereof, including inhalation or aspiration.

**[0096]** In still further embodiments, a composition comprises a recombinant nucleic acid molecule encoding a polypeptide described herein or segments/fragments thereof. Typically a recombinant nucleic acid molecule encoding a polypeptide described herein contains a heterologous promoter. In certain aspects, a recombinant nucleic acid molecule of the invention is a vector, in still other aspects the vector is a plasmid. In certain embodiments the vector is a viral vector. In certain aspects a composition includes a recombinant, non-*staphylococcus* bacterium containing or expressing a polypeptide described herein. In particular aspects the recombinant non-*staphylococcus* bacteria is *Salmonella* or another gram-positive bacteria. A composition is typically administered to mammals, such as human subjects, but administration to other animals that are capable of eliciting an immune response is contemplated. In further aspects the *staphylococcus* bacterium containing or expressing the polypeptide is *Staphylococcus aureus*. In further embodiments the immune response is a protective immune response.

**[0097]** In further embodiments a composition comprises a recombinant nucleic acid molecule encoding all or part of one or more of a Eap, Ehb, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, SpA, vWbp, or vWh protein or peptide or variant thereof. Additional staphylococcal antigens that can be used in combination with the polypeptides described herein include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein. In particular aspects, a bacteria is a recombinant non-*staphylococcus* bacteria, such as a *Salmonella* or other gram-positive bacteria.

**[0098]** Compositions of the invention are typically administered to human subjects, but administration to other animals that are capable of eliciting an immune response to a *staphylococcus* bacterium is contemplated, particularly cattle, horses, goats, sheep and other domestic animals, i.e., mammals.

**[0099]** In certain aspects the *staphylococcus* bacterium is a *Staphylococcus aureus*. In further embodiments the immune response is a protective immune response. In still further aspects, the methods and compositions of the invention can be used to prevent, ameliorate, reduce, or treat infection of tissues or glands, e.g., mammary glands, particularly mastitis and other infections. Other methods include, but are not limited to prophylactically reducing bacterial burden in a subject not exhibiting signs of infection, particularly those subjects suspected of or at risk of being colonized by a target bacteria,

e.g., patients that are or will be at risk or susceptible to infection during a hospital stay, treatment, and/or recovery.

**[0100]** Any embodiment discussed with respect to one aspect of the invention applies to other aspects of the invention as well. In particular, any embodiment discussed in the context of a SpA variant polypeptide or peptide or nucleic acid may be implemented with respect to other antigens, such as Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein (or nucleic acids), and vice versa. It is also understood that any one or more of Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein can be specifically excluded from a claimed composition.

**[0101]** Embodiments of the invention include compositions that contain or do not contain a bacterium. A composition may or may not include an attenuated or viable or intact staphylococcal bacterium. In certain aspects, the composition comprises a bacterium that is not a staphylococcal bacterium or does not contain staphylococcal bacteria. In certain embodiments a bacterial composition comprises an isolated or recombinantly expressed staphylococcal Protein A variant or a nucleotide encoding the same. The composition may be or include a recombinantly engineered *staphylococcus* bacterium that has been altered in a way that comprises specifically altering the bacterium with respect to a secreted virulence factor or cell surface protein. For example, the bacteria may be recombinantly modified to express more of the virulence factor or cell surface protein than it would express if unmodified.

**[0102]** The term “isolated” can refer to a nucleic acid or polypeptide that is substantially free of cellular material, bacterial material, viral material, or culture medium (when produced by recombinant DNA techniques) of their source of

origin, or chemical precursors or other chemicals (when chemically synthesized). Moreover, an isolated compound refers to one that can be administered to a subject as an isolated compound; in other words, the compound may not simply be considered “isolated” if it is adhered to a column or embedded in an agarose gel. Moreover, an “isolated nucleic acid fragment” or “isolated peptide” is a nucleic acid or protein fragment that is not naturally occurring as a fragment and/or is not typically in the functional state.

**[0103]** Moieties of the invention, such as polypeptides, peptides, antigens, or immunogens, may be conjugated or linked covalently or noncovalently to other moieties such as adjuvants, proteins, peptides, supports, fluorescence moieties, or labels. The term “conjugate” or “immunoconjugate” is broadly used to define the operative association of one moiety with another agent and is not intended to refer solely to any type of operative association, and is particularly not limited to chemical “conjugation.” Recombinant fusion proteins are particularly contemplated. Compositions of the invention may further comprise an adjuvant or a pharmaceutically acceptable excipient. An adjuvant may be covalently or non-covalently coupled to a polypeptide or peptide of the invention. In certain aspects, the adjuvant is chemically conjugated to a protein, polypeptide, or peptide.

**[0104]** The term “providing” is used according to its ordinary meaning to indicate “to supply or furnish for use.” In some embodiments, the protein is provided directly by administering the protein, while in other embodiments, the protein is effectively provided by administering a nucleic acid that encodes the protein. In certain aspects the invention contemplates compositions comprising various combinations of nucleic acid, antigens, peptides, and/or epitopes.

**[0105]** The subject will have (e.g., are diagnosed with a staphylococcal infection), will be suspected of having, or will be at risk of developing a staphylococcal infection. Compositions of the present invention include immunogenic compositions wherein the antigen(s) or epitope(s) are contained in an amount effective to achieve the intended purpose. More specifically, an effective amount means an amount of active ingredients necessary to stimulate or elicit an immune response, or provide resistance to, amelioration of, or mitigation of infection. In more specific aspects, an effective amount prevents, alleviates or ameliorates symptoms of disease or infection, or prolongs the survival of the subject being treated. Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any preparation used in the methods of the invention, an effective amount or dose can be estimated initially from *in vitro* studies, cell culture, and/or animal model assays. For example, a dose can be formulated in animal models to achieve a desired immune response or circulating antibody concentration or titer. Such information can be used to more accurately determine useful doses in humans.

**[0106]** The embodiments in the Example section are understood to be embodiments of the invention that are applicable to all aspects of the invention.

**[0107]** The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” It is also contemplated that anything listed using the term “or” may also be specifically excluded.

**[0108]** Throughout this application, the term “about” is used to indicate that a value includes the standard deviation of error for the device or method being employed to determine the value.

**[0109]** Following long-standing patent law, the words “a” and “an,” when used in conjunction with the word “comprising” in the claims or specification, denotes one or more, unless specifically noted.

**[0110]** Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

#### DESCRIPTION OF THE DRAWINGS

**[0111]** So that the matter in which the above-recited features, advantages and objects of the invention as well as others which will become clear are attained and can be understood in detail, more particular descriptions and certain embodiments of the invention briefly summarized above are illustrated in the appended drawings. These drawings form a part of the specification. It is to be noted, however, that the appended drawings illustrate certain embodiments of the invention and therefore are not to be considered limiting in their scope.

**[0112]** FIGS. 1A-1B. (FIG. 1A) Primary structure of the Protein A precursor with an N-terminal YSIRK motif signal peptide, five immunoglobulin binding domains as tandem repeats designated E, D, A, B, C, region X, and the LPXTG sorting signal. (FIG. 1B) Following synthesis of the Protein A precursor, staphylococci secrete this product via the Sec pathway, and sortase A cleaves the LPXTG sorting signal between the T and G residues. Nucleophilic attack of the amino group within lipid II at the sortase-Protein A thioester-linked intermediate forms the amide bond that links Protein A to the cell wall envelope and enables its display on the bacterial surface.

**[0113]** FIG. 2. Three dimensional model of the molecular interactions between the SpA-domain D of Protein A, the VH3 Fab domain of the B cell receptor, and of the Fc $\gamma$  domain of immunoglobulin. The model is derived from two crystal structures (Graillie et al., 2000 and Gouda et al., 1992) that revealed side chain residues involved in the formation of ionic bonds that enable these complexes. Gln-9 and Gln-10 of SpA-D promote binding to Fc $\gamma$ , whereas Asp-36 and Asp-37 enable complex formation with VH3 Fab.

**[0114]** FIG. 3. Left panel—Coomassie Blue stained SDS-PAGE reveals the migrational position of purified His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub>, human IgG, and sortase A (SrtA), a control protein. Right panel—Coomassie Blue stained SDS-PAGE to reveal the elution of Protein A immunoglobulin complexes eluted following affinity chromatography of human IgG on Ni-NTA columns pre-charged with His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or SrtA.

**[0115]** FIG. 4. ELISA assays to quantify human immunoglobulin (hIgG), human F(ab)<sub>2</sub> IgG fragments and human Fc fragments of immunoglobulin (hFc). Plates were coated with equal amounts of His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or SrtA. hIgG-HRP, F(ab)<sub>2</sub>-HRP and hFc-HRP were added onto the plates and incubated for an hour. Absorbance at 450 nm was recorded and plotted to determine the half maximal titers.

**[0116]** FIG. 5. Purified SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or a PBS mock control were injected into the peritoneum of mice and analyzed for their ability to reduce the B cell population in the spleen of experimental BALB/c mice. Animals were killed 4 hours following injection, their spleen removed, tissue homogenized and stained with CD19 antibodies directed against B cells. The number of B cells was quantified by FACS sorting.

**[0117]** FIG. 6 Generation of a non-toxicigenic protein A vaccine. a, Translational protein A (SpA) product of *S. aureus* Newman and USA300 LAC with an N-terminal signal peptide (white box), five immunoglobulin binding domains (IgBDs designated E, D, A, B and C), variable region X and C-terminal sorting signal (black box). b, Amino acid sequence of the five IgBDs as well as nontoxicigenic SpA-D<sub>KKAA</sub>, with the positions of triple  $\alpha$ -helical bundles (H1, H2 and H3) as well as glutamine (Q) 9, 10 and aspartate (D) 36, 37 indicated. c, Coomassie Blue-stained SDS-PAGE of SpA, SpA-D, SpA-D<sub>KKAA</sub> or SrtA purified on Ni-NTA sepharose in the presence or absence of human immunoglobulin (hIgG). d, ELISA examining the association of immobilized SpA, SpA-D or SpA-D<sub>KKAA</sub> with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments and von Willebrand factor (vWF). e, CD19+ B lymphocytes in splenic tissue of BALB/c mice that had been mock immunized or treated with SpA-D or SpA-D<sub>KKAA</sub> were quantified by FACS.

**[0118]** FIG. 7 Non-toxicigenic protein A vaccine prevents abscess formation. Histopathology of renal tissue isolated during necropsy of BALB/c mice that had been mock immunized (PBS) or vaccinated with SpA, SpA-D as well as SpA-D<sub>KKAA</sub> and challenged with *S. aureus* Newman. Thin sectioned tissues were stained with hematoxylin-eosin. White arrows identify polymorphonuclear leukocyte (PMN) infiltrates. Dark arrows identify staphylococcal abscess communities.

**[0119]** FIG. 8 Antibodies raised by the non-toxicigenic protein A vaccine block the B cell superantigen function of SpA. a, Rabbit antibodies raised against SpA-D<sub>KKAA</sub> were purified on a matrix with immobilized antigen and analyzed by Coomassie Blue-stained SDS-PAGE. Antibodies were cleaved with pepsin and F(ab)<sub>2</sub> fragments were purified by a second round of affinity chromatography on SpA-D<sub>KKAA</sub> matrix. b, SpA-D<sub>KKAA</sub> specific F(ab)<sub>2</sub> interfere with the binding of SpA or SpA-D to human immunoglobulin (hIgG) or, c, to von Willebrand Factor (vWF).

**[0120]** FIG. 9 Full-length non-toxicigenic protein A generates improved immune responses. a, Full-length SpA<sub>KKAA</sub> was purified on Ni-NTA sepharose and analyzed by Coomassie-Blue stained SDS-PAGE. b, CD19+ B lymphocytes in splenic tissue of BALB/c mice that had been mock immunized or treated with SpA or SpA<sub>KKAA</sub> were quantified by FACS. c, ELISA examining the association of immobilized SpA or SpA<sub>KKAA</sub> with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments or von Willebrand factor (vWF). d, Human or mouse serum antibody titers to diphtheria toxoid (CRM197) and non-toxicigenic SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub>. Human volunteers with a history of DTaP immunization and staphylococcal infection (n=16) as well as mice (n=20) that had been infected with *S. aureus* Newman or USA 300 LAC or immunized with SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub> were examined by quantitative dot blot.

**[0121]** FIG. 10 Staphylococcal infection does not generate protective immunity. BALB/c mice (n=20) were infected with *S. aureus* Newman or mock challenged (PBS) for thirty

days and infection cleared with chloramphenicol treatment. Both cohorts of animals were then challenged with *S. aureus* Newman and bacterial load (CFU) in kidney tissue homogenate analyzed following necropsy on day 4.

**[0122]** FIG. 11 Comparison of abscess formation in mice treated with PBS, SpA, SpA-D, and SpA-D<sub>KKAA</sub>.

**[0123]** FIGS. 12A-12C (A) ELISA examining the association of immobilized SpA, SpA-D, SpA-DKKAA or SpA-DGGSS with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments and IgM. Statistical significance of SpA-DKKAA and SpA-DGGSS binding to each ligand was compared against SpA-D; SpA-D binding was compared against SpA (n=3); \* signifies P<0.05; \*\* signifies P<0.01. (B) ELISA examining the level of cross-reactive antibodies of hyper-immune sera samples collected from actively immunized mice (n=5) with SpA-D, SpA-DKKAA and SpA-DGGSS. (C) Abscess formation in mice treated with PBS, SpA-D, SpA-D<sub>KKAA</sub> and SpA-D<sub>GGSS</sub>.

**[0124]** FIGS. 13A-13B BALB/c mice (n=18-20) were either mock immunized with PBS/adjuvant or injected with 25 µg of each antigen (Combo 1, ClfA+SdrD+FnBPB; Combo 2, Combo 1+SpA<sub>KKAA</sub>). Immunized mice were challenged by intravenous inoculation with 1×10<sup>7</sup> CFU *S. aureus* Newman. Bacterial loads in kidney tissues were examined at A, day 4 and B, day 18 post challenge. Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values <0.05 were deemed significant.

**[0125]** FIGS. 14A-14H. Active Immunization with Antigens Revealed by Genetic Vaccinology Elicits Protection in Mice against Staphylococcal Abscess Formation. Cohorts of BALB/c mice (n=18-20) were actively immunized with mock (PBS), Combo 1 (ClfA, FnBPB and SdrD) or Combo 2 (ClfA, FnBPB, SdrD and SpA<sub>KKAA</sub>) at day 0 and 11. On day 21, animals were challenged by retro-orbital injection with 1×10<sup>7</sup> CFU *S. aureus* Newman. On days 4 (A) and 18 (B) post challenge, animals were killed to enumerate staphylococcal burden in renal tissues. (C—H) Representative thin-sectioned, hematoxylin-eosin stained histopathology slides from each cohort (n=10, 4 days post challenge) are shown. White arrowheads identify polymorphonuclear leukocyte (PMN) infiltrates. Dark arrowheads identify staphylococcal abscess communities. Animal data are representative of two independent experiments.

**[0126]** FIG. 15 Active Immunization with Antigens Revealed by Genetic Vaccinology Elicits Protection in Mice against Staphylococcal Sepsis. Cohorts of BALB/c mice (n=20) were actively immunized with mock (PBS), Combo 1 (ClfA, FnBPB and SdrD) or Combo 2 (ClfA, FnBPB, SdrD and SpA<sub>KKAA</sub>) at day 0 and 11. On day 21, animals were challenged by retro-orbital injection with 1×10<sup>8</sup> CFU *S. aureus* Newman and monitored for survival. Animal data are representative of two independent experiments.

#### DETAILED DESCRIPTION

**[0127]** *Staphylococcus aureus* is a commensal of the human skin and nares, and the leading cause of bloodstream, skin and soft tissue infections (Klevens et al., 2007). Recent dramatic increases in the mortality of staphylococcal diseases are attributed to the spread of methicillin-resistant *S. aureus* (MRSA) strains often not susceptible to antibiotics (Kennedy et al., 2008). In a large retrospective study, the incidence of MRSA infections was 4.6% of all hospital admissions in the United States (Klevens et al., 2007). The annual health care

costs for 94,300 MRSA infected individuals in the United States exceed \$2.4 billion (Klevens et al., 2007). The current MRSA epidemic has precipitated a public health crisis that needs to be addressed by development of a preventive vaccine (Boucher and Corey, 2008). To date, an FDA licensed vaccine that prevents *S. aureus* diseases is not available.

**[0128]** The inventors describe here the use of Protein A, a cell wall anchored surface protein of staphylococci, for the generation of variants that can serve as subunit vaccines. The pathogenesis of staphylococcal infections is initiated as bacteria invade the skin or blood stream via trauma, surgical wounds, or medical devices (Lowy, 1998). Although the invading pathogen may be phagocytosed and killed, staphylococci can also escape innate immune defenses and seed infections in organ tissues, inducing inflammatory responses that attract macrophages, neutrophils, and other phagocytes (Lowy, 1998). The responsive invasion of immune cells to the site of infection is accompanied by liquefaction necrosis as the host seeks to prevent staphylococcal spread and allow for removal of necrotic tissue debris (Lam et al., 1963). Such lesions can be observed by microscopy as hypercellular areas containing necrotic tissue, leukocytes, and a central nidus of bacteria (Lam et al., 1963). Unless staphylococcal abscesses are surgically drained and treated with antibiotics, disseminated infection and septicemia produce a lethal outcome (Sheagren, 1984).

#### I. Staphylococcal Antigens

**[0129]** A. Staphylococcal Protein A (SpA)

**[0130]** All *Staphylococcus aureus* strains express the structural gene for Protein A (spa) (Jensen, 1958; Said-Salim et al., 2003), a well characterized virulence factor whose cell wall anchored surface protein product (SpA) encompasses five highly homologous immunoglobulin binding domains designated E, D, A, B, and C (Sjodahl, 1977). These domains display ~80% identity at the amino acid level, are 56 to 61 residues in length, and are organized as tandem repeats (Uhlen et al., 1984). SpA is synthesized as a precursor protein with an N-terminal YSIRK/GS signal peptide and a C-terminal LPXTG motif sorting signal (DeDent et al., 2008; Schneewind et al., 1992). Cell wall anchored Protein A is displayed in great abundance on the staphylococcal surface (DeDent et al., 2007; Sjoquist et al., 1972). Each of its immunoglobulin binding domains is composed of anti-parallel α-helices that assemble into a three helix bundle and bind the Fc domain of immunoglobulin G (IgG) (Deisenhofer, 1981; Deisenhofer et al., 1978), the VH3 heavy chain (Fab) of IgM (i.e., the B cell receptor) (Graille et al., 2000), the von Willibrand factor at its A1 domain [vWF A1 is a ligand for platelets] (O'Seaghdha et al., 2006) and the tumor necrosis factor α (TNF-α) receptor I (TNFRI) (Gomez et al., 2006), which is displayed on surfaces of airway epithelia (Gomez et al., 2004; Gomez et al., 2007).

**[0131]** SpA impedes neutrophil phagocytosis of staphylococci through its attribute of binding the Fc component of IgG (Jensen, 1958; Uhlen et al., 1984). Moreover, SpA is able to activate intravascular clotting via its binding to von Willibrand factor A1 domains (Hartleib et al., 2000). Plasma proteins such as fibrinogen and fibronectin act as bridges between staphylococci (ClfA and ClfB) and the platelet integrin GPIIb/IIIa (O'Brien et al., 2002), an activity that is supplemented through Protein A association with vWF A1, which allows staphylococci to capture platelets via the GPIb-α platelet receptor (Foster, 2005; O'Seaghdha et al.,

2006). SpA also binds TNFR1 and this interaction contributes to the pathogenesis of staphylococcal pneumonia (Gomez et al., 2004). SpA activates proinflammatory signaling through TNFR1 mediated activation of TRAF2, the p38/c-Jun kinase, mitogen activate protein kinase (MAPK) and the Rel-transcription factor NF-KB. SpA binding further induces TNFR1 shedding, an activity that appears to require the TNF-converting enzyme (TACE) (Gomez et al., 2007). All of the aforementioned SpA activities are mediated through its five IgG binding domains and can be perturbed by the same amino acid substitutions, initially defined by their requirement for the interaction between Protein A and human IgG1 (Cedergren et al., 1993).

**[0132]** SpA also functions as a B cell superantigen by capturing the Fab region of VH3 bearing IgM, the B cell receptor (Gomez et al., 2007; Goodyear et al., 2003; Goodyear and Silverman, 2004; Roben et al., 1995). Following intravenous challenge, staphylococcal Protein A (SpA) mutations show a reduction in staphylococcal load in organ tissues and dramatically diminished ability to form abscesses (described herein). During infection with wildtype *S. aureus*, abscesses are formed within forty-eight hours and are detectable by light microscopy of hematoxylin-eosin stained, thin-sectioned kidney tissue, initially marked by an influx of polymorphonuclear leukocytes (PMNs). On day 5 of infection, abscesses increase in size and enclosed a central population of staphylococci, surrounded by a layer of eosinophilic, amorphous material and a large cuff of PMNs. Histopathology revealed massive necrosis of PMNs in proximity to the staphylococcal nidus at the center of abscess lesions as well as a mantle of healthy phagocytes. The inventors also observed a rim of necrotic PMNs at the periphery of abscess lesions, bordering the eosinophilic pseudocapsule that separated healthy renal tissue from the infectious lesion. Staphylococcal variants lacking Protein A are unable to establish the histopathology features of abscesses and are cleared during infection.

**[0133]** In previous studies, Cedergren et al. (1993) engineered five individual substitutions in the Fc fragment binding sub-domain of the B domain of SpA, L17D, N28A, I31A and K35A. These authors created these proteins to test data gathered from a three dimensional structure of a complex between one domain of SpA and Fc<sub>1</sub>. Cedergren et al. determined the effects of these mutations on stability and binding, but did not contemplate use of such substitutions for the production of a vaccine antigen.

**[0134]** Brown et al. (1998) describe studies designed to engineer new proteins based on SpA that allow the use of more favorable elution conditions when used as affinity ligands. The mutations studied included single mutations of Q13A, Q 14H, N15A, N15H, F 17H, Y18F, L21H, N32H, or K39H. Brown et al. report that Q13A, N15A, N15H, and N32H substitutions made little difference to the dissociation constant values and that the Y18F substitution resulted in a 2 fold decrease in binding affinity as compared to wild type SpA. Brown et al. also report that L21H and F 17H substitutions decrease the binding affinity by five-fold and a hundred-fold respectively. The authors also studied analogous substitutions in two tandem domains. Thus, the Brown et al. studies were directed to generating a SpA with a more favorable elution profile, hence the use of H is substitutions to provide a pH sensitive alteration in the binding affinity. Brown et al. is silent on the use of SpA as a vaccine antigen.

**[0135]** Graille et al. (2000) describe a crystal structure of domain D of SpA and the Fab fragment of a human IgM

antibody. Graille et al. define by analysis of a crystal structure the D domain amino acid residues that interact with the Fab fragment as residues Q26, G29, F30, Q32, S33, D36, D37, Q40, N43, E47, or L51, as well as the amino acid residues that form the interface between the domain D sub-domains. Graille et al. define the molecular interactions of these two proteins, but is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

**[0136]** O'Seaghda et al. (2006) describe studies directed at elucidating which sub-domain of domain D binds vWF. The authors generated single mutations in either the Fc or VH3 binding sub-domains, i.e., amino acid residues F5A, Q9A, Q10A, F13A, Y14A, L17A, N28A, I31A, K35A, G29A, F30A, S33A, D36A, D37A, Q40A, E47A, or Q32A. The authors discovered that vWF binds the same sub-domain that binds Fc. O'Seaghda et al. define the sub-domain of domain D responsible for binding vWF, but is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

**[0137]** Gomez et al. (2006) describe the identification of residues responsible for activation of the TNFR1 by using single mutations of F5A, F13A, Y14A, L17A, N21A, I31A, Q32A, and K35A. Gomez et al. is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

**[0138]** Recombinant affinity tagged Protein A, a polypeptide encompassing the five IgG domains (EDCAB) (Sjodahl, 1977) but lacking the C-terminal Region X (Guss et al., 1984), was purified from recombinant *E. coli* and used as a vaccine antigen (Stranger-Jones et al., 2006). Because of the attributes of SpA in binding the Fc portion of IgG, a specific humoral immune response to Protein A could not be measured (Stranger-Jones et al., 2006). The inventors have overcome this obstacle through the generation of SpA-DQ9,10K; D36,37A. BALB/c mice immunized with recombinant Protein A (SpA) displayed significant protection against intravenous challenge with *S. aureus* strains: a 2.951 log reduction in staphylococcal load as compared to the wild-type (P>0.005; Student's t-test) (Stranger-Jones et al., 2006). SpA specific antibodies may cause phagocytic clearance prior to abscess formation and/or impact the formation of the aforementioned eosinophilic barrier in abscesses that separate staphylococcal communities from immune cells since these do not form during infection with Protein A mutant strains. Each of the five SpA domains (i.e., domains formed from three helix bundles designated E, D, A, B, and C) exerts similar binding properties (Jansson et al., 1998). The solution and crystal structure of the domain D has been solved both with and without the Fc and VH3 (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille et al., 2000). Mutations in residues known to be involved in IgG binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF AI and TNFR1 binding (Cedergren et al., 1993; Gomez et al., 2006; O'Seaghda et al., 2006), whereas residues important for the VH3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) appear to have no impact on the other binding activities (Graille et al., 2000; Jansson et al., 1998). SpA specifically targets a subset of B cells that express VH3 family related IgM on their surface, i.e., VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells proliferate and commit to apoptosis, leading to preferential and pro-

longed deletion of innate-like B lymphocytes (i.e., marginal zone B cells and follicular B2 cells)(Goodyear et al., 2003; Goodyear et al., 2004).

**[0139]** Molecular Basis of Protein A Surface Display and Function.

**[0140]** Protein A is synthesized as a precursor in the bacterial cytoplasm and secreted via its YSIRK signal peptide at the cross wall, i.e. the cell division septum of staphylococci (FIG. 1) (DeDent et al., 2007; DeDent et al., 2008). Following cleavage of the C-terminal LPXTG sorting signal, Protein A is anchored to bacterial peptidoglycan crossbridges by sortase A (Mazmanian et al., 1999; Schneewind et al., 1995; Mazmanian et al., 2000). Protein A is the most abundant surface protein of staphylococci; the molecule is expressed by virtually all *S. aureus* strains (Cespedes et al., 2005; Kennedy et al., 2008; Said-Salim et al., 2003). Staphylococci turn over 15-20% of their cell wall per division cycle (Navarre and Schneewind, 1999). Murine hydrolases cleave the glycan strands and wall peptides of peptidoglycan, thereby releasing Protein A with its attached C-terminal cell wall disaccharide tetrapeptide into the extracellular medium (Ton-That et al., 1999). Thus, by physiological design, Protein A is both anchored to the cell wall and displayed on the bacterial surface but also released into surrounding tissues during host infection (Marraffini et al., 2006).

**[0141]** Protein A captures immunoglobulins on the bacterial surface and this biochemical activity enables staphylococcal escape from host innate and acquired immune responses (Jensen, 1958; Goodyear et al., 2004). Interestingly, region X of Protein A (Guss et al., 1984), a repeat domain that tethers the IgG binding domains to the LPXTG sorting signal/cell wall anchor, is perhaps the most variable portion of the staphylococcal genome (Said-Salim, 2003; Schneewind et al., 1992). Each of the five immunoglobulin binding domains of Protein A (SpA), formed from three helix bundles and designated E, D, A, B, and C, exerts similar structural and functional properties (Sjodahl, 1977; Jansson et al., 1998). The solution and crystal structure of the domain D has been solved both with and without the Fc and  $V_H3$  (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille 2000).

**[0142]** In the crystal structure complex, the Fab interacts with helix II and helix III of domain D via a surface composed of four VH region  $\beta$ -strands (Graille 2000). The major axis of helix II of domain D is approximately 50° to the orientation of the strands, and the interhelical portion of domain D is most proximal to the C0 strand. The site of interaction on Fab is remote from the Ig light chain and the heavy chain constant region. The interaction involves the following domain D residues: Asp-36 of helix II, Asp-37 and Gln-40 in the loop between helix II and helix III and several other residues (Graille 2000). Both interacting surfaces are composed predominantly of polar side chains, with three negatively charged residues on domain D and two positively charged residues on the 2A2 Fab buried by the interaction, providing an overall electrostatic attraction between the two molecules. Of the five polar interactions identified between Fab and domain D, three are between side chains. A salt bridge is formed between Arg-H19 and Asp-36 and two hydrogen bonds are made between Tyr-H59 and Asp-37 and between Asn-H82a and Ser-33. Because of the conservation of Asp-36 and Asp-37 in all five IgG binding domains of Protein A, the inventors mutated these residues.

**[0143]** The SpA-D sites responsible for Fab binding are structurally separate from the domain surface that mediates Fc $\gamma$  binding. The interaction of Fc $\gamma$  with domain D primarily involves residues in helix I with lesser involvement of helix II (Gouda et al., 1992; Deisenhofer, 1981). With the exception of the Gln-32, a minor contact in both complexes, none of the residues that mediate the Fc $\gamma$  interaction are involved in Fab binding. To examine the spatial relationship between these different Ig-binding sites, the SpA domains in these complexes have been superimposed to construct a model of a complex between Fab, the SpA-domain D, and the Fc $\gamma$  molecule. In this ternary model, Fab and Fc $\gamma$  form a sandwich about opposite faces of the helix II without evidence of steric hindrance of either interaction. These findings illustrate how, despite its small size (i.e., 56-61 aa), an SpA domain can simultaneously display both activities, explaining experimental evidence that the interactions of Fab with an individual domain are noncompetitive. Residues for the interaction between SpA-D and Fc $\gamma$  are Gln-9 and Gln-10.

**[0144]** In contrast, occupancy of the Fc portion of IgG on the domain D blocks its interaction with vWF A1 and probably also TNFR1 (O'Seaghda et al., 2006). Mutations in residues essential for IgG Fc binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF A1 and TNFR1 binding (O'Seaghda et al., 2006; Cedergren et al., 1993; Gomez et al., 2006), whereas residues critical for the VH3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) have no impact on the binding activities of IgG Fc, vWF A1 or TNFR1 (Jansson et al., 1998; Graille et al., 2000). The Protein A immunoglobulin Fab binding activity targets a subset of B cells that express  $V_H3$  family related IgM on their surface, i.e., these molecules function as VH3type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells rapidly proliferate and then commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e., marginal zone B cells and follicular B2 cells) (Goodyear and Silverman, 2004; Goodyear and Silverman, 2003). More than 40% of circulating B cells are targeted by the Protein A interaction and the  $V_H3$  family represents the largest family of human B cell receptors to impart protective humoral responses against pathogens (Goodyear and Silverman, 2004; Goodyear and Silverman, 2003). Thus, Protein A functions analogously to staphylococcal superantigens (Roben et al., 1995), albeit that the latter class of molecules, for example SEB, TSST-1, TSST-2, form complexes with the T cell receptor to inappropriately stimulate host immune responses and thereby precipitating characteristic disease features of staphylococcal infections (Roben et al., 1995; Tiedemann et al., 1995). Together these findings document the contributions of Protein A in establishing staphylococcal infections and in modulating host immune responses.

**[0145]** In sum, Protein A domains can viewed as displaying two different interfaces for binding with host molecules and any development of Protein A based vaccines must consider the generation of variants that do not perturb host cell signaling, platelet aggregation, sequestration of immunoglobulins or the induction of B cell proliferation and apoptosis. Such Protein A variants should also be useful in analyzing vaccines for the ability of raising antibodies that block the aforementioned SpA activities and occupy the five repeat domains at their dual binding interfaces. This goal is articulated and pursued here for the first time and methods are described in detail for the generation of Protein A variants that can be used as a safe vaccine for humans. To perturb IgG Fc $\gamma$ , vWF A1 and

TNFR1 binding, glutamine (Q) 9 and 10 [numbering derived from the SpA domain D as described in Uhlen et al., 1984] were mutated, and generated lysine substitutions for both glutamines with the expectation that these abolish the ligand attributes at the first binding interface. To perturb IgM Fab VH3 binding, aspartate (D) 36 and 37 were mutated, each of which is required for the association with the B cell receptor. D36 and D37 were both substituted with alanine. Q9,10K and D36,37A mutations are here combined in the recombinant molecule SpA-DQ9,10K;D36,37A and tested for the binding attributes of Protein A. Further, SpA-D and SpA-DQ9,10K; D36,37A are subjected to immunization studies in mice and rabbits and analyzed for [1] the production of specific antibodies (SpA-D Ab); [2] the ability of SpA-D Ab to block the association between Protein A and its four different ligands; and, [3] the attributes of SpA-D Ab to generate protective immunity against staphylococcal infections. (See Examples section below).

#### [0146] B. Staphylococcal Coagulases

[0147] Coagulases are enzymes produced by *Staphylococcus* bacteria that convert fibrinogen to fibrin. Coa and vW<sub>h</sub> activate prothrombin without proteolysis (Friedrich et al., 2003). The coagulase•prothrombin complex recognizes fibrinogen as a specific substrate, converting it directly into fibrin. The crystal structure of the active complex revealed binding of the D1 and D2 domains to prothrombin and insertion of its Ile<sup>1</sup>-Val<sup>2</sup> N-terminus into the Ile<sup>16</sup> pocket, inducing a functional active site in the zymogen through conformational change (Friedrich et al., 2003). Exosite I of α-thrombin, the fibrinogen recognition site, and proexosite I on prothrombin are blocked by the D2 of Coa (Friedrich et al., 2003). Nevertheless, association of the tetrameric (Coa•prothrombin)<sub>2</sub> complex binds fibrinogen at a new site with high affinity (Panizzi et al., 2006). This model explains the coagulant properties and efficient fibrinogen conversion by coagulase (Panizzi et al., 2006).

[0148] Fibrinogen is a large glycoprotein (Mr ~340,000), formed by three pairs of Aα-, Bβ-, and γ-chains covalently linked to form a “dimer of trimers,” where A and B designate the fibrinopeptides released by thrombin cleavage (Panizzi et al., 2006). The elongated molecule folds into three separate domains, a central fragment E that contains the N-termini of all six chains and two flanking fragments D formed mainly by the C-termini of the Bβ- and γ-chains. These globular domains are connected by long triple-helical structures. Coagulase-prothrombin complexes, which convert human fibrinogen to the self-polymerizing fibrin, are not targeted by circulating thrombin inhibitors (Panizzi et al., 2006). Thus, staphylococcal coagulases bypass the physiological blood coagulation pathway.

[0149] All *S. aureus* strains secrete coagulase and vWbp (Bjerketorp et al., 2004; Field and Smith, 1945). Although early work reported important contributions of coagulase to the pathogenesis of staphylococcal infections (Ekstedt and Yotis, 1960; Smith et al., 1947), more recent investigations with molecular genetics tools challenged this view by observing no virulence phenotypes with endocarditis, skin abscess and mastitis models in mice (Moreillon et al., 1995; Phonimdaeng et al., 1990). Generating isogenic variants of *S. aureus* Newman, a fully virulent clinical isolate (Duthie et al., 1952), it is described herein that coa mutants indeed display virulence defects in a lethal bacteremia and renal abscess model in mice. In the inventors experience, *S. aureus* 8325-4 is not fully virulent and it is presumed that mutational lesions in this

strain may not be able to reveal virulence defects in vivo. Moreover, antibodies raised against Coa or vWbp perturb the pathogenesis of *S. aureus* Newman infections to a degree mirroring the impact of gene deletions. Coa and vWbp contribute to staphylococcal abscess formation and lethal bacteremia and may also function as protective antigens in subunit vaccines.

[0150] Biochemical studies document the biological value of antibodies against Coa and vWbp. By binding to antigen and blocking its association with clotting factors, the antibodies prevent the formation of Coa•prothrombin and vWbp•prothrombin complexes. Passive transfer studies revealed protection of experimental animals against staphylococcal abscess formation and lethal challenge by Coa and vWbp antibodies. Thus, Coa and vWbp neutralizing antibodies generate immune protection against staphylococcal disease.

[0151] Earlier studies revealed a requirement of coagulase for resisting phagocytosis in blood (Smith et al., 1947) and the inventors observed a similar phenotype for Δcoa mutants in lepirudin-treated mouse blood (see Example 3 below). As vWbp displays higher affinity for human prothrombin than the mouse counterpart, it is suspected the same may be true for ΔvWbp variants in human blood. Further, expression of Coa and vWbp in abscess lesions as well as their striking distribution in the eosinophilic pseudocapsule surrounding (staphylococcal abscess communities (SACs) or the peripheral fibrin wall, suggest that secreted coagulases contribute to the establishment of these lesions. This hypothesis was tested and, indeed, Δcoa mutants were defective in the establishment of abscesses. A corresponding test, blocking Coa function with specific antibodies, produced the same effect. Consequently, it is proposed that the clotting of fibrin is a critical event in the establishment of staphylococcal abscesses that can be targeted for the development of protective vaccines. Due to their overlapping function on human prothrombin, both Coa and vWbp are considered excellent candidates for vaccine development.

#### [0152] C. Other Staphylococcal Antigens

[0153] Research over the past several decades identified *S. aureus* exotoxins, surface proteins and regulatory molecules as important virulence factors (Foster, 2005; Mazmanian et al., 2001; Novick, 2003). Much progress has been achieved regarding the regulation of these genes. For example, staphylococci perform a bacterial census via the secretion of auto-inducing peptides that bind to a cognate receptor at threshold concentration, thereby activating phospho-relay reactions and transcriptional activation of many of the exotoxin genes (Novick, 2003). The pathogenesis of staphylococcal infections relies on these virulence factors (secreted exotoxins, exopolysaccharides, and surface adhesins). The development of staphylococcal vaccines is hindered by the multifaceted nature of staphylococcal invasion mechanisms. It is well established that live attenuated micro-organisms are highly effective vaccines; immune responses elicited by such vaccines are often of greater magnitude and of longer duration than those produced by non-replicating immunogens. One explanation for this may be that live attenuated strains establish limited infections in the host and mimic the early stages of natural infection. Embodiments of the invention are directed to compositions and methods including variant SpA polypeptides and peptides, as well as other immunogenic extracellular proteins, polypeptides, and peptides (including both secreted and cell surface proteins or peptides) of gram



positive bacteria for the use in mitigating or immunizing against infection. In particular embodiments the bacteria is a *staphylococcus* bacteria. Extracellular proteins, polypeptides, or peptides include, but are not limited to secreted and cell surface proteins of the targeted bacteria.

**[0154]** The human pathogen *S. aureus* secretes EsxA and EsxB, two ESAT-6 like proteins, across the bacterial envelope (Burts et al., 2005, which is incorporated herein by reference). Staphylococcal esxA and esxB are clustered with six other genes in the order of transcription: esxA esaA essA esaB essB essC esaC esxB. The acronyms esa, ess, and esx stand for ESAT-6 secretion accessory, system, and extracellular, respectively, depending whether the encoded proteins play an accessory (esa) or direct (ess) role for secretion, or are secreted (esx) in the extracellular milieu. The entire cluster of eight genes is herein referred to as the Ess cluster. EsxA, esxB, esaA, esaB, and essC are all required for synthesis or secretion of EsxA and EsxB. Mutants that fail to produce EsxA, EsxB, and EssC display defects in the pathogenesis of *S. aureus* murine abscesses, suggesting that this specialized secretion system may be a general strategy of human bacterial pathogenesis. Secretion of non-WXG100 substrates by the ESX-1 pathway has been reported for several antigens including EspA, EspB, Rv3483c, and Rv3615c (Fortune et al., 2005; MacGum et al., 2005; McLaughlin et al., 2007; Xu et al., 2007). The alternate ESX-5 pathway has also been shown to secrete both WXG100 and non-WXG100 proteins in pathogenic mycobacteria (Abdallah et al., 2007; Abdallah et al., 2006).

**[0155]** The *Staphylococcus aureus* Ess pathway can be viewed as a secretion module equipped with specialized transport components (Ess), accessory factors (Esa) and cognate secretion substrates (Esx). EssA, EssB and EssC are required for EsxA and EsxB secretion. Because EssA, EssB and EssC are predicted to be transmembrane proteins, it is contemplated that these proteins form a secretion apparatus. Some of the proteins in the ess gene cluster may actively transport secreted substrates (acting as motor) while others may regulate transport (regulator). Regulation may be achieved, but need not be limited to, transcriptional or post-translational mechanisms for secreted polypeptides, sorting of specific substrates to defined locations (e.g., extracellular medium or host cells), or timing of secretion events during infection. At this point, it is unclear whether all secreted Esx proteins function as toxins or contribute indirectly to pathogenesis.

**[0156]** Staphylococci rely on surface protein mediated-adhesion to host cells or invasion of tissues as a strategy for escape from immune defenses. Furthermore, *S. aureus* utilize surface proteins to sequester iron from the host during infection. The majority of surface proteins involved in staphylococcal pathogenesis carry C-terminal sorting signals, i.e., they are covalently linked to the cell wall envelope by sortase. Further, staphylococcal strains lacking the genes required for surface protein anchoring, i.e., sortase A and B, display a dramatic defect in the virulence in several different mouse models of disease. Thus, surface protein antigens represent a validated vaccine target as the corresponding genes are essential for the development of staphylococcal disease and can be exploited in various embodiments of the invention. The sortase enzyme superfamily are Gram-positive transpeptidases responsible for anchoring surface protein virulence factors to the peptidoglycan cell wall layer. Two sortase isoforms have been identified in *Staphylococcus aureus*, SrtA and SrtB.

These enzymes have been shown to recognize a LPXTG motif in substrate proteins. The SrtB isoform appears to be important in heme iron acquisition and iron homeostasis, whereas the SrtA isoform plays a critical role in the pathogenesis of Gram-positive bacteria by modulating the ability of the bacterium to adhere to host tissue via the covalent anchoring of adhesins and other proteins to the cell wall peptidoglycan. In certain embodiments the SpA variants described herein can be used in combination with other staphylococcal proteins such as Coa, Eap, Ebh, Emp, EsaC, EsaB, EsxA, EsxB, Hla, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, IsdC, SasF, vWbp, and/or vWh proteins.

**[0157]** Certain aspects of the invention include methods and compositions concerning proteinaceous compositions including polypeptides, peptides, or nucleic acid encoding SpA variant(s) and other staphylococcal antigens such as other proteins transported by the Ess pathway, or sortase substrates. These proteins may be modified by deletion, insertion, and/or substitution.

**[0158]** The Esx polypeptides include the amino acid sequence of Esx proteins from bacteria in the *Staphylococcus* genus. The Esx sequence may be from a particular *staphylococcus* species, such as *Staphylococcus aureus*, and may be from a particular strain, such as Newman. In certain embodiments, the EsxA sequence is SAV0282 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number Q99WU4 (gi|68565539), which is hereby incorporated by reference. In other embodiments, the EsxB sequence is SAV0290 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number Q99WT7 (gi|68565532), which is hereby incorporated by reference. In further embodiments, other polypeptides transported by the Ess pathway may be used, the sequences of which may be identified by one of skill in the art using databases and interne accessible resources.

**[0159]** The sortase substrate polypeptides include, but are not limited to the amino acid sequence of SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, IsdC or SasF proteins from bacteria in the *Staphylococcus* genus. The sortase substrate polypeptide sequence may be from a particular *staphylococcus* species, such as *Staphylococcus aureus*, and may be from a particular strain, such as Newman. In certain embodiments, the SdrD sequence is from strain N315 and can be accessed using Genbank Accession Number NP\_373773.1 (gi|15926240), which is incorporated by reference. In other embodiments, the SdrE sequence is from strain N315 and can be accessed using Genbank Accession Number NP\_373774.1 (gi|15926241), which is incorporated by reference. In other embodiments, the IsdA sequence is SAV 1130 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number NP\_371654.1 (gi|15924120), which is incorporated by reference. In other embodiments, the IsdB sequence is SAV1129 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number NP\_371653.1 (gi|15924119), which is incorporated by reference. In further embodiments, other polypeptides transported by the Ess pathway or processed by sortase may be used, the sequences of which may be identified by one of skill in the art using databases and interne accessible resources.

**[0160]** In certain embodiments, fibronectin binding protein B sequence can include all or part of the precursor or mature

foam of FnbpB. FnbpB sequence can be found in SEQ ID NO:64 or in GenBank entries having accession numbers NC\_009641.1, AAW37288. (GI:57285194), ZP\_07362431 (GI:304379700), EEV81932 (GI:257859074), NP\_373026 (GI:15925492) or other FnbpB amino acid sequences identified in GenBank.

**[0161]** Examples of various proteins that can be used in the context of the present invention can be identified by analysis of database submissions of bacterial genomes, including but not limited to accession numbers NC\_002951 (GI:57650036 and GenBank CP000046), NC\_002758 (GI:57634611 and GenBank BA000017), NC\_002745 (GI:29165615 and GenBank BA000018), NC\_003923 (GI:21281729 and GenBank BA000033), NC\_002952 (GI:49482253 and GenBank BX571856), NC\_002953 (GI:49484912 and GenBank BX571857), NC\_007793 (GI:87125858 and GenBank CP000255), NC\_007795 (GI:87201381 and GenBank CP000253) each of which are incorporated by reference.

**[0162]** As used herein, a “protein” or “polypeptide” refers to a molecule comprising at least ten amino acid residues. In some embodiments, a wild-type version of a protein or polypeptide are employed, however, in many embodiments of the invention, a modified protein or polypeptide is employed to generate an immune response. The terms described above may be used interchangeably. A “modified protein” or “modified polypeptide” or a “variant” refers to a protein or polypeptide whose chemical structure, particularly its amino acid sequence, is altered with respect to the wild-type protein or polypeptide. In some embodiments, a modified/variant protein or polypeptide has at least one modified activity or function (recognizing that proteins or polypeptides may have multiple activities or functions). It is specifically contemplated that a modified/variant protein or polypeptide may be altered with respect to one activity or function yet retain a wild-type activity or function in other respects, such as immunogenicity.

**[0163]** In certain embodiments the size of a protein or polypeptide (wild-type or modified) may comprise, but is not limited to, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 875, 900, 925, 950, 975, 1000, 1100, 1200, 1300, 1400, 1500, 1750, 2000, 2250, 2500 amino molecules or greater, and any range derivable therein, or derivative of a corresponding amino sequence described or referenced herein. It is contemplated that polypeptides may be mutated by truncation, rendering them shorter than their corresponding wild-type form, but also they might be altered by fusing or conjugating a heterologous protein sequence with a particular function (e.g., for targeting or localization, for enhanced immunogenicity, for purification purposes, etc.).

**[0164]** As used herein, an “amino molecule” refers to any amino acid, amino acid derivative, or amino acid mimic known in the art. In certain embodiments, the residues of the proteinaceous molecule are sequential, without any non-amino molecule interrupting the sequence of amino molecule residues. In other embodiments, the sequence may comprise one or more non-amino molecule moieties. In particular

embodiments, the sequence of residues of the proteinaceous molecule may be interrupted by one or more non-amino molecule moieties.

**[0165]** Accordingly, the term “proteinaceous composition” encompasses amino molecule sequences comprising at least one of the 20 common amino acids in naturally synthesized proteins, or at least one modified or unusual amino acid.

**[0166]** Proteinaceous compositions may be made by any technique known to those of skill in the art, including (i) the expression of proteins, polypeptides, or peptides through standard molecular biological techniques, (ii) the isolation of proteinaceous compounds from natural sources, or (iii) the chemical synthesis of proteinaceous materials. The nucleotide as well as the protein, polypeptide, and peptide sequences for various genes have been previously disclosed, and may be found in the recognized computerized databases. One such database is the National Center for Biotechnology Information’s Genbank and GenPept databases (on the World Wide Web at [ncbi.nlm.nih.gov/](http://ncbi.nlm.nih.gov/)). The coding regions for these genes may be amplified and/or expressed using the techniques disclosed herein or as would be known to those of ordinary skill in the art.

**[0167]** Amino acid sequence variants of SpA, coagulases and other polypeptides of the invention can be substitutional, insertional, or deletion variants. A variation in a polypeptide of the invention may affect 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, or more non-contiguous or contiguous amino acids of the polypeptide, as compared to wild-type. A variant can comprise an amino acid sequence that is at least 50%, 60%, 70%, 80%, or 90%, including all values and ranges there between, identical to any sequence provided or referenced herein, e.g., SEQ ID NO:2-8 or SEQ ID NO:11-30. A variant can include 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more substitute amino acids. A polypeptide processed or secreted by the Ess pathway or other surface proteins (see Table 2) or sortase substrates from any *staphylococcus* species and strain are contemplated for use in compositions and methods described herein.

**[0168]** Deletion variants typically lack one or more residues of the native or wild-type protein. Individual residues can be deleted or a number of contiguous amino acids can be deleted. A stop codon may be introduced (by substitution or insertion) into an encoding nucleic acid sequence to generate a truncated protein. Insertional mutants typically involve the addition of material at a non-terminal point in the polypeptide. This may include the insertion of one or more residues. Terminal additions, called fusion proteins, may also be generated. These fusion proteins include multimers or concatamers of one or more peptide or polypeptide described or referenced herein.

**[0169]** Substitutional variants typically contain the exchange of one amino acid for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide, with or without the loss of other functions or properties. Substitutions may be conservative, that is, one amino acid is replaced with one of similar shape and charge. Conservative substitutions are well known in the art and include, for example, the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or

valine; leucine to valine or isoleucine; lysine to arginine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; and valine to isoleucine or leucine. Alternatively, substitutions may be non-conservative such that a function or activity of the polypeptide is affected. Non-conservative changes typically involve substituting a residue with one that is chemically dissimilar, such as a polar or charged amino acid for a nonpolar or uncharged amino acid, and vice versa.

TABLE 2

Exemplary surface proteins of <i>S. aureus</i> strains.								
SAV #	SA#	Surface	MW2	Mu50	N315	Newman	MRSA252*	MSSA476*
SAV0111	SA0107	Spa	492	450	450	520	516	492
SAV2503	SA2291	FnBPA	1015	1038	1038	741	—	1015
SAV2502	SA2290	FnBPB	943	961	961	677	965	957
SAV0811	SA0742	CifA	946	935	989	933	1029	928
SAV2630	SA2423	CifB	907	877	877	913	873	905
Np	Np	Cna	1183	—	—	—	1183	1183
SAV0561	SA0519	SdrC	955	953	953	947	906	957
SAV0562	SA0520	SdrD	1347	1385	1385	1315	—	1365
SAV0563	SA0521	SdrE	1141	1141	1141	1166	1137	1141
Np	Np	Pls	—	—	—	—	—	—
SAV2654	SA2447	SasA	2275	2271	2271	2271	1351	2275
SAV2160	SA1964	SasB	686	2481	2481	2481	2222	685
	SA1577	SasC	2186	213	2186	2186	2189	2186
SAV0134	SA0129	SasD	241	241	241	241	221	241
SAV1130	SA0977	SasE/IsdA	350	350	350	350	354	350
SAV2646	SA2439	SasF	635	635	635	635	627	635
SAV2496		SasG	1371	525	927	—	—	1371
SAV0023	SA0022	SasH	772	—	772	772	786	786
SAV1731	SA1552	SasI	895	891	891	891	534	895
SAV1129	SA0976	SasJ/IsdB	645	645	645	645	652	645
	SA2381	SasK	198	211	211	—	—	197
	Np	SasL	—	232	—	—	—	—
SAV1131	SA0978	IsdC	227	227	227	227	227	227

[0170] Proteins of the invention may be recombinant, or synthesized in vitro. Alternatively, a non-recombinant or recombinant protein may be isolated from bacteria. It is also contemplated that a bacteria containing such a variant may be implemented in compositions and methods of the invention. Consequently, a protein need not be isolated.

[0171] The term “functionally equivalent codon” is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine, and also refers to codons that encode biologically equivalent amino acids (see Table 3, below).

TABLE 3

Codon Table	
Amino Acids	Codons
Alanine	Ala A GCA GCC GCG GCU
Cysteine	Cys C UGC UGU
Aspartic acid	Asp D GAC GAU
Glutamic acid	Glu E GAA GAG
Phenylalanine	Phe F UUC UUU
Glycine	Gly G GGA GGC GGG GGU

TABLE 3-continued

Codon Table	
Amino Acids	Codons
Histidine	His H CAC CAU
Isoleucine	Ile I AUA AUC AUU
Lysine	Lys K AAA AAG

TABLE 3-continued

Codon Table	
Amino Acids	Codons
Leucine	Leu L UUA UUG CUA CUC CUG CUU
Methionine	Met M AUG
Asparagine	Asn N AAC AAU
Proline	Pro P CCA CCC CCG CCU
Glutamine	Gln Q CAA CAG
Arginine	Arg R AGA AGG CGA CGC CGG CGU
Serine	Ser S AGC AGU UCA UCC UCG UCU
Threonine	Thr T ACA ACC ACG ACU
Valine	Val V GUA GUC GUG GUU
Tryptophan	Trp W UGG
Tyrosine	Tyr Y UAC UAU

[0172] It also will be understood that amino acid and nucleic acid sequences may include additional residues, such as additional N- or C-terminal amino acids, or 5' or 3'

sequences, respectively, and yet still be essentially as set forth in one of the sequences disclosed herein, so long as the sequence meets the criteria set forth above, including the maintenance of biological protein activity (e.g., immunogenicity) where protein expression is concerned. The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region.

**[0173]** The following is a discussion based upon changing of the amino acids of a protein to create a variant polypeptide or peptide. For example, certain amino acids may be substituted for other amino acids in a protein structure with or without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's functional activity, certain amino acid substitutions can be made in a protein sequence, and in its underlying DNA coding sequence, and nevertheless produce a protein with a desirable property. It is thus contemplated by the inventors that various changes may be made in the DNA sequences of genes.

**[0174]** It is contemplated that in compositions of the invention, there is between about 0.001 mg and about 10 mg of total polypeptide, peptide, and/or protein per ml. The concentration of protein in a composition can be about, at least about or at most about 0.001, 0.010, 0.050, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5, 10.0 mg/ml or more (or any range derivable therein). Of this, about, at least about, or at most about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100% may be an SpA variant or a coagulase, and may be used in combination with other peptides or polypeptides, such as other bacterial peptides and/or antigens.

**[0175]** The present invention contemplates the administration of variant SpA polypeptides or peptides to effect a preventative therapy or therapeutic effect against the development of a disease or condition associated with infection by a *staphylococcus* pathogen.

**[0176]** In certain aspects, combinations of staphylococcal antigens are used in the production of an immunogenic composition that is effective at treating or preventing staphylococcal infection. Staphylococcal infections progress through several different stages. For example, the staphylococcal life cycle involves commensal colonization, initiation of infection by accessing adjoining tissues or the bloodstream, and/or anaerobic multiplication in the blood. The interplay between *S. aureus* virulence determinants and the host defense mechanisms can induce complications such as endocarditis, metastatic abscess formation, and sepsis syndrome. Different molecules on the surface of the bacterium are involved in different steps of the infection cycle. Combinations of certain antigens can elicit an immune response which protects against multiple stages of staphylococcal infection. The effectiveness of the immune response can be measured either in animal model assays and/or using an opsonophagocytic assay.

**[0177]** D. Polypeptides and Polypeptide Production

**[0178]** The present invention describes polypeptides, peptides, and proteins and immunogenic fragments thereof for use in various embodiments of the present invention. For example, specific polypeptides are assayed for or used to elicit an immune response. In specific embodiments, all or part of the proteins of the invention can also be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, (1984); Tam et al., (1983); Merrifield, (1986); and Barany and Merrifield (1979), each incorporated herein by reference.

**[0179]** Alternatively, recombinant DNA technology may be employed wherein a nucleotide sequence which encodes a peptide of the invention is inserted into an expression vector, transformed or transfected into an appropriate host cell and cultivated under conditions suitable for expression.

**[0180]** One embodiment of the invention includes the use of gene transfer to cells, including microorganisms, for the production and/or presentation of polypeptides or peptides. The gene for the polypeptide or peptide of interest may be transferred into appropriate host cells followed by culture of cells under the appropriate conditions. The generation of recombinant expression vectors, and the elements included therein, are well known in the art and briefly discussed herein. Alternatively, the protein to be produced may be an endogenous protein normally synthesized by the cell that is isolated and purified.

**[0181]** Another embodiment of the present invention uses autologous B lymphocyte cell lines, which are transfected with a viral vector that expresses an immunogen product, and more specifically, a protein having immunogenic activity. Other examples of mammalian host cell lines include, but are not limited to Vero and HeLa cells, other B- and T-cell lines, such as CEM, 721.221, H9, Jurkat, Raji, as well as cell lines of Chinese hamster ovary, W138, BHK, COS-7, 293, HepG2, 3T3, RIN and MDCK cells. In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or that modifies and processes the gene product in the manner desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed.

**[0182]** A number of selection systems may be used including, but not limited to HSV thymidine kinase, hypoxanthine-guanine phosphoribosyltransferase, and adenine phosphoribosyltransferase genes, in tk-, hgppt- or apt- cells, respectively. Also, anti-metabolite resistance can be used as the basis of selection: for dhfr, which confers resistance to trimethoprim and methotrexate; gpt, which confers resistance to mycophenolic acid; neo, which confers resistance to the aminoglycoside G418; and hyg, which confers resistance to hygromycin.

**[0183]** Animal cells can be propagated in vitro in two modes: as non-anchorage-dependent cells growing in suspension throughout the bulk of the culture or as anchorage-dependent cells requiring attachment to a solid substrate for their propagation (i.e., a monolayer type of cell growth).

**[0184]** Non-anchorage dependent or suspension cultures from continuous established cell lines are the most widely

used means of large scale production of cells and cell products. However, suspension cultured cells have limitations, such as tumorigenic potential and lower protein production than adherent cells.

**[0185]** Where a protein is specifically mentioned herein, it is preferably a reference to a native or recombinant protein or optionally a protein in which any signal sequence has been removed. The protein may be isolated directly from the staphylococcal strain or produced by recombinant DNA techniques. Immunogenic fragments of the protein may be incorporated into the immunogenic composition of the invention. These are fragments comprising at least 10 amino acids, 20 amino acids, 30 amino acids, 40 amino acids, 50 amino acids, or 100 amino acids, including all values and ranges there between, taken contiguously from the amino acid sequence of the protein. In addition, such immunogenic fragments are immunologically reactive with antibodies generated against the Staphylococcal proteins or with antibodies generated by infection of a mammalian host with Staphylococci. Immunogenic fragments also include fragments that when administered at an effective dose, (either alone or as a hapten bound to a carrier), elicit a protective or therapeutic immune response against Staphylococcal infection, in certain aspects it is protective against *S. aureus* and/or *S. epidermidis* infection. Such an immunogenic fragment may include, for example, the protein lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment according to the invention comprises substantially all of the extracellular domain of a protein which has at least 80% identity, at least 85% identity, at least 90% identity, at least 95% identity, or at least 97-99% identity, including all values and ranges there between, to a sequence selected segment of a polypeptide described or referenced herein.

**[0186]** Also included in immunogenic compositions of the invention are fusion proteins composed of one or more Staphylococcal proteins, or immunogenic fragments of staphylococcal proteins. Such fusion proteins may be made recombinantly and may comprise one portion of at least 1, 2, 3, 4, 5, or 6 staphylococcal proteins or segments. Alternatively, a fusion protein may comprise multiple portions of at least 1, 2, 3, 4 or 5 staphylococcal proteins. These may combine different Staphylococcal proteins and/or multiples of the same protein or protein fragment, or immunogenic fragments in the same protein (forming a multimer or a concatamer). Alternatively, the invention also includes individual fusion proteins of Staphylococcal proteins or immunogenic fragments thereof, as a fusion protein with heterologous sequences such as a provider of T-cell epitopes or purification tags, for example:  $\beta$ -galactosidase, glutathione-S-transferase, green fluorescent proteins (GFP), epitope tags such as FLAG, myc tag, poly histidine, or viral surface proteins such as influenza virus haemagglutinin, or bacterial proteins such as tetanus toxoid, diphtheria toxoid, or CRM197.

## II. Nucleic Acids

**[0187]** In certain embodiments, the present invention concerns recombinant polynucleotides encoding the proteins, polypeptides, peptides of the invention. The nucleic acid sequences for SpA, coagulases and other bacterial proteins are included, all of which are incorporated by reference, and can be used to prepare peptides or polypeptides.

**[0188]** As used in this application, the term "polynucleotide" refers to a nucleic acid molecule that either is recom-

binant or has been isolated free of total genomic nucleic acid. Included within the term "polynucleotide" are oligonucleotides (nucleic acids of 100 residues or less in length), recombinant vectors, including, for example, plasmids, cosmids, phage, viruses, and the like. Polynucleotides include, in certain aspects, regulatory sequences, isolated substantially away from their naturally occurring genes or protein encoding sequences. Polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be RNA, DNA (genomic, cDNA or synthetic), analogs thereof, or a combination thereof. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide.

**[0189]** In this respect, the term "gene," "polynucleotide," or "nucleic acid" is used to refer to a nucleic acid that encodes a protein, polypeptide, or peptide (including any sequences required for proper transcription, post-translational modification, or localization). As will be understood by those in the art, this term encompasses genomic sequences, expression cassettes, cDNA sequences, and smaller engineered nucleic acid segments that express, or may be adapted to express, proteins, polypeptides, domains, peptides, fusion proteins, and mutants. A nucleic acid encoding all or part of a polypeptide may contain a contiguous nucleic acid sequence of: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 441, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, 1000, 1010, 1020, 1030, 1040, 1050, 1060, 1070, 1080, 1090, 1095, 1100, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 9000, 10000, or more nucleotides, nucleosides, or base pairs, including all values and ranges therebetween, of a polynucleotide encoding one or more amino acid sequence described or referenced herein. It also is contemplated that a particular polypeptide may be encoded by nucleic acids containing variations having slightly different nucleic acid sequences but, nonetheless, encode the same or substantially similar protein (see Table 3 above).

**[0190]** In particular embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a variant SpA or coagulase. The term "recombinant" may be used in conjunction with a polynucleotide or polypeptide and generally refers to a polypeptide or polynucleotide produced and/or manipulated in vitro or that is a replication product of such a molecule.

**[0191]** In other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a variant SpA or coagulase polypeptide or peptide to generate an immune response in a subject. In various embodiments the nucleic acids of the invention may be used in genetic vaccines.

**[0192]** The nucleic acid segments used in the present invention can be combined with other nucleic acid sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and

use in the intended recombinant nucleic acid protocol. In some cases, a nucleic acid sequence may encode a polypeptide sequence with additional heterologous coding sequences, for example to allow for purification of the polypeptide, transport, secretion, post-translational modification, or for therapeutic benefits such as targeting or efficacy. As discussed above, a tag or other heterologous polypeptide may be added to the modified polypeptide-encoding sequence, wherein "heterologous" refers to a polypeptide that is not the same as the modified polypeptide.

**[0193]** In certain other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors that include within their sequence a contiguous nucleic acid sequence from SEQ ID NO:1 (SpA domain D) or SEQ ID NO:3 (SpA) or any other nucleic acid sequences encoding coagulases or other secreted virulence factors and/or surface proteins including proteins transported by the Ess pathway, processed by sortase, or proteins incorporated herein by reference.

**[0194]** In certain embodiments, the present invention provides polynucleotide variants having substantial identity to the sequences disclosed herein; those comprising at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher sequence identity, including all values and ranges there between, compared to a polynucleotide sequence of this invention using the methods described herein (e.g., BLAST analysis using standard parameters).

**[0195]** The invention also contemplates the use of polynucleotides which are complementary to all the above described polynucleotides.

**[0196]** A. Vectors

**[0197]** Polypeptides of the invention may be encoded by a nucleic acid molecule comprised in a vector. The term "vector" is used to refer to a carrier nucleic acid molecule into which a heterologous nucleic acid sequence can be inserted for introduction into a cell where it can be replicated and expressed. A nucleic acid sequence can be "heterologous," which means that it is in a context foreign to the cell in which the vector is being introduced or to the nucleic acid in which is incorporated, which includes a sequence homologous to a sequence in the cell or nucleic acid but in a position within the host cell or nucleic acid where it is ordinarily not found. Vectors include DNAs, RNAs, plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (e.g., YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (for example Sambrook et al., 2001; Ausubel et al., 1996, both incorporated herein by reference). In addition to encoding a variant SpA polypeptide the vector can encode other polypeptide sequences such as a one or more other bacterial peptide, a tag, or an immunogenicity enhancing peptide. Useful vectors encoding such fusion proteins include pIN vectors (Inouye et al., 1985), vectors encoding a stretch of histidines, and pGEX vectors, for use in generating glutathione S-transferase (GST) soluble fusion proteins for later purification and separation or cleavage.

**[0198]** The term "expression vector" refers to a vector containing a nucleic acid sequence coding for at least part of a gene product capable of being transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. Expression vectors can contain a variety of "control sequences," which refer to nucleic acid sequences necessary for the transcription and possibly translation of an operably linked coding sequence in a particular host organ-

ism. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described herein.

**[0199]** 1. Promoters and Enhancers

**[0200]** A "promoter" is a control sequence. The promoter is typically a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind such as RNA polymerase and other transcription factors. The phrases "operatively positioned," "operatively linked," "under control," and "under transcriptional control" mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and expression of that sequence. A promoter may or may not be used in conjunction with an "enhancer," which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

**[0201]** Naturally, it may be important to employ a promoter and/or enhancer that effectively directs the expression of the DNA segment in the cell type or organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression (see Sambrook et al., 2001, incorporated herein by reference). The promoters employed may be constitutive, tissue-specific, or inducible and in certain embodiments may direct high level expression of the introduced DNA segment under specified conditions, such as large-scale production of recombinant proteins or peptides.

**[0202]** Various elements/promoters may be employed in the context of the present invention to regulate the expression of a gene. Examples of such inducible elements, which are regions of a nucleic acid sequence that can be activated in response to a specific stimulus, include but are not limited to Immunoglobulin Heavy Chain (Banerji et al., 1983; Gilles et al., 1983; Grosschedl et al., 1985; Atchinson et al., 1986, 1987; Imler et al., 1987; Weinberger et al., 1984; Kiledjian et al., 1988; Porton et al., 1990), Immunoglobulin Light Chain (Queen et al., 1983; Picard et al., 1984), T Cell Receptor (Luria et al., 1987; Winoto et al., 1989; Redondo et al., 1990), HLA DQ a and/or DQ (3 (Sullivan et al., 1987),  $\beta$  Interferon (Goodbourn et al., 1986; Fujita et al., 1987; Goodbourn et al., 1988), Interleukin-2 (Greene et al., 1989), Interleukin-2 Receptor (Greene et al., 1989; Lin et al., 1990), MHC Class II 5 (Koch et al., 1989), MHC Class II HLA-DRa (Sherman et al., 1989),  $\beta$ -Actin (Kawamoto et al., 1988; Ng et al., 1989), Muscle Creatine Kinase (MCK) (Jaynes et al., 1988; Horlick et al., 1989; Johnson et al., 1989), Prealbumin (Transthyretin) (Costa et al., 1988), Elastase I (Ornitz et al., 1987), Metallothionein (MTII) (Karin et al., 1987; Culotta et al., 1989), Collagenase (Pinkert et al., 1987; Angel et al., 1987), Albumin (Pinkert et al., 1987; Tronche et al., 1989, 1990),  $\alpha$ -Fetoprotein (Godbout et al., 1988; Campere et al., 1989),  $\gamma$ -Globin (Bodine et al., 1987; Perez-Stable et al., 1990),  $\beta$ -Globin (Trudel et al., 1987), c-fos (Cohen et al., 1987), c-Ha-Ras (Triesman, 1986; Deschamps et al., 1985), Insulin (Edlund et al., 1985), Neural Cell Adhesion Molecule (NCAM) (Hirsh et al., 1990),  $\alpha$ 1-Antitrypsin (Latimer et al., 1990), H<sub>2</sub>B (TH2B) Histone (Hwang et al., 1990), Mouse and/or Type I Collagen (Ripe et al., 1989), Glucose-Regulated Proteins (GRP94 and GRP78) (Chang et al., 1989), Rat Growth Hormone (Larsen et al., 1986), Human Serum Amyloid A (SAA) (Edbrooke et al., 1989), Troponin I (TN I)

(Yützey et al., 1989), Platelet-Derived Growth Factor (PDGF) (Pech et al., 1989), Duchenne Muscular Dystrophy (Klamut et al., 1990), SV40 (Banerji et al., 1981; Moreau et al., 1981; Sleigh et al., 1985; Firak et al., 1986; Herr et al., 1986; Imbra et al., 1986; Kadesch et al., 1986; Wang et al., 1986; Ondek et al., 1987; Kuhl et al., 1987; Schaffner et al., 1988), Polyoma (Swartzendruber et al., 1975; Vasseur et al., 1980; Katinka et al., 1980, 1981; Tyndell et al., 1981; Dandolo et al., 1983; de Villiers et al., 1984; Hen et al., 1986; Satake et al., 1988; Campbell et al., 1988), Retroviruses (Kriegler et al., 1982, 1983; Levinson et al., 1982; Kriegler et al., 1983, 1984a, b, 1988; Bosze et al., 1986; Miksicek et al., 1986; Celander et al., 1987; Thiesen et al., 1988; Celander et al., 1988; Choi et al., 1988; Reisman et al., 1989), Papilloma Virus (Campo et al., 1983; Lusk et al., 1983; Spandidos and Wilkie, 1983; Spalholz et al., 1985; Lusk et al., 1986; Cripe et al., 1987; Gloss et al., 1987; Hirochika et al., 1987; Stephens et al., 1987), Hepatitis B Virus (Bulla et al., 1986; Jameel et al., 1986; Shaul et al., 1987; Spandau et al., 1988; Vannice et al., 1988), Human Immunodeficiency Virus (Muesing et al., 1987; Hauber et al., 1988; Jakobovits et al., 1988; Feng et al., 1988; Takebe et al., 1988; Rosen et al., 1988; Berkhout et al., 1989; Laspia et al., 1989; Sharp et al., 1989; Braddock et al., 1989), Cytomegalovirus (CMV) IE (Weber et al., 1984; Boshart et al., 1985; Foecking et al., 1986), Gibbon Ape Leukemia Virus (Holbrook et al., 1987; Quinn et al., 1989).

**[0203]** Inducible elements include, but are not limited to MT II—Phorbol Ester (TFA)/Heavy metals (Palmiter et al., 1982; Haslinger et al., 1985; Searle et al., 1985; Stuart et al., 1985; Imagawa et al., 1987; Karin et al., 1987; Angel et al., 1987b; McNeall et al., 1989); MMTV (mouse mammary tumor virus)—Glucocorticoids (Huang et al., 1981; Lee et al., 1981; Majors et al., 1983; Chandler et al., 1983; Lee et al., 1984; Ponta et al., 1985; Sakai et al., 1988);  $\beta$ -Interferon—poly(rI)/poly(rc) (Tavernier et al., 1983); Adenovirus 5 E2—E1A (Imperiale et al., 1984); Collagenase—Phorbol Ester (TPA) (Angel et al., 1987a); Stromelysin—Phorbol Ester (TPA) (Angel et al., 1987b); SV40—Phorbol Ester (TPA) (Angel et al., 1987b); Murine MX Gene—Interferon, Newcastle Disease Virus (Hug et al., 1988); GRP78 Gene—A23187 (Resendez et al., 1988);  $\alpha$ -2-Macroglobulin—IL-6 (Kunz et al., 1989); Vimentin—Serum (Rittling et al., 1989); MHC Class I Gene H-2kb—Interferon (Blonar et al., 1989); HSP70—E1A/SV40 Large T Antigen (Taylor et al., 1989, 1990a, 1990b); Proliferin—Phorbol Ester/TPA (Mordacq et al., 1989); Tumor Necrosis Factor—PMA (Hensel et al., 1989); and Thyroid Stimulating Hormone  $\alpha$  Gene—Thyroid Hormone (Chatterjee et al., 1989).

**[0204]** The particular promoter that is employed to control the expression of peptide or protein encoding polynucleotide of the invention is not believed to be critical, so long as it is capable of expressing the polynucleotide in a targeted cell, preferably a bacterial cell. Where a human cell is targeted, it is preferable to position the polynucleotide coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a bacterial, human or viral promoter.

**[0205]** In embodiments in which a vector is administered to a subject for expression of the protein, it is contemplated that a desirable promoter for use with the vector is one that is not down-regulated by cytokines or one that is strong enough that even if down-regulated, it produces an effective amount of a variant SpA for eliciting an immune response. Non-limiting

examples of these are CMV IE and RSV LTR. Tissue specific promoters can be used, particularly if expression is in cells in which expression of an antigen is desirable, such as dendritic cells or macrophages. The mammalian MHC I and MHC II promoters are examples of such tissue-specific promoters.

**[0206]** 2. Initiation Signals and Internal Ribosome Binding Sites (IRES)

**[0207]** A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals.

**[0208]** In certain embodiments of the invention, the use of internal ribosome entry sites (IRES) elements are used to create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, 1988; Macejak and Sarnow, 1991). IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be transcribed together, each separated by an IRES, creating polycistronic messages. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message (see U.S. Pat. Nos. 5,925,565 and 5,935,819, herein incorporated by reference).

**[0209]** 3. Selectable and Screenable Markers

**[0210]** In certain embodiments of the invention, cells containing a nucleic acid construct of the present invention may be identified in vitro or in vivo by encoding a screenable or selectable marker in the expression vector. When transcribed and translated, a marker confers an identifiable change to the cell permitting easy identification of cells containing the expression vector. Generally, a selectable marker is one that confers a property that allows for selection. A positive selectable marker is one in which the presence of the marker allows for its selection, while a negative selectable marker is one in which its presence prevents its selection. An example of a positive selectable marker is a drug resistance marker.

**[0211]** B. Host Cells

**[0212]** As used herein, the terms “cell,” “cell line,” and “cell culture” may be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. In the context of expressing a heterologous nucleic acid sequence, “host cell” refers to a prokaryotic or eukaryotic cell, and it includes any transformable organism that is capable of replicating a vector or expressing a heterologous gene encoded by a vector. A host cell can, and has been, used as a recipient for vectors or viruses. A host cell may be “transfected” or “transformed,” which refers to a process by which exogenous nucleic acid, such as a recombinant protein-encoding sequence, is transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny.

**[0213]** Host cells may be derived from prokaryotes or eukaryotes, including bacteria, yeast cells, insect cells, and mammalian cells for replication of the vector or expression of part or all of the nucleic acid sequence(s). Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection

(ATCC), which is an organization that serves as an archive for living cultures and genetic materials (www.atcc.org).

#### [0214] C. Expression Systems

[0215] Numerous expression systems exist that comprise at least a part or all of the compositions discussed above. Prokaryote- and/or eukaryote-based systems can be employed for use with the present invention to produce nucleic acid sequences, or their cognate polypeptides, proteins and peptides. Many such systems are commercially and widely available.

[0216] The insect cell/baculovirus system can produce a high level of protein expression of a heterologous nucleic acid segment, such as described in U.S. Pat. Nos. 5,871,986, 4,879,236, both herein incorporated by reference, and which can be bought, for example, under the name MAXBAC® 2.0 from INVITROGEN® and BACPACK™ BACULOVIRUS EXPRESSION SYSTEM FROM CLONTECH®.

[0217] In addition to the disclosed expression systems of the invention, other examples of expression systems include STRATAGENE®'s COMPLETE CONTROL™ Inducible Mammalian Expression System, which involves a synthetic ecdysone-inducible receptor, or its pET Expression System, an *E. coli* expression system. Another example of an inducible expression system is available from INVITROGEN®, which carries the T-REX™ (tetracycline-regulated expression) System, an inducible mammalian expression system that uses the full-length CMV promoter. INVITROGEN® also provides a yeast expression system called the *Pichia methanolica* Expression System, which is designed for high-level production of recombinant proteins in the methylotrophic yeast *Pichia methanolica*. One of skill in the art would know how to express a vector, such as an expression construct, to produce a nucleic acid sequence or its cognate polypeptide, protein, or peptide.

### III. Polysaccharides

[0218] The immunogenic compositions of the invention may further comprise capsular polysaccharides including one or more of PIA (also known as PNAG) and/or *S. aureus* Type V and/or type VIII capsular polysaccharide and/or *S. epidermidis* Type I, and/or Type II and/or Type III capsular polysaccharide.

#### [0219] A. PIA (PNAG)

[0220] It is now clear that the various forms of staphylococcal surface polysaccharides identified as PS/A, PIA and SAA are the same chemical entity—PNAG (Maira-Litran et al., 2004). Therefore the term PIA or PNAG encompasses all these polysaccharides or oligosaccharides derived from them.

[0221] PIA is a polysaccharide intercellular adhesin and is composed of a polymer of (β-(1→6)-linked glucosamine substituted with N-acetyl and O-succinyl constituents). This polysaccharide is present in both *S. aureus* and *S. epidermidis* and can be isolated from either source (Joyce et al., 2003; Maira-Litran et al., 2002). For example, PNAG may be isolated from *S. aureus* strain MN8m (WO04/43407). PIA isolated from *S. epidermidis* is an integral constituent of biofilm. It is responsible for mediating cell-cell adhesion and probably also functions to shield the growing colony from the host's immune response. The polysaccharide previously known as poly-N-succinyl-β-(1→6)-glucosamine (PNSG) was recently shown not to have the expected structure since the identification of N-succinylation was incorrect (Maira-Litran

et al., 2002). Therefore the polysaccharide formally known as PNSG and now found to be PNAG is also encompassed by the term PIA.

[0222] PIA (or PNAG) may be of different sizes varying from over 400 kDa to between 75 and 400 kDa to between 10 and 75 kDa to oligosaccharides composed of up to 30 repeat units (of β-(1→6)-linked glucosamine substituted with N-acetyl and O-succinyl constituents). Any size of PIA polysaccharide or oligosaccharide may be used in an immunogenic composition of the invention, in one aspect the polysaccharide is over 40 kDa. Sizing may be achieved by any method known in the art, for instance by microfluidization, ultrasonic irradiation or by chemical cleavage (WO 03/53462, EP497524, EP497525). In certain aspects PIA (PNAG) is at least or at most 40-400 kDa, 40-300 kDa, 50-350 kDa, 60-300 kDa, 50-250 kDa and 60-200 kDa.

[0223] PIA (PNAG) can have different degree of acetylation due to substitution on the amino groups by acetate. PIA produced in vitro is almost fully substituted on amino groups (95-100%). Alternatively, a deacetylated PIA (PNAG) can be used having less than 60%, 50%, 40%, 30%, 20%, 10% acetylation. Use of a deacetylated PIA (PNAG) is preferred since non-acetylated epitopes of PNAG are efficient at mediating opsonic killing of Gram positive bacteria, preferably *S. aureus* and/or *S. epidermidis*. In certain aspects, the PIA (PNAG) has a size between 40 kDa and 300 kDa and is deacetylated so that less than 60%, 50%, 40%, 30% or 20% of amino groups are acetylated.


[0224] The term deacetylated PNAG (dPNAG) refers to a PNAG polysaccharide or oligosaccharide in which less than 60%, 50%, 40%, 30%, 20% or 10% of the amino groups are acetylated. In certain aspects, PNAG is deacetylated to form dPNAG by chemically treating the native polysaccharide. For example, the native PNAG is treated with a basic solution such that the pH rises to above 10. For instance the PNAG is treated with 0.1-5 M, 0.2-4 M, 0.3-3 M, 0.5-2 M, 0.75-1.5 M or 1 M NaOH, KOH or NH<sub>4</sub>OH. Treatment is for at least 10 to 30 minutes, or 1, 2, 3, 4, 5, 10, 15 or 20 hours at a temperature of 20-100, 25-80, 30-60 or 30-50 or 35-45° C. dPNAG may be prepared as described in WO 04/43405.

[0225] The polysaccharide(s) can be conjugated or unconjugated to a carrier protein.


#### [0226] B. Type 5 and Type 8 Polysaccharides from *S. aureus*

[0227] Most strains of *S. aureus* that cause infection in man contain either Type 5 or Type 8 polysaccharides. Approximately 60% of human strains are Type 8 and approximately 30% are Type 5. The structures of Type 5 and Type 8 capsular polysaccharide antigens are described in Moreau et al., (1990) and Fournier et al., (1984). Both have FucNAcp in their repeat unit as well as ManNAcA which can be used to introduce a sulfhydryl group. The structures are:

##### Type 5



##### Type 8





## Type 5

[0230]  $\rightarrow 4$ )- $\beta$ -D-ManNAcA-(1 $\rightarrow$ 4)- $\alpha$ -L-FucNAc  
(3OAc)-(1-6)43-D-FucNAc-(1 $\rightarrow$

## Type 8

[0231]  $\rightarrow 3$ )- $\beta$ -D-ManNAcA(4OAc)-(1 $\rightarrow$ 3)- $\alpha$ -L-Fuc-  
NAc(1 $\rightarrow$ 3)- $\alpha$ -D-FucNAc(1 $\rightarrow$

[0232] Polysaccharides may be extracted from the appropriate strain of *S. aureus* using method well known to of skill in the art, See U.S. Pat. No. 6,294,177. For example, ATCC 12902 is a Type 5 *S. aureus* strain and ATCC 12605 is a Type 8 *S. aureus* strain.

[0233] Polysaccharides are of native size or alternatively may be sized, for instance by microfluidisation, ultrasonic irradiation, or by chemical treatment. The invention also covers oligosaccharides derived from the type 5 and 8 polysaccharides from *S. aureus*. The type 5 and 8 polysaccharides included in the immunogenic composition of the invention are preferably conjugated to a carrier protein as described below or are alternatively unconjugated. The immunogenic compositions of the invention alternatively contains either type 5 or type 8 polysaccharide.

[0234] *C. S. aureus* 336 Antigen

[0235] In an embodiment, the immunogenic composition of the invention comprises the *S. aureus* 336 antigen described in U.S. Pat. No. 6,294,177. The 336 antigen comprises n-linked hexosamine, contains no O-acetyl groups, and specifically binds to antibodies to *S. aureus* Type 336 deposited under ATCC 55804. In an embodiment, the 336 antigen is a polysaccharide which is of native size or alternatively may be sized, for instance by microfluidisation, ultrasonic irradiation, or by chemical treatment. The invention also covers oligosaccharides derived from the 336 antigen. The 336 antigen can be unconjugated or conjugated to a carrier protein.

[0236] D. Type I, II and III Polysaccharides from *S. epidermidis*

[0237] Amongst the problems associated with the use of polysaccharides in vaccination, is the fact that polysaccharides per se are poor immunogens. It is preferred that the polysaccharides utilized in the invention are linked to a protein carrier which provide bystander T-cell help to improve immunogenicity. Examples of such carriers which may be conjugated to polysaccharide immunogens include the Diphtheria and Tetanus toxoids (DT, DT CRM197 and TT respectively), Keyhole Limpet Haemocyanin (KLH), and the purified protein derivative of Tuberculin (PPD), *Pseudomonas aeruginosa* exoprotein A (rEPA), protein D from *Haemophilus influenzae*, pneumolysin or fragments of any of the above. Fragments suitable for use include fragments encompassing T-helper epitopes. In particular the protein D fragment from *H. influenzae* will preferably contain the N-terminal  $\frac{1}{3}$  of the protein. Protein D is an IgD-binding protein from *Haemophilus influenzae* (EP 0 594 610 B1) and is a potential immunogen. In addition, staphylococcal proteins may be used as a carrier protein in the polysaccharide conjugates of the invention.

[0238] A carrier protein that would be particularly advantageous to use in the context of a staphylococcal vaccine is staphylococcal alpha toxin. The native form may be conjugated to a polysaccharide since the process of conjugation reduces toxicity. Preferably genetically detoxified alpha toxins such as the His35Leu or His35Arg variants are used as carriers since residual toxicity is lower. Alternatively the

alpha toxin is chemically detoxified by treatment with a cross-linking reagent, formaldehyde or glutaraldehyde. A genetically detoxified alpha toxin is optionally chemically detoxified, preferably by treatment with a cross-linking reagent, formaldehyde or glutaraldehyde to further reduce toxicity.

[0239] The polysaccharides may be linked to the carrier protein(s) by any known method (for example those methods described in U.S. Pat. Nos. 4,372,945, 4,474,757, and 4,356,170). Preferably, CDAP conjugation chemistry is carried out (see WO95/08348). In CDAP, the cyanating reagent 1-cyano-dimethylaminopyridinium tetrafluoroborate (CDAP) is preferably used for the synthesis of polysaccharide-protein conjugates. The cyanilation reaction can be performed under relatively mild conditions, which avoids hydrolysis of the alkaline sensitive polysaccharides. This synthesis allows direct coupling to a carrier protein.

[0240] Conjugation preferably involves producing a direct linkage between the carrier protein and polysaccharide. Optionally a spacer (such as adipic dihydride (ADH)) may be introduced between the carrier protein and the polysaccharide.

## IV. Immune Response and Assays

[0241] As discussed above, the invention concerns evoking or inducing an immune response in a subject against a variant SpA or coagulase peptide. In one embodiment, the immune response can protect against or treat a subject having, suspected of having, or at risk of developing an infection or related disease, particularly those related to staphylococci. One use of the immunogenic compositions of the invention is to prevent nosocomial infections by inoculating a subject prior to undergoing procedures in a hospital or other environment having an increased risk of infection.

[0242] A. Immunoassays

[0243] The present invention includes the implementation of serological assays to evaluate whether and to what extent an immune response is induced or evoked by compositions of the invention. There are many types of immunoassays that can be implemented. Immunoassays encompassed by the present invention include, but are not limited to, those described in U.S. Pat. No. 4,367,110 (double monoclonal antibody sandwich assay) and U.S. Pat. No. 4,452,901 (western blot). Other assays include immunoprecipitation of labeled ligands and immunocytochemistry, both in vitro and in vivo.

[0244] Immunoassays generally are binding assays. Certain preferred immunoassays are the various types of enzyme linked immunosorbent assays (ELISAs) and radioimmunoassays (RIA) known in the art. Immunohistochemical detection using tissue sections is also particularly useful. In one example, antibodies or antigens are immobilized on a selected surface, such as a well in a polystyrene microtiter plate, dipstick, or column support. Then, a test composition suspected of containing the desired antigen or antibody, such as a clinical sample, is added to the wells. After binding and washing to remove non specifically bound immune complexes, the bound antigen or antibody may be detected. Detection is generally achieved by the addition of another antibody, specific for the desired antigen or antibody, that is linked to a detectable label. This type of ELISA is known as a "sandwich ELISA." Detection also may be achieved by the addition of a second antibody specific for the desired antigen,

followed by the addition of a third antibody that has binding affinity for the second antibody, with the third antibody being linked to a detectable label.

**[0245]** Competition ELISAs are also possible implementations in which test samples compete for binding with known amounts of labeled antigens or antibodies. The amount of reactive species in the unknown sample is determined by mixing the sample with the known labeled species before or during incubation with coated wells. The presence of reactive species in the sample acts to reduce the amount of labeled species available for binding to the well and thus reduces the ultimate signal. Irrespective of the format employed, ELISAs have certain features in common, such as coating, incubating or binding, washing to remove non specifically bound species, and detecting the bound immune complexes.

**[0246]** Antigen or antibodies may also be linked to a solid support, such as in the form of plate, beads, dipstick, membrane, or column matrix, and the sample to be analyzed is applied to the immobilized antigen or antibody. In coating a plate with either antigen or antibody, one will generally incubate the wells of the plate with a solution of the antigen or antibody, either overnight or for a specified period. The wells of the plate will then be washed to remove incompletely-adsorbed material. Any remaining available surfaces of the wells are then "coated" with a nonspecific protein that is antigenically neutral with regard to the test antisera. These include bovine serum albumin (BSA), casein, and solutions of milk powder. The coating allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific binding of antisera onto the surface.

**[0247]** B. Diagnosis of Bacterial Infection

**[0248]** In addition to the use of proteins, polypeptides, and/or peptides, as well as antibodies binding these polypeptides, proteins, and/or peptides, to treat or prevent infection as described above, the present invention contemplates the use of these polypeptides, proteins, peptides, and/or antibodies in a variety of ways, including the detection of the presence of Staphylococci to diagnose an infection, whether in a patient or on medical equipment which may also become infected. In accordance with the invention, a preferred method of detecting the presence of infections involves the steps of obtaining a sample suspected of being infected by one or more staphylococcal bacteria species or strains, such as a sample taken from an individual, for example, from one's blood, saliva, tissues, bone, muscle, cartilage, or skin. Following isolation of the sample, diagnostic assays utilizing the polypeptides, proteins, peptides, and/or antibodies of the present invention may be carried out to detect the presence of staphylococci, and such assay techniques for determining such presence in a sample are well known to those skilled in the art and include methods such as radioimmunoassay, western blot analysis and ELISA assays. In general, in accordance with the invention, a method of diagnosing an infection is contemplated wherein a sample suspected of being infected with staphylococci has added to it the polypeptide, protein, peptide, antibody, or monoclonal antibody in accordance with the present invention, and staphylococci are indicated by antibody binding to the polypeptides, proteins, and/or peptides, or polypeptides, proteins, and/or peptides binding to the antibodies in the sample.

**[0249]** Accordingly, antibodies in accordance with the invention may be used for the prevention of infection from staphylococcal bacteria (i.e., passive immunization), for the

treatment of an ongoing infection, or for use as research tools. The term "antibodies" as used herein includes monoclonal, polyclonal, chimeric, single chain, bispecific, simianized, and humanized or primatized antibodies as well as Fab fragments, such as those fragments which maintain the binding specificity of the antibodies, including the products of an Fab immunoglobulin expression library. Accordingly, the invention contemplates the use of single chains such as the variable heavy and light chains of the antibodies. Generation of any of these types of antibodies or antibody fragments is well known to those skilled in the art. Specific examples of the generation of an antibody to a bacterial protein can be found in U.S. Patent Application Pub. No. 20030153022, which is incorporated herein by reference in its entirety.

**[0250]** Any of the above described polypeptides, proteins, peptides, and/or antibodies may be labeled directly with a detectable label for identification and quantification of staphylococcal bacteria. Labels for use in immunoassays are generally known to those skilled in the art and include enzymes, radioisotopes, and fluorescent, luminescent and chromogenic substances, including colored particles such as colloidal gold or latex beads. Suitable immunoassays include enzyme-linked immunosorbent assays (ELISA).

**[0251]** C. Protective Immunity

**[0252]** In some embodiments of the invention, proteinaceous compositions confer protective immunity to a subject. Protective immunity refers to a body's ability to mount a specific immune response that protects the subject from developing a particular disease or condition that involves the agent against which there is an immune response. An immunogenically effective amount is capable of conferring protective immunity to the subject.

**[0253]** As used herein in the specification and in the claims section that follows, the term polypeptide or peptide refer to a stretch of amino acids covalently linked there amongst via peptide bonds. Different polypeptides have different functionalities according to the present invention. While according to one aspect, a polypeptide is derived from an immunogen designed to induce an active immune response in a recipient, according to another aspect of the invention, a polypeptide is derived from an antibody which results following the elicitation of an active immune response in, for example, an animal, and which can serve to induce a passive immune response in the recipient. In both cases, however, the polypeptide is encoded by a polynucleotide according to any possible codon usage.

**[0254]** As used herein the phrase "immune response" or its equivalent "immunological response" refers to the development of a humoral (antibody mediated), cellular (mediated by antigen-specific T cells or their secretion products) or both humoral and cellular response directed against a protein, peptide, carbohydrate, or polypeptide of the invention in a recipient patient. Such a response can be an active response induced by administration of immunogen or a passive response induced by administration of antibody, antibody containing material, or primed T-cells. A cellular immune response is elicited by the presentation of polypeptide epitopes in association with Class I or Class II MHC molecules, to activate antigen-specific CD4 (+) T helper cells and/or CD8 (+) cytotoxic T cells. The response may also involve activation of monocytes, macrophages, NK cells, basophils, dendritic cells, astrocytes, microglia cells, eosinophils or other compo-

nents of innate immunity. As used herein “active immunity” refers to any immunity conferred upon a subject by administration of an antigen.

**[0255]** As used herein “passive immunity” refers to any immunity conferred upon a subject without administration of an antigen to the subject. “Passive immunity” therefore includes, but is not limited to, administration of activated immune effectors including cellular mediators or protein mediators (e.g., monoclonal and/or polyclonal antibodies) of an immune response. A monoclonal or polyclonal antibody composition may be used in passive immunization for the prevention or treatment of infection by organisms that carry the antigen recognized by the antibody. An antibody composition may include antibodies that bind to a variety of antigens that may in turn be associated with various organisms. The antibody component can be a polyclonal antiserum. In certain aspects the antibody or antibodies are affinity purified from an animal or second subject that has been challenged with an antigen(s). Alternatively, an antibody mixture may be used, which is a mixture of monoclonal and/or polyclonal antibodies to antigens present in the same, related, or different microbes or organisms, such as gram-positive bacteria, gram-negative bacteria, including but not limited to *staphylococcus* bacteria.

**[0256]** Passive immunity may be imparted to a patient or subject by administering to the patient immunoglobulins (Ig) and/or other immune factors obtained from a donor or other non-patient source having a known immunoreactivity. In other aspects, an antigenic composition of the present invention can be administered to a subject who then acts as a source or donor for globulin, produced in response to challenge with the antigenic composition (“hyperimmune globulin”), that contains antibodies directed against *Staphylococcus* or other organism. A subject thus treated would donate plasma from which hyperimmune globulin would then be obtained, via conventional plasma-fractionation methodology, and administered to another subject in order to impart resistance against or to treat *staphylococcus* infection. Hyperimmune globulins according to the invention are particularly useful for immune-compromised individuals, for individuals undergoing invasive procedures or where time does not permit the individual to produce their own antibodies in response to vaccination. See U.S. Pat. Nos. 6,936,258, 6,770,278, 6,756,361, 5,548,066, 5,512,282, 4,338,298, and 4,748,018, each of which is incorporated herein by reference in its entirety, for exemplary methods and compositions related to passive immunity.

**[0257]** For purposes of this specification and the accompanying claims the terms “epitope” and “antigenic determinant” are used interchangeably to refer to a site on an antigen to which B and/or T cells respond or recognize. B-cell epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, e.g., Epitope Mapping Protocols (1996). Antibodies that recognize the same epitope can be identified in a simple immunoassay showing the ability of one antibody to block the binding of another antibody to a target antigen.

T-cells recognize continuous epitopes of about nine amino acids for CD8 cells or about 13-15 amino acids for CD4 cells. T cells that recognize the epitope can be identified by in vitro assays that measure antigen-dependent proliferation, as determined by <sup>3</sup>H-thymidine incorporation by primed T cells in response to an epitope (Burke et al., 1994), by antigen-dependent killing (cytotoxic T lymphocyte assay, Tigges et al., 1996) or by cytokine secretion.

**[0258]** The presence of a cell-mediated immunological response can be determined by proliferation assays (CD4 (+) T cells) or CTL (cytotoxic T lymphocyte) assays. The relative contributions of humoral and cellular responses to the protective or therapeutic effect of an immunogen can be distinguished by separately isolating IgG and T-cells from an immunized syngeneic animal and measuring protective or therapeutic effect in a second subject.

**[0259]** As used herein and in the claims, the terms “antibody” or “immunoglobulin” are used interchangeably and refer to any of several classes of structurally related proteins that function as part of the immune response of an animal or recipient, which proteins include IgG, IgD, IgE, IgA, IgM and related proteins.

**[0260]** Under normal physiological conditions antibodies are found in plasma and other body fluids and in the membrane of certain cells and are produced by lymphocytes of the type denoted B cells or their functional equivalent. Antibodies of the IgG class are made up of four polypeptide chains linked together by disulfide bonds. The four chains of intact IgG molecules are two identical heavy chains referred to as H-chains and two identical light chains referred to as L-chains.

**[0261]** In order to produce polyclonal antibodies, a host, such as a rabbit or goat, is immunized with the antigen or antigen fragment, generally with an adjuvant and, if necessary, coupled to a carrier. Antibodies to the antigen are subsequently collected from the sera of the host. The polyclonal antibody can be affinity purified against the antigen rendering it monospecific.

**[0262]** Monoclonal antibodies can be produced by hyperimmunization of an appropriate donor with the antigen or ex-vivo by use of primary cultures of splenic cells or cell lines derived from spleen (Anavi, 1998; Huston et al., 1991; Johnson et al., 1991; Mernaugh et al., 1995).

**[0263]** As used herein and in the claims, the phrase “an immunological portion of an antibody” includes a Fab fragment of an antibody, a Fv fragment of an antibody, a heavy chain of an antibody, a light chain of an antibody, a heterodimer consisting of a heavy chain and a light chain of an antibody, a variable fragment of a light chain of an antibody, a variable fragment of a heavy chain of an antibody, and a single chain variant of an antibody, which is also known as scFv. In addition, the term includes chimeric immunoglobulins which are the expression products of fused genes derived from different species, one of the species can be a human, in which case a chimeric immunoglobulin is said to be humanized. Typically, an immunological portion of an antibody competes with the intact antibody from which it was derived for specific binding to an antigen.

**[0264]** Optionally, an antibody or preferably an immunological portion of an antibody, can be chemically conjugated to, or expressed as, a fusion protein with other proteins. For purposes of this specification and the accompanying claims, all such fused proteins are included in the definition of antibodies or an immunological portion of an antibody.

[0265] As used herein the terms “immunogenic agent” or “immunogen” or “antigen” are used interchangeably to describe a molecule capable of inducing an immunological response against itself on administration to a recipient, either alone, in conjunction with an adjuvant, or presented on a display vehicle.

[0266] D. Treatment Methods

[0267] A method of the present invention includes treatment for a disease or condition caused by a *staphylococcus* pathogen. An immunogenic polypeptide of the invention can be given to induce an immune response in a person infected with *staphylococcus* or suspected of having been exposed to *staphylococcus*. Methods may be employed with respect to individuals who have tested positive for exposure to *staphylococcus* or who are deemed to be at risk for infection based on possible exposure.

[0268] In particular, the invention encompasses a method of treatment for staphylococcal infection, particularly hospital acquired nosocomial infections. The immunogenic compositions and vaccines of the invention are particularly advantageous to use in cases of elective surgery. Such patients will know the date of surgery in advance and could be inoculated in advance. The immunogenic compositions and vaccines of the invention are also advantageous to use to inoculate health care workers.

[0269] In some embodiments, the treatment is administered in the presence of adjuvants or carriers or other staphylococcal antigens. Furthermore, in some examples, treatment comprises administration of other agents commonly used against bacterial infection, such as one or more antibiotics.

[0270] The use of peptides for vaccination can require, but not necessarily, conjugation of the peptide to an immunogenic carrier protein, such as hepatitis B surface antigen, keyhole limpet hemocyanin, or bovine serum albumin. Methods for performing this conjugation are well known in the art.

#### V. Vaccine and other pharmaceutical compositions and Administration

[0271] A. Vaccines

[0272] The present invention includes methods for preventing or ameliorating staphylococcal infections, particularly hospital acquired nosocomial infections. As such, the invention contemplates vaccines for use in both active and passive immunization embodiments. Immunogenic compositions, proposed to be suitable for use as a vaccine, may be prepared from immunogenic SpA polypeptide(s), such as a SpA domain D variant, or immunogenic coagulases. In other embodiments SpA or coagulases can be used in combination with other secreted virulence proteins, surface proteins or immunogenic fragments thereof. In certain aspects, antigenic material is extensively dialyzed to remove undesired small molecular weight molecules and/or lyophilized for more ready formulation into a desired vehicle.

[0273] Other options for a protein/peptide-based vaccine involve introducing nucleic acids encoding the antigen(s) as DNA vaccines. In this regard, recent reports described construction of recombinant vaccinia viruses expressing either 10 contiguous minimal CTL epitopes (Thomson, 1996) or a combination of B cell, cytotoxic T-lymphocyte (CTL), and T-helper (Th) epitopes from several microbes (An, 1997), and successful use of such constructs to immunize mice for priming protective immune responses. Thus, there is ample evidence in the literature for successful utilization of peptides, peptide-pulsed antigen presenting cells (APCs), and peptide-

encoding constructs for efficient in vivo priming of protective immune responses. The use of nucleic acid sequences as vaccines is exemplified in U.S. Pat. Nos. 5,958,895 and 5,620,896.

[0274] The preparation of vaccines that contain polypeptide or peptide sequence(s) as active ingredients is generally well understood in the art, as exemplified by U.S. Pat. Nos. 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all of which are incorporated herein by reference. Typically, such vaccines are prepared as injectables either as liquid solutions or suspensions: solid forms suitable for solution in or suspension in liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients that are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants that enhance the effectiveness of the vaccines. In specific embodiments, vaccines are formulated with a combination of substances, as described in U.S. Pat. Nos. 6,793,923 and 6,733,754, which are incorporated herein by reference.

[0275] Vaccines may be conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkalene glycols or triglycerides: such suppositories may be formed from mixtures containing the active ingredient in the range of about 0.5% to about 10%, preferably about 1% to about 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 10% to about 95% of active ingredient, preferably about 25% to about 70%.

[0276] The polypeptides and polypeptide-encoding DNA constructs may be formulated into a vaccine as neutral or salt forms. Pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the peptide) and those that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like.

[0277] Typically, vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, including the capacity of the individual's immune system to synthesize antibodies and the degree of protection desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are of the order of several hundred micrograms of active ingredient per vaccination. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed by subsequent inoculations or other administrations.

[0278] The manner of application may be varied widely. Any of the conventional methods for administration of a

vaccine are applicable. These are believed to include oral application within a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection and the like. The dosage of the vaccine will depend on the route of administration and will vary according to the size and health of the subject.

[0279] In certain instances, it will be desirable to have multiple administrations of the vaccine, e.g., 2, 3, 4, 5, 6 or more administrations. The vaccinations can be at 1, 2, 3, 4, 5, 6, 7, 8, to 5, 6, 7, 8, 9, 10, 11, 12 twelve week intervals, including all ranges there between. Periodic boosters at intervals of 1-5 years will be desirable to maintain protective levels of the antibodies. The course of the immunization may be followed by assays for antibodies against the antigens, as described in U.S. Pat. Nos. 3,791,932; 4,174,384 and 3,949,064.

#### [0280] 1. Carriers

[0281] A given composition may vary in its immunogenicity. It is often necessary therefore to boost the host immune system, as may be achieved by coupling a peptide or polypeptide to a carrier. Exemplary and preferred carriers are keyhole limpet hemocyanin (KLH) and bovine serum albumin (BSA). Other albumins such as ovalbumin, mouse serum albumin, or rabbit serum albumin can also be used as carriers. Means for conjugating a polypeptide to a carrier protein are well known in the art and include glutaraldehyde, m-maleimidobenzoyl-N-hydroxysuccinimide ester, carbodiimide, and bis-biaozitized benzidine.

#### [0282] 2. Adjuvants

[0283] The immunogenicity of polypeptide or peptide compositions can be enhanced by the use of non-specific stimulators of the immune response, known as adjuvants. Suitable adjuvants include all acceptable immunostimulatory compounds, such as cytokines, toxins, or synthetic compositions. A number of adjuvants can be used to enhance an antibody response against a variant SpA polypeptide or coagulase, or any other bacterial protein or combination contemplated herein. Adjuvants can (1) trap the antigen in the body to cause a slow release; (2) attract cells involved in the immune response to the site of administration; (3) induce proliferation or activation of immune system cells; or (4) improve the spread of the antigen throughout the subject's body.

[0284] Adjuvants include, but are not limited to, oil-in-water emulsions, water-in-oil emulsions, mineral salts, polynucleotides, and natural substances. Specific adjuvants that may be used include IL-1, IL-2, IL-4, IL-7, IL-12,  $\gamma$ -interferon, GMCSF, BCG, aluminum salts, such as aluminum hydroxide or other aluminum compound, MDP compounds, such as thur-MDP and nor-MDP, CGP (MTP-PE), lipid A, and monophosphoryl lipid A (MPL). RIBI, which contains three components extracted from bacteria, MPL, trehalose dimycolate (TDM), and cell wall skeleton (CWS) in a 2% squalene/Tween 80 emulsion. MHC antigens may even be used. Others adjuvants or methods are exemplified in U.S. Pat. Nos. 6,814,971, 5,084,269, 6,656,462, each of which is incorporated herein by reference).

[0285] Various methods of achieving adjuvant affect for the vaccine includes use of agents such as aluminum hydroxide or phosphate (alum), commonly used as about 0.05 to about 0.1% solution in phosphate buffered saline, admixture with synthetic polymers of sugars (Carbopol®) used as an about 0.25% solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between about 70°

to about 101° C. for a 30-second to 2-minute period, respectively. Aggregation by reactivating with pepsin-treated (Fab) antibodies to albumin; mixture with bacterial cells (e.g., *C. parvum*), endotoxins or lipopolysaccharide components of Gram-negative bacteria; emulsion in physiologically acceptable oil vehicles (e.g., mannide mono-oleate (Aracel A)); or emulsion with a 20% solution of a perfluorocarbon (Fluosol-DA®) used as a block substitute may also be employed to produce an adjuvant effect.

[0286] Examples of and often preferred adjuvants include complete Freund's adjuvant (a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*), incomplete Freund's adjuvants, and aluminum hydroxide.

[0287] In some aspects, it is preferred that the adjuvant be selected to be a preferential inducer of either a Th1 or a Th2 type of response. High levels of Th1-type cytokines tend to favor the induction of cell mediated immune responses to a given antigen, while high levels of Th2-type cytokines tend to favor the induction of humoral immune responses to the antigen.

[0288] The distinction of Th1 and Th2-type immune response is not absolute. In reality an individual will support an immune response which is described as being predominantly Th1 or predominantly Th2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4+ T cell clones by Mosmann and Coffman (Mosmann, and Coffman, 1989). Traditionally, Th1-type responses are associated with the production of the INF- $\gamma$  and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of Th1-type immune responses are not produced by T-cells, such as IL-12. In contrast, Th2-type responses are associated with the secretion of IL-4, IL-5, IL-6, IL-10.

[0289] In addition to adjuvants, it may be desirable to co-administer biologic response modifiers (BRM) to enhance immune responses. BRMs have been shown to upregulate T cell immunity or downregulate suppresser cell activity. Such BRMs include, but are not limited to, Cimetidine (CIM; 1200 mg/d) (Smith/Kline, PA); or low-dose Cyclophosphamide (CYP; 300 mg/m<sup>2</sup>) (Johnson/Mead, NJ) and cytokines such as  $\gamma$ -interferon, IL-2, or IL-12 or genes encoding proteins involved in immune helper functions, such as B-7.

#### [0290] B. Lipid Components and Moieties

[0291] In certain embodiments, the present invention concerns compositions comprising one or more lipids associated with a nucleic acid or a polypeptide/peptide. A lipid is a substance that is insoluble in water and extractable with an organic solvent. Compounds other than those specifically described herein are understood by one of skill in the art as lipids, and are encompassed by the compositions and methods of the present invention. A lipid component and a non-lipid may be attached to one another, either covalently or non-covalently.

[0292] A lipid may be a naturally occurring lipid or a synthetic lipid. However, a lipid is usually a biological substance. Biological lipids are well known in the art, and include for example, neutral fats, phospholipids, phosphoglycerides, steroids, terpenes, lysolipids, glycosphingolipids, glucolipids, sulphatides, lipids with ether and ester-linked fatty acids and polymerizable lipids, and combinations thereof.

[0293] A nucleic acid molecule or a polypeptide/peptide, associated with a lipid may be dispersed in a solution containing a lipid, dissolved with a lipid, emulsified with a lipid,

mixed with a lipid, combined with a lipid, covalently bonded to a lipid, contained as a suspension in a lipid or otherwise associated with a lipid. A lipid or lipid-poxvirus-associated composition of the present invention is not limited to any particular structure. For example, they may also simply be interspersed in a solution, possibly forming aggregates which are not uniform in either size or shape. In another example, they may be present in a bilayer structure, as micelles, or with a "collapsed" structure. In another non-limiting example, a lipofectamine(Gibco BRL)-poxvirus or Superfect (Qiagen)-poxvirus complex is also contemplated.

**[0294]** In certain embodiments, a composition may comprise about 1%, about 2%, about 3%, about 4%, about 5%, about 6%, about 7%, about 8%, about 9%, about 10%, about 11%, about 12%, about 13%, about 14%, about 15%, about 16%, about 17%, about 18%, about 19%, about 20%, about 21%, about 22%, about 23%, about 24%, about 25%, about 26%, about 27%, about 28%, about 29%, about 30%, about 31%, about 32%, about 33%, about 34%, about 35%, about 36%, about 37%, about 38%, about 39%, about 40%, about 41%, about 42%, about 43%, about 44%, about 45%, about 46%, about 47%, about 48%, about 49%, about 50%, about 51%, about 52%, about 53%, about 54%, about 55%, about 56%, about 57%, about 58%, about 59%, about 60%, about 61%, about 62%, about 63%, about 64%, about 65%, about 66%, about 67%, about 68%, about 69%, about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or any range therebetween, of a particular lipid, lipid type, or non-lipid component such as an adjuvant, antigen, peptide, polypeptide, sugar, nucleic acid or other material disclosed herein or as would be known to one of skill in the art. In a non-limiting example, a composition may comprise about 10% to about 20% neutral lipids, and about 33% to about 34% of a cerebroside, and about 1% cholesterol. In another non-limiting example, a liposome may comprise about 4% to about 12% terpenes, wherein about 1% of the micelle is specifically lycopene, leaving about 3% to about 11% of the liposome as comprising other terpenes; and about 10% to about 35% phosphatidyl choline, and about 1% of a non-lipid component. Thus, it is contemplated that compositions of the present invention may comprise any of the lipids, lipid types or other components in any combination or percentage range.

#### **[0295]** C. Combination Therapy

**[0296]** The compositions and related methods of the present invention, particularly administration of a secreted virulence factor or surface protein, including a variant SpA polypeptide or peptide, and/or other bacterial peptides or proteins to a patient/subject, may also be used in combination with the administration of traditional therapies. These include, but are not limited to, the administration of antibiotics such as streptomycin, ciprofloxacin, doxycycline, gentamycin, chloramphenicol, trimethoprim, sulfamethoxazole, ampicillin, tetracycline or various combinations of antibiotics.

**[0297]** In one aspect, it is contemplated that a polypeptide vaccine and/or therapy is used in conjunction with antibacterial treatment. Alternatively, the therapy may precede or follow the other agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agents and/or

proteins or polynucleotides are administered separately, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and antigenic composition would still be able to exert an advantageously combined effect on the subject. In such instances, it is contemplated that one may administer both modalities within about 12-24 h of each other or within about 6-12 h of each other. In some situations, it may be desirable to extend the time period for administration significantly, where several days (2, 3, 4, 5, 6 or 7) to several weeks (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

**[0298]** Various combinations may be employed, for example antibiotic therapy is "A" and the immunogenic molecule given as part of an immune therapy regime, such as an antigen, is "B":

**[0299]** A/B/A B/A/B B/B/A A/A/B A/B/B B/A/A A/B/B B/A/B/A

**[0300]** B/B/B/A B/B/A/B A/A/B/B A/B/A/B A/B/B/A B/B/A/A

**[0301]** B/A/B/A B/A/A/B A/A/A/B B/A/A/A A/B/A/A A/A/B/A

**[0302]** Administration of the immunogenic compositions of the present invention to a patient/subject will follow general protocols for the administration of such compounds, taking into account the toxicity, if any, of the SpA composition, or other compositions described herein. It is expected that the treatment cycles would be repeated as necessary. It also is contemplated that various standard therapies, such as hydration, may be applied in combination with the described therapy.

#### **[0303]** D. General Pharmaceutical Compositions

**[0304]** In some embodiments, pharmaceutical compositions are administered to a subject. Different aspects of the present invention involve administering an effective amount of a composition to a subject. In some embodiments of the present invention, staphylococcal antigens, members of the Ess pathway, including polypeptides or peptides of the Esa or Esx class, and/or members of sortase substrates may be administered to the patient to protect against infection by one or more *staphylococcus* pathogens. Alternatively, an expression vector encoding one or more such polypeptides or peptides may be given to a patient as a preventative treatment. Additionally, such compounds can be administered in combination with an antibiotic or an antibacterial. Such compositions will generally be dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium.

**[0305]** In addition to the compounds formulated for parenteral administration, such as those for intravenous or intramuscular injection, other pharmaceutically acceptable forms include, e.g., tablets or other solids for oral administration; time release capsules; and any other form currently used, including creams, lotions, mouthwashes, inhalants and the like.

**[0306]** The active compounds of the present invention can be formulated for parenteral administration, e.g., formulated for injection via the intravenous, intramuscular, sub-cutaneous, or even intraperitoneal routes. The preparation of an aqueous composition that contains a compound or compounds that increase the expression of an MHC class 1 molecule will be known to those of skill in the art in light of the present disclosure. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for use to prepare solutions or suspen-

sions upon the addition of a liquid prior to injection can also be prepared; and, the preparations can also be emulsified.

**[0307]** Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

**[0308]** The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil, or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that it may be easily injected. It also should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

**[0309]** The proteinaceous compositions may be formulated into a neutral or salt form. Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

**[0310]** The carrier also can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion, and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

**[0311]** Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques, which yield a powder of the active ingredient, plus any additional desired ingredient from a previously sterile-filtered solution thereof.

**[0312]** Administration of the compositions according to the present invention will typically be via any common route. This includes, but is not limited to oral, nasal, or buccal administration. Alternatively, administration may be by

orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal, intranasal, or intravenous injection. In certain embodiments, a vaccine composition may be inhaled (e.g., U.S. Pat. No. 6,651,655, which is specifically incorporated by reference). Such compositions would normally be administered as pharmaceutically acceptable compositions that include physiologically acceptable carriers, buffers or other excipients. As used herein, the term "pharmaceutically acceptable" refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem complications commensurate with a reasonable benefit/risk ratio. The term "pharmaceutically acceptable carrier," means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting a chemical agent.

**[0313]** For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous, and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in isotonic NaCl solution and either added to hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, Remington's Pharmaceutical Sciences, 1990). Some variation in dosage will necessarily occur depending on the condition of the subject. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

**[0314]** An effective amount of therapeutic or prophylactic composition is determined based on the intended goal. The term "unit dose" or "dosage" refers to physically discrete units suitable for use in a subject, each unit containing a predetermined quantity of the composition calculated to produce the desired responses discussed above in association with its administration, i.e., the appropriate route and regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the protection desired.

**[0315]** Precise amounts of the composition also depend on the judgment of the practitioner and are peculiar to each individual. Factors affecting dose include physical and clinical state of the subject, route of administration, intended goal of treatment (alleviation of symptoms versus cure), and potency, stability, and toxicity of the particular composition.

**[0316]** Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically or prophylactically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above.

**[0317]** E. In Vitro, Ex Vivo, or In Vivo Administration

**[0318]** As used herein, the term in vitro administration refers to manipulations performed on cells removed from or outside of a subject, including, but not limited to cells in culture. The term ex vivo administration refers to cells which have been manipulated in vitro, and are subsequently admin-

istered to a subject. The term *in vivo* administration includes all manipulations performed within a subject.

**[0319]** In certain aspects of the present invention, the compositions may be administered either *in vitro*, *ex vivo*, or *in vivo*. In certain *in vitro* embodiments, autologous B-lymphocyte cell lines are incubated with a virus vector of the instant invention for 24 to 48 hours or with a variant SpA and/or coagulase and/or any other composition described herein for two hours. The transduced cells can then be used for *in vitro* analysis, or alternatively for *ex vivo* administration. U.S. Pat. Nos. 4,690,915 and 5,199,942, both incorporated herein by reference, disclose methods for *ex vivo* manipulation of blood mononuclear cells and bone marrow cells for use in therapeutic applications.

**[0320]** F. Antibodies And Passive Immunization

**[0321]** Another aspect of the invention is a method of preparing an immunoglobulin for use in prevention or treatment of staphylococcal infection comprising the steps of immunizing a recipient or donor with the vaccine of the invention and isolating immunoglobulin from the recipient or donor. An immunoglobulin prepared by this method is a further aspect of the invention. A pharmaceutical composition comprising the immunoglobulin of the invention and a pharmaceutically acceptable carrier is a further aspect of the invention which could be used in the manufacture of a medicament for the treatment or prevention of staphylococcal disease. A method for treatment or prevention of staphylococcal infection comprising a step of administering to a patient an effective amount of the pharmaceutical preparation of the invention is a further aspect of the invention.

**[0322]** Inocula for polyclonal antibody production are typically prepared by dispersing the antigenic composition in a physiologically tolerable diluent such as saline or other adjuvants suitable for human use to form an aqueous composition. An immunostimulatory amount of inoculum is administered to a mammal and the inoculated mammal is then maintained for a time sufficient for the antigenic composition to induce protective antibodies.

**[0323]** The antibodies can be isolated to the extent desired by well known techniques such as affinity chromatography (Harlow and Lane, 1988). Antibodies can include antiserum preparations from a variety of commonly used animals, e.g. goats, primates, donkeys, swine, horses, guinea pigs, rats or man.

**[0324]** An immunoglobulin produced in accordance with the present invention can include whole antibodies, antibody fragments or subfragments. Antibodies can be whole immunoglobulins of any class (e.g., IgG, IgM, IgA, IgD or IgE), chimeric antibodies or hybrid antibodies with dual specificity to two or more antigens of the invention. They may also be fragments (e.g., F(ab')<sub>2</sub>, Fab', Fab, Fv and the like) including hybrid fragments. An immunoglobulin also includes natural, synthetic, or genetically engineered proteins that act like an antibody by binding to specific antigens to form a complex.

**[0325]** A vaccine of the present invention can be administered to a recipient who then acts as a source of immunoglobulin, produced in response to challenge from the specific vaccine. A subject thus treated would donate plasma from which hyperimmune globulin would be obtained via conventional plasma fractionation methodology. The hyperimmune globulin would be administered to another subject in order to impart resistance against or treat staphylococcal infection. Hyperimmune globulins of the invention are particularly useful for treatment or prevention of staphylococcal disease in

infants, immune compromised individuals, or where treatment is required and there is no time for the individual to produce antibodies in response to vaccination.

**[0326]** An additional aspect of the invention is a pharmaceutical composition comprising two or more monoclonal antibodies (or fragments thereof; preferably human or humanised) reactive against at least two constituents of the immunogenic composition of the invention, which could be used to treat or prevent infection by Gram positive bacteria, preferably staphylococci, more preferably *S. aureus* or *S. epidermidis*. Such pharmaceutical compositions comprise monoclonal antibodies that can be whole immunoglobulins of any class, chimeric antibodies, or hybrid antibodies with specificity to two or more antigens of the invention. They may also be fragments (e.g., F(ab')<sub>2</sub>, Fab', Fab, Fv and the like) including hybrid fragments.

**[0327]** Methods of making monoclonal antibodies are well known in the art and can include the fusion of splenocytes with myeloma cells (Kohler and Milstein, 1975; Harlow and Lane, 1988). Alternatively, monoclonal Fv fragments can be obtained by screening a suitable phage display library (Vaughan et al., 1998). Monoclonal antibodies may be humanized or part humanized by known methods.

## VI. Examples

**[0328]** The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion. One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the methods described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses which are encompassed within the spirit of the invention as defined by the scope of the claims will occur to those skilled in the art.

### Example 1

#### Non-Toxicigenic Protein a Variants as Subunit Vaccines to Prevent *Staphylococcus aureus* Infections

**[0329]** A. Results

**[0330]** An Animal Model for *S. aureus* Infection

**[0331]** BALB/c mice were infected by intravenous injection with  $1 \times 10^7$  CFU of the human clinical isolate *S. aureus* Newman (Baba et al., 2007). Within 6 hours following infection, 99.999% of staphylococci disappeared from the blood stream and were distributed via the vasculature. Staphylococcal dissemination to peripheral tissues occurred rapidly, as the bacterial load in kidney and other peripheral organ tissues reached  $1 \times 10^5$  CFU g<sup>-1</sup> within the first three hours. The staphylococcal load in kidney tissues increased by 1.5 log CFU within twenty-four hours. Forty-eight hours following infection, mice developed disseminated abscesses in multiple organs, detectable by light microscopy of hematoxylin-eosin stained, thin-sectioned kidney tissue. The initial abscess diameter was 524 μM (±65 μM); lesions were initially marked by an influx of polymorphonuclear leukocytes (PMNs) and harbored no discernable organization of staphylococci, most of which appeared to reside within PMNs. On day 5 of infec-



tion, abscesses increased in size and enclosed a central population of staphylococci, surrounded by a layer of eosinophilic, amorphous material and a large cuff of PMNs. Histopathology revealed massive necrosis of PMNs in proximity to the staphylococcal nidus at the center of abscess lesions as well as a mantle of healthy phagocytes. A rim of necrotic PMNs were

of the appearance of infectious lesions and the morphological attributes of abscesses Ruined by *S. aureus* Newman were similar to those observed following mouse infection with *S. aureus* USA300 (LAC), the current epidemic community-acquired methicillin-resistant *S. aureus* (CA-MRSA) clone in the United States (Diep et al., 2006).

TABLE 4

Genotype	Genetic requirements for <i>S. aureus</i> Newman abscess formation in mice			Abscess formation in kidney tissue		
	Staphylococcal load in kidney tissue			<sup>d</sup> Surface	<sup>e</sup> Number of	<sup>f</sup> Significance (P-value)
	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup> tissue	<sup>b</sup> Significance (P-value)	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	abscesses (%)	abscesses per kidney	
wild-type	6.141 ± 0.192	—	—	70	4.364 ± 0.889	—
ΔsrtA	4.095 ± 0.347	6.7 × 10 <sup>-6</sup>	2.046	0	0.000 ± 0.000	0.0216
spa	5.137 ± 0.374	0.0144	1.004	13	0.375 ± 0.374	0.0356

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 5 days following infection in cohorts of fifteen BALB/c mice per challenge strain. Standard error of the means (±SEM) is indicated.

<sup>b</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values <0.05 were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Abscess formation in kidney tissues five days following infection was measured by macroscopic inspection (% positive)

<sup>e</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys from eight to ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

<sup>f</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values <0.05 were deemed significant.

observed at the periphery of abscess lesions, bordering eosinophilic, amorphous material that separates healthy renal tissue from lesions. Abscesses eventually reached a diameter of 1,524 μm on day 15 or 36. At later time intervals, the staphylococcal load was increased to 10<sup>4</sup>-10<sup>6</sup> CFU g<sup>-1</sup> and growing abscess lesions migrated towards the organ capsule. Peripheral lesions were prone to rupture, thereby releasing necrotic material and staphylococci into the peritoneal cavity or the retroperitoneal space. These events resulted in bacteremia as well as a secondary wave of abscesses, eventually precipitating a lethal outcome.

**[0332]** To enumerate staphylococcal load in renal tissue, animals were killed, their kidneys excised and tissue homogenate spread on agar media for colony formation. On day 5 of infection, a mean of 1×10<sup>6</sup> CFU g<sup>-1</sup> renal tissue for *S. aureus* Newman was observed. To quantify abscess formation, kidneys were visually inspected, and each individual organ was given a score of one or zero. The final sum was divided by the total number of kidneys to calculate percent surface abscesses (Table 4). In addition, randomly chosen kidneys were fixed in formalin, embedded, thin sectioned, and stained with hematoxylin-eosin. For each kidney, four sagittal sections at 200 μm intervals were viewed by microscopy. The numbers of lesions were counted for each section and averaged to quantify the number of abscesses within the kidneys. *S. aureus* Newman caused 4.364±0.889 abscesses per kidney, and surface abscesses were observed on 14 out of 20 kidneys (70%) (Table 4).

**[0333]** When examined by scanning electron microscopy, *S. aureus* Newman was located in tightly associated lawns at the center of abscesses. Staphylococci were contained by an amorphous pseudocapsule that separated bacteria from the cuff of abscesses leukocytes. No immune cells were observed in these central nests of staphylococci, however occasional red blood cells were located among the bacteria. Bacterial populations at the abscess center, designated staphylococcal abscess communities (SAC), appeared homogenous and coated by an electron-dense, granular material. The kinetics

**[0334]** *S. aureus* Protein A (Spa) Mutants are Avirulent and Cannot Form Abscesses

**[0335]** Sortase A is a transpeptidase that immobilizes nineteen surface proteins in the envelope of *S. aureus* strain Newman (Mazmanian et al., 1999; Mazmanian et al., 2000). Earlier work identified sortase A as a virulence factor in multiple animal model systems, however the contributions of this enzyme and its anchored surface proteins to abscess formation or persistence have not yet been revealed (Jonsson et al., 2002; Weiss et al., 2004). Compared to the wild-type parent (Baba et al., 2007), an isogenic srtA variant (ΔsrtA) failed to form abscess lesions on either macroscopic or histopathology examination on days 2, 5, or 15. In mice infected with the strA mutant, only 1×10<sup>4</sup> CFU g<sup>-1</sup> was recovered from kidney tissue on day 5 of infection, which is a 2.046 log<sub>10</sub> CFU g<sup>-1</sup> reduction compared to the wild-type parent strain (P=6.73×10<sup>-6</sup>). A similar defect was observed for the srtA mutant of MRSA strain USA300 (data not shown). Scanning electron microscopy showed that srtA mutants were highly dispersed and often associated with leukocytes in otherwise healthy renal tissue. On day fifteen following infection, srtA mutants were cleared from renal tissues, a ≥3.5 log<sub>10</sub> CFU g<sup>-1</sup> reduction compared to the wild-type (Table 4). Thus, sortase A anchored surface proteins enable the formation of abscess lesions and the persistence of bacteria in host tissues, wherein staphylococci replicate as communities embedded in an extracellular matrix and shielded from surrounding leukocytes by an amorphous pseudocapsule.

**[0336]** Sortase A anchors a large spectrum of proteins with LPXTG motif sorting signals to the cell wall envelope, thereby providing for the surface display of many virulence factors (Mazmanian et al., 2002). To identify surface proteins required for staphylococcal abscess foimation, *bursa aurealis* insertions were introduced in 5' coding sequences of genes that encode polypeptides with LPXTG motif proteins (Bae et al., 2004) and these mutations were transduced into *S. aureus* Newman. Mutations in the structural gene for Protein A (spa) reduced the staphylococcal load in infected mouse kidney

tissues by  $1.004 \log_{10}$  ( $P=0.0144$ ). When analyzed for their ability to form abscesses in kidney tissues by histopathology, we observed that the spa mutants were unable to form abscesses as compared with the wild-type parent strain *S. aureus* Newman (wild-type *S. aureus* Newman  $4.364 \pm 0.889$  abscesses per kidney vs. the isogenic spa mutant with  $0.375 \pm 0.374$  lesions;  $P=0.0356$ ).

**[0337]** Protein A Blocks Innate and Adaptive Immune Responses.

**[0338]** Studies identified Protein A as a critical virulence factor during the pathogenesis of *S. aureus* infections. Earlier work demonstrated that Protein A impedes phagocytosis of staphylococci by binding the Fc component of immunoglobulin (Jensen 1958; Uhlén et al., 1984), activates platelet aggregation via the von Willebrand factor (Hartleib et al., 2000), functions as a B cell superantigen by capturing the F(ab)<sub>2</sub> region of VH3 bearing IgM (Roben et al., 1995), and, through its activation of TNFR1, can initiate staphylococcal pneumonia (Gomez et al., 2004). Due to the fact that Protein A captures immunoglobulin and displays toxic attributes, the possibility that this surface molecule may function as a vaccine in humans has not been rigorously pursued. The inventors demonstrate for the first time that Protein A variants no longer able to bind to immunoglobulins, vWF and TNFR-1 are removed of their toxigenic potential and are able to stimulate humoral immune responses that protect against staphylococcal disease.

**[0339]** Molecular Basis of Protein A Surface Display and Function.

**[0340]** Protein A is synthesized as a precursor in the bacterial cytoplasm and secreted via its YSIRK signal peptide at the cross wall, i.e., the cell division septum of staphylococci (FIG. 1). (DeDent et al., 2007; DeDent et al., 2008). Following cleavage of the C-terminal LPXTG sorting signal, Protein A is anchored to bacterial peptidoglycan crossbridges by sortase A (Schneewind et al., 1995; Mazmanian et al., 1999; Mazmanian et al., 2000). Protein A is the most abundant surface protein of staphylococci; the molecule is expressed by virtually all *S. aureus* strains (Saïd-Salim et al., 2003; Cespedes et al., 2005; Kennedy et al., 2008). Staphylococci turn over 15-20% of their cell wall per division cycle (Navarre and Schneewind 1999). Murine hydrolases cleave the glycan strands and wall peptides of peptidoglycan, thereby releasing Protein A with its attached C-terminal cell wall disaccharide tetrapeptide into the extracellular medium (Ton-That et al., 1999). Thus, by physiological design, Protein A is both anchored to the cell wall and displayed on the bacterial surface but also released into surrounding tissues during host infection (Marraffini et al., 2006).

**[0341]** Protein A captures immunoglobulins on the bacterial surface and this biochemical activity enables staphylococcal escape from host innate and acquired immune responses (Jensen 1958; Goodyear and Silvean 2004). Interestingly, region X of Protein A (Guss et al., 1984), a repeat domain that tethers the IgG binding domains to the LPXTG sorting signal/cell wall anchor, is perhaps the most variable portion of the staphylococcal genome (Schneewind et al., 1992; Sad-Salim et al., 2003). Each of the five immunoglobulin binding domains of Protein A (SpA), formed from three helix bundles and designated E, D, A, B, and C, exerts similar structural and functional properties (Sjodahl 1977; Jansson et al., 1998). The solution and crystal structure of domain D has been solved both with and without the Fc and

V<sub>H</sub>3 (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille et al., 2000).

**[0342]** In the crystal structure complex, the Fab interacts with helix II and helix III of domain D via a surface composed of four VH region  $\beta$ -strands (Graille et al., 2000). The major axis of helix II of domain D is approximately 50° to the orientation of the strands, and the interhelical portion of domain D is most proximal to the C0 strand. The site of interaction on Fab is remote from the Ig light chain and the heavy chain constant region. The interaction involves the following domain D residues: Asp-36 of helix II as well as Asp-37 and Gln-40 in the loop between helix II and helix III, in addition to several other residues with SpA-D (Graille et al., 2000). Both interacting surfaces are composed predominantly of polar side chains, with three negatively charged residues on domain D and two positively charged residues on the 2A2 Fab buried by the interaction, providing an overall electrostatic attraction between the two molecules. Of the five polar interactions identified between Fab and domain D, three are between side chains. A salt bridge is formed between Arg-H19 and Asp-36 and two hydrogen bonds are made between Tyr-H59 and Asp-37 and between Asn-H82a and Ser-33. Because of the conservation of Asp-36 and Asp-37 in all five IgG binding domains of Protein A, these residues were selected for mutagenesis.

**[0343]** The SpA-D sites responsible for Fab binding are structurally separate from the domain surface that mediates Fc $\gamma$  binding. The interaction of Fc $\gamma$  with domain B primarily involves residues in helix I with lesser involvement of helix II (Deisenhofer 1981; Gouda et al., 1992). With the exception of the Gln-32, a minor contact in both complexes, none of the residues that mediate the Fc $\gamma$  interaction are involved in Fab binding. To examine the spatial relationship between these different Ig-binding sites, the SpA domains in these complexes have been superimposed to construct a model of a complex between Fab, the SpA-domain D, and the Fc $\gamma$  molecule. In this ternary model, Fab and Fc $\gamma$  form a sandwich about opposite faces of the helix II without evidence of steric hindrance of either interaction. These findings illustrate how, despite its small size (i.e., 56-61 aa), a SpA domain can simultaneously display both activities, explaining experimental evidence that the interactions of Fab with an individual domain are noncompetitive. Residues for the interaction between SpA-D and Fc $\gamma$  are Gln-9 and Gln-10.

**[0344]** In contrast, occupancy of the Fc portion of IgG on the domain D blocks its interaction with vWF A1 and probably also TNFR1 (O'Seaghda et al., 2006). Mutations in residues essential for IgG Fc binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF A1 and TNFR1 binding (Cedergren et al., 1993; Gomez et al., 2006; O'Seaghda et al. 2006), whereas residues critical for the V<sub>H</sub>3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) have no impact on the binding activities of IgG Fc, vWF A1 or TNFR1 (Jansson et al., 1998; Graille et al., 2000). The Protein A immunoglobulin Fab binding activity targets a subset of B cells that express VH3 family related IgM on their surface, i.e. these molecules function as VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells rapidly proliferate and then commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e. marginal zone B cells and follicular B2 cells) (Goodyear and Silverman 2003; Goodyear and Silverman 2004). It is important to note that more than 40% of circulating B cells are targeted by the Protein A interaction

and the VH3 family represents the largest family of human B cell receptors to impart protective humoral responses against pathogens (Goodyear and Silverman 2003; Goodyear and Silverman 2004). Thus, Protein A functions analogously to staphylococcal superantigens (Roben et al., 1995), albeit that the latter class of molecules, for example SEB, TSST-1, TSST-2, form complexes with the T cell receptor to inappropriately stimulate host immune responses and thereby precipitating characteristic disease features of staphylococcal infections (Roben et al., 1995; Tiedemann et al., 1995). Together these findings document the contributions of Protein A in establishing staphylococcal infections and in modulating host immune responses.

**[0345]** Non-Toxicogenic Variant of Protein A.

**[0346]** The inventors have developed a non-toxicogenic variant of staphylococcal Protein A and, with this reagent in hand, aimed for the first time to measure the immune response of animals to Protein A immunization. Further, the inventors address whether immunization of animals with a non-toxicogenic variant of Protein A could generate immune responses that raise protective immunity against staphylococcal infection.

**[0347]** To perturb the IgG Fc, vWF A1 and TNFR1 binding activities of Protein A, glutamine (Q) residues 9 and 10 [the numbering here is derived from that established for the SpA domain D] were modified generating lysine or glycine substitutions for both glutamines with the expectation that these substitutions abolish the ion bonds formed between wild-type Protein A and its ligands. The added effect of the dual lysine substitutions may be that these positively charged residues institute a repellent charge for immunoglobulins. To perturb IgM Fab VH3 binding, the inventors selected the aspartate (D) residues 36 and 37 of SpA-D, each of which is required for the association of Protein A with the B cell receptor. D36 and D37 were both substituted with alanine. The Q9,10K and D36,37A mutations were combined in the recombinant molecule SpA-D<sub>Q9,10K;D36,37A</sub> and examined for the binding attributes of Protein A.

**[0348]** In brief, the Protein A (spa) genomic sequence of *Staphylococcus aureus* N315 was PCR amplified with the primers (GCTGCACATATGGCGCAACACGATGAAGCTCAAC [5' primer](SEQ ID NO:35) and AGTG-GATCCTTATGCTTTGTTAGCATCTGC [3' primer] (SEQ ID NO:36)), cloned into the pET15b vector (pYSJ1, codons 48-486) (Stranger-Jones, et al., 2006) and recombinant plasmid transformed into *E. coli* BL21(DE3) (Studier et al., 1990). The Protein A product derived from pYSJ1 harbors SpA residues 36-265 fused to the N-terminal His tag (MGSSHHHHHSSGLVPRGS (SEQ ID NO:37)). Following IPTG inducible expression, recombinant N-terminal His<sub>6</sub>-tagged SpA was purified by affinity chromatography on Ni-NTA resin (Stranger-Jones et al., 2006). The domain D of SpA (SpA-D) was PCR amplified with a pair of specific primers (AACATATGTTCAACAAAGATCAACAAAGC [5' primer](SEQ ID NO:38) and AAGGATCCAGATTTCGTTAATTTTTTAGC [3' primer] (SEQ ID NO:39)), sub-cloned into the pET15b vector (pHAN 1, spa codons 212-261) and recombinant plasmid transformed into *E. coli* BL21(DE3) to express and purify recombinant N-terminal His<sub>6</sub>-tagged protein. To generate mutations in the SpA-D coding sequence, sets of two pairs of primers were synthesized (for D to A substitutions: CTTCAATCAAAGTCTTAAAGCCGC-CCCAAGCAAAGCACTAAC [5' primer] (SEQ ID NO:40) and GTTAGTGCTTTGGCTTGGGGCGGCTT-

TAAGACTTTGAATGAAG [3' primer] (SEQ ID NO:41); for Q to K substitutions CATATGTTCAACAAA-GATAAAAAAGCGCCTTCTATGAAATC [5' primer] (SEQ ID NO:42) and GATTCATAGAAGGCGCTTTTTT-TATCTTTGTTGAACATATG [3' primer] (SEQ ID NO:43); for Q to G substitutions CATATGTTCAACAAAGATG-GAGGAAGCGCCTTCTATGAAATC [5' primer] (SEQ ID NO:44) and GATTCATAGAAGGCGCTTCTC-CATCTTTGTTGAACATATG [3' primer] (SEQ ID NO:45). Primers were used for quick-change mutagenesis protocols. Following mutagenesis, DNA sequences were confirmed for each of the recombinant proteins: SpA, SpA-D and SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10K;D36,37A</sub>. All proteins were purified from lysates of recombinant *E. coli* using Ni-NTA chromatography and subsequently dialyzed against PBS and stored at 4° C.

**[0349]** To measure binding of immunoglobulin to Protein A and its variants, 200 µg of purified protein was diluted into a 1 ml volume using column buffer (50 mM Tris-HCl, 150 mM NaCl, pH7.5) and then loaded onto a pre-equilibrated Ni-NTA column (1 ml bed volume). Columns were washed with 10 ml of column buffer. 200 µg of purified human IgG was diluted in a total volume of 1 ml column buffer and then applied to each of the columns charged with Protein A and its variants. The columns were subsequently washed with 5 ml wash buffer (10 mM imidazole in column buffer) and 5 ml column buffer. Protein samples were eluted with 2 ml elution buffer (500 mM imidazole in column buffer), fractions collected and aliquots subjected to SDS-PAGE gel electrophoresis, followed by Coomassie-Blue staining. As shown in FIG. 3, wild-type Protein A (SpA) and its SpA-domain D both retained immunoglobulin during chromatography. In contrast, the SpA-D<sub>Q9,10K;D36,37A</sub> variant did not bind to immunoglobulin.

**[0350]** To quantify the binding of Protein A and its variants to the Fc portion of immunoglobulin and the VH3 domain of Fab, HRP conjugated human immunoglobulin G [hIgG], the Fc portion of human IgG [hFc] and the F(ab)<sub>2</sub> portion of human IgG [hF(ab)<sub>2</sub>] as well as ELISA assays were used to quantify the relative amount binding to Protein A and its variants. The data in FIG. 4 demonstrate the binding of SpA and SpA-D to hIgG and hFc, whereas SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10K;D36,37A</sub> displayed only background binding activities. SpA bound similar amounts of hFc and hF(ab)<sub>2</sub>, however the binding of SpA-D to hF(ab)<sub>2</sub> was reduced compared to full length SpA. This result suggests that the presence of multiple IgG binding domains may cooperatively increase the ability of Protein A to bind to the B cell receptor. When compared with the reduced binding power of SpA-D for hF(ab)<sub>2</sub>, of the two variants only SpA-D<sub>Q9,10K;D36,37A</sub> displayed a significant reduction in the ability to bind the VH3 domain of immunoglobulin. To examine the toxicogenic attributes of SpA-D and its variants, purified proteins were injected into mice, which were sacrificed after 4 hours to remove their spleens. Organ tissue was homogenized, capsular material removed and B cells stained with fluorescent CD19 antibodies. Following FACS analysis to quantify the abundance of B cells in splenic tissues, it was observed that SpA-D caused a 5% drop in the B cell count compared to a mock (PBS) control (FIG. 5). In contrast, SpA-D<sub>Q9,10K;D36,37A</sub> did not cause a reduction in B-cell counts, indicating that the mutant molecule had lost its toxicogenic attributes of stimulating B cell proliferation and death (FIG. 5). In summary, amino acid substitutions in the SpA-D residues Q9, Q10,

D36, and D37 abolished the ability of Protein A domains to bind immunoglobulins or exert toxic functions in human and animal tissues.

**[0351]** Non-Toxicigenic Protein A Variants Elicit Vaccine Protection.

**[0352]** To test whether or not Protein A and its variants can function as vaccine antigens, SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub>, and SpA-D<sub>Q9,10K;D36,37A</sub> were emulsified with complete or incomplete Freund's adjuvant and immunized 4 week old BALB/c mice on day 1 and day 11 with 50 µg of purified protein. Cohort of animals (n=5) were analyzed for humoral immune responses to immunization by bleeding the animals before (day 0) and after the immunization schedule (day 21). Table 5 indicates that immunized mice generated only a modest humoral immune response directed at wild-type Protein A or its SpA-D module, whereas the amount of antibody raised following immunization with SpA-D<sub>Q9,10K;D36,37A</sub> or SpA-D<sub>Q9,10K;D36,37A</sub> was increased four to five fold. Following intravenous challenge with 1×10<sup>7</sup> CFU *S. aureus* Newman, animals were killed on day 4, their kidneys removed and either analyzed for staphylococcal load (by plating tissue

lesions per organ (P=0.02 and P=0.04) were identified. Thus, immunization with non-toxicigenic Protein A variants generates increased humoral immune responses for Protein A and provides protective immunity against staphylococcal challenge. These data indicate that Protein A is an ideal candidate for a human vaccine that prevents *S. aureus* disease.

**[0353]** These exciting results have several implications for the design of a human vaccine. First, the generation of substitution mutations that affect the ability of the immunoglobulin binding domains of Protein A, either alone or in combination of two or more domains, can generate non-toxicigenic variants suitable for vaccine development. It seems likely that a combination of mutant IgG binding domains closely resembling the structure of Protein A can generate even better humoral immune responses as is reported here for the SpA-domain D alone. Further, a likely attribute of Protein A specific antibodies may be that the interaction of antigen binding sites with the microbial surface can neutralize the ability of staphylococci to capture immunoglobulins via their Fc portion or to stimulate the B cell receptor via the VH3 binding activities.

TABLE 5

Non-toxicigenic Protein A variants as vaccine antigens that prevent <i>S. aureus</i> disease									
Antigen	Bacterial load in kidney (n = number of mice)			IgG titer	<sup>d</sup> Surface abscess	Abscess formation in mice (n = number of mice)			
	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> Reduction	<sup>c</sup> P value			Reduction	<sup>e</sup> Histopathology	Reduction	<sup>f</sup> P value
Mock	6.46 ± 0.25 (n = 19)	—	—	<100	14/19 (70%)	—	3.7 ± 1.2 (n = 10)	—	—
SpA	3.95 ± 0.56 (n = 20)	2.51	0.0003	1706 ± 370	10/20 (50%)	32%	2.1 ± 1.2 (n = 10)	2.2	0.35
SpA-D	4.43 ± 0.41 (n = 18)	2.03	0.0001	381 ± 27	10/18 (55%)	25%	1.5 ± 0.8 (n = 10)	2.2	0.15
SpA-D1	3.39 ± 0.50 (n = 19)	3.07	<0.0001	5600 ± 801	6/20 (30%)	59%	0.5 ± 0.4 (n = 10)	3.2	0.02
SpA-D2	3.43 ± 0.46 (n = 19)	3.03	<0.0001	3980 ± 676	6/19 (32%)	57%	0.8 ± 0.5 (n = 10)	2.9	0.04

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of 18 to 20 BALB/c mice. Standard error of the means (±SEM) is indicated.

<sup>c</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values <0.05 were deemed significant.

<sup>b</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Abscess formation in kidney tissues four days following infection was measured by macroscopic inspection (% positive)

<sup>e</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys from ten animals, the number of abscesses per kidney was recorded and averaged for the final mean (±SEM).

<sup>f</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values <0.05 were deemed significant.

SpA-D1 and SpA-D2 represent SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10G;D36,37A</sub>, respectively.

homogenate on agar plates and enumerating colony forming units, CFU) or histopathology. As expected, mock (PBS) immunized mice (n=19) harbored 6.46 log<sub>10</sub> (±0.25) CFU in kidney tissue and infectious lesions were organized into 3.7 (±1.2) abscesses per organ (n=10) (Table 5). Immunization of animals with SpA led to a 2.51 log<sub>10</sub> CFU reduction on day 5 (P=0.0003) with 2.1 (±1.2) abscesses per organ. The latter data indicate that there was no significant reduction in abscess formation (P=0.35). Immunization with SpA-D generated similar results: a 2.03 log<sub>10</sub> CFU reduction on day 5 (P=0.0001) with 1.5 (±0.8) abscesses per organ (P=0.15). In contrast, immunization with SpA-D<sub>Q9,10K;D36,37A</sub> or SpA-D<sub>Q9,10G;D36,37A</sub> created increased protection, with 3.07 log<sub>10</sub> and 3.03 log<sub>10</sub> CFU reduction on day 4, respectively (statistical significance P<0.0001 for both observations). Further, immunization with both SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10G;D36,37A</sub> generated significant protection from staphylococcal abscess formation, as only 0.5 (±0.4) and 0.8 (±0.5) infectious

**[0354]** Vaccine Protection in Murine Abscess, Murine Lethal Infection, and Murine Pneumonia Models.

**[0355]** Three animal models have been established for the study of *S. aureus* infectious disease. These models are used here to examine the level of protective immunity provided via the generation of Protein A specific antibodies.

**[0356]** Murine Abscess

**[0357]** BALB/c mice (24-day-old female, 8-10 mice per group, Charles River Laboratories, Wilmington, Mass.) are immunized by intramuscular injection into the hind leg with purified protein (Chang et al., 2003; Schneewind et al., 1992). Purified SpA, SpA-D or SpA-DQ9,10K;D36,37A (50 µg protein) is administered on days 0 (emulsified 1:1 with complete Freund's adjuvant) and 11 (emulsified 1:1 with incomplete Freund's adjuvant). Blood samples are drawn by retroorbital bleeding on days 0, 11, and 20. Sera are examined by ELISA for IgG titers for specific SpA-D and SpA-DQ9,10K;D36,

37A binding activity. Immunized animals are challenged on day 21 by retroorbital injection of 100  $\mu$ l of *S. aureus* Newman or *S. aureus* USA300 suspension ( $1 \times 10^7$  cfu). For this, overnight cultures of *S. aureus* Newman are diluted 1:100 into fresh tryptic soy broth and grown for 3 h at 37° C. Staphylococci are centrifuged, washed twice, and diluted in PBS to yield an  $A_{600}$  of 0.4 ( $1 \times 10^8$  cfu per ml). Dilutions are verified experimentally by agar plating and colony formation. Mice are anesthetized by intraperitoneal injection of 80-120 mg of ketamine and 3-6 mg of xylazine per kilogram of body weight and infected by retroorbital injection. On day 5 or 15 following challenge, mice are euthanized by compressed CO<sub>2</sub> inhalation. Kidneys are removed and homogenized in 1% Triton X-100. Aliquots are diluted and plated on agar medium for triplicate determination of cfu. For histology, kidney tissue is incubated at room temperature in 10% formalin for 24 h. Tissues are embedded in paraffin, thin-sectioned, stained with hematoxylin-eosin, and examined by microscopy.

**[0358]** Murine Lethal Infection

**[0359]** BALB/c mice (24-day-old female, 8-10 mice per group, Charles River Laboratories, Wilmington, Mass.) are immunized by intramuscular injection into the hind leg with purified SpA, SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> (50  $\mu$ g protein). Vaccine is administered on days 0 (emulsified 1:1 with complete Freund's adjuvant) and 11 (emulsified 1:1 with incomplete Freund's adjuvant). Blood samples are drawn by retroorbital bleeding on days 0, 11, and 20. Sera are examined by ELISA for IgG titers with specific SpA-D and SpA-D<sub>Q9,10K;D36,37A</sub> binding activity. Immunized animals are challenged on day 21 by retroorbital injection of 100  $\mu$ l of *S. aureus* Newman or *S. aureus* USA300 suspension ( $15 \times 10^7$  cfu) (34). For this, overnight cultures of *S. aureus* Newman are diluted 1:100 into fresh tryptic soy broth and grown for 3 h at 37° C. Staphylococci are centrifuged, washed twice, diluted in PBS to yield an  $A_{600}$  of 0.4 ( $1 \times 10^8$  cfu per ml) and concentrated. Dilutions are verified experimentally by agar plating and colony formation. Mice are anesthetized by intraperitoneal injection of 80-120 mg of ketamine and 3-6 mg of xylazine per kilogram of body weight. Immunized animals are challenged on day 21 by intraperitoneal inject with  $2 \times 10^{10}$  cfu of *S. aureus* Newman or  $3-10 \times 10^9$  cfu of clinical *S. aureus* isolates. Animals are monitored for 14 days, and lethal disease is recorded.

**[0360]** Murine Pneumonia Model

**[0361]** *S. aureus* strains Newman or USA300 (LAC) are grown at 37° C. in tryptic soy broth/agar to OD<sub>660</sub> 0.5. 50-ml culture aliquots are centrifuged, washed in PBS, and suspended in 750  $\mu$ l PBS for mortality studies ( $3-4 \times 10^8$  CFU per 30- $\mu$ l volume), or 1,250  $\mu$ l PBS ( $2 \times 10^8$  CFU per 30- $\mu$ l volume) for bacterial load and histopathology experiments (2, 3). For lung infection, 7-wk-old C57BL/6J mice (The Jackson Laboratory) are anesthetized before inoculation of 30  $\mu$ l of *S. aureus* suspension into the left nare. Animals are placed into the cage in a supine position for recovery and observed for 14 days. For active immunization, 4-wk-old mice receive 20  $\mu$ g SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> in CFA on day 0 via the i.m. route, followed by a boost with 20  $\mu$ g SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> in incomplete Freund's adjuvant (IFA) on day 10. Animals are challenged with *S. aureus* on day 21. Sera are collected before immunization and on day 20 to assess specific antibody production. For passive immunization studies, 7-wk-old mice receive 100  $\mu$ l of either NRS (normal rabbit serum) or SpA-D-specific rabbit antisera via i.p. injection 24 h before challenge. To assess the pathological corre-

lates of pneumonia, infected animals are killed via forced CO<sub>2</sub> inhalation before removal of both lungs. The right lung is homogenized for enumeration of lung bacterial load. The left lung is placed in 1% formalin and paraffin embedded, thin sectioned, stained with hematoxylin-eosin, and analyzed by microscopy.

**[0362]** Rabbit Antibodies

**[0363]** Purified 200  $\mu$ g SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> is used as an immunogen for the production of rabbit antisera. 200  $\mu$ g protein is emulsified with CFA for injection at day 0, followed by booster injections with 200  $\mu$ g protein emulsified with IFA on days 21 and 42. Rabbit antibody titers are determined by ELISA. Purified antibodies are obtained by affinity chromatography of rabbit serum on SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> sepharose. The concentration of eluted antibodies is measured by absorbance at  $A_{280}$  and specific antibody titers are determined by ELISA.

**[0364]** Active Immunization with SpA-Domain D Variants.

**[0365]** To determine vaccine efficacy, animals are actively immunized with purified SpA-D or SpAD<sub>Q9,10K;D36,37A</sub>. As a control, animals are immunized with adjuvant alone. Antibody titers against Protein A preparations are determined using SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> as antigens; note that the SpA-D<sub>Q9,10K;D36,37A</sub> variant cannot bind the Fc or Fab portion of IgG. Using infectious disease models described above, any reduction in bacterial load (murine abscess and pneumonia), histopathology evidence of staphylococcal disease (murine abscess and pneumonia) and protection from lethal disease (murine lethal challenge and pneumonia) is measured.

**[0366]** Passive Immunization with Affinity Purified Rabbit Polyclonal Antibodies Generated Against SpA-Domain D Variants.

**[0367]** To determine protective immunity of Protein A specific rabbit antibodies, mice are passively immunized with 5 mg/kg of purified SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> derived rabbit antibodies. Both of these antibody preparations are purified by affinity chromatography using immobilized SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub>. As a control, animals are passively immunized with rV 10 antibodies (a plague protective antigen that has no impact on the outcome of staphylococcal infections). Antibody titers against all Protein A preparations are determined using SpA-D<sub>Q9,10K;D36,37A</sub> as an antigen, as this variant cannot bind the Fc or Fab portion of IgG. Using the infectious disease models described above, the reduction in bacterial load (murine abscess and pneumonia), histopathology evidence of staphylococcal disease (murine abscess and pneumonia), and the protection from lethal disease (murine lethal challenge and pneumonia) is measured.

Example 2

**[0368]** Non-Toxicogenic Protein a Vaccine for Methicillin-Resistant *Staphylococcus aureus* infection

**[0369]** Clinical isolates of *S. aureus* express protein A (Shopsin et al., 1999, whose primary translational product is comprised of an N-terminal signal peptide (DeDent et al., 2008), five Ig-BDs (designated E, D, A, B and C) (Sjodahl, 1977), region X with variable repeats of an eight residue peptide (Guss et al., 1984), and C-terminal sorting signal for the cell wall anchoring of SpA (Schneewind et al., 1992; Schneewind et al., 1995) (FIG. 6). Guided by amino acid homology (Uhlen et al., 1984), the triple  $\alpha$ -helical bundle

structure of IgBDs (Deisenhofer et al., 1978; Deisenhofer et al., 1981) and their atomic interactions with Fab  $V_H3$  (Graille et al., 2000) or Fc $\gamma$  (Gouda et al., 1998), glutamine 9 and 10 were selected as well as aspartate 36 and 37 as critical for the association of SpA with antibodies or B cell receptor, respectively. Substitutions Gln9Lys, Gln10Lys, Asp36Ala and Asp37Ala were introduced into the D domain to generate SpA-D<sub>KKAA</sub> (FIG. 6). The ability of isolated SpA-D or SpA-D<sub>KKAA</sub> to bind human IgG was analyzed by affinity chromatography (FIG. 6). Polyhistidine tagged SpA-D as well as full-length SpA retained human IgG on Ni-NTA, whereas SpA-D<sub>KKAA</sub> and a negative control (SrtA) did not (FIG. 6). A similar result was observed with von Willebrand factor (Hartleib et al., 2000), which, along with tumor necrosis factor receptor 1 (TNFR1)(Gomez et al., 2004), can also bind protein A via glutamine 9 and 10 (FIG. 6). Human immunoglobulin encompasses 60-70%  $V_H3$ -type IgG. The inventors distinguish between Fc domain and B cell receptor activation of Igs and measured association of human Fc $\gamma$  and F(ab)<sub>2</sub> fragments, both of which bound to full-length SpA or SpA-D, but not to SpA-D<sub>KKAA</sub> (FIG. 6). Injection of SpA-D into the peritoneal cavity of mice resulted in B cell expansion followed by apoptotic collapse of CD 19+ lymphocytes in spleen tissue of BALB/c mice (Goodyear and Silverman, 2003)(FIG. 6). B cell superantigen activity was not observed following injection with SpA-D<sub>KKAA</sub>, and TUNEL-staining of splenic tissue failed to detect the increase in apoptotic cells that follows injection of SpA or SpA-D (FIG. 6).

**[0370]** Naive six week old BALB/c mice were injected with 5014 each of purified SpA, SpA-D or SpA-D<sub>KKAA</sub> emulsified in CFA and boosted with the same antigen emulsified in IFA. In agreement with the hypothesis that SpA-D promotes the apoptotic collapse of activated clonal B cell populations, the inventors observed a ten-fold higher titer of SpA-D<sub>KKAA</sub> specific antibodies following immunization of mice with the non-toxicigenic variant as compared to the B cell superantigen (SpA-D vs. SpA-D<sub>KKAA</sub> P<0.0001, Table 6). Antibody titers raised by immunization with full-length SpA were higher than those elicited by SpA-D (P=0.0022), which is likely due to the larger size and reiterative domain structure of this antigen (Table 6). Nevertheless, even SpA elicited lower antibody titers than SpA-D<sub>KKAA</sub> (P=0.0003), which encompasses only 50 amino acids of protein A (520 residues, SEQ ID NO:33). Immunized mice were challenged by intravenous inoculation with *S. aureus* Newman and the ability of staphylococci to seed abscesses in renal tissues was examined by necropsy four days after challenge. In homogenized renal tissue of mock (PBS/adjuvant) immunized mice, an average staphylococcal load of 6.46 log<sub>10</sub> CFU g<sup>-1</sup> was enumerated (Table 6). Immunization of mice with SpA or SpA-D led to a reduction in staphylococcal load, however SpA-D<sub>KKAA</sub> vaccinated animals displayed an even greater, 3.07 log<sub>10</sub> CFU g<sup>-1</sup> reduction of *S. aureus* Newman in renal tissues (P<0.0001, Table 6). Abscess formation in kidneys was analyzed by histopathology (FIG. 7). Mock immunized animals harbored an average of 3.7 (±1.2) abscesses per kidney (Table 6). Vaccination with SpA-D<sub>KKAA</sub> reduced the average number of abscesses to 0.5 (±0.4) (P=0.0204), whereas immunization with SpA or SpA-D did not cause a significant reduction in the number of abscess lesions (Table 6). Lesions from SpA-D<sub>KKAA</sub> vaccinated animals were smaller in size, with fewer infiltrating PMNs and characteristically lacked staphylococcal abscess communities (Cheng et al., 2009)(FIG. 7). Abscesses in animals that had been immunized with SpA or

SpA-D displayed the same overall structure of lesions in mock immunized animals (FIG. 7).

**[0371]** The inventors examined whether SpA-D<sub>KKAA</sub> immunization can protect mice against MRSA strains and selected the USA300 LAC isolate for animal challenge (Diep et al., 2006). This highly virulent CA-MRSA strain spread rapidly throughout the United States, causing significant human morbidity and mortality (Kennedy et al., 2008). Compared to adjuvant control mice, SpA-D<sub>KKAA</sub> immunized animals harbored a 1.07 log<sub>10</sub> CFU g<sup>-1</sup> reduction in bacterial load of infected kidney tissues. Histopathology examination of renal tissue following *S. aureus* USA300 challenge revealed that the average number of abscesses was reduced from 4.04 (±0.8) to 1.6 (±0.6) (P=0.02774). In contrast, SpA or SpA-D immunization did not cause a significant reduction in bacterial load or abscess formation (Table 6).

**[0372]** Rabbits were immunized with SpA-D<sub>KKAA</sub> and specific antibodies were purified on SpA-D<sub>KKAA</sub> affinity column followed by SDS-PAGE (FIG. 8). SpA-D<sub>KKAA</sub> specific IgG was cleaved with pepsin to generate Fc $\gamma$  and F(ab)<sub>2</sub> fragments, the latter of which were purified by chromatography on SpA-D<sub>KKAA</sub> column (FIG. 8). Binding of human IgG or vWF to SpA or SpA-D was perturbed by SpA-D<sub>KKAA</sub> specific F(ab)<sub>2</sub>, indicating that SpA-D<sub>KKAA</sub> derived antibodies neutralize the B cell superantigen function of protein A as well as its interactions with Ig (FIG. 8).

**[0373]** To further improve the vaccine properties for non-toxicigenic protein A, the inventors generated SpA<sub>KKAA</sub>, which includes all five IgBDs with four amino acid substitutions—substitutions corresponding to Gln9Lys, Gln10Lys, Asp36Ala and Asp37Ala of domain D—in each of its five domains (E, D, A, B and C). Polyhistidine tagged SpA<sub>KKAA</sub> was purified by affinity chromatography and analyzed by Coomassie Blue-stained SDS-PAGE (FIG. 9). Unlike full-length SpA, SpA<sub>KKAA</sub> did not bind human IgG, Fe and F(ab)<sub>2</sub> or vWF (FIG. 9). SpA<sub>KKAA</sub> failed to display B cell superantigen activity, as injection of the variant into BALB/c mice did not cause a depletion of CD19+ B cells in splenic tissue (FIG. 9). SpA<sub>KKAA</sub> vaccination generated higher specific antibody titers than SpA-D<sub>KKAA</sub> immunization and provided mice with elevated protection against *S. aureus* USA300 challenge (Table 6). Four days following challenge, SpA<sub>KKAA</sub> vaccinated animals harbored 3.54 log<sub>10</sub> CFU g<sup>-1</sup> fewer staphylococci in renal tissues (P=0.0001) and also caused a greater reduction in the number of abscess lesions (P=0.0109) (Table 6).

**[0374]** SpA<sub>KKAA</sub> was used to immunize rabbits. Rabbit antibodies specific for SpA-D<sub>KKAA</sub> or SpA<sub>KKAA</sub> were affinity purified on matrices with immobilized cognate antigen and injected at a concentration of 5 mg kg<sup>-1</sup> body weight into the peritoneal cavity of BALB/c mice (Table 7). Twenty-four hours later, specific antibody titers were determined in serum and animals challenged by intravenous inoculation with *S. aureus* Newman. Passive transfer reduced the staphylococcal load in kidney tissues for SpA-D<sub>KKAA</sub> (P=0.0016) or SpA<sub>KKAA</sub> (P=0.0005) specific antibodies. On histopathology examination, both antibodies reduced the abundance of abscess lesions in the kidneys of mice challenged with *S. aureus* Newman (Table 7). Together these data reveal that vaccine protection following immunization with SpA-D<sub>KKAA</sub> or SpA<sub>KKAA</sub> is conferred by antibodies that neutralize protein A.

TABLE 6

Immunization of mice with protein A vaccines.						
Staphylococcal load and abscess formation in renal tissue						
Antigen	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> P-value	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>d</sup> IgG Titer	<sup>e</sup> Number of abscesses	<sup>f</sup> P-value
<i>S. aureus</i> Newman challenge						
Mock	6.46 ± 0.25	—	—	<100	3.7 ± 1.2	—
SpA	3.95 ± 0.56	0.0003	2.51	1706 ± 370	2.1 ± 1.2	0.3581
SpA-D	4.43 ± 0.41	0.0001	2.03	381 ± 27	1.5 ± 0.8	0.1480
SpA D <sub>IKAA</sub>	3.39 ± 0.50	<0.0001	3.07	5600 ± 801	0.5 ± 0.4	0.0204
<i>S. aureus</i> USA300 (LAC) challenge						
Mock	7.20 ± 0.24	—	—	<100	4.0 ± 0.8	—
SpA	6.81 ± 0.26	0.2819	0.39	475 ± 60	3.3 ± 1.0	0.5969
SpA-D	6.34 ± 0.52	0.1249	0.85	358 ± 19	2.2 ± 0.6	0.0912
SpA-D <sub>KKAA</sub>	6.00 ± 0.42	0.0189	1.20	3710 ± 1147	1.6 ± 0.6	0.0277
SpA <sub>KKAA</sub>	3.66 ± 0.76	0.0001	3.54	10200 ± 2476	1.2 ± 0.5	0.0109

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of fifteen to twenty BALB/c mice per immunization. Representative of two independent and reproducible animal experiments is shown. Standard error of the means (±SEM) is indicated.

<sup>b</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values <0.05 were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Means of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA

<sup>e</sup>Histopathology of hematoxyline-eosin stained, thin sectioned kidneys from ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

TABLE 7

Passive immunization of mice with antibodies against protein A.						
Staphylococcal load and abscess formation in renal tissue						
<sup>a</sup> Antibody	<sup>b</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>c</sup> P-value	<sup>d</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>e</sup> IgG Titer	<sup>f</sup> Number of abscesses	<sup>g</sup> P-value
Mock	7.10 ± 0.14	—	—	<100	4.5 ± 0.8	—
α-SpA-D <sub>KKAA</sub>	5.53 ± 0.43	0.0016	1.57	466 ± 114	1.9 ± 0.7	0.0235
α-SpA <sub>KKAA</sub>	5.69 ± 0.34	0.0005	1.41	1575 ± 152	1.6 ± 0.5	0.0062

<sup>a</sup>Affinity purified antibodies were injected into the peritoneal cavity of BALB/c mice at a concentration of 5 mg · kg<sup>-1</sup> twenty-four hours prior to intravenous challenge with 1 × 10<sup>7</sup> CFU *S. aureus* Newman.

<sup>b</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of fifteen BALB/c mice per immunization. Representative of two independent and reproducible animal experiments is shown. Standard error of the means (±SEM) is indicated.

<sup>c</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values <0.05 were deemed significant.

<sup>d</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>e</sup>Means of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA

<sup>f</sup>Histopathology of hematoxyline-eosin stained, thin sectioned kidneys from ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

**[0375]** Following infection with virulent *S. aureus*, mice do not develop protective immunity against subsequent infection with the same strain (Burts et al., 2008) (FIG. 10). The average abundance of SpA-D<sub>KKAA</sub> specific IgG in these animals was determined by dot blot as 0.20 μg ml<sup>-1</sup> (±0.04) and 0.14 μg ml<sup>-1</sup> (±0.01) for strains Newman and USA300 LAC, respectively (FIG. 9). The minimal concentration of protein A-specific IgG required for disease protection in SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub> vaccinated animals (P .05 log<sub>10</sub> reduction in staphylococcal CFU g<sup>-1</sup> renal tissue) was calculated as 4.05 μg ml<sup>-1</sup> (±0.88). Average serum concentration of SpA-specific IgG in adult healthy human volunteers (n=16) was 0.21 μg ml<sup>-1</sup> (±0.02). Thus, *S. aureus* infections in mice or humans are not associated with immune responses that raise significant levels of neutralizing antibodies directed against protein A, which is likely due to the B cell superantigen attributes of this molecule. In contrast, the average serum concentration of IgG specific for diphtheria toxin in human volunteers, 0.068 ml<sup>-1</sup> (±0.20), was within range for protective immunity against diphtheria (Behring, 1890; Lagergard et al., 1992).

**[0376]** Clinical *S. aureus* isolates express protein A, an essential virulence factor whose B cell superantigen activity and evasive attributes towards opsono-phagocytic clearance are absolutely required for staphylococcal abscess formation (Palmqvist et al., 2005; Cheng et al., 2009; Silverman and Goodyear, 2006). Protein A can thus be thought of as a toxin, essential for pathogenesis, whose molecular attributes must be neutralized in order to achieve protective immunity. By generating non-toxic variants unable to bind Igs via Fcγ or VH<sub>3</sub>-Fab domains, the inventors measure here for the first time protein A neutralizing immune responses as a correlate for protective immunity against *S. aureus* infection. In contrast to many methicillin-sensitive strains, CA-MRSA isolate USA300 LAC is significantly more virulent (Cheng et al., 2009). For example, immunization of experimental animals with the surface protein IsdB (Kuklin et al., 2006; Stranger-Jones et al., 2006) raises antibodies that confer protection against *S. aureus* Newman (Stranger-Jones et al., 2009) but not against USA300 challenge.

[0377] The methods utilized include:

[0378] Bacterial Strains and Growth.

[0379] *Staphylococcus aureus* strains Newman and USA300 were grown in tryptic soy broth (TSB) at 37° C. *Escherichia coli* strains DH5 $\alpha$  and BL21 (DE3) were grown in Luria-Bertani (LB) broth with 100  $\mu$ g mY<sup>1</sup> ampicillin at 37° C.

[0380] Rabbit Antibodies.

[0381] The coding sequence for SpA was PCR-amplified with two primers, gctgcacatatggcgcaacacgatgaagctcaac (SEQ ID NO:35) and agtggatccttatgcttgagattgttagcatctgc (SEQ ID NO:36) using *S. aureus* Newman template DNA. SpA-D was PCR-amplified with two primers, aacatattgcaacaagatcaacaaagc (SEQ ID NO:38) and aaggatccagatgctt-taatttttagc (SEQ ID NO:39). The sequence for SpA-D<sub>KKAA</sub> was mutagenized with two sets of primers catatgttcaacaaagataaaaaagcgctctatgaaatc (SEQ ID NO:42) and gatttcatagaaggcgtttttttatctgttgaacatag (SEQ ID NO:43) for Q9K, Q10K as well as cttcattcaaatcttaagccgc-cccaagcacaagcactaac (SEQ ID NO:40), and gttagtgtggct-tggggcgattaagacttgaatgaag (SEQ ID NO:41) for D36A, D37A. The sequence of SpA<sub>KKAA</sub> was synthesized by Integrated DNA Technologies, Inc. PCR products were cloned into pET-15b generating N-terminal His<sub>6</sub> tagged recombinant protein. Plasmids were transformed into BL21 (DE3). Overnight cultures of transformants were diluted 1:100 into fresh media and grown at 37° C. to an OD<sub>600</sub> 0.5, at which point cultures were induced with 1 mM isopropyl  $\beta$ -D-1-thiogalactopyranoside (IPTG) and grown for an additional three hours. Bacterial cells were sedimented by centrifugation, suspended in column buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl) and disrupted with a French pressure cell at 14,000 psi. Lysates were cleared of membrane and insoluble components by ultracentrifugation at 40,000 $\times$ g. Proteins in the soluble lysate were subjected to nickel-nitrilotriacetic acid (Ni-NTA, Qiagen) affinity chromatography. Proteins were eluted in column buffer containing successively higher concentrations of imidazole (100-500 mM). Protein concentrations were determined by bicinchonic acid (BCA) assay (Thermo Scientific). For antibody generation, rabbits (6 month old New-Zealand white, female, Charles River Laboratories) were immunized with 500  $\mu$ g protein emulsified in Complete Freund's Adjuvant (Difco) by subscapular injection. For booster immunizations, proteins emulsified in Incomplete Freund's Adjuvant and injected 24 or 48 days following the initial immunization. On day 60, rabbits were bled and serum recovered.

[0382] Purified antigen (5 mg protein) was covalently linked to HiTrap NHS-activated HP columns (GE Healthcare). Antigen-matrix was used for affinity chromatography of 10-20 ml of rabbit serum at 4° C. Charged matrix was washed with 50 column volumes of PBS, antibodies eluted with elution buffer (1 M glycine, pH 2.5, 0.5 M NaCl) and immediately neutralized with 1M Tris-HCl, pH 8.5. Purified antibodies were dialyzed overnight against PBS at 4° C.

[0383] F(ab)<sub>2</sub> Fragments.

[0384] Affinity purified antibodies were mixed with 3 mg of pepsin at 37° C. for 30 minutes. The reaction was quenched

with 1 M Tris-HCl, pH 8.5 and F(ab)<sub>2</sub> fragments were affinity purified with specific antigen-conjugated HiTrap NHS-activated HP columns. Purified antibodies were dialyzed overnight against PBS at 4° C., loaded onto SDS-PAGE gel and visualized with Coomassie Blue staining.

[0385] Active and Passive Immunization.

[0386] BALB/c mice (3 week old, female, Charles River Laboratories) were immunized with 50  $\mu$ g protein emulsified in Complete Freund's Adjuvant (Difco) by intramuscular injection. For booster immunizations, proteins were emulsified in Incomplete Freund's Adjuvant and injected 11 days following the initial immunization. On day 20 following immunization, 5 mice were bled to obtain sera for specific antibody titers by enzyme-linked immunosorbent assay (ELISA).

[0387] Affinity purified antibodies in PBS were injected at a concentration 5 mg kg<sup>-1</sup> of experimental animal weight into the peritoneal cavity of BALB/c mice (6 week old, female, Charles River Laboratories) 24 hours prior to challenge with *S. aureus*. Animal blood was collected via periorbital vein puncture. Blood cells were removed with heparinized micro-hematocrit capillary tubes (Fisher) and Z-gel serum separation micro tubes (Sarstedt) were used to collect and measure antigen specific antibody titers by ELISA.

[0388] Mouse Renal Abscess.

[0389] Overnight cultures of *S. aureus* Newman or USA300 (LAC) were diluted 1:100 into fresh TSB and grown for 2 hours at 37° C. Staphylococci were sedimented, washed and suspended PBS at OD<sub>600</sub> of 0.4 (~1 $\times$ 10<sup>8</sup> CFU ml<sup>-1</sup>). Inocula were quantified by spreading sample aliquots on TSA and enumerating colonies formed. BALB/c mice (6 week old, female, Charles River Laboratories) were anesthetized via intraperitoneal injection with 100 mg ml<sup>-1</sup> ketamine and 20 mg ml<sup>-1</sup> xylazine per kilogram of body weight. Mice were infected by retro-orbital injection with 1 $\times$ 10<sup>7</sup> CFU of *S. aureus* Newman or 5 $\times$ 10<sup>6</sup> CFU of *S. aureus* USA300. On day 4 following challenge, mice were killed by CO<sub>2</sub> inhalation. Both kidneys were removed, and the staphylococcal load in one organ was analyzed by homogenizing renal tissue with PBS, 1% Triton X-100. Serial dilutions of homogenate were spread on TSA and incubated for colony formation. The remaining organ was examined by histopathology. Briefly, kidneys were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with hematoxylin-eosin, and inspected by light microscopy to enumerate abscess lesions. All mouse experiments were performed in accordance with the institutional guidelines following experimental protocol review and approval by the Institutional Biosafety Committee (IBC) and the Institutional Animal Care and Use Committee (IACUC) at the University of Chicago.

[0390] Protein A Binding.

[0391] For human IgG binding, Ni-NTA affinity columns were pre-charged with 200  $\mu$ g of purified proteins (SpA, SpA-D, SpA-D<sub>KKAA</sub>, and SrtA) in column buffer. After washing, 200  $\mu$ g of human IgG (Sigma) was loaded onto the column. Protein samples were collected from washes and elutions and subjected to SDS-PAGE gel electrophoresis,



followed by Coomassie Blue staining. Purified proteins (SpA, SpA<sub>KKAA</sub>, SpA-D and SpA-D<sub>KKAA</sub>) were coated onto MaxiSorp ELISA plates (NUNC) in 0.1M carbonate buffer (pH 9.5) at 1  $\mu\text{g ml}^{-1}$  concentration overnight at 4° C. Plates were next blocked with 5% whole milk followed by incubation with serial dilutions of peroxidase-conjugated human IgG, Fc or F(ab)<sub>2</sub> fragments for one hour. Plates were washed and developed using OptEIA ELISA reagents (BD). Reactions were quenched with 1 M phosphoric acid and A<sub>450</sub> readings were used to calculate half maximal titer and percent binding.

**[0392]** von Willebrand Factor (vWF) Binding Assays.

**[0393]** Purified proteins (SpA, SpA<sub>KKAA</sub>, SpA D and SpA-D<sub>KKAA</sub>) were coated and blocked as described above. Plates were incubated with human vWF at 1  $\mu\text{g ml}^{-1}$  concentration for two hours, then washed and blocked with human IgG for another hour. After washing, plates were incubated with serial dilution of peroxidase-conjugated antibody directed against human vWF for one hour. Plates were washed and developed using OptEIA ELISA reagents (BD). Reactions were quenched with 1 M phosphoric acid and A<sub>450</sub> readings were used to calculate half maximal titer and percent binding. For inhibition assays, plates were incubated with affinity purified F(ab)<sub>2</sub> fragments specific for SpA-D<sub>KKAA</sub> at 10  $\mu\text{g ml}^{-1}$  concentration for one hour prior to ligand binding assays.

**[0394]** Splenocyte Apoptosis.

**[0395]** Affinity purified proteins (150  $\mu\text{g}$  of SpA, SpA-D, SpA<sub>KKAA</sub>, and SpA-D<sub>KKAA</sub>) were injected into the peritoneal cavity of BALB/c mice (6 week old, female, Charles River Laboratories). Four hours following injection, animals were killed by CO<sub>2</sub> inhalation. Their spleens were removed and homogenized. Cell debris were removed using cell strainer and suspended cells were transferred to ACK lysis buffer (0.15 M NH<sub>4</sub>Cl, 10 mM KHCO<sub>3</sub>, 0.1 mM EDTA) to lyse red blood cells. White blood cells were sedimented by centrifugation, suspended in PBS and stained with 1:250 diluted R-PE conjugated anti-CD19 monoclonal antibody (Invitrogen) on ice and in the dark for one hour. Cells were washed with 1% FBS and fixed with 4% formalin overnight at 4° C. The following day, cells were diluted in PBS and analyzed by flow cytometry. The remaining organ was examined for histopathology. Briefly, spleens were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with the Apoptosis detection kit (Millipore), and inspected by light microscopy.

**[0396]** Antibody Quantification.

**[0397]** Sera were collected from healthy human volunteers or BALB/c mice that had been either infected with *S. aureus* Newman or USA300 for 30 days or that had been immunized with SpA-D<sub>KKAA</sub>/SpA<sub>KKAA</sub> as described above. Human/mouse IgG (Jackson Immunology Laboratory), SpA<sub>KKAA</sub>, and CRM<sub>197</sub> were blotted onto nitrocellulose membrane. Membranes were blocked with 5% whole milk, followed by incubation with either human or mouse sera. IRDye 700DX conjugated affinity purified anti-human/mouse IgG (Rockland) was used to quantify signal intensities using the Odyssey™ infrared imaging system (Li-cor). Experiments with blood from human volunteers involved protocols that were reviewed, approved and performed under regulatory supervision of The University of Chicago's Institutional Review Board (IRB).

**[0398]** Statistical Analysis.

**[0399]** Two tailed Student's t tests were performed to analyze the statistical significance of renal abscess, ELISA, and B cell superantigen data.

### Example 3

#### Active Immunization Using Subunit Vaccine Including Multiple Antigens

**[0400]** BALB/c mice (n=18-20) were either mock immunized with PBS/adjuvant or injected with 25  $\mu\text{g}$  of each antigen (Combo 1, ClfA+SdrD+FnBPB; Combo 2, Combo 1+SpA<sub>KKAA</sub>). Immunized mice were challenged by intravenous inoculation with  $1 \times 10^7$  CFU *S. aureus* Newman. Bacterial loads in kidney tissues were examined at day 4 (FIG. 13A) and day 18 (FIG. 13B) post challenge. Statistical significance was calculated with the unpaired two-tailed Student's t-test and P-values recorded; P-values <0.05 were deemed significant. Combo 1 and Combo 2 showed significant reduction in bacterial load at 4 and 18 days post challenge.

**[0401]** Genetic Vaccinology Identifies Protective Antigens of *S. aureus*.

**[0402]** The putative protective antigens identified by genetic vaccinology are sortase A-anchored surface proteins with C-terminal LPXTG sorting signals. Previous work assessed the contribution of surface proteins to disease pathogenesis and vaccine protection in the murine abscess model. Mutations in sdrD or clfA, but not fnbpB or sasF, reduced the staphylococcal load in infected renal tissues. When used as a single subunit vaccine antigen, purified SdrD or ClfA, not SasF or FnBPB, elicited IgG immune responses that conferred significant reduction in staphylococcal load. FnBPB is a homolog of FnBPA (60% sequence identity) and both polypeptides are known to bind fibronectin as well as fibrinogen. The contribution of both surface proteins to disease pathogens and protective immunity has not yet been assessed and this prompted the inclusion of FnBPB into a combination vaccine with ClfA and SdrD (Combo 1). Previous work identified non-toxicogenic protein A (SpA<sub>KKAA</sub>) as a protective antigen, which elicits neutralizing IgG responses for the Fc $\gamma$  and Fab VH3 binding B cell superantigen attributes of SpA. The inventors included SpA<sub>KKAA</sub> to the antigen mixture with ClfA, FnBPB and SdrD (Combo 2).

**[0403]** Immunization of animals with Combo 1 or 2 emulsified in complete Freund adjuvant and boosted with the same antigen mixture emulsified in incomplete Freund adjuvant, raised specific IgG responses. Following intravenous challenge with *S. aureus* Newman, a significant reduction in bacterial load for both vaccines on day four after challenge with the wild-type strain *S. aureus* Newman was observed (FIG. 14; Table 8). To monitor the ability of vaccine formulations to prevent staphylococcal persistence, immunized animals were also analyzed eighteen days after challenge (FIG. 14; Table 8). Again, immunization with either Combo 1 or 2 conferred protection against persistent *S. aureus* Newman infection. Post vaccination antibody titers were also assessed and the results of these analyses are shown in Table 9 below.

TABLE 8

Active immunization with antigen combinations prevents staphylococcal abscess formation					
Staphylococcal load and abscess formation in renal tissue					
Vaccine	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> P-value	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>d</sup> Number of abscesses	<sup>b</sup> P-value
<i>S. aureus</i> Newman challenge at day 4					
Mock	4.56 ± 0.51 (n = 20)	—	—	2.1 ± 0.7 (n = 10)	—
Combo 1	2.74 ± 0.47 (n = 20)	0.0125	1.82	0.4 ± 0.3 (n = 10)	0.0471
Combo 2	1.65 ± 0.59 (n = 20)	0.0005	2.91	0.3 ± 0.3 (n = 10)	0.0363
<i>S. aureus</i> Newman challenge at day 18					
Mock	3.86 ± 0.58 (n = 18)	—	—	1.9 ± 0.8 (n = 10)	—
Combo 1	1.10 ± 0.48 (n = 19)	0.0012	2.76	0.1 ± 0.1 (n = 10)	0.0404
Combo 2	0.26 ± 0.26 (n = 20)	<0.0001	3.60	0.0 ± 0.0 (n = 10)	0.0304

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 or 18 days following infection in cohorts of twenty BALB/c mice per immunization. Combo 1 is composed of affinity-purified, recombinant ClfA, SdrD, and FnBPB. Combo 2 contains one additional antigen, SpA<sub>KKAA</sub>. Representative data of two independent animal experiments are shown. Standard error of the means (±SEM) is indicated.

<sup>b</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

TABLE 9

Humoral immune responses to staphylococcal subunit vaccines					
Antigen specific IgG titer <sup>a</sup>					
Vaccine	ClfA	FnbpB	SdrD	SdrE	SpA <sub>KKAA</sub>
Mock	<100	<100	<100	<100	<100
Combo 1	2975 ± 396	6351 ± 1981	7569 ± 1405	2297 ± 538	<100
Combo 2	3457 ± 887	5539 ± 1292	4716 ± 870	3128 ± 1813	6667 ± 1980

<sup>a</sup>Means (±SEM) of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA using individual antigens.

**[0404]** Vaccine Protection against Staphylococcal Sepsis.

**[0405]** The mortality of *S. aureus* infections increases dramatically when the pathogen replicates in blood or on endocardial tissue. The inventors conducted studies to determine if combo 1 and 2 protect animals against lethal *S. aureus* Newman challenge. All of the mock immunized animals succumbed to challenge within four days (FIG. 15). In contrast, Combo 1 immunized mice displayed either a delayed time to death or survived the lethal challenge (FIG. 15). Mice immunized with Combo 2 displayed a further increase in protective immunity and delayed time-to-death (FIG. 15). Thus, the combination of antibodies against ClfA, FnBPB, SdrD and SpA generates significant protection from staphylococcal abscess formation and lethal challenge.

**[0406]** Bacterial Strains and Culturing Conditions.

**[0407]** Staphylococci were cultured on tryptic soy agar or broth at 37° C. *E. coli* strains DH5α and BL21(DE3) (Studier et al., (1990) Methods Enzymol. 185, 60-89) were cultured on Luria agar or broth at 37° C. Ampicillin (100 μg erythromycin (200 μg ml<sup>-1</sup>) and spectinomycin (200 μg ml<sup>-1</sup>) were used for pET15b (Studier et al., (1990) Methods Enzymol. 185, 60-89), transposon mutant (Bae et al., (2004) Proc. Natl. Acad. Sci. USA 101, 12312-12317) and protein A mutant (Kim et al., J Exp Med 207, 1863-70) selection, respectively.

**[0408]** Mutagenesis.

**[0409]** *Bursa aurealis* mini-transposon insertions from the Phoenix library were transduced into *S. aureus* Newman. The

spa gene on the chromosome of *S. aureus* Newman was deleted by allelic replacement as described previously.

**[0410]** Cloning and Purification.

**[0411]** Coding sequences for ClfA, SdrD, and FnBPB were PCR amplified using *S. aureus* Newman template DNA (Stranger-Jones et al., (2006) Proc. Nat. Acad. Sci. USA 103, 16942-16947). PCR products were cloned into pET15b to express recombinant proteins with N-terminal His<sub>6</sub>-tag fusion. Cloning of non-toxicogenic protein A was described previously (Kim et al., J Exp Med 207, 1863-70). Plasmids were transformed into BL21(DE3). Overnight cultures of transformants were diluted 1:100 into fresh media and grown at 37° C. to an OD<sub>600</sub> 0.5, at which point cultures were induced with 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) and grown for an additional three hours. Bacterial cells were sedimented by centrifugation, suspended in column buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl) and disrupted with a French pressure cell at 14,000 psi. Lysates were cleared of membrane and insoluble components by ultracentrifugation at 40,000×g. Proteins in the soluble lysate were subjected to nickel-nitrilotriacetic acid (Ni-NTA, Qiagen) affinity chromatography. Proteins were eluted in column buffer containing successively higher concentrations of imidazole (100-500 mM). Protein concentrations were determined by bicinchoninic acid (BCA) assay (Thermo Scientific).

**[0412]** Live-Attenuated Vaccine and Renal Abscess Model.

**[0413]** Overnight cultures of *S. aureus* Newman and its isogenic mutants were diluted 1:100 into fresh TSB and grown for 2 hours at 37° C. Staphylococci were sedimented, washed and suspended PBS at OD600 of 0.4 (~1×10<sup>8</sup> CFU ml<sup>-1</sup>). Inocula were quantified by spreading sample aliquots on TSA and enumerating colony formation. BALB/c mice (4 week old, female, Charles River Laboratories) were anesthetized via intraperitoneal injection with 100 mg ml<sup>-1</sup> ketamine and 20 mg ml<sup>-1</sup> xylazine per kilogram of body weight. Mice were infected with 100 µl of bacterial suspension (1×10<sup>7</sup> CFU) by retro-orbital injection. On day 19 following infection, cohorts of mice were treated with antibiotics, a mixture of ampicillin (1 mg ml<sup>-1</sup>) and chloramphenicol (1 mg ml<sup>-1</sup>) in water for 3 days. On day 26, mice were challenged with 100 µl of *S. aureus* Newman (1×10<sup>7</sup> CFU) by retro-orbital injection or bled to analyze adaptive immune response towards components of the antigen matrix. Animals were killed by CO<sub>2</sub> inhalation on day 18 and 30 post initial infection. Both kidneys were removed, and the staphylococcal load in right kidney was analyzed by homogenizing renal tissue with PBS, 0.1% Triton X-100. Serial dilutions of homogenate were spread on TSA or TSA containing antibiotics and incubated for colony formation. The left kidney was examined by histopathology. Briefly, kidneys were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with hematoxylin-eosin, and inspected by light microscopy to enumerate abscess lesions. Also, hyper-immune sera were collected via cardiac puncture and analyzed against components of the antigen matrix. All mouse experiments were performed in accordance with the institutional guidelines following experimental protocol review and approval by the Institutional Biosafety Committee (IBC) and the Institutional Animal Care and Use Committee (IACUC) at the University of Chicago.

**[0414]** Active Immunization.

**[0415]** BALB/c mice (3 week old, female, Charles River Laboratories) were immunized with 25 µg protein emulsified in Complete Freund's Adjuvant (Difco) by intramuscular injection. For booster immunizations, proteins were emulsified in Incomplete Freund's Adjuvant and injected 11 days following the initial immunization. On day 20 following immunization, 5 mice were bled to obtain sera for specific antibody titers by enzyme-linked immunosorbent assay (ELISA). On day 21, all mice were challenged with 1×10<sup>7</sup> CFU *S. aureus* Newman. Four and eighteen days following challenge, kidneys were removed during necropsy, and renal tissue was analyzed for staphylococcal load or histopathology. Also, hyper-immune sera were collected via cardiac puncture and analyzed against components of the staphylococcal antigen matrix.

**[0416]** Antibody Quantification.

**[0417]** For the antigen matrix, nitrocellulose membrane was blotted with 2 µg of a collection of Ni-NTA affinity purified recombinant His6 tagged staphylococcal proteins. Signal intensities in mouse sera were quantified and normalized using anti-His6 antibody with the Odyssey™.

**[0418]** Statistical Analysis.

**[0419]** Unpaired two-tailed Student's t tests were performed to analyze the statistical significance. Linear regression analysis was performed using Graphpad Prism.

## REFERENCES

**[0420]** The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- [0421]** U.S. Pat. No. 3,791,932
- [0422]** U.S. Pat. No. 3,949,064
- [0423]** U.S. Pat. No. 4,174,384
- [0424]** U.S. Pat. No. 4,338,298
- [0425]** U.S. Pat. No. 4,356,170
- [0426]** U.S. Pat. No. 4,367,110
- [0427]** U.S. Pat. No. 4,372,945
- [0428]** U.S. Pat. No. 4,452,901
- [0429]** U.S. Pat. No. 4,474,757
- [0430]** U.S. Pat. No. 4,554,101
- [0431]** U.S. Pat. No. 4,578,770
- [0432]** U.S. Pat. No. 4,596,792
- [0433]** U.S. Pat. No. 4,599,230
- [0434]** U.S. Pat. No. 4,599,231
- [0435]** U.S. Pat. No. 4,601,903
- [0436]** U.S. Pat. No. 4,608,251
- [0437]** U.S. Pat. No. 4,683,195
- [0438]** U.S. Pat. No. 4,683,202
- [0439]** U.S. Pat. No. 4,684,611
- [0440]** U.S. Pat. No. 4,690,915
- [0441]** U.S. Pat. No. 4,690,915
- [0442]** U.S. Pat. No. 4,748,018
- [0443]** U.S. Pat. No. 4,800,159
- [0444]** U.S. Pat. No. 4,879,236
- [0445]** U.S. Pat. No. 4,952,500
- [0446]** U.S. Pat. No. 5,084,269
- [0447]** U.S. Pat. No. 5,199,942
- [0448]** U.S. Pat. No. 5,221,605
- [0449]** U.S. Pat. No. 5,238,808
- [0450]** U.S. Pat. No. 5,302,523
- [0451]** U.S. Pat. No. 5,310,687
- [0452]** U.S. Pat. No. 5,322,783
- [0453]** U.S. Pat. No. 5,384,253
- [0454]** U.S. Pat. No. 5,464,765
- [0455]** U.S. Pat. No. 5,512,282
- [0456]** U.S. Pat. No. 5,512,282
- [0457]** U.S. Pat. No. 5,538,877
- [0458]** U.S. Pat. No. 5,538,880
- [0459]** U.S. Pat. No. 5,548,066
- [0460]** U.S. Pat. No. 5,550,318
- [0461]** U.S. Pat. No. 5,563,055
- [0462]** U.S. Pat. No. 5,580,859
- [0463]** U.S. Pat. No. 5,589,466
- [0464]** U.S. Pat. No. 5,591,616
- [0465]** U.S. Pat. No. 5,610,042
- [0466]** U.S. Pat. No. 5,620,896
- [0467]** U.S. Pat. No. 5,648,240
- [0468]** U.S. Pat. No. 5,656,610
- [0469]** U.S. Pat. No. 5,702,932
- [0470]** U.S. Pat. No. 5,736,524
- [0471]** U.S. Pat. No. 5,780,448
- [0472]** U.S. Pat. No. 5,789,215
- [0473]** U.S. Pat. No. 5,801,234
- [0474]** U.S. Pat. No. 5,840,846
- [0475]** U.S. Pat. No. 5,843,650
- [0476]** U.S. Pat. No. 5,846,709
- [0477]** U.S. Pat. No. 5,846,783
- [0478]** U.S. Pat. No. 5,849,497
- [0479]** U.S. Pat. No. 5,849,546

- [0480] U.S. Pat. No. 5,849,547  
 [0481] U.S. Pat. No. 5,858,652  
 [0482] U.S. Pat. No. 5,866,366  
 [0483] U.S. Pat. No. 5,871,986  
 [0484] U.S. Pat. No. 5,916,776  
 [0485] U.S. Pat. No. 5,922,574  
 [0486] U.S. Pat. No. 5,925,565  
 [0487] U.S. Pat. No. 5,925,565  
 [0488] U.S. Pat. No. 5,928,905  
 [0489] U.S. Pat. No. 5,928,906  
 [0490] U.S. Pat. No. 5,932,451  
 [0491] U.S. Pat. No. 5,935,819  
 [0492] U.S. Pat. No. 5,935,825  
 [0493] U.S. Pat. No. 5,939,291  
 [0494] U.S. Pat. No. 5,942,391  
 [0495] U.S. Pat. No. 5,945,100  
 [0496] U.S. Pat. No. 5,958,895  
 [0497] U.S. Pat. No. 5,981,274  
 [0498] U.S. Pat. No. 5,994,624  
 [0499] U.S. Pat. No. 6,00,8341  
 [0500] U.S. Pat. No. 6,288,214  
 [0501] U.S. Pat. No. 6,294,177  
 [0502] U.S. Pat. No. 6,651,655  
 [0503] U.S. Pat. No. 6,656,462  
 [0504] U.S. Pat. No. 6,733,754  
 [0505] U.S. Pat. No. 6,756,361  
 [0506] U.S. Pat. No. 6,770,278  
 [0507] U.S. Pat. No. 6,793,923  
 [0508] U.S. Pat. No. 6,814,971  
 [0509] U.S. Pat. No. 6,936,258  
 [0510] U.S. Patent Appln. 2002/0169288  
 [0511] U.S. Patent Appln. 2003/0153022  
 [0512] Abdallah et al., *Mol. Microbiol.*, 62, 667-679, 2006.  
 [0513] Abdallah et al., *Nat. Rev. Microbiol.*, 5, 883-891, 2007.  
 [0514] Albus et al., *Infect. Immun.*, 59:1008-1014, 1991.  
 [0515] An, *J. Virol.*, 71(3):2292-302, 1997.  
 [0516] Anavi, Sc. thesis from the department of Molecular Microbiology and Biotechnology of the Tel-Aviv University, Israel, 1998.  
 [0517] Andersen et al., *J. Immunol.*, 154, 3359-3372, 1995.  
 [0518] Angel et al., *Cell*, 49:729, 1987b.  
 [0519] Angel et al., *Mol. Cell. Biol.*, 7:2256, 1987a.  
 [0520] Archer, *Clin. Infect. Dis.*, 26, 1179-1181, 1998.  
 [0521] Atchison and Perry, *Cell*, 46:253, 1986.  
 [0522] Atchison and Perry, *Cell*, 48:121, 1987.  
 [0523] Ausubel et al., In: *Current Protocols in Molecular Biology*, John, Wiley & Sons, Inc, New York, 1996.  
 [0524] Baba et al., *J. Bacteriol.* 190:300-310, 2007.  
 [0525] Bae and Schneewind, *Plasmid*, 55:58-63, 2006.  
 [0526] Bae et al., *Proc. Natl. Acad. Sci. USA*, 101, 12312-12317, 2004.  
 [0527] Banerji et al., *Cell*, 27(2 Pt 1):299-308, 1981.  
 [0528] Banerji et al., *Cell*, 33(3):729-740, 1983.  
 [0529] Barany and Merrifield, In: *The Peptides*, Gross and Meienhofer (Eds.), Academic Press, NY, 1-284, 1979.  
 [0530] Behring EA. Über das Zustandekommen der Diphtherie—Immunität bei Thieren. *Deutsche Medizinische Wochenschrift*, 16:1145-8, 1890.  
 [0531] Bellus, *J. Macromol. Sci. Pure Appl. Chem.*, A31(1): 1355-1376, 1994.  
 [0532] Berkhout et al., *Cell*, 59:273-282, 1989.  
 [0533] Birch-Hirschfeld, L. 1934. Über die Agglutination von Staphylokokken durch Bestandteile des Säugetierblutplasmas. *Klinische Wochenschrift* 13:331.  
 [0534] Bjerketorp et al., *FEMS Microbiol. Lett.*, 234:309-314, 2004.  
 [0535] Blonar et al., *EMBO J.*, 8:1139, 1989.  
 [0536] Bodine and Ley, *EMBO J.* 6:2997, 1987.  
 [0537] Borrebaeck, In: *Antibody Engineering—A Practical Guide*, W.H. Freeman and Co., 1992.  
 [0538] Boshart et al., *Cell*, 41:521, 1985.  
 [0539] Bosze et al., *EMBO* 1, 5(7):1615-1623, 1986.  
 [0540] Boucher and Corey. *Clin. Infect. Dis.* 46:S334-S349, 2008.  
 [0541] Braddock et al., *Cell*, 58:269, 1989.  
 [0542] Brown et al., *Biochemistry*, 37:4397-4406, 1998.  
 [0543] Bubeck Wardenburg and Schneewind. *J. Exp. Med.* 205:287-294, 2008.  
 [0544] Bubeck-Wardenburg et al., *Infect. Immun.* 74:1040-1044, 2007.  
 [0545] Bubeck-Wardenburg et al., *Proc. Natl. Acad. Sci. USA*, 103:13831-13836, 2006.  
 [0546] Bulla and Siddiqui, *J. Virol.*, 62:1437, 1986.  
 [0547] Burke et al., *J. Inf. Dis.*, 170:1110-1119, 1994.  
 [0548] Burlak et al., *Cell Microbiol.*, 9:1172-1190, 2007.  
 [0549] Burts and Missiakas, *Mol. Microbiol.*, 69:736-46, 2008.  
 [0550] Burts et al., *Proc. Natl. Acad. Sci. USA*, 102:1169-1174, 2005.  
 [0551] Campbell and Villarreal, *Mol. Cell. Biol.*, 8:1993, 1988.  
 [0552] Campere and Tilghman, *Genes and Dev.*, 3:537, 1989.  
 [0553] Campo et al., *Nature*, 303:77, 1983.  
 [0554] Carbonelli et al., *FEMS Microbiol. Lett.*, 177(1):75-82, 1999.  
 [0555] Cedergren et al., *Protein Eng.*, 6:441-448, 1993.  
 [0556] Celandier and Haseltine, *J. Virology*, 61:269, 1987.  
 [0557] Celandier et al., *J. Virology*, 62:1314, 1988.  
 [0558] Cespedes et al., *J. Infect. Dis.*, 191(3):444-52, 2005.  
 [0559] Champion et al., *Science*, 313:1632-1636, 2006.  
 [0560] Chandler et al., *Cell*, 33:489, 1983.  
 [0561] Chandler et al., *Proc. Natl. Acad. Sci. USA*, 94(8): 3596-601, 1997.  
 [0562] Chang et al., *Lancet.*, 362(9381):362-369, 2003.  
 [0563] Chang et al., *Mol. Cell. Biol.*, 9:2153, 1989.  
 [0564] Chatterjee et al., *Proc. Natl. Acad. Sci. USA*, 86:9114, 1989.  
 [0565] Chen and Okayama, *Mol. Cell. Biol.*, 7(8):2745-2752, 1987.  
 [0566] Cheng et al., *FASEB J.*, 23:1-12, 2009.  
 [0567] Cheng et al., *PLoS Pathog* 6.  
 [0568] Choi et al., *Cell*, 53:519, 1988.  
 [0569] Cocea, *Biotechniques*, 23(5):814-816, 1997.  
 [0570] Cohen et al., *J. Cell. Physiol.*, 5:75, 1987.  
 [0571] Cosgrove et al., *Infect. Control Hosp. Epidemiol.* 26:166-174, 2005.  
 [0572] Costa et al., *Mol. Cell. Biol.*, 8:81, 1988.  
 [0573] Cripe et al., *EMBO J.*, 6:3745, 1987.  
 [0574] Culotta and Hamer, *Mol. Cell. Biol.*, 9:1376, 1989.  
 [0575] Dalbey and Wickner, *J. Biol. Chem.*, 260:15925-15931, 1985.  
 [0576] Dandolo et al., *J. Virology*, 47:55-64, 1983.  
 [0577] De Villiers et al., *Nature*, 312(5991):242-246, 1984.  
 [0578] DeBord et al., *Infect. Immun.*, 74:4910-4914, 2006.

- [0579] DeDent et al., *EMBO J.* 27:2656-2668, 2008.
- [0580] DeDent et al., *J. Bacteriol.* 189:4473-4484, 2007.
- [0581] Deisenhofer et al., *Hoppe-Seyh Zeitsch. Physiol. Chem.* 359:975-985, 1978.
- [0582] Deisenhofer, *Biochemistry* 20:2361-2370, 1981.
- [0583] Deschamps et al., *Science*, 230:1174-1177, 1985.
- [0584] Devereux et al., *Nucl. Acid Res.*, 12:387-395, 1984.
- [0585] Diep et al., *J. Infect. Dis.*, 193:1495-1503, 2006a.
- [0586] Diep et al., *Lancet.*, 367:731-739, 2006b.
- [0587] Dinges et al., *Clin. Microbiol. Rev.*, 13:16-34, 2000.
- [0588] Duthie and Lorenz, *J. Gen. Microbiol.*, 6:95-107, 1952.
- [0589] Edbrooke et al., *Mol. Cell. Biol.*, 9:1908, 1989.
- [0590] Edlund et al., *Science*, 230:912-916, 1985.
- [0591] Ekstedt and Yotis, *Ann. N.Y. Acad. Sci.*, 80:496-500, 1960.
- [0592] Emorl and Gaynes, *Clin. Microbiol. Rev.*, 6:428-442, 1993.
- [0593] EP 0786519
- [0594] EP 497524
- [0595] EP 497525
- [0596] Epitope Mapping Protocols In: *Methods in Molecular Biology*, Vol. 66, Morris (Ed.), 1996.
- [0597] Fehheimer, et al., *Proc Natl. Acad. Sci. USA*, 84:8463-8467, 1987.
- [0598] Feng and Holland, *Nature*, 334:6178, 1988.
- [0599] Field and Smith, *J. Comp. Pathol.*, 55:63, 1945.
- [0600] Firak and Subramanian, *Mol. Cell. Biol.*, 6:3667, 1986.
- [0601] Foecking and Hofstetter, *Gene*, 45(1):101-105, 1986.
- [0602] Fortune et al., *Proc Natl. Acad. Sci. USA*, 102:10676-10681, 2005.
- [0603] Foster, *Nat. Rev. Microbiol.*, 3:948-958, 2005.
- [0604] Fournier et al., *Infect. Immun.*, 45:87-93, 1984.
- [0605] Fraley et al., *Proc. Natl. Acad. Sci. USA*, 76:3348-3352, 1979.
- [0606] Friedrich et al., *Nature*, 425:535-539, 2003.
- [0607] Fujita et al., *Cell*, 49:357, 1987.
- [0608] GB Appln. 2 202 328
- [0609] Gilles et al., *Cell*, 33:717, 1983.
- [0610] Gloss et al., *EMBO J.*, 6:3735, 1987.
- [0611] Godbout et al., *Mol. Cell. Biol.*, 8:1169, 1988.
- [0612] Gomez et al., *EMBO J.* 26:701-709, 2007.
- [0613] Gomez et al., *J. Biol. Chem.* 281:20190-20196, 2006.
- [0614] Gomez et al., *Nature Med.* 10:842-8, 2004.
- [0615] Goodbourn and Maniatis, *Proc. Natl. Acad. Sci. USA*, 85:1447, 1988.
- [0616] Goodbourn et al., *Cell*, 45:601, 1986.
- [0617] Goodyear and Silverman, *J. Exp. Med.*, 197:1125-1139, 2003.
- [0618] Goodyear and Silverman, *Proc. Nat. Acad. Sci. USA*, 101:11392-11397, 2004.
- [0619] Gopal, *Mol. Cell. Biol.*, 5:1188-1190, 1985.
- [0620] Gouda et al., *Biochemistry*, 31(40):9665-72, 1992.
- [0621] Gouda et al., *Biochemistry*, 37:129-36, 1998. Graham and Van Der Eb, *Virology*, 52:456-467, 1973.
- [0622] Graille et al., *Proc. Nat. Acad. Sci. USA* 97:5399-5404, 2000.
- [0623] Greene et al., *Immunology Today*, 10:272, 1989
- [0624] Grosschedl and Baltimore, *Cell*, 41:885, 1985.
- [0625] Guinn et al., *Mol. Microbiol.*, 51:359-370, 2004.
- [0626] Guss et al., *Eur. J. Biochem.* 138:413-420, 1984.
- [0627] Harland and Weintraub, *J. Cell Biol.*, 101(3):1094-1099, 1985.
- [0628] Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., Chapter 8, 1988.
- [0629] Hartleib et al., *Blood* 96:2149-2156, 2000.
- [0630] Harvey et al., *Proc. Natl. Acad. Sci. USA*, 83:1084-1088, 1986.
- [0631] Haslinger and Karin, *Proc. Natl. Acad. Sci. USA*, 82:8572, 1985.
- [0632] Hauber and Cullen, *J. Virology*, 62:673, 1988.
- [0633] Hen et al., *Nature*, 321:249, 1986.
- [0634] Hensel et al., *Lymphokine Res.*, 8:347, 1989.
- [0635] Herr and Clarke, *Cell*, 45:461, 1986.
- [0636] Hirochika et al., *J. Virol.*, 61:2599, 1987.
- [0637] Hirsch et al., *Mol. Cell. Biol.*, 10:1959, 1990.
- [0638] Holbrook et al., *Virology*, 157:211, 1987.
- [0639] Horlick and Benfield, *Mol. Cell. Biol.*, 9:2396, 1989.
- [0640] Hsu et al., *Proc. Natl. Acad. Sci. USA*, 100:12420-12425, 2003.
- [0641] Huang et al., *Cell*, 27:245, 1981.
- [0642] Hug et al., *Mol. Cell. Biol.*, 8:3065, 1988.
- [0643] Huston et al., In: *Methods in Enzymology*, Langone (Ed.), Academic Press, NY, 203:46-88, 1991.
- [0644] Hwang et al., *Mol. Cell. Biol.*, 10:585, 1990.
- [0645] Imagawa et al., *Cell*, 51:251, 1987.
- [0646] Imbra and Karin, *Nature*, 323:555, 1986.
- [0647] Imler et al., *Mol. Cell. Biol.*, 7:2558, 1987.
- [0648] Imperiale and Nevins, *Mol. Cell. Biol.*, 4:875, 1984.
- [0649] Innis et al., *Proc Natl Acad Sci USA*, 85(24):9436-9440, 1988.
- [0650] Inouye and Inouye, *Nucleic Acids Res.*, 13: 3101-3109, 1985.
- [0651] Jakobovits et al., *Mol. Cell. Biol.*, 8:2555, 1988.
- [0652] Jameel and Siddiqui, *Mol. Cell. Biol.*, 6:710, 1986.
- [0653] Jansson et al., *FEMS Immunol. Med. Microbiol.* 20:69-78 1998.
- [0654] Jaynes et al., *Mol. Cell. Biol.*, 8:62, 1988.
- [0655] Jensen, *Acta Path. Microbiol. Scand.* 44:421-428, 1958.
- [0656] Johnson et al., *Methods in Enzymol.*, 203:88-99, 1991.
- [0657] Johnson et al., *Mol. Cell. Biol.*, 9:3393, 1989.
- [0658] Jones, *Carb. Research*, 340:1097-1106, 2005.
- [0659] Jonsson et al., *Oral Dis.*, 8(3):130-140, 2002.
- [0660] Joyce et al., *Carbohydrate Research* 338:903-922 (2003)
- [0661] Kadesch and Berg, *Mol. Cell. Biol.*, 6:2593, 1986.
- [0662] Kaeppler et al., *Plant Cell Rep.*, 8:415-418, 1990.
- [0663] Kaneda et al., *Science*, 243:375-378, 1989.
- [0664] Karin et al., *Mol. Cell. Biol.*, 7:606, 1987.
- [0665] Katinka et al., *Cell*, 20:393, 1980.
- [0666] Kato et al., *J. Biol. Chem.*, 266:3361-3364, 1991.
- [0667] Kawamoto et al., *Mol. Cell. Biol.*, 8:267, 1988.
- [0668] Kennedy et al., *Proc. Natl. Acad. Sci. USA* 105:1327-1332, 2008.
- [0669] Kiledjian et al., *Mol. Cell. Biol.*, 8:145, 1988.
- [0670] Kim et al., *J Exp Med* 207, 1863-70, 2010.
- [0671] Kinoshita, M., N. Kobayashi, S. Nagashima, M. Ishino, S. Otokozawa, K. Mise, A. Sumi, H. Tsutsumi, N. Uehara, N. Watanabe, and M. Endo. 2008. Diversity of

- staphylocoagulase and identification of novel variants of staphylocoagulase gene in *Staphylococcus aureus*. *Microbiol. Immunols.* 52:334-348.
- [0672] Klamut et al., *Mol. Cell. Biol.*, 10:193, 1990.
- [0673] Klevens et al., *Clin. Infect. Dis.*, 2008; 47:927-30, 2008.
- [0674] Klevens et al., *JAMA*, 298:1763-1771, 2007.
- [0675] Koch et al., *Mol. Cell. Biol.*, 9:303, 1989.
- [0676] Kohler and Milstein, *Nature* 256:495-497 (1975)
- [0677] Kriegler and Botchan, In: *Eukaryotic Viral Vectors*, Gluzman (Ed.), Cold Spring Harbor: Cold Spring Harbor Laboratory, NY, 1982.
- [0678] Kriegler and Botchan, *Mol. Cell. Biol.*, 3:325, 1983.
- [0679] Kriegler et al., *Cell*, 38:483, 1984a.
- [0680] Kriegler et al., *Cell*, 53:45, 1988.
- [0681] Kriegler et al., In: *Cancer Cells 2/Oncogenes and Viral Genes*, Van de Woude et al. eds, Cold Spring Harbor, Cold Spring Harbor Laboratory, 1984b.
- [0682] Kroh et al., *Proc. Natl. Acad. Sci. USA*, 106:7786-7791, 2009.
- [0683] Kuhl et al., *Cell*, 50:1057, 1987.
- [0684] Kuklin et al., *Infect. Immun.*, 74:2215-23, 2006.
- [0685] Kunz et al., *Nucl. Acids Res.*, 17:1121, 1989.
- [0686] Kuroda et al., *Lancet.*, 357:1225-1240, 2001.
- [0687] Kyte and Doolittle, *J. Mol. Biol.*, 157(1):105-132, 1982.
- [0688] Lagergard et al., *Eur. J. Clin. Microbiol. Infect. Dis.*, 11:341-5, 1992.
- [0689] Lam et al., *J. Bacteriol.*, 86:87-91, 1963.
- [0690] Larsen et al., *Proc Natl. Acad. Sci. USA.*, 83:8283, 1986, 1963.
- [0691] Laspia et al., *Cell*, 59:283, 1989.
- [0692] Latimer et al., *Mol. Cell. Biol.*, 10:760, 1990.
- [0693] Lee et al., *Nature*, 294:228, 1981.
- [0694] Lee et al., *Nucleic Acids Res.*, 12:4191-206, 1984.
- [0695] Lee, *Trends Microbiol.* 4(4):162-166, 1996.
- [0696] Levenson et al., *Hum. Gene Ther.*, 9(8):1233-1236, 1998.
- [0697] Levinson et al., *Nature*, 295:79, 1982.
- [0698] Lin et al., *Mol. Cell. Biol.*, 10:850, 1990.
- [0699] Lowy, *New Engl. J. Med.*, 339:520-532, 1998.
- [0700] Luria et al., *EMBO J.*, 6:3307, 1987.
- [0701] Lusky and Botchan, *Proc. Natl. Acad. Sci. USA*, 83:3609, 1986.
- [0702] Lusky et al., *Mol. Cell. Biol.*, 3:1108, 1983.
- [0703] Macejak and Sarnow, *Nature*, 353:90-94, 1991.
- [0704] MacGum et al., *Mol. Microbiol.*, 57:1653-1663, 2005.
- [0705] Maira-Litran et al., *Infect. Immun.*, 70:4433-4440, 2002.
- [0706] Maira-Litran et al., *Vaccine*, 22:872-879, 2004.
- [0707] Majors and Varmus, *Proc. Natl. Acad. Sci. USA*, 80:5866, 1983.
- [0708] Markwardt, *Untersuchungen über Hirudin. Naturwissenschaften*, 41:537-538, 1955.
- [0709] Mazmanian et al., *Mol. Microbiol.* 40, 1049-1057, 2001.
- [0710] Mazmanian et al., *Mol. Microbiol.*, 40(5):1049-1057, 2001.
- [0711] Mazmanian et al., *Proc. Natl. Acad. Sci. USA*, 97:5510-5515, 2000.
- [0712] Mazmanian et al., *Science*, 285(5428):760-3, 1999.
- [0713] McLaughlin et al., *PLoS Pathog.*, 3:e105, 2007.
- [0714] McNeill et al., *Gene*, 76:81, 1989.
- [0715] Memaugh et al., In: *Molecular Methods in Plant Pathology*, Singh et al. (Eds.), CRC Press Inc., Boca Raton, Fla., 359-365, 1995.
- [0716] Merrifield, *Science*, 232(4748):341-347, 1986.
- [0717] Miksicek et al., *Cell*, 46:203, 1986.
- [0718] Mordacq and Linzer, *Genes and Dev.*, 3:760, 1989.
- [0719] Moreau et al., *Carbohydrate Res.*, 201:285-297, 1990.
- [0720] Moreau et al., *Nucl. Acids Res.*, 9:6047, 1981.
- [0721] Moreillon et al., *Infect. Immun.*, 63:4738-4743, 1995.
- [0722] Moreillon et al., *Infect. Immun.*, 63:4738-4743, 1995.
- [0723] Mosmann and Coffman, *Ann. Rev. Immunol.*, 7:145-173, 1989.
- [0724] Muesing et al., *Cell*, 48:691, 1987.
- [0725] Musher et al., *Medicine (Baltimore)*, 73:186-208, 1994.
- [0726] Navarre and Schneewind, *J. Biol. Chem.*, 274:15847-15856, 1999.
- [0727] Needleman & Wunsch, *J. Mol. Biol.*, 48:443, 1970.
- [0728] Ng et al., *Nuc. Acids Res.*, 17:601, 1989.
- [0729] Nicolau and Sene, *Biochim. Biophys. Acta*, 721:185-190, 1982.
- [0730] Nicolau et al., *Methods Enzymol.*, 149:157-176, 1987.
- [0731] Novick, *Mol. Microbiol.*, 48:1429-1449, 2003.
- [0732] O'Brien et al., *Mol. Microbiol.* 44:1033-1044, 2002.
- [0733] O'Seaghda et al., *FEBS J.* 273:4831-4841, 2006.
- [0734] Omirulleh et al., *Plant Mol. Biol.*, 21(3):415-28, 1993.
- [0735] Ondek et al., *EMBO J.*, 6:1017, 1987.
- [0736] Ornitz et al., *Mol. Cell. Biol.*, 7:3466, 1987.
- [0737] Pallen, *Trends Microbiol.*, 10:209-212, 2002.
- [0738] Palmiter et al., *Nature*, 300:611, 1982.
- [0739] Palmqvist et al., *Microbes. Infect.*, 7:1501-11, 2005.
- [0740] Panizzi et al., *J. Biol. Chem.*, 281:1179-1187, 2006.
- [0741] PCT Appln. PCT/US89/01025
- [0742] PCT Appln. WO 00/02523
- [0743] PCT Appln. WO 00/12132
- [0744] PCT Appln. WO 00/12689
- [0745] PCT Appln. WO 00/15238
- [0746] PCT Appln. WO 01/34809
- [0747] PCT Appln. WO 01/60852
- [0748] PCT Appln. WO 01/98499
- [0749] PCT Appln. WO 02/059148
- [0750] PCT Appln. WO 02/094868
- [0751] PCT Appln. WO 03/53462
- [0752] PCT Appln. WO 04/43407
- [0753] PCT Appln. WO 06/032472
- [0754] PCT Appln. WO 06/032475
- [0755] PCT Appln. WO 06/032500
- [0756] PCT Appln. WO 07/113,222
- [0757] PCT Appln. WO 07/113,223
- [0758] PCT Appln. WO 10/119,343
- [0759] PCT Appln. WO 94/09699
- [0760] PCT Appln. WO 95/06128
- [0761] PCT Appln. WO 95/08348
- [0762] PCT Appln. WO 98/57994
- [0763] Pearson & Lipman, *Proc. Natl. Acad. Sci. USA*, 85:2444, 1988.
- [0764] Pech et al., *Mol. Cell. Biol.*, 9:396, 1989.
- [0765] Pelletier and Sonenberg, *Nature*, 334(6180):320-325, 1988.

- [0766] Perez-Stable and Constantini, *Mol. Cell. Biol.*, 10:1116, 1990.
- [0767] Phonimdaeng et al., *Mol. Microbiol.*, 4:393-404, 1990.
- [0768] Picard and Schaffner, *Nature*, 307:83, 1984.
- [0769] Pinkert et al., *Genes and Dev.*, 1:268, 1987.
- [0770] Ponta et al., *Proc. Natl. Acad. Sci. USA*, 82:1020, 1985.
- [0771] Porton et al., *Mol. Cell. Biol.*, 10:1076, 1990.
- [0772] Potrykus et al., *Mol. Gen. Genet.*, 199(2):169-177, 1985.
- [0773] Pugsley, *Microbiol. Rev.*, 57:50-108, 1993.
- [0774] Pym et al., *Mol. Microbiol.*, 46:709-717, 2002.
- [0775] Pym et al., *Nat. Med.*, 9:533-539, 2003.
- [0776] Queen and Baltimore, *Cell*, 35:741, 1983.
- [0777] Quinn et al., *Mol. Cell. Biol.*, 9:4713, 1989.
- [0778] Redondo et al., *Science*, 247:1225, 1990.
- [0779] Reisman and Rotter, *Mol. Cell. Biol.*, 9:3571, 1989.
- [0780] Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1289-1329, 1990.
- [0781] Resendez Jr. et al., *Mol. Cell. Biol.*, 8:4579, 1988.
- [0782] Ripe et al., *Mol. Cell. Biol.*, 9:2224, 1989.
- [0783] Rippe, et al., *Mol. Cell. Biol.*, 10:689-695, 1990.
- [0784] Riffing et al., *Nuc. Acids Res.*, 17:1619, 1989.
- [0785] Roben et al., *J. Immunol.* 154:6437-6445, 1995.
- [0786] Rosen et al., *Cell*, 41:813, 1988.
- [0787] Sakai et al., *Genes and Dev.*, 2:1144, 1988.
- [0788] Salid-Salim et al., *Infect. Control Hosp. Epidemiol.* 24:451-455, 2003.
- [0789] Sambrook et al., *In: Molecular cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001.
- [0790] Schaffner et al., *J. Mol. Biol.*, 201:81, 1988.
- [0791] Schneewind et al., *Cell* 70:267-281, 1992.
- [0792] Schneewind et al., *EMBO*, 12:4803-4811, 1993.
- [0793] Schneewind et al., *Science*, 268:103-6, 1995.
- [0794] Searle et al., *Mol. Cell. Biol.*, 5:1480, 1985.
- [0795] Sharp and Marciniak, *Cell*, 59:229, 1989.
- [0796] Shaul and Ben-Levy, *EMBO J.*, 6:1913, 1987.
- [0797] Shaw et al., *Microbiology*, 150:217-228, 2004.
- [0798] Sheagren, *N. Engl. J. Med.* 310:1368-1373, 1984.
- [0799] Sherman et al., *Mol. Cell. Biol.*, 9:50, 1989.
- [0800] Shopsin et al., *J. Clin. Microbiol.*, 37:3556-63, 1999.
- [0801] Sibbald et al., *Microbiol. Mol. Biol. Rev.*, 70:755-788, 2006.
- [0802] Silverman and Goodyear. *Nat. Rev. Immunol.*, 6:465-75, 2006.
- [0803] Sjudahl, *Eur. J. Biochem.* 73:343-351, 1977.
- [0804] Sjoquist et al., *Eur. J. Biochem.* 30:190-194, 1972.
- [0805] Sleight and Lockett, *J. EMBO*, 4:3831, 1985.
- [0806] Smith & Waterman, *Adv. Appl. Math.*, 2:482, 1981.
- [0807] Smith et al., *Brit. J. Exp. Pathol.*, 28:57, 1947.
- [0808] Sorensen et al., *Infect. Immun.*, 63:1710-1717, 1995.
- [0809] Spalholz et al., *Cell*, 42:183, 1985.
- [0810] Spandau and Lee, *J. Virology*, 62:427, 1988.
- [0811] Spandidos and Wilkie, *EMBO J.*, 2:1193, 1983.
- [0812] Stanley et al., *Proc. Natl. Acad. Sci. USA*, 100:13001-13006, 2003.
- [0813] Stephens and Hentschel, *Biochem. J.*, 248:1, 1987.
- [0814] Stewart and Young, In: *Solid Phase Peptide Synthesis*, 2d. ed., Pierce Chemical Co., 1984.
- [0815] Stranger-Jones et al., *Proc. Nat. Acad. Sci. USA* 103, 16942-16947, 2006.
- [0816] Stranger-Jones et al., *Proc. Nat. Acad. Sci. USA*, 103:16942-16947, 2006.
- [0817] Stuart et al., *Nature*, 317:828, 1985.
- [0818] Studier et al., *Methods Enzymol.* 185, 60-89, 1990.
- [0819] Studier et al., *Methods Enzymol.* 185:60-89 1990.
- [0820] Sullivan and Peterlin, *Mol. Cell. Biol.*, 7:3315, 1987.
- [0821] Swartzendruber and Lehman, *J. Cell. Physiology*, 85:179, 1975.
- [0822] Takebe et al., *Mol. Cell. Biol.*, 8:466, 1988.
- [0823] Tam et al., *J. Am. Chem. Soc.*, 105:6442, 1983.
- [0824] Tavernier et al., *Nature*, 301:634, 1983.
- [0825] Taylor and Kingston, *Mol. Cell. Biol.*, 10:165, 1990a.
- [0826] Taylor and Kingston, *Mol. Cell. Biol.*, 10:176, 1990b.
- [0827] Taylor et al., *J. Biol. Chem.*, 264:15160, 1989.
- [0828] Thiesen et al., *J. Virology*, 62:614, 1988.
- [0829] Thomson et al., *J. Immunol.*, 157(2):822-826, 1996.
- [0830] Tigges et al., *J. Immunol.*, 156(10):3901-3910, 1996.
- [0831] Tigges et al., *J. Immunol.*, 156(10):3901-3910, 1996.
- [0832] Ton-That et al., *Proc. Natl. Acad. Sci. USA*, 96(22):12424-9, 1999.
- [0833] Treisman, *Cell*, 42:889, 1985.
- [0834] Tronche et al., *Mol. Biol. Med.*, 7:173, 1990.
- [0835] Trudel and Constantini, *Genes and Dev.*, 6:954, 1987.
- [0836] Tyndell et al., *Nuc. Acids Res.*, 9:6231, 1981.
- [0837] Uhlen et al., *J. Biol. Chem.* 259:1695-1702 and 13628 (Corr.) 1984.
- [0838] van den Ent and Lowe, *FEBS Lett.*, 579:3837-3841, 2005.
- [0839] van Wely et al., *FEMS Microbiol. Rev.*, 25:437-454, 2001.
- [0840] Vannice and Levinson, *J. Virology*, 62:1305, 1988.
- [0841] Vasseur et al., *Proc Natl. Acad. Sci. USA*, 77:1068, 1980.
- [0842] Vaughan, et al., *Nat. Biotech.* 16; 535-539, 1998.
- [0843] Wang and Calame, *Cell*, 47:241, 1986.
- [0844] Weber et al., *Cell*, 36:983, 1984.
- [0845] Weinberger et al. *Mol. Cell. Biol.*, 8:988, 1984.
- [0846] Weiss et al., *J. Antimicrob. Chemother.*, 53(3):480-6, 2004.
- [0847] Winoto and Baltimore, *Cell*, 59:649, 1989.
- [0848] Wong et al., *Gene*, 10:87-94, 1980.
- [0849] Xu et al., *J. Infect. Dis.*, 189:2323-2333, 2004.
- [0850] Xu et al., *Mol. Microbiol.*, 66(3):787-800, 2007.
- [0851] Yutzey et al. *Mol. Cell. Biol.*, 9:1397, 1989.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 67

<210> SEQ ID NO 1

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 1

```

ttcaacaaag atcaacaaag cgccttctat gaaatcttga acatgcctaa cttaaacgaa      60
gcgcaacgta acgggttcat tcaaagtctt aaagacgacc caagccaaag cactaatggt      120
ttaggtgaag ctaaaaaatt aaacgaatct                                     150

```

<210> SEQ ID NO 2

<211> LENGTH: 54

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 2

```

Gln Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile
 1          5          10          15
Leu Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln
          20          25          30
Ser Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala
          35          40          45
Lys Lys Leu Asn Glu Ser
          50

```

<210> SEQ ID NO 3

<211> LENGTH: 51

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 3

```

Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr Gln Val Leu Asn Met
 1          5          10          15
Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys
          20          25          30
Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys Leu
          35          40          45
Asn Asp Ser
          50

```

<210> SEQ ID NO 4

<211> LENGTH: 52

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 4

```

Asn Asn Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn
 1          5          10          15
Met Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
          20          25          30
Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys
          35          40          45
Leu Asn Glu Ser
          50

```



---

-continued

---

<210> SEQ ID NO 5  
<211> LENGTH: 52  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 5

Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His  
1 5 10 15  
Leu Pro Asn Leu Thr Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu  
20 25 30  
Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys  
35 40 45  
Leu Asn Asp Ala  
50

<210> SEQ ID NO 6  
<211> LENGTH: 52  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 6

Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His  
1 5 10 15  
Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu  
20 25 30  
Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys  
35 40 45  
Leu Asn Asp Ala  
50

<210> SEQ ID NO 7  
<211> LENGTH: 52  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (7)..(8)  
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (34)..(35)  
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 7

Asn Asn Phe Asn Lys Asp Xaa Xaa Ser Ala Phe Tyr Glu Ile Leu Asn  
1 5 10 15  
Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu  
20 25 30  
Lys Xaa Xaa Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys  
35 40 45  
Leu Asn Glu Ser  
50

<210> SEQ ID NO 8  
<211> LENGTH: 52  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.  
<220> FEATURE:  
<221> NAME/KEY: MISC\_FEATURE  
<222> LOCATION: (7)..(8)

-continued

<223> OTHER INFORMATION: where X is any amino acid other than Q  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (12)..(35)  
 <223> OTHER INFORMATION: where Y is any amion acid other than D

<400> SEQUENCE: 8

Asn Asn Phe Asn Lys Asp Xaa Xaa Ser Ala Phe Tyr Glu Ile Leu Asn  
 1 5 10 15  
 Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu  
 20 25 30  
 Lys Tyr Tyr Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys  
 35 40 45  
 Leu Asn Glu Ser  
 50

<210> SEQ ID NO 9  
 <211> LENGTH: 450  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 9

Met Lys Lys Lys Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile  
 1 5 10 15  
 Ala Ser Val Thr Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro  
 20 25 30  
 Ala Ala Asn Ala Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr  
 35 40 45  
 Gln Val Leu Asn Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe  
 50 55 60  
 Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly  
 65 70 75 80  
 Glu Ala Gln Lys Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln  
 85 90 95  
 Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu  
 100 105 110  
 Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser  
 115 120 125  
 Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys  
 130 135 140  
 Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys  
 145 150 155 160  
 Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn  
 165 170 175  
 Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser  
 180 185 190  
 Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln  
 195 200 205  
 Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe  
 210 215 220  
 Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly  
 225 230 235 240  
 Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu  
 245 250 255  
 Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Glu Glu Asp



-continued

165					170					175					
Glu	Glu	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser
			180					185					190		
Gln	Ser	Ala	Asn	Leu	Leu	Ser	Glu	Ala	Lys	Lys	Leu	Asn	Glu	Ser	Gln
		195					200					205			
Ala	Pro	Lys	Ala	Asp	Asn	Lys	Phe	Asn	Lys	Glu	Gln	Gln	Asn	Ala	Phe
		210					215					220			
Tyr	Glu	Ile	Leu	His	Leu	Pro	Asn	Leu	Asn	Glu	Glu	Gln	Arg	Asn	Gly
		225					230					235			240
Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser	Val	Ser	Lys	Glu	Ile	Leu
				245					250					255	
Ala	Glu	Ala	Lys	Lys	Leu	Asn	Asp	Ala	Gln	Ala	Pro	Lys	Glu	Glu	Asp
			260					265					270		
Asn	Lys	Lys	Pro	Gly	Lys	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp
		275					280					285			
Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Lys	Lys	Pro	Gly	Lys	Glu	Asp
		290					295					300			
Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp
		305					310					315			320
Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp
			325						330					335	
Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp
			340					345						350	
Gly	Asn	Gly	Val	His	Val	Val	Lys	Pro	Gly	Asp	Thr	Val	Asn	Asp	Ile
		355					360					365			
Ala	Lys	Ala	Asn	Gly	Thr	Thr	Ala	Asp	Lys	Ile	Ala	Ala	Asp	Asn	Lys
		370					375					380			
Leu	Ala	Asp	Lys	Asn	Met	Ile	Lys	Pro	Gly	Gln	Glu	Leu	Val	Val	Asp
		385					390					395			400
Lys	Lys	Gln	Pro	Ala	Asn	His	Ala	Asp	Ala	Asn	Lys	Ala	Gln	Ala	Leu
				405					410					415	
Pro	Glu	Thr	Gly	Glu	Glu	Asn	Pro	Phe	Ile	Gly	Thr	Thr	Val	Phe	Gly
			420					425					430		
Gly	Leu	Ser	Leu	Ala	Leu	Gly	Ala	Ala	Leu	Leu	Ala	Gly	Arg	Arg	Arg
		435					440					445			
Glu	Leu														
	450														

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 97

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 11

Met	Ala	Met	Ile	Lys	Met	Ser	Pro	Glu	Glu	Ile	Arg	Ala	Lys	Ser	Gln
1				5					10					15	
Ser	Tyr	Gly	Gln	Gly	Ser	Asp	Gln	Ile	Arg	Gln	Ile	Leu	Ser	Asp	Leu
		20					25						30		
Thr	Arg	Ala	Gln	Gly	Glu	Ile	Ala	Ala	Asn	Trp	Glu	Gly	Gln	Ala	Phe
		35					40						45		
Ser	Arg	Phe	Glu	Glu	Gln	Phe	Gln	Gln	Leu	Ser	Pro	Lys	Val	Glu	Lys
		50				55					60				
Phe	Ala	Gln	Leu	Leu	Glu	Glu	Ile	Lys	Gln	Gln	Leu	Asn	Ser	Thr	Ala

-continued

---

65		70		75		80									
Asp	Ala	Val	Gln	Glu	Gln	Asp	Gln	Gln	Leu	Ser	Asn	Asn	Phe	Gly	Leu
				85					90					95	

Gln

<210> SEQ ID NO 12  
 <211> LENGTH: 102  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 12

Met	Gly	Gly	Tyr	Lys	Gly	Ile	Lys	Ala	Asp	Gly	Gly	Lys	Val	Asn	Gln
1				5					10					15	
Ala	Lys	Gln	Leu	Ala	Ala	Lys	Ile	Ala	Lys	Asp	Ile	Glu	Ala	Cys	Gln
			20					25					30		
Lys	Gln	Thr	Gln	Gln	Leu	Ala	Glu	Tyr	Ile	Glu	Gly	Ser	Asp	Trp	Glu
		35					40					45			
Gly	Gln	Phe	Ala	Asn	Lys	Val	Lys	Asp	Val	Leu	Leu	Ile	Met	Ala	Lys
	50					55					60				
Phe	Gln	Glu	Glu	Leu	Val	Gln	Pro	Met	Ala	Asp	His	Gln	Lys	Ala	Ile
65					70					75					80
Asp	Asn	Leu	Ser	Gln	Asn	Leu	Ala	Lys	Tyr	Asp	Thr	Leu	Ser	Ile	Lys
				85					90					95	
Gln	Gly	Leu	Asp	Arg	Val										
				100											

<210> SEQ ID NO 13  
 <211> LENGTH: 1385  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 13

Met	Leu	Asn	Arg	Glu	Asn	Lys	Thr	Ala	Ile	Thr	Arg	Lys	Gly	Met	Val
1				5					10					15	
Ser	Asn	Arg	Leu	Asn	Lys	Phe	Ser	Ile	Arg	Lys	Tyr	Thr	Val	Gly	Thr
			20					25					30		
Ala	Ser	Ile	Leu	Val	Gly	Thr	Thr	Leu	Ile	Phe	Gly	Leu	Gly	Asn	Gln
		35					40					45			
Glu	Ala	Lys	Ala	Ala	Glu	Ser	Thr	Asn	Lys	Glu	Leu	Asn	Glu	Ala	Thr
	50					55					60				
Thr	Ser	Ala	Ser	Asp	Asn	Gln	Ser	Ser	Asp	Lys	Val	Asp	Met	Gln	Gln
65					70					75					80
Leu	Asn	Gln	Glu	Asp	Asn	Thr	Lys	Asn	Asp	Asn	Gln	Lys	Glu	Met	Val
				85					90					95	
Ser	Ser	Gln	Gly	Asn	Glu	Thr	Thr	Ser	Asn	Gly	Asn	Lys	Ser	Ile	Glu
			100					105					110		
Lys	Glu	Ser	Val	Gln	Ser	Thr	Thr	Gly	Asn	Lys	Val	Glu	Val	Ser	Thr
		115					120					125			
Ala	Lys	Ser	Asp	Glu	Gln	Ala	Ser	Pro	Lys	Ser	Thr	Asn	Glu	Asp	Leu
		130				135						140			
Asn	Thr	Lys	Gln	Thr	Ile	Ser	Asn	Gln	Glu	Gly	Leu	Gln	Pro	Asp	Leu
145					150					155					160
Leu	Glu	Asn	Lys	Ser	Val	Val	Asn	Val	Gln	Pro	Thr	Asn	Glu	Glu	Asn
				165					170						175



-continued

---

Val Gly Asn Val Thr Val Thr Val Phe Asp Asn Asn Thr Asn Thr Lys  
595 600 605

Val Gly Glu Ala Val Thr Lys Glu Asp Gly Ser Tyr Leu Ile Pro Asn  
610 615 620

Leu Pro Asn Gly Asp Tyr Arg Val Glu Phe Ser Asn Leu Pro Lys Gly  
625 630 635 640

Tyr Glu Val Thr Pro Ser Lys Gln Gly Asn Asn Glu Glu Leu Asp Ser  
645 650 655

Asn Gly Leu Ser Ser Val Ile Thr Val Asn Gly Lys Asp Asn Leu Ser  
660 665 670

Ala Asp Leu Gly Ile Tyr Lys Pro Lys Tyr Asn Leu Gly Asp Tyr Val  
675 680 685

Trp Glu Asp Thr Asn Lys Asn Gly Ile Gln Asp Gln Asp Glu Lys Gly  
690 695 700

Ile Ser Gly Val Thr Val Thr Leu Lys Asp Glu Asn Gly Asn Val Leu  
705 710 715 720

Lys Thr Val Thr Thr Asp Ala Asp Gly Lys Tyr Lys Phe Thr Asp Leu  
725 730 735

Asp Asn Gly Asn Tyr Lys Val Glu Phe Thr Thr Pro Glu Gly Tyr Thr  
740 745 750

Pro Thr Thr Val Thr Ser Gly Ser Asp Ile Glu Lys Asp Ser Asn Gly  
755 760 765

Leu Thr Thr Thr Gly Val Ile Asn Gly Ala Asp Asn Met Thr Leu Asp  
770 775 780

Ser Gly Phe Tyr Lys Thr Pro Lys Tyr Asn Leu Gly Asn Tyr Val Trp  
785 790 795 800

Glu Asp Thr Asn Lys Asp Gly Lys Gln Asp Ser Thr Glu Lys Gly Ile  
805 810 815

Ser Gly Val Thr Val Thr Leu Lys Asn Glu Asn Gly Glu Val Leu Gln  
820 825 830

Thr Thr Lys Thr Asp Lys Asp Gly Lys Tyr Gln Phe Thr Gly Leu Glu  
835 840 845

Asn Gly Thr Tyr Lys Val Glu Phe Glu Thr Pro Ser Gly Tyr Thr Pro  
850 855 860

Thr Gln Val Gly Ser Gly Thr Asp Glu Gly Ile Asp Ser Asn Gly Thr  
865 870 875 880

Ser Thr Thr Gly Val Ile Lys Asp Lys Asp Asn Asp Thr Ile Asp Ser  
885 890 895

Gly Phe Tyr Lys Pro Thr Tyr Asn Leu Gly Asp Tyr Val Trp Glu Asp  
900 905 910

Thr Asn Lys Asn Gly Val Gln Asp Lys Asp Glu Lys Gly Ile Ser Gly  
915 920 925

Val Thr Val Thr Leu Lys Asp Glu Asn Asp Lys Val Leu Lys Thr Val  
930 935 940

Thr Thr Asp Glu Asn Gly Lys Tyr Gln Phe Thr Asp Leu Asn Asn Gly  
945 950 955 960

Thr Tyr Lys Val Glu Phe Glu Thr Pro Ser Gly Tyr Thr Pro Thr Ser  
965 970 975

Val Thr Ser Gly Asn Asp Thr Glu Lys Asp Ser Asn Gly Leu Thr Thr  
980 985 990

Thr Gly Val Ile Lys Asp Ala Asp Asn Met Thr Leu Asp Ser Gly Phe

-continued

995				1000				1005						
Tyr	Lys	Thr	Pro	Lys	Tyr	Ser	Leu	Gly	Asp	Tyr	Val	Trp	Tyr	Asp
1010						1015					1020			
Ser	Asn	Lys	Asp	Gly	Lys	Gln	Asp	Ser	Thr	Glu	Lys	Gly	Ile	Lys
1025						1030					1035			
Asp	Val	Lys	Val	Ile	Leu	Leu	Asn	Glu	Lys	Gly	Glu	Val	Ile	Gly
1040						1045					1050			
Thr	Thr	Lys	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	Arg	Phe	Asp	Asn	Leu
1055						1060					1065			
Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu	Lys	Pro	Thr	Gly	Leu
1070						1075					1080			
Thr	Gln	Thr	Gly	Thr	Asn	Thr	Thr	Glu	Asp	Asp	Lys	Asp	Ala	Asp
1085						1090					1095			
Gly	Gly	Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp	Asp	Phe	Thr
1100						1105					1110			
Leu	Asp	Asn	Gly	Tyr	Tyr	Glu	Glu	Glu	Thr	Ser	Asp	Ser	Asp	Ser
1115						1120					1125			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
1130						1135					1140			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
1145						1150					1155			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
1160						1165					1170			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
1175						1180					1185			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
1190						1195					1200			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
1205						1210					1215			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
1220						1225					1230			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
1235						1240					1245			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
1250						1255					1260			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
1265						1270					1275			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
1280						1285					1290			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
1295						1300					1305			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
1310						1315					1320			
Ser	Asp	Ala	Gly	Lys	His	Thr	Pro	Val	Lys	Pro	Met	Ser	Thr	Thr
1325						1330					1335			
Lys	Asp	His	His	Asn	Lys	Ala	Lys	Ala	Leu	Pro	Glu	Thr	Gly	Asn
1340						1345					1350			
Glu	Asn	Ser	Gly	Ser	Asn	Asn	Ala	Thr	Leu	Phe	Gly	Gly	Leu	Phe
1355						1360					1365			
Ala	Ala	Leu	Gly	Ser	Leu	Leu	Leu	Phe	Gly	Arg	Arg	Lys	Lys	Gln
1370						1375					1380			



-continued

Asn Lys  
1385

<210> SEQ ID NO 14  
<211> LENGTH: 1141  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 14

Met Ile Asn Arg Asp Asn Lys Lys Ala Ile Thr Lys Lys Gly Met Ile  
1 5 10 15

Ser Asn Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Thr Val Gly Thr  
20 25 30

Ala Ser Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Gly Asn Gln  
35 40 45

Glu Ala Lys Ala Ala Glu Asn Thr Ser Thr Glu Asn Ala Lys Gln Asp  
50 55 60

Asp Ala Thr Thr Ser Asp Asn Lys Glu Val Val Ser Glu Thr Glu Asn  
65 70 75 80

Asn Ser Thr Thr Glu Asn Asp Ser Thr Asn Pro Ile Lys Lys Glu Thr  
85 90 95

Asn Thr Asp Ser Gln Pro Glu Ala Lys Glu Glu Ser Thr Thr Ser Ser  
100 105 110

Thr Gln Gln Gln Gln Asn Asn Val Thr Ala Thr Thr Glu Thr Lys Pro  
115 120 125

Gln Asn Ile Glu Lys Glu Asn Val Lys Pro Ser Thr Asp Lys Thr Ala  
130 135 140

Thr Glu Asp Thr Ser Val Ile Leu Glu Glu Lys Lys Ala Pro Asn Tyr  
145 150 155 160

Thr Asn Asn Asp Val Thr Thr Lys Pro Ser Thr Ser Glu Ile Gln Thr  
165 170 175

Lys Pro Thr Thr Pro Gln Glu Ser Thr Asn Ile Glu Asn Ser Gln Pro  
180 185 190

Gln Pro Thr Pro Ser Lys Val Asp Asn Gln Val Thr Asp Ala Thr Asn  
195 200 205

Pro Lys Glu Pro Val Asn Val Ser Lys Glu Glu Leu Lys Asn Asn Pro  
210 215 220

Glu Lys Leu Lys Glu Leu Val Arg Asn Asp Asn Asn Thr Asp Arg Ser  
225 230 235 240

Thr Lys Pro Val Ala Thr Ala Pro Thr Ser Val Ala Pro Lys Arg Leu  
245 250 255

Asn Ala Lys Met Arg Phe Ala Val Ala Gln Pro Ala Ala Val Ala Ser  
260 265 270

Asn Asn Val Asn Asp Leu Ile Thr Val Thr Lys Gln Thr Ile Lys Val  
275 280 285

Gly Asp Gly Lys Asp Asn Val Ala Ala Ala His Asp Gly Lys Asp Ile  
290 295 300

Glu Tyr Asp Thr Glu Phe Thr Ile Asp Asn Lys Val Lys Lys Gly Asp  
305 310 315 320

Thr Met Thr Ile Asn Tyr Asp Lys Asn Val Ile Pro Ser Asp Leu Thr  
325 330 335

Asp Lys Asn Asp Pro Ile Asp Ile Thr Asp Pro Ser Gly Glu Val Ile  
340 345 350

-continued

---

Ala	Lys	Gly	Thr	Phe	Asp	Lys	Ala	Thr	Lys	Gln	Ile	Thr	Tyr	Thr	Phe
		355					360					365			
Thr	Asp	Tyr	Val	Asp	Lys	Tyr	Glu	Asp	Ile	Lys	Ala	Arg	Leu	Thr	Leu
		370				375					380				
Tyr	Ser	Tyr	Ile	Asp	Lys	Gln	Ala	Val	Pro	Asn	Glu	Thr	Ser	Leu	Asn
		385				390					395				400
Leu	Thr	Phe	Ala	Thr	Ala	Gly	Lys	Glu	Thr	Ser	Gln	Asn	Val	Ser	Val
				405					410						415
Asp	Tyr	Gln	Asp	Pro	Met	Val	His	Gly	Asp	Ser	Asn	Ile	Gln	Ser	Ile
				420					425					430	
Phe	Thr	Lys	Leu	Asp	Glu	Asn	Lys	Gln	Thr	Ile	Glu	Gln	Gln	Ile	Tyr
				435				440					445		
Val	Asn	Pro	Leu	Lys	Lys	Thr	Ala	Thr	Asn	Thr	Lys	Val	Asp	Ile	Ala
				450			455					460			
Gly	Ser	Gln	Val	Asp	Asp	Tyr	Gly	Asn	Ile	Lys	Leu	Gly	Asn	Gly	Ser
				465			470				475				480
Thr	Ile	Ile	Asp	Gln	Asn	Thr	Glu	Ile	Lys	Val	Tyr	Lys	Val	Asn	Pro
				485					490						495
Asn	Gln	Gln	Leu	Pro	Gln	Ser	Asn	Arg	Ile	Tyr	Asp	Phe	Ser	Gln	Tyr
				500					505					510	
Glu	Asp	Val	Thr	Ser	Gln	Phe	Asp	Asn	Lys	Lys	Ser	Phe	Ser	Asn	Asn
			515					520					525		
Val	Ala	Thr	Leu	Asp	Phe	Gly	Asp	Ile	Asn	Ser	Ala	Tyr	Ile	Ile	Lys
			530				535					540			
Val	Val	Ser	Lys	Tyr	Thr	Pro	Thr	Ser	Asp	Gly	Glu	Leu	Asp	Ile	Ala
				545			550				555				560
Gln	Gly	Thr	Ser	Met	Arg	Thr	Thr	Asp	Lys	Tyr	Gly	Tyr	Tyr	Asn	Tyr
				565					570						575
Ala	Gly	Tyr	Ser	Asn	Phe	Ile	Val	Thr	Ser	Asn	Asp	Thr	Gly	Gly	Gly
				580					585					590	
Asp	Gly	Thr	Val	Lys	Pro	Glu	Glu	Lys	Leu	Tyr	Lys	Ile	Gly	Asp	Tyr
				595				600					605		
Val	Trp	Glu	Asp	Val	Asp	Lys	Asp	Gly	Val	Gln	Gly	Thr	Asp	Ser	Lys
				610			615					620			
Glu	Lys	Pro	Met	Ala	Asn	Val	Leu	Val	Thr	Leu	Thr	Tyr	Pro	Asp	Gly
				625			630				635				640
Thr	Thr	Lys	Ser	Val	Arg	Thr	Asp	Ala	Asn	Gly	His	Tyr	Glu	Phe	Gly
				645					650						655
Gly	Leu	Lys	Asp	Gly	Glu	Thr	Tyr	Thr	Val	Lys	Phe	Glu	Thr	Pro	Ala
				660					665					670	
Gly	Tyr	Leu	Pro	Thr	Lys	Val	Asn	Gly	Thr	Thr	Asp	Gly	Glu	Lys	Asp
				675				680					685		
Ser	Asn	Gly	Ser	Ser	Ile	Thr	Val	Lys	Ile	Asn	Gly	Lys	Asp	Asp	Met
				690				695				700			
Ser	Leu	Asp	Thr	Gly	Phe	Tyr	Lys	Glu	Pro	Lys	Tyr	Asn	Leu	Gly	Asp
				705				710				715			720
Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Ile	Gln	Asp	Ala	Asn	Glu
				725					730						735
Pro	Gly	Ile	Lys	Asp	Val	Lys	Val	Thr	Leu	Lys	Asp	Ser	Thr	Gly	Lys
				740					745					750	
Val	Ile	Gly	Thr	Thr	Thr	Thr	Asp	Ala	Ser	Gly	Lys	Tyr	Lys	Phe	Thr
				755				760						765	

-continued

---

Asp Leu Asp Asn Gly Asn Tyr Thr Val Glu Phe Glu Thr Pro Ala Gly  
 770 775 780  
 Tyr Thr Pro Thr Val Lys Asn Thr Thr Ala Glu Asp Lys Asp Ser Asn  
 785 790 795 800  
 Gly Leu Thr Thr Thr Gly Val Ile Lys Asp Ala Asp Asn Met Thr Leu  
 805 810 815  
 Asp Ser Gly Phe Tyr Lys Thr Pro Lys Tyr Ser Leu Gly Asp Tyr Val  
 820 825 830  
 Trp Tyr Asp Ser Asn Lys Asp Gly Lys Gln Asp Ser Thr Glu Lys Gly  
 835 840 845  
 Ile Lys Asp Val Lys Val Thr Leu Leu Asn Glu Lys Gly Glu Val Ile  
 850 855 860  
 Gly Thr Thr Lys Thr Asp Glu Asn Gly Lys Tyr Arg Phe Asp Asn Leu  
 865 870 875 880  
 Asp Ser Gly Lys Tyr Lys Val Ile Phe Glu Lys Pro Ala Gly Leu Thr  
 885 890 895  
 Gln Thr Val Thr Asn Thr Thr Glu Asp Asp Lys Asp Ala Asp Gly Gly  
 900 905 910  
 Glu Val Asp Val Thr Ile Thr Asp His Asp Asp Phe Thr Leu Asp Asn  
 915 920 925  
 Gly Tyr Phe Glu Glu Asp Thr Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 930 935 940  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 945 950 955 960  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 965 970 975  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 980 985 990  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 995 1000 1005  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 1010 1015 1020  
 Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 1025 1030 1035  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 1040 1045 1050  
 Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 1055 1060 1065  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ala Gly  
 1070 1075 1080  
 Lys His Thr Pro Val Lys Pro Met Ser Thr Thr Lys Asp His His  
 1085 1090 1095  
 Asn Lys Ala Lys Ala Leu Pro Glu Thr Gly Ser Glu Asn Asn Gly  
 1100 1105 1110  
 Ser Asn Asn Ala Thr Leu Phe Gly Gly Leu Phe Ala Ala Leu Gly  
 1115 1120 1125  
 Ser Leu Leu Leu Phe Gly Arg Arg Lys Lys Gln Asn Lys  
 1130 1135 1140

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 350

&lt;212&gt; TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 15

```

Met Thr Lys His Tyr Leu Asn Ser Lys Tyr Gln Ser Glu Gln Arg Ser
1           5           10           15
Ser Ala Met Lys Lys Ile Thr Met Gly Thr Ala Ser Ile Ile Leu Gly
          20           25           30
Ser Leu Val Tyr Ile Gly Ala Asp Ser Gln Gln Val Asn Ala Ala Thr
          35           40           45
Glu Ala Thr Asn Ala Thr Asn Asn Gln Ser Thr Gln Val Ser Gln Ala
          50           55           60
Thr Ser Gln Pro Ile Asn Phe Gln Val Gln Lys Asp Gly Ser Ser Glu
65           70           75           80
Lys Ser His Met Asp Asp Tyr Met Gln His Pro Gly Lys Val Ile Lys
          85           90           95
Gln Asn Asn Lys Tyr Tyr Phe Gln Thr Val Leu Asn Asn Ala Ser Phe
          100          105          110
Trp Lys Glu Tyr Lys Phe Tyr Asn Ala Asn Asn Gln Glu Leu Ala Thr
          115          120          125
Thr Val Val Asn Asp Asn Lys Lys Ala Asp Thr Arg Thr Ile Asn Val
          130          135          140
Ala Val Glu Pro Gly Tyr Lys Ser Leu Thr Thr Lys Val His Ile Val
145          150          155          160
Val Pro Gln Ile Asn Tyr Asn His Arg Tyr Thr Thr His Leu Glu Phe
          165          170          175
Glu Lys Ala Ile Pro Thr Leu Ala Asp Ala Ala Lys Pro Asn Asn Val
          180          185          190
Lys Pro Val Gln Pro Lys Pro Ala Gln Pro Lys Thr Pro Thr Glu Gln
          195          200          205
Thr Lys Pro Val Gln Pro Lys Val Glu Lys Val Lys Pro Thr Val Thr
          210          215          220
Thr Thr Ser Lys Val Glu Asp Asn His Ser Thr Lys Val Val Ser Thr
225          230          235          240
Asp Thr Thr Lys Asp Gln Thr Lys Thr Gln Thr Ala His Thr Val Lys
          245          250          255
Thr Ala Gln Thr Ala Gln Glu Gln Asn Lys Val Gln Thr Pro Val Lys
          260          265          270
Asp Val Ala Thr Ala Lys Ser Glu Ser Asn Asn Gln Ala Val Ser Asp
          275          280          285
Asn Lys Ser Gln Gln Thr Asn Lys Val Thr Lys His Asn Glu Thr Pro
          290          295          300
Lys Gln Ala Ser Lys Ala Lys Glu Leu Pro Lys Thr Gly Leu Thr Ser
305          310          315          320
Val Asp Asn Phe Ile Ser Thr Val Ala Phe Ala Thr Leu Ala Leu Leu
          325          330          335
Gly Ser Leu Ser Leu Leu Leu Phe Lys Arg Lys Glu Ser Lys
          340          345          350

```

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 645

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 16

-continued

---

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys  
 1 5 10 15  
 Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu  
 20 25 30  
 Met Ser Asn Gly Glu Ala Gln Ala Ala Ala Glu Glu Thr Gly Gly Thr  
 35 40 45  
 Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr  
 50 55 60  
 Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser  
 65 70 75 80  
 Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala  
 85 90 95  
 Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Ala Val Lys  
 100 105 110  
 Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu  
 115 120 125  
 Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser  
 130 135 140  
 Ala Pro Asn Ser Arg Pro Ile Asp Phe Glu Met Lys Lys Glu Asn Gly  
 145 150 155 160  
 Glu Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Lys Pro Ala Arg Val  
 165 170 175  
 Ile Phe Thr Asp Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly  
 180 185 190  
 Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro  
 195 200 205  
 Ile Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg  
 210 215 220  
 Phe Ser Val Ser Asn Gly Thr Lys Ala Val Lys Ile Val Ser Ser Thr  
 225 230 235 240  
 His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Glu Phe  
 245 250 255  
 Ala Gln Pro Ile Tyr Asn Ser Ala Asp Lys Phe Lys Thr Glu Glu Asp  
 260 265 270  
 Tyr Lys Ala Glu Lys Leu Leu Ala Pro Tyr Lys Lys Ala Lys Thr Leu  
 275 280 285  
 Glu Arg Gln Val Tyr Glu Leu Asn Lys Ile Gln Asp Lys Leu Pro Glu  
 290 295 300  
 Lys Leu Lys Ala Glu Tyr Lys Lys Lys Leu Glu Asp Thr Lys Lys Ala  
 305 310 315 320  
 Leu Asp Glu Gln Val Lys Ser Ala Ile Thr Glu Phe Gln Asn Val Gln  
 325 330 335  
 Pro Thr Asn Glu Lys Met Thr Asp Leu Gln Asp Thr Lys Tyr Val Val  
 340 345 350  
 Tyr Glu Ser Val Glu Asn Asn Glu Ser Met Met Asp Thr Phe Val Lys  
 355 360 365  
 His Pro Ile Lys Thr Gly Met Leu Asn Gly Lys Lys Tyr Met Val Met  
 370 375 380  
 Glu Thr Thr Asn Asp Asp Tyr Trp Lys Asp Phe Met Val Glu Gly Gln  
 385 390 395 400  
 Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile

-continued

405					410					415					
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys
			420						425				430		
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile
			435					440					445		
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys
			450					455					460		
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr
				465				470					475		480
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln
				485					490					495	
Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu
			500					505					510		
Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys
			515					520					525		
Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser	Ser	Ser	Thr	Thr	Pro	Thr	Lys	Val
			530					535					540		
Val	Ser	Thr	Thr	Gln	Asn	Val	Ala	Lys	Pro	Thr	Thr	Ala	Ser	Ser	Lys
				545				550					555		560
Thr	Thr	Lys	Asp	Val	Val	Gln	Thr	Ser	Ala	Gly	Ser	Ser	Glu	Ala	Lys
				565					570					575	
Asp	Ser	Ala	Pro	Leu	Gln	Lys	Ala	Asn	Ile	Lys	Asn	Thr	Asn	Asp	Gly
			580					585						590	
His	Thr	Gln	Ser	Gln	Asn	Asn	Lys	Asn	Thr	Gln	Glu	Asn	Lys	Ala	Lys
			595					600					605		
Ser	Leu	Pro	Gln	Thr	Gly	Glu	Glu	Ser	Asn	Lys	Asp	Met	Thr	Leu	Pro
			610					615					620		
Leu	Met	Ala	Leu	Leu	Ala	Leu	Ser	Ser	Ile	Val	Ala	Phe	Val	Leu	Pro
			625						630					635	640
Arg	Lys	Arg	Lys	Asn											
				645											

<210> SEQ ID NO 17  
 <211> LENGTH: 80  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 17

Met	Asn	Gln	His	Val	Lys	Val	Thr	Phe	Asp	Phe	Thr	Asn	Tyr	Asn	Tyr
1				5					10					15	
Gly	Thr	Tyr	Asp	Leu	Ala	Val	Pro	Ala	Tyr	Leu	Pro	Ile	Lys	Asn	Leu
			20					25					30		
Ile	Ala	Leu	Val	Leu	Asp	Ser	Leu	Asp	Ile	Ser	Ile	Phe	Asp	Val	Asn
			35					40					45		
Thr	Gln	Ile	Lys	Val	Met	Thr	Lys	Gly	Gln	Leu	Leu	Val	Glu	Asn	Asp
			50					55					60		
Arg	Leu	Ile	Asp	Tyr	Gln	Ile	Ala	Asp	Gly	Asp	Ile	Leu	Lys	Leu	Leu
			65					70					75		80

<210> SEQ ID NO 18  
 <211> LENGTH: 877  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 18

-continued

---

Met Lys Lys Arg Ile Asp Tyr Leu Ser Asn Lys Gln Asn Lys Tyr Ser  
 1 5 10 15  
 Ile Arg Arg Phe Thr Val Gly Thr Thr Ser Val Ile Val Gly Ala Thr  
 20 25 30  
 Ile Leu Phe Gly Ile Gly Asn His Gln Ala Gln Ala Ser Glu Gln Ser  
 35 40 45  
 Asn Asp Thr Thr Gln Ser Ser Lys Asn Asn Ala Ser Ala Asp Ser Glu  
 50 55 60  
 Lys Asn Asn Met Ile Glu Thr Pro Gln Leu Asn Thr Thr Ala Asn Asp  
 65 70 75 80  
 Thr Ser Asp Ile Ser Ala Asn Thr Asn Ser Ala Asn Val Asp Ser Thr  
 85 90 95  
 Thr Lys Pro Met Ser Thr Gln Thr Ser Asn Thr Thr Thr Thr Glu Pro  
 100 105 110  
 Ala Ser Thr Asn Glu Thr Pro Gln Pro Thr Ala Ile Lys Asn Gln Ala  
 115 120 125  
 Thr Ala Ala Lys Met Gln Asp Gln Thr Val Pro Gln Glu Ala Asn Ser  
 130 135 140  
 Gln Val Asp Asn Lys Thr Thr Asn Asp Ala Asn Ser Ile Ala Thr Asn  
 145 150 155 160  
 Ser Glu Leu Lys Asn Ser Gln Thr Leu Asp Leu Pro Gln Ser Ser Pro  
 165 170 175  
 Gln Thr Ile Ser Asn Ala Gln Gly Thr Ser Lys Pro Ser Val Arg Thr  
 180 185 190  
 Arg Ala Val Arg Ser Leu Ala Val Ala Glu Pro Val Val Asn Ala Ala  
 195 200 205  
 Asp Ala Lys Gly Thr Asn Val Asn Asp Lys Val Thr Ala Ser Asn Phe  
 210 215 220  
 Lys Leu Glu Lys Thr Thr Phe Asp Pro Asn Gln Ser Gly Asn Thr Phe  
 225 230 235 240  
 Met Ala Ala Asn Phe Thr Val Thr Asp Lys Val Lys Ser Gly Asp Tyr  
 245 250 255  
 Phe Thr Ala Lys Leu Pro Asp Ser Leu Thr Gly Asn Gly Asp Val Asp  
 260 265 270  
 Tyr Ser Asn Ser Asn Asn Thr Met Pro Ile Ala Asp Ile Lys Ser Thr  
 275 280 285  
 Asn Gly Asp Val Val Ala Lys Ala Thr Tyr Asp Ile Leu Thr Lys Thr  
 290 295 300  
 Tyr Thr Phe Val Phe Thr Asp Tyr Val Asn Asn Lys Glu Asn Ile Asn  
 305 310 315 320  
 Gly Gln Phe Ser Leu Pro Leu Phe Thr Asp Arg Ala Lys Ala Pro Lys  
 325 330 335  
 Ser Gly Thr Tyr Asp Ala Asn Ile Asn Ile Ala Asp Glu Met Phe Asn  
 340 345 350  
 Asn Lys Ile Thr Tyr Asn Tyr Ser Ser Pro Ile Ala Gly Ile Asp Lys  
 355 360 365  
 Pro Asn Gly Ala Asn Ile Ser Ser Gln Ile Ile Gly Val Asp Thr Ala  
 370 375 380  
 Ser Gly Gln Asn Thr Tyr Lys Gln Thr Val Phe Val Asn Pro Lys Gln  
 385 390 395 400  
 Arg Val Leu Gly Asn Thr Trp Val Tyr Ile Lys Gly Tyr Gln Asp Lys

-continued

405				410				415							
Ile	Glu	Glu	Ser	Ser	Gly	Lys	Val	Ser	Ala	Thr	Asp	Thr	Lys	Leu	Arg
			420								425				430
Ile	Phe	Glu	Val	Asn	Asp	Thr	Ser	Lys	Leu	Ser	Asp	Ser	Tyr	Tyr	Ala
			435												445
Asp	Pro	Asn	Asp	Ser	Asn	Leu	Lys	Glu	Val	Thr	Asp	Gln	Phe	Lys	Asn
			450				455								460
Arg	Ile	Tyr	Tyr	Glu	His	Pro	Asn	Val	Ala	Ser	Ile	Lys	Phe	Gly	Asp
							470								480
Ile	Thr	Lys	Thr	Tyr	Val	Val	Leu	Val	Glu	Gly	His	Tyr	Asp	Asn	Thr
															495
Gly	Lys	Asn	Leu	Lys	Thr	Gln	Val	Ile	Gln	Glu	Asn	Val	Asp	Pro	Val
			500												510
Thr	Asn	Arg	Asp	Tyr	Ser	Ile	Phe	Gly	Trp	Asn	Asn	Glu	Asn	Val	Val
			515				520								525
Arg	Tyr	Gly	Gly	Gly	Ser	Ala	Asp	Gly	Asp	Ser	Ala	Val	Asn	Pro	Lys
			530				535								540
Asp	Pro	Thr	Pro	Gly	Pro	Pro	Val	Asp	Pro	Glu	Pro	Ser	Pro	Asp	Pro
							550								560
Glu	Pro	Glu	Pro	Thr	Pro	Asp	Pro	Glu	Pro	Ser	Pro	Asp	Pro	Glu	Pro
							565								575
Glu	Pro	Ser	Pro	Asp	Pro	Asp	Pro	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			580												590
Gly	Ser	Asp	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser
			595				600								605
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser
			610				615								620
Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
							630								640
Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
							645								655
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Glu	Ser
			660												670
Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			675				680								685
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			690				695								700
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser
							710								720
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
							725								735
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			740												750
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			755				760								765
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			770				775								780
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
							790								800
Asp	Ser	Asp	Ser	Arg	Val	Thr	Pro	Pro	Asn	Asn	Glu	Gln	Lys	Ala	Pro
															815



-continued

---

Ser Asn Pro Lys Gly Glu Val Asn His Ser Asn Lys Val Ser Lys Gln  
820 825 830

His Lys Thr Asp Ala Leu Pro Glu Thr Gly Asp Lys Ser Glu Asn Thr  
835 840 845

Asn Ala Thr Leu Phe Gly Ala Met Met Ala Leu Leu Gly Ser Leu Leu  
850 855 860

Leu Phe Arg Lys Arg Lys Gln Asp His Lys Glu Lys Ala  
865 870 875

<210> SEQ ID NO 19  
<211> LENGTH: 227  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 19

Met Lys Asn Ile Leu Lys Val Phe Asn Thr Thr Ile Leu Ala Leu Ile  
1 5 10 15

Ile Ile Ile Ala Thr Phe Ser Asn Ser Ala Asn Ala Ala Asp Ser Gly  
20 25 30

Thr Leu Asn Tyr Glu Val Tyr Lys Tyr Asn Thr Asn Asp Thr Ser Ile  
35 40 45

Ala Asn Asp Tyr Phe Asn Lys Pro Ala Lys Tyr Ile Lys Lys Asn Gly  
50 55 60

Lys Leu Tyr Val Gln Ile Thr Val Asn His Ser His Trp Ile Thr Gly  
65 70 75 80

Met Ser Ile Glu Gly His Lys Glu Asn Ile Ile Ser Lys Asn Thr Ala  
85 90 95

Lys Asp Glu Arg Thr Ser Glu Phe Glu Val Ser Lys Leu Asn Gly Lys  
100 105 110

Ile Asp Gly Lys Ile Asp Val Tyr Ile Asp Glu Lys Val Asn Gly Lys  
115 120 125

Pro Phe Lys Tyr Asp His His Tyr Asn Ile Thr Tyr Lys Phe Asn Gly  
130 135 140

Pro Thr Asp Val Ala Gly Ala Asn Ala Pro Gly Lys Asp Asp Lys Asn  
145 150 155 160

Ser Ala Ser Gly Ser Asp Lys Gly Ser Asp Gly Thr Thr Thr Gly Gln  
165 170 175

Ser Glu Ser Asn Ser Ser Asn Lys Asp Lys Val Glu Asn Pro Gln Thr  
180 185 190

Asn Ala Gly Thr Pro Ala Tyr Ile Tyr Ala Ile Pro Val Ala Ser Leu  
195 200 205

Ala Leu Leu Ile Ala Ile Thr Leu Phe Val Arg Lys Lys Ser Lys Gly  
210 215 220

Asn Val Glu  
225

<210> SEQ ID NO 20  
<211> LENGTH: 635  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 20

Met Ala Lys Tyr Arg Gly Lys Pro Phe Gln Leu Tyr Val Lys Leu Ser  
1 5 10 15

Cys Ser Thr Met Met Ala Ser Ser Ile Ile Leu Thr Asn Ile Leu Pro

-continued

20				25				30							
Tyr	Asp	Ala	Gln	Ala	Ala	Ser	Glu	Lys	Asp	Thr	Glu	Ile	Ser	Lys	Glu
	35						40					45			
Ile	Leu	Ser	Lys	Gln	Asp	Leu	Leu	Asp	Lys	Val	Asp	Lys	Ala	Ile	Arg
	50					55					60				
Gln	Ile	Glu	Gln	Leu	Lys	Gln	Leu	Ser	Ala	Ser	Ser	Lys	Ala	His	Tyr
	65				70					75				80	
Lys	Ala	Gln	Leu	Asn	Glu	Ala	Lys	Thr	Ala	Ser	Gln	Ile	Asp	Glu	Ile
			85						90					95	
Ile	Lys	Arg	Ala	Asn	Glu	Leu	Asp	Ser	Lys	Glu	Asn	Lys	Ser	Ser	His
			100						105					110	
Thr	Glu	Met	Asn	Gly	Gln	Ser	Asp	Ile	Asp	Ser	Lys	Leu	Asp	Gln	Leu
		115					120					125			
Leu	Lys	Asp	Leu	Asn	Glu	Val	Ser	Ser	Asn	Val	Asp	Arg	Gly	Gln	Gln
	130					135					140				
Ser	Gly	Glu	Asp	Asp	Leu	Asn	Ala	Met	Lys	Asn	Asp	Met	Ser	Gln	Thr
	145				150					155					160
Ala	Thr	Thr	Lys	Tyr	Gly	Glu	Lys	Asp	Asp	Lys	Asn	Asp	Glu	Ala	Met
			165						170					175	
Val	Asn	Lys	Ala	Leu	Glu	Asp	Leu	Asp	His	Leu	Asn	Gln	Gln	Ile	His
			180						185					190	
Lys	Ser	Lys	Asp	Ala	Leu	Lys	Asp	Ala	Ser	Lys	Asp	Pro	Ala	Val	Ser
		195					200					205			
Thr	Thr	Asp	Ser	Asn	His	Glu	Val	Ala	Lys	Thr	Pro	Asn	Asn	Asp	Gly
		210				215					220				
Ser	Gly	His	Val	Val	Leu	Asn	Lys	Phe	Leu	Ser	Asn	Glu	Glu	Asn	Gln
	225				230					235					240
Ser	His	Ser	Asn	Gln	Leu	Thr	Asp	Lys	Leu	Gln	Gly	Ser	Asp	Lys	Ile
			245						250					255	
Asn	His	Ala	Met	Ile	Glu	Lys	Leu	Ala	Lys	Ser	Asn	Ala	Ser	Thr	Gln
			260						265					270	
His	Tyr	Thr	Tyr	His	Lys	Leu	Asn	Thr	Leu	Gln	Ser	Leu	Asp	Gln	Arg
		275				280						285			
Ile	Ala	Asn	Thr	Gln	Leu	Pro	Lys	Asn	Gln	Lys	Ser	Asp	Leu	Met	Ser
		290				295					300				
Glu	Val	Asn	Lys	Thr	Lys	Glu	Arg	Ile	Lys	Ser	Gln	Arg	Asn	Ile	Ile
	305				310					315					320
Leu	Glu	Glu	Leu	Ala	Arg	Thr	Asp	Asp	Lys	Lys	Tyr	Ala	Thr	Gln	Ser
			325						330					335	
Ile	Leu	Glu	Ser	Ile	Phe	Asn	Lys	Asp	Glu	Ala	Asp	Lys	Ile	Leu	Lys
			340						345					350	
Asp	Ile	Arg	Val	Asp	Gly	Lys	Thr	Asp	Gln	Gln	Ile	Ala	Asp	Gln	Ile
		355					360					365			
Thr	Arg	His	Ile	Asp	Gln	Leu	Ser	Leu	Thr	Thr	Ser	Asp	Asp	Leu	Leu
		370				375					380				
Thr	Ser	Leu	Ile	Asp	Gln	Ser	Gln	Asp	Lys	Ser	Leu	Leu	Ile	Ser	Gln
		385			390					395					400
Ile	Leu	Gln	Thr	Lys	Leu	Gly	Lys	Ala	Glu	Ala	Asp	Lys	Leu	Ala	Lys
			405						410					415	
Asp	Trp	Thr	Asn	Lys	Gly	Leu	Ser	Asn	Arg	Gln	Ile	Val	Asp	Gln	Leu
			420						425					430	

-continued

---

Lys Lys His Phe Ala Ser Thr Gly Asp Thr Ser Ser Asp Asp Ile Leu  
 435 440 445  
 Lys Ala Ile Leu Asn Asn Ala Lys Asp Lys Lys Gln Ala Ile Glu Thr  
 450 455 460  
 Ile Leu Ala Thr Arg Ile Glu Arg Gln Lys Ala Lys Leu Leu Ala Asp  
 465 470 475 480  
 Leu Ile Thr Lys Ile Glu Thr Asp Gln Asn Lys Ile Phe Asn Leu Val  
 485 490 495  
 Lys Ser Ala Leu Asn Gly Lys Ala Asp Asp Leu Leu Asn Leu Gln Lys  
 500 505 510  
 Arg Leu Asn Gln Thr Lys Lys Asp Ile Asp Tyr Ile Leu Ser Pro Ile  
 515 520 525  
 Val Asn Arg Pro Ser Leu Leu Asp Arg Leu Asn Lys Asn Gly Lys Thr  
 530 535 540  
 Thr Asp Leu Asn Lys Leu Ala Asn Leu Met Asn Gln Gly Ser Asn Leu  
 545 550 555 560  
 Leu Asp Ser Ile Pro Asp Ile Pro Thr Pro Lys Pro Glu Lys Thr Leu  
 565 570 575  
 Thr Leu Gly Lys Gly Asn Gly Leu Leu Ser Gly Leu Leu Asn Ala Asp  
 580 585 590  
 Gly Asn Val Ser Leu Pro Lys Ala Gly Glu Thr Ile Lys Glu His Trp  
 595 600 605  
 Leu Pro Ile Ser Val Ile Val Gly Ala Met Gly Val Leu Met Ile Trp  
 610 615 620  
 Leu Ser Arg Arg Asn Lys Leu Lys Asn Lys Ala  
 625 630 635

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 953

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 21

Met Asn Asn Lys Lys Thr Ala Thr Asn Arg Lys Gly Met Ile Pro Asn  
 1 5 10 15  
 Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Ser Val Gly Thr Ala Ser  
 20 25 30  
 Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Ser Gly His Glu Ala  
 35 40 45  
 Lys Ala Ala Glu His Thr Asn Gly Glu Leu Asn Gln Ser Lys Asn Glu  
 50 55 60  
 Thr Thr Ala Pro Ser Glu Asn Lys Thr Thr Glu Lys Val Asp Ser Arg  
 65 70 75 80  
 Gln Leu Lys Asp Asn Thr Gln Thr Ala Thr Ala Asp Gln Pro Lys Val  
 85 90 95  
 Thr Met Ser Asp Ser Ala Thr Val Lys Glu Thr Ser Ser Asn Met Gln  
 100 105 110  
 Ser Pro Gln Asn Ala Thr Ala Ser Gln Ser Thr Thr Gln Thr Ser Asn  
 115 120 125  
 Val Thr Thr Asn Asp Lys Ser Ser Thr Thr Tyr Ser Asn Glu Thr Asp  
 130 135 140  
 Lys Ser Asn Leu Thr Gln Ala Lys Asn Val Ser Thr Thr Pro Lys Thr  
 145 150 155 160

-continued

---

Thr	Thr	Ile	Lys	Gln	Arg	Ala	Leu	Asn	Arg	Met	Ala	Val	Asn	Thr	Val
				165					170					175	
Ala	Ala	Pro	Gln	Gln	Gly	Thr	Asn	Val	Asn	Asp	Lys	Val	His	Phe	Thr
			180					185					190		
Asn	Ile	Asp	Ile	Ala	Ile	Asp	Lys	Gly	His	Val	Asn	Lys	Thr	Thr	Gly
		195					200					205			
Asn	Thr	Glu	Phe	Trp	Ala	Thr	Ser	Ser	Asp	Val	Leu	Lys	Leu	Lys	Ala
	210					215					220				
Asn	Tyr	Thr	Ile	Asp	Asp	Ser	Val	Lys	Glu	Gly	Asp	Thr	Phe	Thr	Phe
225					230					235					240
Lys	Tyr	Gly	Gln	Tyr	Phe	Arg	Pro	Gly	Ser	Val	Arg	Leu	Pro	Ser	Gln
				245					250					255	
Thr	Gln	Asn	Leu	Tyr	Asn	Ala	Gln	Gly	Asn	Ile	Ile	Ala	Lys	Gly	Ile
			260					265						270	
Tyr	Asp	Ser	Lys	Thr	Asn	Thr	Thr	Thr	Tyr	Thr	Phe	Thr	Asn	Tyr	Val
		275					280					285			
Asp	Gln	Tyr	Thr	Asn	Val	Ser	Gly	Ser	Phe	Glu	Gln	Val	Ala	Phe	Ala
	290					295					300				
Lys	Arg	Glu	Asn	Ala	Thr	Thr	Asp	Lys	Thr	Ala	Tyr	Lys	Met	Glu	Val
305					310					315					320
Thr	Leu	Gly	Asn	Asp	Thr	Tyr	Ser	Lys	Asp	Val	Ile	Val	Asp	Tyr	Gly
				325					330					335	
Asn	Gln	Lys	Gly	Gln	Gln	Leu	Ile	Ser	Ser	Thr	Asn	Tyr	Ile	Asn	Asn
			340					345					350		
Glu	Asp	Leu	Ser	Arg	Asn	Met	Thr	Val	Tyr	Val	Asn	Gln	Pro	Lys	Lys
		355					360					365			
Thr	Tyr	Thr	Lys	Glu	Thr	Phe	Val	Thr	Asn	Leu	Thr	Gly	Tyr	Lys	Phe
	370					375					380				
Asn	Pro	Asp	Ala	Lys	Asn	Phe	Lys	Ile	Tyr	Glu	Val	Thr	Asp	Gln	Asn
385					390					395					400
Gln	Phe	Val	Asp	Ser	Phe	Thr	Pro	Asp	Thr	Ser	Lys	Leu	Lys	Asp	Val
				405					410					415	
Thr	Gly	Gln	Phe	Asp	Val	Ile	Tyr	Ser	Asn	Asp	Asn	Lys	Thr	Ala	Thr
			420					425					430		
Val	Asp	Leu	Leu	Asn	Gly	Gln	Ser	Ser	Ser	Asp	Lys	Gln	Tyr	Ile	Ile
		435					440					445			
Gln	Gln	Val	Ala	Tyr	Pro	Asp	Asn	Ser	Ser	Thr	Asp	Asn	Gly	Lys	Ile
		450					455				460				
Asp	Tyr	Thr	Leu	Glu	Thr	Gln	Asn	Gly	Lys	Ser	Ser	Trp	Ser	Asn	Ser
465					470					475					480
Tyr	Ser	Asn	Val	Asn	Gly	Ser	Ser	Thr	Ala	Asn	Gly	Asp	Gln	Lys	Lys
				485					490					495	
Tyr	Asn	Leu	Gly	Asp	Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Lys
		500						505					510		
Gln	Asp	Ala	Asn	Glu	Lys	Gly	Ile	Lys	Gly	Val	Tyr	Val	Ile	Leu	Lys
		515					520					525			
Asp	Ser	Asn	Gly	Lys	Glu	Leu	Asp	Arg	Thr	Thr	Thr	Asp	Glu	Asn	Gly
	530					535						540			
Lys	Tyr	Gln	Phe	Thr	Gly	Leu	Ser	Asn	Gly	Thr	Tyr	Ser	Val	Glu	Phe
545					550					555					560
Ser	Thr	Pro	Ala	Gly	Tyr	Thr	Pro	Thr	Thr	Ala	Asn	Ala	Gly	Thr	Asp
				565					570					575	

-continued

---

Asp Ala Val Asp Ser Asp Gly Leu Thr Thr Thr Gly Val Ile Lys Asp  
                   580                                  585                                  590

Ala Asp Asn Met Thr Leu Asp Ser Gly Phe Tyr Lys Thr Pro Lys Tyr  
                   595                                  600                                  605

Ser Leu Gly Asp Tyr Val Trp Tyr Asp Ser Asn Lys Asp Gly Lys Gln  
                   610                                  615                                  620

Asp Ser Thr Glu Lys Gly Ile Lys Gly Val Lys Val Thr Leu Gln Asn  
   625                                  630                                  635                                  640

Glu Lys Gly Glu Val Ile Gly Thr Thr Glu Thr Asp Glu Asn Gly Lys  
                                   645                                  650                                  655

Tyr Arg Phe Asp Asn Leu Asp Ser Gly Lys Tyr Lys Val Ile Phe Glu  
                   660                                  665                                  670

Lys Pro Ala Gly Leu Thr Gln Thr Gly Thr Asn Thr Thr Glu Asp Asp  
                   675                                  680                                  685

Lys Asp Ala Asp Gly Gly Glu Val Asp Val Thr Ile Thr Asp His Asp  
                   690                                  695                                  700

Asp Phe Thr Leu Asp Asn Gly Tyr Tyr Glu Glu Glu Thr Ser Asp Ser  
   705                                  710                                  715                                  720

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   725                                  730                                  735

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   740                                  745                                  750

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   755                                  760                                  765

Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   770                                  775                                  780

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
   785                                  790                                  795                                  800

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   805                                  810                                  815

Asp Ser Asp Ser Asp Ser Asp Asn Asp Ser Asp Ser Asp Ser Asp Ser  
                   820                                  825                                  830

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   835                                  840                                  845

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   850                                  855                                  860

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
   865                                  870                                  875                                  880

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Ala Gly Lys  
                   885                                  890                                  895

His Thr Pro Thr Lys Pro Met Ser Thr Val Lys Asp Gln His Lys Thr  
                   900                                  905                                  910

Ala Lys Ala Leu Pro Glu Thr Gly Ser Glu Asn Asn Asn Ser Asn Asn  
                   915                                  920                                  925

Gly Thr Leu Phe Gly Gly Leu Phe Ala Ala Leu Gly Ser Leu Leu Leu  
                   930                                  935                                  940

Phe Gly Arg Arg Lys Lys Gln Asn Lys  
   945                                  950

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 989

&lt;212&gt; TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 22

Met Asn Met Lys Lys Lys Glu Lys His Ala Ile Arg Lys Lys Ser Ile  
 1 5 10 15  
 Gly Val Ala Ser Val Leu Val Gly Thr Leu Ile Gly Phe Gly Leu Leu  
 20 25 30  
 Ser Ser Lys Glu Ala Asp Ala Ser Glu Asn Ser Val Thr Gln Ser Asp  
 35 40 45  
 Ser Ala Ser Asn Glu Ser Lys Ser Asn Asp Ser Ser Ser Val Ser Ala  
 50 55 60  
 Ala Pro Lys Thr Asp Asp Thr Asn Val Ser Asp Thr Lys Thr Ser Ser  
 65 70 75 80  
 Asn Thr Asn Asn Gly Glu Thr Ser Val Ala Gln Asn Pro Ala Gln Gln  
 85 90 95  
 Glu Thr Thr Gln Ser Ser Ser Thr Asn Ala Thr Thr Glu Glu Thr Pro  
 100 105 110  
 Val Thr Gly Glu Ala Thr Thr Thr Thr Thr Asn Gln Ala Asn Thr Pro  
 115 120 125  
 Ala Thr Thr Gln Ser Ser Asn Thr Asn Ala Glu Glu Leu Val Asn Gln  
 130 135 140  
 Thr Ser Asn Glu Thr Thr Ser Asn Asp Thr Asn Thr Val Ser Ser Val  
 145 150 155 160  
 Asn Ser Pro Gln Asn Ser Thr Asn Ala Glu Asn Val Ser Thr Thr Gln  
 165 170 175  
 Asp Thr Ser Thr Glu Ala Thr Pro Ser Asn Asn Glu Ser Ala Pro Gln  
 180 185 190  
 Asn Thr Asp Ala Ser Asn Lys Asp Val Val Ser Gln Ala Val Asn Pro  
 195 200 205  
 Ser Thr Pro Arg Met Arg Ala Phe Ser Leu Ala Ala Val Ala Ala Asp  
 210 215 220  
 Ala Pro Ala Ala Gly Thr Asp Ile Thr Asn Gln Leu Thr Asp Val Lys  
 225 230 235 240  
 Val Thr Ile Asp Ser Gly Thr Thr Val Tyr Pro His Gln Ala Gly Tyr  
 245 250 255  
 Val Lys Leu Asn Tyr Gly Phe Ser Val Pro Asn Ser Ala Val Lys Gly  
 260 265 270  
 Asp Thr Phe Lys Ile Thr Val Pro Lys Glu Leu Asn Leu Asn Gly Val  
 275 280 285  
 Thr Ser Thr Ala Lys Val Pro Pro Ile Met Ala Gly Asp Gln Val Leu  
 290 295 300  
 Ala Asn Gly Val Ile Asp Ser Asp Gly Asn Val Ile Tyr Thr Phe Thr  
 305 310 315 320  
 Asp Tyr Val Asp Asn Lys Glu Asn Val Thr Ala Asn Ile Thr Met Pro  
 325 330 335  
 Ala Tyr Ile Asp Pro Glu Asn Val Thr Lys Thr Gly Asn Val Thr Leu  
 340 345 350  
 Thr Thr Gly Ile Gly Thr Asn Thr Ala Ser Lys Thr Val Leu Ile Asp  
 355 360 365  
 Tyr Glu Lys Tyr Gly Gln Phe His Asn Leu Ser Ile Lys Gly Thr Ile  
 370 375 380  
 Asp Gln Ile Asp Lys Thr Asn Asn Thr Tyr Arg Gln Thr Ile Tyr Val

-continued

---

385					390						395				400
Asn	Pro	Ser	Gly	Asp	Asn	Val	Val	Leu	Pro	Ala	Leu	Thr	Gly	Asn	Leu
				405						410				415	
Ile	Pro	Asn	Thr	Lys	Ser	Asn	Ala	Leu	Ile	Asp	Ala	Lys	Asn	Thr	Asp
			420					425					430		
Ile	Lys	Val	Tyr	Arg	Val	Asp	Asn	Ala	Asn	Asp	Leu	Ser	Glu	Ser	Tyr
		435					440					445			
Tyr	Val	Asn	Pro	Ser	Asp	Phe	Glu	Asp	Val	Thr	Asn	Gln	Val	Arg	Ile
	450					455					460				
Ser	Phe	Pro	Asn	Ala	Asn	Gln	Tyr	Lys	Val	Glu	Phe	Pro	Thr	Asp	Asp
465					470					475					480
Asp	Gln	Ile	Thr	Thr	Pro	Tyr	Ile	Val	Val	Asn	Gly	His	Ile	Asp	
			485					490					495		
Pro	Ala	Ser	Thr	Gly	Asp	Leu	Ala	Leu	Arg	Ser	Thr	Phe	Tyr	Gly	Tyr
			500					505					510		
Asp	Ser	Asn	Phe	Ile	Trp	Arg	Ser	Met	Ser	Trp	Asp	Asn	Glu	Val	Ala
		515					520					525			
Phe	Asn	Asn	Gly	Ser	Gly	Ser	Gly	Asp	Gly	Ile	Asp	Lys	Pro	Val	Val
	530					535				540					
Pro	Glu	Gln	Pro	Asp	Glu	Pro	Gly	Glu	Ile	Glu	Pro	Ile	Pro	Glu	Asp
545					550					555					560
Ser	Asp	Ser	Asp	Pro	Gly	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asn	Ser	Asp
				565					570					575	
Ser	Gly	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Thr	Ser	Asp	Ser	Gly	Ser	Asp
			580					585					590		
Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp
		595					600					605			
Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala
	610					615					620				
Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp
	625				630					635					640
Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp
				645					650					655	
Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Asp
			660					665					670		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		675					680					685			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	690					695					700				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	705				710					715					720
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
				725					730					735	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			740					745					750		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		755					760					765			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	770					775					780				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	785				790					795					800

-continued

---

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp  
805 810 815

Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Glu  
820 825 830

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp  
835 840 845

Ser Asp Ser Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp  
850 855 860

Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp  
865 870 875 880

Ser Ala Ser Asp Ser Asp Ser Gly Ser Asp Ser Asp Ser Ser Ser Asp  
885 890 895

Ser Asp Ser Asp Ser Thr Ser Asp Thr Gly Ser Asp Asn Asp Ser Asp  
900 905 910

Ser Asp Ser Asn Ser Asp Ser Glu Ser Gly Ser Asn Asn Asn Val Val  
915 920 925

Pro Pro Asn Ser Pro Lys Asn Gly Thr Asn Ala Ser Asn Lys Asn Glu  
930 935 940

Ala Lys Asp Ser Lys Glu Pro Leu Pro Asp Thr Gly Ser Glu Asp Glu  
945 950 955 960

Ala Asn Thr Ser Leu Ile Trp Gly Leu Leu Ala Ser Leu Gly Ser Leu  
965 970 975

Leu Leu Phe Arg Arg Lys Lys Glu Asn Lys Asp Lys Lys  
980 985

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 584

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 23

Met Lys Phe Lys Ser Leu Ile Thr Thr Thr Leu Ala Leu Gly Val Leu  
1 5 10 15

Ala Ser Thr Gly Ala Asn Phe Asn Asn Asn Glu Ala Ser Ala Ala Ala  
20 25 30

Lys Pro Leu Asp Lys Ser Ser Ser Ser Leu His His Gly Tyr Ser Lys  
35 40 45

Val His Val Pro Tyr Ala Ile Thr Val Asn Gly Thr Ser Gln Asn Ile  
50 55 60

Leu Ser Ser Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp  
65 70 75 80

Leu Glu Asp Arg Val Lys Ser Val Leu Lys Ser Asp Arg Gly Ile Ser  
85 90 95

Asp Ile Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Tyr Phe  
100 105 110

Lys Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ala Gly Ile Tyr Thr  
115 120 125

Ala Asp Leu Ile Asn Thr Ser Glu Ile Lys Ala Ile Asn Ile Asn Val  
130 135 140

Asp Thr Lys Lys Gln Val Glu Asp Lys Lys Lys Asp Lys Ala Asn Tyr  
145 150 155 160

Gln Val Pro Tyr Thr Ile Thr Val Asn Gly Thr Ser Gln Asn Ile Leu  
165 170 175







-continued

---

Gln Val Pro Val Gly Arg Thr Asp Phe Val Thr Val Asn Ser Asp Gly  
 370 375 380  
 Thr Asn Val Gln Trp Ser His Gly Ala Gly Ala Gly Ala Asn Lys Pro  
 385 390 395 400  
 Leu Gln Gln Met Trp Glu Tyr Gly Val Asn Asp Pro His Arg Ser His  
 405 410 415  
 Asp Phe Lys Ile Arg Asn Arg Ser Gly Gln Val Ile Tyr Asp Trp Pro  
 420 425 430  
 Thr Val His Ile Tyr Ser Leu Glu Asp Leu Ser Arg Ala Ser Asp Tyr  
 435 440 445  
 Phe Ser Glu Ala Gly Ala Thr Pro Ala Thr Lys Ala Phe Gly Arg Gln  
 450 455 460  
 Asn Phe Glu Tyr Ile Asn Gly Gln Lys Pro Ala Glu Ser Pro Gly Val  
 465 470 475 480  
 Pro Lys Val Tyr Thr Phe Ile Gly Gln Gly Asp Ala Ser Tyr Thr Ile  
 485 490 495  
 Ser Phe Lys Thr Gln Gly Pro Thr Val Asn Lys Leu Tyr Tyr Ala Ala  
 500 505 510  
 Gly Gly Arg Ala Leu Glu Tyr Asn Gln Leu Phe Met Tyr Ser Gln Leu  
 515 520 525  
 Tyr Val Glu Ser Thr Gln Asp His Gln Gln Arg Leu Asn Gly Leu Arg  
 530 535 540  
 Gln Val Val Asn Arg Thr Tyr Arg Ile Gly Thr Thr Lys Arg Val Glu  
 545 550 555 560  
 Val Ser Gln Gly Asn Val Gln Thr Lys Lys Val Leu Glu Ser Thr Asn  
 565 570 575  
 Leu Asn Ile Asp Asp Phe Val Asp Asp Pro Leu Ser Tyr Val Lys Thr  
 580 585 590  
 Pro Ser Asn Lys Val Leu Gly Phe Tyr Ser Asn Asn Ala Asn Thr Asn  
 595 600 605  
 Ala Phe Arg Pro Gly Gly Ala Gln Gln Leu Asn Glu Tyr Gln Leu Ser  
 610 615 620  
 Gln Leu Phe Thr Asp Gln Lys Leu Gln Glu Ala Ala Arg Thr Arg Asn  
 625 630 635 640  
 Pro Ile Arg Leu Met Ile Gly Phe Asp Tyr Pro Asp Ala Tyr Gly Asn  
 645 650 655  
 Ser Glu Thr Leu Val Pro Val Asn Leu Thr Val Leu Pro Glu Ile Gln  
 660 665 670  
 His Asn Ile Lys Phe Phe Lys Asn Asp Asp Thr Gln Asn Ile Ala Glu  
 675 680 685  
 Lys Pro Phe Ser Lys Gln Ala Gly His Pro Val Phe Tyr Val Tyr Ala  
 690 695 700  
 Gly Asn Gln Gly Asn Ala Ser Val Asn Leu Gly Gly Ser Val Thr Ser  
 705 710 715 720  
 Ile Gln Pro Leu Arg Ile Asn Leu Thr Ser Asn Glu Asn Phe Thr Asp  
 725 730 735  
 Lys Asp Trp Gln Ile Thr Gly Ile Pro Arg Thr Leu His Ile Glu Asn  
 740 745 750  
 Ser Thr Asn Arg Pro Asn Asn Ala Arg Glu Arg Asn Ile Glu Leu Val  
 755 760 765  
 Gly Asn Leu Leu Pro Gly Asp Tyr Phe Gly Thr Ile Arg Phe Gly Arg

-continued

770	775	780
Lys Glu Gln Leu Phe 785	Glu Ile Arg Val 790	Lys Pro His Thr Pro Thr Ile 795 800
Thr Thr Thr Ala 805	Glu Gln Leu Arg Gly Thr 810	Ala Leu Gln Lys Val Pro 815
Val Asn Ile Ser Gly 820	Ile Pro Leu Asp Pro Ser 825	Ala Leu Val Tyr Leu 830
Val Ala Pro Thr Asn 835	Gln Thr Thr Asn Gly Gly 840	Ser Glu Ala Asp Gln 845
Ile Pro Ser Gly Tyr Thr 850	Ile Leu Ala Thr Gly Thr 855 860	Pro Asp Gly Val
His Asn Thr Ile Thr 865	Ile Arg Pro Gln Asp Tyr 870 875	Val Val Phe Ile Pro 880
Pro Val Gly Lys Gln 885	Ile Arg Ala Val Val Tyr 890	Tyr Tyr Asn Lys Val Val 895
Ala Ser Asn Met Ser 900	Asn Ala Val Thr Ile Leu 905	Pro Asp Asp Ile Pro 910
Pro Thr Ile Asn Asn Pro 915	Val Gly Ile Asn Ala Lys 920 925	Tyr Tyr Arg Gly
Asp Glu Val Asn Phe Thr 930	Met Gly Val Ser Asp Arg 935 940	His Ser Gly Ile
Lys Asn Thr Thr Ile Thr 945	Thr Leu Pro Asn Gly Trp 950 955	Thr Ser Asn Leu 960
Thr Lys Ala Asp Lys 965	Asn Asn Gly Ser Leu Ser 970	Ile Thr Gly Arg Val 975
Ser Met Asn Gln Ala Phe 980	Asn Ser Asp Ile Thr Phe 985	Lys Val Ser Ala 990
Thr Asp Asn Val Asn Asn Thr 995	Thr Asn Asp Ser Gln Ser 1000	Lys His Val 1005
Ser Ile His Val Gly Lys 1010	Ile Ser Glu Asp Ala His 1015 1020	Pro Ile Val
Leu Gly Asn Thr Glu Lys Val 1025	Val Val Val Asn Pro Thr 1030 1035	Ala Val
Ser Asn Asp Glu Lys Gln Ser 1040	Ile Ile Thr Ala Phe Met 1045 1050	Asn Lys
Asn Gln Asn Ile Arg Gly Tyr 1055	Leu Ala Ser Thr Asp Pro 1060 1065	Val Thr
Val Asp Asn Asn Gly Asn Val 1070	Thr Leu His Tyr Arg Asp 1075 1080	Gly Ser
Ser Thr Thr Leu Asp Ala Thr 1085	Asn Val Met Thr Tyr Glu 1090 1095	Pro Val
Val Lys Pro Glu Tyr Gln Thr 1100	Val Asn Ala Ala Lys Thr 1105 1110	Ala Thr
Val Thr Ile Ala Lys Gly Gln 1115	Ser Phe Ser Ile Gly Asp 1120 1125	Ile Lys
Gln Tyr Phe Thr Leu Ser Asn 1130	Gly Gln Pro Ile Pro Ser 1135 1140	Gly Thr
Phe Thr Asn Ile Thr Ser Asp 1145	Arg Thr Ile Pro Thr Ala 1150 1155	Gln Glu
Val Ser Gln Met Asn Ala Gly 1160	Thr Gln Leu Tyr His Ile 1165 1170	Thr Ala

-continued

---

Thr	Asn	Ala	Tyr	His	Lys	Asp	Ser	Glu	Asp	Phe	Tyr	Ile	Ser	Leu
	1175					1180						1185		
Lys	Ile	Ile	Asp	Val	Lys	Gln	Pro	Glu	Gly	Asp	Gln	Arg	Val	Tyr
	1190					1195					1200			
Arg	Thr	Ser	Thr	Tyr	Asp	Leu	Thr	Thr	Asp	Glu	Ile	Ser	Lys	Val
	1205					1210					1215			
Lys	Gln	Ala	Phe	Ile	Asn	Ala	Asn	Arg	Asp	Val	Ile	Thr	Leu	Ala
	1220					1225					1230			
Glu	Gly	Asp	Ile	Ser	Val	Thr	Asn	Thr	Pro	Asn	Gly	Ala	Asn	Val
	1235					1240					1245			
Ser	Thr	Ile	Thr	Val	Asn	Ile	Asn	Lys	Gly	Arg	Leu	Thr	Lys	Ser
	1250					1255					1260			
Phe	Ala	Ser	Asn	Leu	Ala	Asn	Met	Asn	Phe	Leu	Arg	Trp	Val	Asn
	1265					1270					1275			
Phe	Pro	Gln	Asp	Tyr	Thr	Val	Thr	Trp	Thr	Asn	Ala	Lys	Ile	Ala
	1280					1285					1290			
Asn	Arg	Pro	Thr	Asp	Gly	Gly	Leu	Ser	Trp	Ser	Asp	Asp	His	Lys
	1295					1300					1305			
Ser	Leu	Ile	Tyr	Arg	Tyr	Asp	Ala	Thr	Leu	Gly	Thr	Gln	Ile	Thr
	1310					1315					1320			
Thr	Asn	Asp	Ile	Leu	Thr	Met	Leu	Lys	Ala	Thr	Thr	Thr	Val	Pro
	1325					1330					1335			
Gly	Leu	Arg	Asn	Asn	Ile	Thr	Gly	Asn	Glu	Lys	Ser	Gln	Ala	Glu
	1340					1345					1350			
Ala	Gly	Gly	Arg	Pro	Asn	Phe	Arg	Thr	Thr	Gly	Tyr	Ser	Gln	Ser
	1355					1360					1365			
Asn	Ala	Thr	Thr	Asp	Gly	Gln	Arg	Gln	Phe	Thr	Leu	Asn	Gly	Gln
	1370					1375					1380			
Val	Ile	Gln	Val	Leu	Asp	Ile	Ile	Asn	Pro	Ser	Asn	Gly	Tyr	Gly
	1385					1390					1395			
Gly	Gln	Pro	Val	Thr	Asn	Ser	Asn	Thr	Arg	Ala	Asn	His	Ser	Asn
	1400					1405					1410			
Ser	Thr	Val	Val	Asn	Val	Asn	Glu	Pro	Ala	Ala	Asn	Gly	Ala	Gly
	1415					1420					1425			
Ala	Phe	Thr	Ile	Asp	His	Val	Val	Lys	Ser	Asn	Ser	Thr	His	Asn
	1430					1435					1440			
Ala	Ser	Asp	Ala	Val	Tyr	Lys	Ala	Gln	Leu	Tyr	Leu	Thr	Pro	Tyr
	1445					1450					1455			
Gly	Pro	Lys	Gln	Tyr	Val	Glu	His	Leu	Asn	Gln	Asn	Thr	Gly	Asn
	1460					1465					1470			
Thr	Thr	Asp	Ala	Ile	Asn	Ile	Tyr	Phe	Val	Pro	Ser	Asp	Leu	Val
	1475					1480					1485			
Asn	Pro	Thr	Ile	Ser	Val	Gly	Asn	Tyr	Thr	Asn	His	Gln	Val	Phe
	1490					1495					1500			
Ser	Gly	Glu	Thr	Phe	Thr	Asn	Thr	Ile	Thr	Ala	Asn	Asp	Asn	Phe
	1505					1510					1515			
Gly	Val	Gln	Ser	Val	Thr	Val	Pro	Asn	Thr	Ser	Gln	Ile	Thr	Gly
	1520					1525					1530			
Thr	Val	Asp	Asn	Asn	His	Gln	His	Val	Ser	Ala	Thr	Ala	Pro	Asn
	1535					1540					1545			
Val	Thr	Ser	Ala	Thr	Asn	Lys	Thr	Ile	Asn	Leu	Leu	Ala	Thr	Asp
	1550					1555					1560			

-continued

---

Thr Ser	Gly Asn	Thr Ala	Thr	Thr Ser	Phe Asn	Val	Thr Val	Lys	
1565			1570			1575			
Pro Leu	Arg Asp	Lys Tyr	Arg	Val Gly	Thr Ser	Ser	Thr Ala	Ala	
1580			1585			1590			
Asn Pro	Val Arg	Ile Ala	Asn	Ile Ser	Asn Asn	Ala	Thr Val	Ser	
1595			1600			1605			
Gln Ala	Asp Gln	Thr Thr	Ile	Ile Asn	Ser Leu	Thr	Phe Thr	Glu	
1610			1615			1620			
Thr Val	Pro Asn	Arg Ser	Tyr	Ala Arg	Ala Ser	Ala	Asn Glu	Ile	
1625			1630			1635			
Thr Ser	Lys Thr	Val Ser	Asn	Val Ser	Arg Thr	Gly	Asn Asn	Ala	
1640			1645			1650			
Asn Val	Thr Val	Thr Val	Thr	Tyr Gln	Asp Gly	Thr	Thr Ser	Thr	
1655			1660			1665			
Val Thr	Val Pro	Val Lys	His	Val Ile	Pro Glu	Ile	Val Ala	His	
1670			1675			1680			
Ser His	Tyr Thr	Val Gln	Gly	Gln Asp	Phe Pro	Ala	Gly Asn	Gly	
1685			1690			1695			
Ser Ser	Ala Ser	Asp Tyr	Phe	Lys Leu	Ser Asn	Gly	Ser Asp	Ile	
1700			1705			1710			
Ala Asp	Ala Thr	Ile Thr	Trp	Val Ser	Gly Gln	Ala	Pro Asn	Lys	
1715			1720			1725			
Asp Asn	Thr Arg	Ile Gly	Glu	Asp Ile	Thr Val	Thr	Ala His	Ile	
1730			1735			1740			
Leu Ile	Asp Gly	Glu Thr	Thr	Pro Ile	Thr Lys	Thr	Ala Thr	Tyr	
1745			1750			1755			
Lys Val	Val Arg	Thr Val	Pro	Lys His	Val Phe	Glu	Thr Ala	Arg	
1760			1765			1770			
Gly Val	Leu Tyr	Pro Gly	Val	Ser Asp	Met Tyr	Asp	Ala Lys	Gln	
1775			1780			1785			
Tyr Val	Lys Pro	Val Asn	Asn	Ser Trp	Ser Thr	Asn	Ala Gln	His	
1790			1795			1800			
Met Asn	Phe Gln	Phe Val	Gly	Thr Tyr	Gly Pro	Asn	Lys Asp	Val	
1805			1810			1815			
Val Gly	Ile Ser	Thr Arg	Leu	Ile Arg	Val Thr	Tyr	Asp Asn	Arg	
1820			1825			1830			
Gln Thr	Glu Asp	Leu Thr	Ile	Leu Ser	Lys Val	Lys	Pro Asp	Pro	
1835			1840			1845			
Pro Arg	Ile Asp	Ala Asn	Ser	Val Thr	Tyr Lys	Ala	Gly Leu	Thr	
1850			1855			1860			
Asn Gln	Glu Ile	Lys Val	Asn	Asn Val	Leu Asn	Asn	Ser Ser	Val	
1865			1870			1875			
Lys Leu	Phe Lys	Ala Asp	Asn	Thr Pro	Leu Asn	Val	Thr Asn	Ile	
1880			1885			1890			
Thr His	Gly Ser	Gly Phe	Ser	Ser Val	Val Thr	Val	Ser Asp	Ala	
1895			1900			1905			
Leu Pro	Asn Gly	Gly Ile	Lys	Ala Lys	Ser Ser	Ile	Ser Met	Asn	
1910			1915			1920			
Asn Val	Thr Tyr	Thr Thr	Gln	Asp Glu	His Gly	Gln	Val Val	Thr	
1925			1930			1935			
Val Thr	Arg Asn	Glu Ser	Val	Asp Ser	Asn Asp	Ser	Ala Thr	Val	



-continued

Gln Ala Thr Thr Lys Ile Trp	Gln Asn Gly His Ile Asp Ile Thr
2330	2335 2340
Pro Asn Asn Pro Ser Gly His	Leu Ile Asn Pro Thr Gln Ala Met
2345	2350 2355
Asp Ile Ala Tyr Thr Glu Lys	Val Gly Asn Gly Ala Glu His Ser
2360	2365 2370
Lys Thr Ile Asn Val Val Arg	Gly Gln Asn Asn Gln Trp Thr Ile
2375	2380 2385
Ala Asn Lys Pro Asp Tyr Val	Thr Leu Asp Ala Gln Thr Gly Lys
2390	2395 2400
Val Thr Phe Asn Ala Asn Thr	Ile Lys Pro Asn Ser Ser Ile Thr
2405	2410 2415
Ile Thr Pro Lys Ala Gly Thr	Gly His Ser Val Ser Ser Asn Pro
2420	2425 2430
Ser Thr Leu Thr Ala Pro Ala	Ala His Thr Val Asn Thr Thr Glu
2435	2440 2445
Ile Val Lys Asp Tyr Gly Ser	Asn Val Thr Ala Ala Glu Ile Asn
2450	2455 2460
Asn Ala Val Gln Val Ala Asn	Lys Arg Thr Ala Thr Ile Lys Asn
2465	2470 2475
Gly Thr Ala Met Pro Thr Asn	Leu Ala Gly Gly Ser Thr Thr Thr
2480	2485 2490
Ile Pro Val Thr Val Thr Tyr	Asn Asp Gly Ser Thr Glu Glu Val
2495	2500 2505
Gln Glu Ser Ile Phe Thr Lys	Ala Asp Lys Arg Glu Leu Ile Thr
2510	2515 2520
Ala Lys Asn His Leu Asp Asp	Pro Val Ser Thr Glu Gly Lys Lys
2525	2530 2535
Pro Gly Thr Ile Thr Gln Tyr	Asn Asn Ala Met His Asn Ala Gln
2540	2545 2550
Gln Gln Ile Asn Thr Ala Lys	Thr Glu Ala Gln Gln Val Ile Asn
2555	2560 2565
Asn Glu Arg Ala Thr Pro Gln	Gln Val Ser Asp Ala Leu Thr Lys
2570	2575 2580
Val Arg Ala Ala Gln Thr Lys	Ile Asp Gln Ala Lys Ala Leu Leu
2585	2590 2595
Gln Asn Lys Glu Asp Asn Ser	Gln Leu Val Thr Ser Lys Asn Asn
2600	2605 2610
Leu Gln Ser Ser Val Asn Gln	Val Pro Ser Thr Ala Gly Met Thr
2615	2620 2625
Gln Gln Ser Ile Asp Asn Tyr	Asn Ala Lys Lys Arg Glu Ala Glu
2630	2635 2640
Thr Glu Ile Thr Ala Ala Gln	Arg Val Ile Asp Asn Gly Asp Ala
2645	2650 2655
Thr Ala Gln Gln Ile Ser Asp	Glu Lys His Arg Val Asp Asn Ala
2660	2665 2670
Leu Thr Ala Leu Asn Gln Ala	Lys His Asp Leu Thr Ala Asp Thr
2675	2680 2685
His Ala Leu Glu Gln Ala Val	Gln Gln Leu Asn Arg Thr Gly Thr
2690	2695 2700
Thr Thr Gly Lys Lys Pro Ala	Ser Ile Thr Ala Tyr Asn Asn Ser
2705	2710 2715



-continued

---

Ile Arg 2720	Ala Leu Gln Ser 2725	Asp 2725	Leu Thr Ser Ala 2730	Lys 2730	Asn Ser Ala 2730
Asn Ala 2735	Ile Ile Gln Lys 2740	Pro 2740	Ile Arg Thr Val 2745	Gln 2745	Glu Val Gln 2745
Ser Ala 2750	Leu Thr Asn Val 2755	Asn 2755	Arg Val Asn Glu 2760	Arg 2760	Leu Thr Gln 2760
Ala Ile 2765	Asn Gln Leu Val 2770	Pro 2770	Leu Ala Asp Asn 2775	Ser 2775	Ala Leu Lys 2775
Thr Ala 2780	Lys Thr Lys Leu 2785	Asp 2785	Glu Glu Ile Asn 2790	Lys 2790	Ser Val Thr 2790
Thr Asp 2795	Gly Met Thr Gln 2800	Ser 2800	Ser Ile Gln Ala 2805	Tyr 2805	Glu Asn Ala 2805
Lys Arg 2810	Ala Gly Gln Thr 2815	Glu 2815	Ser Thr Asn Ala 2820	Gln 2820	Asn Val Ile 2820
Asn Asn 2825	Gly Asp Ala Thr 2830	Asp 2830	Gln Gln Ile Ala 2835	Ala 2835	Glu Lys Thr 2835
Lys Val 2840	Glu Glu Lys Tyr 2845	Asn 2845	Ser Leu Lys Gln 2850	Ala 2850	Ile Ala Gly 2850
Leu Thr 2855	Pro Asp Leu Ala 2860	Pro 2860	Leu Gln Thr Ala 2865	Lys 2865	Thr Gln Leu 2865
Gln Asn 2870	Asp Ile Asp Gln 2875	Pro 2875	Thr Ser Thr Thr 2880	Gly 2880	Met Thr Ser 2880
Ala Ser 2885	Ile Ala Ala Phe 2890	Asn 2890	Glu Lys Leu Ser 2895	Ala 2895	Ala Arg Thr 2895
Lys Ile 2900	Gln Glu Ile Asp 2905	Arg 2905	Val Leu Ala Ser 2910	His 2910	Pro Asp Val 2910
Ala Thr 2915	Ile Arg Gln Asn 2920	Val 2920	Thr Ala Ala Asn 2925	Ala 2925	Ala Lys Ser 2925
Ala Leu 2930	Asp Gln Ala Arg 2935	Asn 2935	Gly Leu Thr Val 2940	Asp 2940	Lys Ala Pro 2940
Leu Glu 2945	Asn Ala Lys Asn 2950	Gln 2950	Leu Gln His Ser 2955	Ile 2955	Asp Thr Gln 2955
Thr Ser 2960	Thr Thr Gly Met 2965	Thr 2965	Gln Asp Ser Ile 2970	Asn 2970	Ala Tyr Asn 2970
Ala Lys 2975	Leu Thr Ala Ala 2980	Arg 2980	Asn Lys Ile Gln 2985	Gln 2985	Ile Asn Gln 2985
Val Leu 2990	Ala Gly Ser Pro 2995	Thr 2995	Val Glu Gln Ile 3000	Asn 3000	Thr Asn Thr 3000
Ser Thr 3005	Ala Asn Gln Ala 3010	Lys 3010	Ser Asp Leu Asp 3015	His 3015	Ala Arg Gln 3015
Ala Leu 3020	Thr Pro Asp Lys 3025	Ala 3025	Pro Leu Gln Thr 3030	Ala 3030	Lys Thr Gln 3030
Leu Glu 3035	Gln Ser Ile Asn 3040	Gln 3040	Pro Thr Asp Thr 3045	Thr 3045	Gly Met Thr 3045
Thr Ala 3050	Ser Leu Asn Ala 3055	Tyr 3055	Asn Gln Lys Leu 3060	Gln 3060	Ala Ala Arg 3060
Gln Lys 3065	Leu Thr Glu Ile 3070	Asn 3070	Gln Val Leu Asn 3075	Gly 3075	Asn Pro Thr 3075
Val Gln 3080	Asn Ile Asn Asp 3085	Lys 3085	Val Thr Glu Ala 3090	Asn 3090	Gln Ala Lys 3090
Asp Gln 3095	Leu Asn Thr Ala 3100	Arg 3100	Gln Gly Leu Thr 3105	Leu 3105	Asp Arg Gln 3105

-continued

3095	3100	3105
Pro Ala Leu Thr Thr Leu His Gly Ala Ser Asn Leu Asn Gln Ala 3110 3115 3120		
Gln Gln Asn Asn Phe Thr Gln Gln Ile Asn Ala Ala Gln Asn His 3125 3130 3135		
Ala Ala Leu Glu Thr Ile Lys Ser Asn Ile Thr Ala Leu Asn Thr 3140 3145 3150		
Ala Met Thr Lys Leu Lys Asp Ser Val Ala Asp Asn Asn Thr Ile 3155 3160 3165		
Lys Ser Asp Gln Asn Tyr Thr Asp Ala Thr Pro Ala Asn Lys Gln 3170 3175 3180		
Ala Tyr Asp Asn Ala Val Asn Ala Ala Lys Gly Val Ile Gly Glu 3185 3190 3195		
Thr Thr Asn Pro Thr Met Asp Val Asn Thr Val Asn Gln Lys Ala 3200 3205 3210		
Ala Ser Val Lys Ser Thr Lys Asp Ala Leu Asp Gly Gln Gln Asn 3215 3220 3225		
Leu Gln Arg Ala Lys Thr Glu Ala Thr Asn Ala Ile Thr His Ala 3230 3235 3240		
Ser Asp Leu Asn Gln Ala Gln Lys Asn Ala Leu Thr Gln Gln Val 3245 3250 3255		
Asn Ser Ala Gln Asn Val Gln Ala Val Asn Asp Ile Lys Gln Thr 3260 3265 3270		
Thr Gln Ser Leu Asn Thr Ala Met Thr Gly Leu Lys Arg Gly Val 3275 3280 3285		
Ala Asn His Asn Gln Val Val Gln Ser Asp Asn Tyr Val Asn Ala 3290 3295 3300		
Asp Thr Asn Lys Lys Asn Asp Tyr Asn Asn Ala Tyr Asn His Ala 3305 3310 3315		
Asn Asp Ile Ile Asn Gly Asn Ala Gln His Pro Val Ile Thr Pro 3320 3325 3330		
Ser Asp Val Asn Asn Ala Leu Ser Asn Val Thr Ser Lys Glu His 3335 3340 3345		
Ala Leu Asn Gly Glu Ala Lys Leu Asn Ala Ala Lys Gln Glu Ala 3350 3355 3360		
Asn Thr Ala Leu Gly His Leu Asn Asn Leu Asn Asn Ala Gln Arg 3365 3370 3375		
Gln Asn Leu Gln Ser Gln Ile Asn Gly Ala His Gln Ile Asp Ala 3380 3385 3390		
Val Asn Thr Ile Lys Gln Asn Ala Thr Asn Leu Asn Ser Ala Met 3395 3400 3405		
Gly Asn Leu Arg Gln Ala Val Ala Asp Lys Asp Gln Val Lys Arg 3410 3415 3420		
Thr Glu Asp Tyr Ala Asp Ala Asp Thr Ala Lys Gln Asn Ala Tyr 3425 3430 3435		
Asn Ser Ala Val Ser Ser Ala Glu Thr Ile Ile Asn Gln Thr Thr 3440 3445 3450		
Asn Pro Thr Met Ser Val Asp Asp Val Asn Arg Ala Thr Ser Ala 3455 3460 3465		
Val Thr Ser Asn Lys Asn Ala Leu Asn Gly Tyr Glu Lys Leu Ala 3470 3475 3480		

-continued

---

Gln Ser	Lys Thr Asp Ala Ala	Arg Ala Ile Asp Ala	Leu Pro His
3485	3490	3495	
Leu Asn	Asn Ala Gln Lys Ala	Asp Val Lys Ser Lys	Ile Asn Ala
3500	3505	3510	
Ala Ser	Asn Ile Ala Gly Val	Asn Thr Val Lys Gln	Gln Gly Thr
3515	3520	3525	
Asp Leu	Asn Thr Ala Met Gly	Asn Leu Gln Gly Ala	Ile Asn Asp
3530	3535	3540	
Glu Gln	Thr Thr Leu Asn Ser	Gln Asn Tyr Gln Asp	Ala Thr Pro
3545	3550	3555	
Ser Lys	Lys Thr Ala Tyr Thr	Asn Ala Val Gln Ala	Ala Lys Asp
3560	3565	3570	
Ile Leu	Asn Lys Ser Asn Gly	Gln Asn Lys Thr Lys	Asp Gln Val
3575	3580	3585	
Thr Glu	Ala Met Asn Gln Val	Asn Ser Ala Lys Asn	Asn Leu Asp
3590	3595	3600	
Gly Thr	Arg Leu Leu Asp Gln	Ala Lys Gln Thr Ala	Lys Gln Gln
3605	3610	3615	
Leu Asn	Asn Met Thr His Leu	Thr Thr Ala Gln Lys	Thr Asn Leu
3620	3625	3630	
Thr Asn	Gln Ile Asn Ser Gly	Thr Thr Val Ala Gly	Val Gln Thr
3635	3640	3645	
Val Gln	Ser Asn Ala Asn Thr	Leu Asp Gln Ala Met	Asn Thr Leu
3650	3655	3660	
Arg Gln	Ser Ile Ala Asn Lys	Asp Ala Thr Lys Ala	Ser Glu Asp
3665	3670	3675	
Tyr Val	Asp Ala Asn Asn Asp	Lys Gln Thr Ala Tyr	Asn Asn Ala
3680	3685	3690	
Val Ala	Ala Ala Glu Thr Ile	Ile Asn Ala Asn Ser	Asn Pro Glu
3695	3700	3705	
Met Asn	Pro Ser Thr Ile Thr	Gln Lys Ala Glu Gln	Val Asn Ser
3710	3715	3720	
Ser Lys	Thr Ala Leu Asn Gly	Asp Glu Asn Leu Ala	Ala Ala Lys
3725	3730	3735	
Gln Asn	Ala Lys Thr Tyr Leu	Asn Thr Leu Thr Ser	Ile Thr Asp
3740	3745	3750	
Ala Gln	Lys Asn Asn Leu Ile	Ser Gln Ile Thr Ser	Ala Thr Arg
3755	3760	3765	
Val Ser	Gly Val Asp Thr Val	Lys Gln Asn Ala Gln	His Leu Asp
3770	3775	3780	
Gln Ala	Met Ala Ser Leu Gln	Asn Gly Ile Asn Asn	Glu Ser Gln
3785	3790	3795	
Val Lys	Ser Ser Glu Lys Tyr	Arg Asp Ala Asp Thr	Asn Lys Gln
3800	3805	3810	
Gln Glu	Tyr Asp Asn Ala Ile	Thr Ala Ala Lys Ala	Ile Leu Asn
3815	3820	3825	
Lys Ser	Thr Gly Pro Asn Thr	Ala Gln Asn Ala Val	Glu Ala Ala
3830	3835	3840	
Leu Gln	Arg Val Asn Asn Ala	Lys Asp Ala Leu Asn	Gly Asp Ala
3845	3850	3855	
Lys Leu	Ile Ala Ala Gln Asn	Ala Ala Lys Gln His	Leu Gly Thr
3860	3865	3870	

-continued

---

Leu Thr	His Ile	Thr Thr	Ala	Gln Arg	Asn Asp	Leu Thr	Asn Gln		
3875			3880			3885			
Ile Ser	Gln Ala	Thr Asn	Leu	Ala Gly	Val Glu	Ser	Val Lys	Gln	
3890			3895			3900			
Asn Ala	Asn Ser	Leu Asp	Gly	Ala Met	Gly Asn	Leu	Gln Thr	Ala	
3905			3910			3915			
Ile Asn	Asp Lys	Ser Gly	Thr	Leu Ala	Ser Gln	Asn	Phe Leu	Asp	
3920			3925			3930			
Ala Asp	Glu Gln	Lys Arg	Asn	Ala Tyr	Asn Gln	Ala	Val Ser	Ala	
3935			3940			3945			
Ala Glu	Thr Ile	Leu Asn	Lys	Gln Thr	Gly Pro	Asn	Thr Ala	Lys	
3950			3955			3960			
Thr Ala	Val Glu	Gln Ala	Leu	Asn Asn	Val Asn	Asn	Ala Lys	His	
3965			3970			3975			
Ala Leu	Asn Gly	Thr Gln	Asn	Leu Asn	Asn Ala	Lys	Gln Ala	Ala	
3980			3985			3990			
Ile Thr	Ala Ile	Asn Gly	Ala	Ser Asp	Leu Asn	Gln	Lys Gln	Lys	
3995			4000			4005			
Asp Ala	Leu Lys	Ala Gln	Ala	Asn Gly	Ala Gln	Arg	Val Ser	Asn	
4010			4015			4020			
Ala Gln	Asp Val	Gln His	Asn	Ala Thr	Glu Leu	Asn	Thr Ala	Met	
4025			4030			4035			
Gly Thr	Leu Lys	His Ala	Ile	Ala Asp	Lys Thr	Asn	Thr Leu	Ala	
4040			4045			4050			
Ser Ser	Lys Tyr	Val Asn	Ala	Asp Ser	Thr Lys	Gln	Asn Ala	Tyr	
4055			4060			4065			
Thr Thr	Lys Val	Thr Asn	Ala	Glu His	Ile Ile	Ser	Gly Thr	Pro	
4070			4075			4080			
Thr Val	Val Thr	Thr Pro	Ser	Glu Val	Thr Ala	Ala	Ala Asn	Gln	
4085			4090			4095			
Val Asn	Ser Ala	Lys Gln	Glu	Leu Asn	Gly Asp	Glu	Arg Leu	Arg	
4100			4105			4110			
Glu Ala	Lys Gln	Asn Ala	Asn	Thr Ala	Ile Asp	Ala	Leu Thr	Gln	
4115			4120			4125			
Leu Asn	Thr Pro	Gln Lys	Ala	Lys Leu	Lys Glu	Gln	Val Gly	Gln	
4130			4135			4140			
Ala Asn	Arg Leu	Glu Asp	Val	Gln Thr	Val Gln	Thr	Asn Gly	Gln	
4145			4150			4155			
Ala Leu	Asn Asn	Ala Met	Lys	Gly Leu	Arg Asp	Ser	Ile Ala	Asn	
4160			4165			4170			
Glu Thr	Thr Val	Lys Thr	Ser	Gln Asn	Tyr Thr	Asp	Ala Ser	Pro	
4175			4180			4185			
Asn Asn	Gln Ser	Thr Tyr	Asn	Ser Ala	Val Ser	Asn	Ala Lys	Gly	
4190			4195			4200			
Ile Ile	Asn Gln	Thr Asn	Asn	Pro Thr	Met Asp	Thr	Ser Ala	Ile	
4205			4210			4215			
Thr Gln	Ala Thr	Thr Gln	Val	Asn Asn	Ala Lys	Asn	Gly Leu	Asn	
4220			4225			4230			
Gly Ala	Glu Asn	Leu Arg	Asn	Ala Gln	Asn Thr	Ala	Lys Gln	Asn	
4235			4240			4245			
Leu Asn	Thr Leu	Ser His	Leu	Thr Asn	Asn Gln	Lys	Ser Ala	Ile	

-continued

4250	4255	4260
Ser Ser Gln Ile Asp Arg Ala Gly His Val Ser Glu Val Thr Ala 4265	4270	4275
Thr Lys Asn Ala Ala Thr Glu Leu Asn Thr Gln Met Gly Asn Leu 4280	4285	4290
Glu Gln Ala Ile His Asp Gln Asn Thr Val Lys Gln Ser Val Lys 4295	4300	4305
Phe Thr Asp Ala Asp Lys Ala Lys Arg Asp Ala Tyr Thr Asn Ala 4310	4315	4320
Val Ser Arg Ala Glu Ala Ile Leu Asn Lys Thr Gln Gly Ala Asn 4325	4330	4335
Thr Ser Lys Gln Asp Val Glu Ala Ala Ile Gln Asn Val Ser Ser 4340	4345	4350
Ala Lys Asn Ala Leu Asn Gly Asp Gln Asn Val Thr Asn Ala Lys 4355	4360	4365
Asn Ala Ala Lys Asn Ala Leu Asn Asn Leu Thr Ser Ile Asn Asn 4370	4375	4380
Ala Gln Lys Arg Asp Leu Thr Thr Lys Ile Asp Gln Ala Thr Thr 4385	4390	4395
Val Ala Gly Val Glu Ala Val Ser Asn Thr Ser Thr Gln Leu Asn 4400	4405	4410
Thr Ala Met Ala Asn Leu Gln Asn Gly Ile Asn Asp Lys Thr Asn 4415	4420	4425
Thr Leu Ala Ser Glu Asn Tyr His Asp Ala Asp Ser Asp Lys Lys 4430	4435	4440
Thr Ala Tyr Thr Gln Ala Val Thr Asn Ala Glu Asn Ile Leu Asn 4445	4450	4455
Lys Asn Ser Gly Ser Asn Leu Asp Lys Thr Ala Val Glu Asn Ala 4460	4465	4470
Leu Ser Gln Val Ala Asn Ala Lys Gly Ala Leu Asn Gly Asn His 4475	4480	4485
Asn Leu Glu Gln Ala Lys Ser Asn Ala Asn Thr Thr Ile Asn Gly 4490	4495	4500
Leu Gln His Leu Thr Thr Ala Gln Lys Asp Lys Leu Lys Gln Gln 4505	4510	4515
Val Gln Gln Ala Gln Asn Val Ala Gly Val Asp Thr Val Lys Ser 4520	4525	4530
Ser Ala Asn Thr Leu Asn Gly Ala Met Gly Thr Leu Arg Asn Ser 4535	4540	4545
Ile Gln Asp Asn Thr Ala Thr Lys Asn Gly Gln Asn Tyr Leu Asp 4550	4555	4560
Ala Thr Glu Arg Asn Lys Thr Asn Tyr Asn Asn Ala Val Asp Ser 4565	4570	4575
Ala Asn Gly Val Ile Asn Ala Thr Ser Asn Pro Asn Met Asp Ala 4580	4585	4590
Asn Ala Ile Asn Gln Ile Ala Thr Gln Val Thr Ser Thr Lys Asn 4595	4600	4605
Ala Leu Asp Gly Thr His Asn Leu Thr Gln Ala Lys Gln Thr Ala 4610	4615	4620
Thr Asn Ala Ile Asp Gly Ala Thr Asn Leu Asn Lys Ala Gln Lys 4625	4630	4635

-continued

Asp	Ala	Leu	Lys	Ala	Gln	Val	Thr	Ser	Ala	Gln	Arg	Val	Ala	Asn
4640						4645					4650			
Val	Thr	Ser	Ile	Gln	Gln	Thr	Ala	Asn	Glu	Leu	Asn	Thr	Ala	Met
4655						4660					4665			
Gly	Gln	Leu	Gln	His	Gly	Ile	Asp	Asp	Glu	Asn	Ala	Thr	Lys	Gln
4670						4675					4680			
Thr	Gln	Lys	Tyr	Arg	Asp	Ala	Glu	Gln	Ser	Lys	Lys	Thr	Ala	Tyr
4685						4690					4695			
Asp	Gln	Ala	Val	Ala	Ala	Ala	Lys	Ala	Ile	Leu	Asn	Lys	Gln	Thr
4700						4705					4710			
Gly	Ser	Asn	Ser	Asp	Lys	Ala	Ala	Val	Asp	Arg	Ala	Leu	Gln	Gln
4715						4720					4725			
Val	Thr	Ser	Thr	Lys	Asp	Ala	Leu	Asn	Gly	Asp	Ala	Lys	Leu	Ala
4730						4735					4740			
Glu	Ala	Lys	Ala	Ala	Ala	Lys	Gln	Asn	Leu	Gly	Thr	Leu	Asn	His
4745						4750					4755			
Ile	Thr	Asn	Ala	Gln	Arg	Thr	Asp	Leu	Glu	Gly	Gln	Ile	Asn	Gln
4760						4765					4770			
Ala	Thr	Thr	Val	Asp	Gly	Val	Asn	Thr	Val	Lys	Thr	Asn	Ala	Asn
4775						4780					4785			
Thr	Leu	Asp	Gly	Ala	Met	Asn	Ser	Leu	Gln	Gly	Ser	Ile	Asn	Asp
4790						4795					4800			
Lys	Asp	Ala	Thr	Leu	Arg	Asn	Gln	Asn	Tyr	Leu	Asp	Ala	Asp	Glu
4805						4810					4815			
Ser	Lys	Arg	Asn	Ala	Tyr	Thr	Gln	Ala	Val	Thr	Ala	Ala	Glu	Gly
4820						4825					4830			
Ile	Leu	Asn	Lys	Gln	Thr	Gly	Gly	Asn	Thr	Ser	Lys	Ala	Asp	Val
4835						4840					4845			
Asp	Asn	Ala	Leu	Asn	Ala	Val	Thr	Arg	Ala	Lys	Ala	Ala	Leu	Asn
4850						4855					4860			
Gly	Ala	Asp	Asn	Leu	Arg	Asn	Ala	Lys	Thr	Ser	Ala	Thr	Asn	Thr
4865						4870					4875			
Ile	Asp	Gly	Leu	Pro	Asn	Leu	Thr	Gln	Leu	Gln	Lys	Asp	Asn	Leu
4880						4885					4890			
Lys	His	Gln	Val	Glu	Gln	Ala	Gln	Asn	Val	Ala	Gly	Val	Asn	Gly
4895						4900					4905			
Val	Lys	Asp	Lys	Gly	Asn	Thr	Leu	Asn	Thr	Ala	Met	Gly	Ala	Leu
4910						4915					4920			
Arg	Thr	Ser	Ile	Gln	Asn	Asp	Asn	Thr	Thr	Lys	Thr	Ser	Gln	Asn
4925						4930					4935			
Tyr	Leu	Asp	Ala	Ser	Asp	Ser	Asn	Lys	Asn	Asn	Tyr	Asn	Thr	Ala
4940						4945					4950			
Val	Asn	Asn	Ala	Asn	Gly	Val	Ile	Asn	Ala	Thr	Asn	Asn	Pro	Asn
4955						4960					4965			
Met	Asp	Ala	Asn	Ala	Ile	Asn	Gly	Met	Ala	Asn	Gln	Val	Asn	Thr
4970						4975					4980			
Thr	Lys	Ala	Ala	Leu	Asn	Gly	Ala	Gln	Asn	Leu	Ala	Gln	Ala	Lys
4985						4990					4995			
Thr	Asn	Ala	Thr	Asn	Thr	Ile	Asn	Asn	Ala	His	Asp	Leu	Asn	Gln
5000						5005					5010			
Lys	Gln	Lys	Asp	Ala	Leu	Lys	Thr	Gln	Val	Asn	Asn	Ala	Gln	Arg
5015						5020					5025			

-continued

---

Val	Ser	Asp	Ala	Asn	Asn	Val	Gln	His	Thr	Ala	Thr	Glu	Leu	Asn
5030						5035						5040		
Ser	Ala	Met	Thr	Ala	Leu	Lys	Ala	Ala	Ile	Ala	Asp	Lys	Glu	Arg
5045						5050						5055		
Thr	Lys	Ala	Ser	Gly	Asn	Tyr	Val	Asn	Ala	Asp	Gln	Glu	Lys	Arg
5060						5065						5070		
Gln	Ala	Tyr	Asp	Ser	Lys	Val	Thr	Asn	Ala	Glu	Asn	Ile	Ile	Ser
5075						5080						5085		
Gly	Thr	Pro	Asn	Ala	Thr	Leu	Thr	Val	Asn	Asp	Val	Asn	Ser	Ala
5090						5095						5100		
Ala	Ser	Gln	Val	Asn	Ala	Ala	Lys	Thr	Ala	Leu	Asn	Gly	Asp	Asn
5105						5110						5115		
Asn	Leu	Arg	Val	Ala	Lys	Glu	His	Ala	Asn	Asn	Thr	Ile	Asp	Gly
5120						5125						5130		
Leu	Ala	Gln	Leu	Asn	Asn	Ala	Gln	Lys	Ala	Lys	Leu	Lys	Glu	Gln
5135						5140						5145		
Val	Gln	Ser	Ala	Thr	Thr	Leu	Asp	Gly	Val	Gln	Thr	Val	Lys	Asn
5150						5155						5160		
Ser	Ser	Gln	Thr	Leu	Asn	Thr	Ala	Met	Lys	Gly	Leu	Arg	Asp	Ser
5165						5170						5175		
Ile	Ala	Asn	Glu	Ala	Thr	Ile	Lys	Ala	Gly	Gln	Asn	Tyr	Thr	Asp
5180						5185						5190		
Ala	Ser	Pro	Asn	Asn	Arg	Asn	Glu	Tyr	Asp	Ser	Ala	Val	Thr	Ala
5195						5200						5205		
Ala	Lys	Ala	Ile	Ile	Asn	Gln	Thr	Ser	Asn	Pro	Thr	Met	Glu	Pro
5210						5215						5220		
Asn	Thr	Ile	Thr	Gln	Val	Thr	Ser	Gln	Val	Thr	Thr	Lys	Glu	Gln
5225						5230						5235		
Ala	Leu	Asn	Gly	Ala	Arg	Asn	Leu	Ala	Gln	Ala	Lys	Thr	Thr	Ala
5240						5245						5250		
Lys	Asn	Asn	Leu	Asn	Asn	Leu	Thr	Ser	Ile	Asn	Asn	Ala	Gln	Lys
5255						5260						5265		
Asp	Ala	Leu	Thr	Arg	Ser	Ile	Asp	Gly	Ala	Thr	Thr	Val	Ala	Gly
5270						5275						5280		
Val	Asn	Gln	Glu	Thr	Ala	Lys	Ala	Thr	Glu	Leu	Asn	Asn	Ala	Met
5285						5290						5295		
His	Ser	Leu	Gln	Asn	Gly	Ile	Asn	Asp	Glu	Thr	Gln	Thr	Lys	Gln
5300						5305						5310		
Thr	Gln	Lys	Tyr	Leu	Asp	Ala	Glu	Pro	Ser	Lys	Lys	Ser	Ala	Tyr
5315						5320						5325		
Asp	Gln	Ala	Val	Asn	Ala	Ala	Lys	Ala	Ile	Leu	Thr	Lys	Ala	Ser
5330						5335						5340		
Gly	Gln	Asn	Val	Asp	Lys	Ala	Ala	Val	Glu	Gln	Ala	Leu	Gln	Asn
5345						5350						5355		
Val	Asn	Ser	Thr	Lys	Thr	Ala	Leu	Asn	Gly	Asp	Ala	Lys	Leu	Asn
5360						5365						5370		
Glu	Ala	Lys	Ala	Ala	Ala	Lys	Gln	Thr	Leu	Gly	Thr	Leu	Thr	His
5375						5380						5385		
Ile	Asn	Asn	Ala	Gln	Arg	Thr	Ala	Leu	Asp	Asn	Glu	Ile	Thr	Gln
5390						5395						5400		
Ala	Thr	Asn	Val	Glu	Gly	Val	Asn	Thr	Val	Lys	Ala	Lys	Ala	Gln

-continued

---

5405	5410	5415
Gln Leu Asp Gly Ala Met 5420	Gly Gln Leu Glu Thr 5425	Ser Ile Arg Asp 5430
Lys Asp Thr Thr Leu Gln 5435	Ser Gln Asn Tyr Gln 5440	Asp Ala Asp Asp 5445
Ala Lys Arg Thr Ala Tyr 5450	Ser Gln Ala Val Asn 5455	Ala Ala Ala Thr 5460
Ile Leu Asn Lys Thr Ala 5465	Gly Gly Asn Thr Pro 5470	Lys Ala Asp Val 5475
Glu Arg Ala Met Gln Ala 5480	Val Thr Gln Ala Asn 5485	Thr Ala Leu Asn 5490
Gly Ile Gln Asn Leu Asp 5495	Arg Ala Lys Gln Ala 5500	Ala Asn Thr Ala 5505
Ile Thr Asn Ala Ser Asp 5510	Leu Asn Thr Lys Gln 5515	Lys Glu Ala Leu 5520
Lys Ala Gln Val Thr Ser 5525	Ala Gly Arg Val Ser 5530	Ala Ala Asn Gly 5535
Val Glu His Thr Ala Thr 5540	Glu Leu Asn Thr Ala 5545	Met Thr Ala Leu 5550
Lys Arg Ala Ile Ala Asp 5555	Lys Ala Glu Thr Lys 5560	Ala Ser Gly Asn 5565
Tyr Val Asn Ala Asp Ala 5570	Asn Lys Arg Gln Ala 5575	Tyr Asp Glu Lys 5580
Val Thr Ala Ala Glu Asn 5585	Ile Val Ser Gly Thr 5590	Pro Thr Pro Thr 5595
Leu Thr Pro Ala Asp Val 5600	Thr Asn Ala Ala Thr 5605	Gln Val Thr Asn 5610
Ala Lys Thr Gln Leu Asn 5615	Gly Asn His Asn Leu 5620	Glu Val Ala Lys 5625
Gln Asn Ala Asn Thr Ala 5630	Ile Asp Gly Leu Thr 5635	Ser Leu Asn Gly 5640
Pro Gln Lys Ala Lys Leu 5645	Lys Glu Gln Val Gly 5650	Gln Ala Thr Thr 5655
Leu Pro Asn Val Gln Thr 5660	Val Arg Asp Asn Ala 5665	Gln Thr Leu Asn 5670
Thr Ala Met Lys Gly Leu 5675	Arg Asp Ser Ile Ala 5680	Asn Glu Ala Thr 5685
Ile Lys Ala Gly Gln Asn 5690	Tyr Thr Asp Ala Ser 5695	Gln Asn Lys Gln 5700
Thr Asp Tyr Asn Ser Ala 5705	Val Thr Ala Ala Lys 5710	Ala Ile Ile Gly 5715
Gln Thr Thr Ser Pro Ser 5720	Met Asn Ala Gln Glu 5725	Ile Asn Gln Ala 5730
Lys Asp Gln Val Thr Ala 5735	Lys Gln Gln Ala Leu 5740	Asn Gly Gln Glu 5745
Asn Leu Arg Thr Ala Gln 5750	Thr Asn Ala Lys Gln 5755	His Leu Asn Gly 5760
Leu Ser Asp Leu Thr Asp 5765	Ala Gln Lys Asp Ala 5770	Val Lys Arg Gln 5775
Ile Glu Gly Ala Thr His 5780	Val Asn Glu Val Thr 5785	Gln Ala Gln Asn 5790





-continued

---

Lys	His	Ala	Leu	Ala	Glu	Lys	Asp	Asn	Thr	Lys	Arg	Ser	Val	Asn
6185						6190					6195			
Tyr	Thr	Asp	Ala	Asp	Gln	Pro	Lys	Gln	Gln	Ala	Tyr	Asp	Thr	Ala
6200						6205					6210			
Val	Thr	Gln	Ala	Glu	Ala	Ile	Thr	Asn	Ala	Asn	Gly	Ser	Asn	Ala
6215						6220					6225			
Asn	Glu	Thr	Gln	Val	Gln	Ala	Ala	Leu	Asn	Gln	Leu	Asn	Gln	Ala
6230						6235					6240			
Lys	Asn	Asp	Leu	Asn	Gly	Asp	Asn	Lys	Val	Ala	Gln	Ala	Lys	Glu
6245						6250					6255			
Ser	Ala	Lys	Arg	Ala	Leu	Ala	Ser	Tyr	Ser	Asn	Leu	Asn	Asn	Ala
6260						6265					6270			
Gln	Ser	Thr	Ala	Ala	Ile	Ser	Gln	Ile	Asp	Asn	Ala	Thr	Thr	Val
6275						6280					6285			
Ala	Gly	Val	Thr	Ala	Ala	Gln	Asn	Thr	Ala	Asn	Glu	Leu	Asn	Thr
6290						6295					6300			
Ala	Met	Gly	Gln	Leu	Gln	Asn	Gly	Ile	Asn	Asp	Gln	Asn	Thr	Val
6305						6310					6315			
Lys	Gln	Gln	Val	Asn	Phe	Thr	Asp	Ala	Asp	Gln	Gly	Lys	Lys	Asp
6320						6325					6330			
Ala	Tyr	Thr	Asn	Ala	Val	Thr	Asn	Ala	Gln	Gly	Ile	Leu	Asp	Lys
6335						6340					6345			
Ala	His	Gly	Gln	Asn	Met	Thr	Lys	Ala	Gln	Val	Glu	Ala	Ala	Leu
6350						6355					6360			
Asn	Gln	Val	Thr	Thr	Ala	Lys	Asn	Ala	Leu	Asn	Gly	Asp	Ala	Asn
6365						6370					6375			
Val	Arg	Gln	Ala	Lys	Ser	Asp	Ala	Lys	Ala	Asn	Leu	Gly	Thr	Leu
6380						6385					6390			
Thr	His	Leu	Asn	Asn	Ala	Gln	Lys	Gln	Asp	Leu	Thr	Ser	Gln	Ile
6395						6400					6405			
Glu	Gly	Ala	Thr	Thr	Val	Asn	Gly	Val	Asn	Gly	Val	Lys	Thr	Lys
6410						6415					6420			
Ala	Gln	Asp	Leu	Asp	Gly	Ala	Met	Gln	Arg	Leu	Gln	Ser	Ala	Ile
6425						6430					6435			
Ala	Asn	Lys	Asp	Gln	Thr	Lys	Ala	Ser	Glu	Asn	Tyr	Ile	Asp	Ala
6440						6445					6450			
Asp	Pro	Thr	Lys	Lys	Thr	Ala	Phe	Asp	Asn	Ala	Ile	Thr	Gln	Ala
6455						6460					6465			
Glu	Ser	Tyr	Leu	Asn	Lys	Asp	His	Gly	Ala	Asn	Lys	Asp	Lys	Gln
6470						6475					6480			
Ala	Val	Glu	Gln	Ala	Ile	Gln	Ser	Val	Thr	Ser	Thr	Glu	Asn	Ala
6485						6490					6495			
Leu	Asn	Gly	Asp	Ala	Asn	Leu	Gln	Arg	Ala	Lys	Thr	Glu	Ala	Ile
6500						6505					6510			
Gln	Ala	Ile	Asp	Asn	Leu	Thr	His	Leu	Asn	Thr	Pro	Gln	Lys	Thr
6515						6520					6525			
Ala	Leu	Lys	Gln	Gln	Val	Asn	Ala	Ala	Gln	Arg	Val	Ser	Gly	Val
6530						6535					6540			
Thr	Asp	Leu	Lys	Asn	Ser	Ala	Thr	Ser	Leu	Asn	Asn	Ala	Met	Asp
6545						6550					6555			
Gln	Leu	Lys	Gln	Ala	Ile	Ala	Asp	His	Asp	Thr	Ile	Val	Ala	Ser

-continued

6560	6565	6570
Gly Asn Tyr Thr Asn Ala Ser Pro Asp Lys Gln Gly Ala Tyr Thr 6575 6580		6585
Asp Ala Tyr Asn Ala Ala Lys Asn Ile Val Asn Gly Ser Pro Asn 6590 6595		6600
Val Ile Thr Asn Ala Ala Asp Val Thr Ala Ala Thr Gln Arg Val 6605 6610		6615
Asn Asn Ala Glu Thr Gly Leu Asn Gly Asp Thr Asn Leu Ala Thr 6620 6625		6630
Ala Lys Gln Gln Ala Lys Asp Ala Leu Arg Gln Met Thr His Leu 6635 6640		6645
Ser Asp Ala Gln Lys Gln Ser Ile Thr Gly Gln Ile Asp Ser Ala 6650 6655		6660
Thr Gln Val Thr Gly Val Gln Ser Val Lys Asp Asn Ala Thr Asn 6665 6670		6675
Leu Asp Asn Ala Met Asn Gln Leu Arg Asn Ser Ile Ala Asn Lys 6680 6685		6690
Asp Asp Val Lys Ala Ser Gln Pro Tyr Val Asp Ala Asp Arg Asp 6695 6700		6705
Lys Gln Asn Ala Tyr Asn Thr Ala Val Thr Asn Ala Glu Asn Ile 6710 6715		6720
Ile Asn Ala Thr Ser Gln Pro Thr Leu Asp Pro Ser Ala Val Thr 6725 6730		6735
Gln Ala Ala Asn Gln Val Ser Thr Asn Lys Thr Ala Leu Asn Gly 6740 6745		6750
Ala Gln Asn Leu Ala Asn Lys Lys Gln Glu Thr Thr Ala Asn Ile 6755 6760		6765
Asn Gln Leu Ser His Leu Asn Asn Ala Gln Lys Gln Asp Leu Asn 6770 6775		6780
Thr Gln Val Thr Asn Ala Pro Asn Ile Ser Thr Val Asn Gln Val 6785 6790		6795
Lys Thr Lys Ala Glu Gln Leu Asp Gln Ala Met Glu Arg Leu Ile 6800 6805		6810
Asn Gly Ile Gln Asp Lys Asp Gln Val Lys Gln Ser Val Asn Phe 6815 6820		6825
Thr Asp Ala Asp Pro Glu Lys Gln Thr Ala Tyr Asn Asn Ala Val 6830 6835		6840
Thr Ala Ala Glu Asn Ile Ile Asn Gln Ala Asn Gly Thr Asn Ala 6845 6850		6855
Asn Gln Ser Gln Val Glu Ala Ala Leu Ser Thr Val Thr Thr Thr 6860 6865		6870
Lys Gln Ala Leu Asn Gly Asp Arg Lys Val Thr Asp Ala Lys Asn 6875 6880		6885
Asn Ala Asn Gln Thr Leu Ser Thr Leu Asp Asn Leu Asn Asn Ala 6890 6895		6900
Gln Lys Gly Ala Val Thr Gly Asn Ile Asn Gln Ala His Thr Val 6905 6910		6915
Ala Glu Val Thr Gln Ala Ile Gln Thr Ala Gln Glu Leu Asn Thr 6920 6925		6930
Ala Met Gly Asn Leu Lys Asn Ser Leu Asn Asp Lys Asp Thr Thr 6935 6940		6945

-continued

---

Leu Gly	Ser Gln Asn Phe	Ala Asp Ala Asp Pro	Glu Lys Lys Asn	
6950		6955	6960	
Ala Tyr	Asn Glu Ala Val	His Asn Ala Glu Asn	Ile Leu Asn Lys	
6965		6970	6975	
Ser Thr	Gly Thr Asn Val	Pro Lys Asp Gln Val	Glu Ala Ala Met	
6980		6985	6990	
Asn Gln	Val Asn Ala Thr	Lys Ala Ala Leu Asn	Gly Thr Gln Asn	
6995		7000	7005	
Leu Glu	Lys Ala Lys Gln	His Ala Asn Thr Ala	Ile Asp Gly Leu	
7010		7015	7020	
Ser His	Leu Thr Asn Ala	Gln Lys Glu Ala Leu	Lys Gln Leu Val	
7025		7030	7035	
Gln Gln	Ser Thr Thr Val	Ala Glu Ala Gln Gly	Asn Glu Gln Lys	
7040		7045	7050	
Ala Asn	Asn Val Asp Ala	Ala Met Asp Lys Leu	Arg Gln Ser Ile	
7055		7060	7065	
Ala Asp	Asn Ala Thr Thr	Lys Gln Asn Gln Asn	Tyr Thr Asp Ala	
7070		7075	7080	
Ser Gln	Asn Lys Lys Asp	Ala Tyr Asn Asn Ala	Val Thr Thr Ala	
7085		7090	7095	
Gln Gly	Ile Ile Asp Gln	Thr Thr Ser Pro Thr	Leu Asp Pro Thr	
7100		7105	7110	
Val Ile	Asn Gln Ala Ala	Gly Gln Val Ser Thr	Thr Lys Asn Ala	
7115		7120	7125	
Leu Asn	Gly Asn Glu Asn	Leu Glu Ala Ala Lys	Gln Gln Ala Ser	
7130		7135	7140	
Gln Ser	Leu Gly Ser Leu	Asp Asn Leu Asn Asn	Ala Gln Lys Gln	
7145		7150	7155	
Thr Val	Thr Asp Gln Ile	Asn Gly Ala His Thr	Val Asp Glu Ala	
7160		7165	7170	
Asn Gln	Ile Lys Gln Asn	Ala Gln Asn Leu Asn	Thr Ala Met Gly	
7175		7180	7185	
Asn Leu	Lys Gln Ala Ile	Ala Asp Lys Asp Ala	Thr Lys Ala Thr	
7190		7195	7200	
Val Asn	Phe Thr Asp Ala	Asp Gln Ala Lys Gln	Gln Ala Tyr Asn	
7205		7210	7215	
Thr Ala	Val Thr Asn Ala	Glu Asn Ile Ser Lys	Ala Asn Gly Asn	
7220		7225	7230	
Ala Thr	Gln Ala Glu Val	Glu Gln Ala Ile Lys	Gln Val Asn Ala	
7235		7240	7245	
Ala Lys	Gln Ala Leu Asn	Gly Asn Ala Asn Val	Gln His Ala Lys	
7250		7255	7260	
Asp Glu	Ala Thr Ala Leu	Ile Asn Ser Ser Asn	Asp Leu Asn Gln	
7265		7270	7275	
Ala Gln	Lys Asp Ala Leu	Lys Gln Gln Val Gln	Asn Ala Thr Thr	
7280		7285	7290	
Val Ala	Gly Val Asn Asn	Val Lys Gln Thr Ala	Gln Glu Leu Asn	
7295		7300	7305	
Asn Ala	Met Thr Gln Leu	Lys Gln Gly Ile Ala	Asp Lys Glu Gln	
7310		7315	7320	
Thr Lys	Ala Asp Gly Asn	Phe Val Asn Ala Asp	Pro Asp Lys Gln	
7325		7330	7335	

-continued

---

Asn Ala	Tyr Asn	Gln Ala	Val	Ala Lys	Ala Glu	Ala	Leu Ile	Ser	
7340			7345			7350			
Ala Thr	Pro Asp	Val Val	Val	Thr Pro	Ser Glu	Ile	Thr Ala	Ala	
7355			7360			7365			
Leu Asn	Lys Val	Thr Gln	Ala	Lys Asn	Asp Leu	Asn	Gly Asn	Thr	
7370			7375			7380			
Asn Leu	Ala Thr	Ala Lys	Gln	Asn Val	Gln His	Ala	Ile Asp	Gln	
7385			7390			7395			
Leu Pro	Asn Leu	Asn Gln	Ala	Gln Arg	Asp Glu	Tyr	Ser Lys	Gln	
7400			7405			7410			
Ile Thr	Gln Ala	Thr Leu	Val	Pro Asn	Val Asn	Ala	Ile Gln	Gln	
7415			7420			7425			
Ala Ala	Thr Thr	Leu Asn	Asp	Ala Met	Thr Gln	Leu	Lys Gln	Gly	
7430			7435			7440			
Ile Ala	Asn Lys	Ala Gln	Ile	Lys Gly	Ser Glu	Asn	Tyr His	Asp	
7445			7450			7455			
Ala Asp	Thr Asp	Lys Gln	Thr	Ala Tyr	Asp Asn	Ala	Val Thr	Lys	
7460			7465			7470			
Ala Glu	Glu Leu	Leu Lys	Gln	Thr Thr	Asn Pro	Thr	Met Asp	Pro	
7475			7480			7485			
Asn Thr	Ile Gln	Gln Ala	Leu	Thr Lys	Val Asn	Asp	Thr Asn	Gln	
7490			7495			7500			
Ala Leu	Asn Gly	Asn Gln	Lys	Leu Ala	Asp Ala	Lys	Gln Asp	Ala	
7505			7510			7515			
Lys Thr	Thr Leu	Gly Thr	Leu	Asp His	Leu Asn	Asp	Ala Gln	Lys	
7520			7525			7530			
Gln Ala	Leu Thr	Thr Gln	Val	Glu Gln	Ala Pro	Asp	Ile Ala	Thr	
7535			7540			7545			
Val Asn	Asn Val	Lys Gln	Asn	Ala Gln	Asn Leu	Asn	Asn Ala	Met	
7550			7555			7560			
Thr Asn	Leu Asn	Asn Ala	Leu	Gln Asp	Lys Thr	Glu	Thr Leu	Asn	
7565			7570			7575			
Ser Ile	Asn Phe	Thr Asp	Ala	Asp Gln	Ala Lys	Lys	Asp Ala	Tyr	
7580			7585			7590			
Thr Asn	Ala Val	Ser His	Ala	Glu Gly	Ile Leu	Ser	Lys Ala	Asn	
7595			7600			7605			
Gly Ser	Asn Ala	Ser Gln	Thr	Glu Val	Glu Gln	Ala	Met Gln	Arg	
7610			7615			7620			
Val Asn	Glu Ala	Lys Gln	Ala	Leu Asn	Gly Asn	Asp	Asn Val	Gln	
7625			7630			7635			
Arg Ala	Lys Asp	Ala Ala	Lys	Gln Val	Ile Thr	Asn	Ala Asn	Asp	
7640			7645			7650			
Leu Asn	Gln Ala	Gln Lys	Asp	Ala Leu	Lys Gln	Gln	Val Asp	Ala	
7655			7660			7665			
Ala Gln	Thr Val	Ala Asn	Val	Asn Thr	Ile Lys	Gln	Thr Ala	Gln	
7670			7675			7680			
Asp Leu	Asn Gln	Ala Met	Thr	Gln Leu	Lys Gln	Gly	Ile Ala	Asp	
7685			7690			7695			
Lys Asp	Gln Thr	Lys Ala	Asn	Gly Asn	Phe Val	Asn	Ala Asp	Thr	
7700			7705			7710			
Asp Lys	Gln Asn	Ala Tyr	Asn	Asn Ala	Val Ala	His	Ala Glu	Gln	

-continued

---

7715		7720		7725
Ile Ile Ser Gly Thr Pro	Asn Ala Asn Val Asp	Pro Gln Gln Val		
7730	7735	7740		
Ala Gln Ala Leu Gln Gln	Val Asn Gln Ala Lys Gly	Asp Leu Asn		
7745	7750	7755		
Gly Asn His Asn Leu Gln	Val Ala Lys Asp Asn Ala	Asn Thr Ala		
7760	7765	7770		
Ile Asp Gln Leu Pro Asn	Leu Asn Gln Pro Gln Lys	Thr Ala Leu		
7775	7780	7785		
Lys Asp Gln Val Ser His	Ala Glu Leu Val Thr Gly	Val Asn Ala		
7790	7795	7800		
Ile Lys Gln Asn Ala Asp	Ala Leu Asn Asn Ala Met	Gly Thr Leu		
7805	7810	7815		
Lys Gln Gln Ile Gln Ala	Asn Ser Gln Val Pro Gln	Ser Val Asp		
7820	7825	7830		
Phe Thr Gln Ala Asp Gln	Asp Lys Gln Gln Ala Tyr	Asn Asn Ala		
7835	7840	7845		
Ala Asn Gln Ala Gln Gln	Ile Ala Asn Gly Ile Pro	Thr Pro Val		
7850	7855	7860		
Leu Thr Pro Asp Thr Val	Thr Gln Ala Val Thr Thr	Met Asn Gln		
7865	7870	7875		
Ala Lys Asp Ala Leu Asn	Gly Asp Glu Lys Leu Ala	Gln Ala Lys		
7880	7885	7890		
Gln Glu Ala Leu Ala Asn	Leu Asp Thr Leu Arg Asp	Leu Asn Gln		
7895	7900	7905		
Pro Gln Arg Asp Ala Leu	Arg Asn Gln Ile Asn Gln	Ala Gln Ala		
7910	7915	7920		
Leu Ala Thr Val Glu Gln	Thr Lys Gln Asn Ala Gln	Asn Val Asn		
7925	7930	7935		
Thr Ala Met Ser Asn Leu	Lys Gln Gly Ile Ala Asn	Lys Asp Thr		
7940	7945	7950		
Val Lys Ala Ser Glu Asn	Tyr His Asp Ala Asp Ala	Asp Lys Gln		
7955	7960	7965		
Thr Ala Tyr Thr Asn Ala	Val Ser Gln Ala Glu Gly	Ile Ile Asn		
7970	7975	7980		
Gln Thr Thr Asn Pro Thr	Leu Asn Pro Asp Glu Ile	Thr Arg Ala		
7985	7990	7995		
Leu Thr Gln Val Thr Asp	Ala Lys Asn Gly Leu Asn	Gly Glu Ala		
8000	8005	8010		
Lys Leu Ala Thr Glu Lys	Gln Asn Ala Lys Asp Ala	Val Ser Gly		
8015	8020	8025		
Met Thr His Leu Asn Asp	Ala Gln Lys Gln Ala Leu	Lys Gly Gln		
8030	8035	8040		
Ile Asp Gln Ser Pro Glu	Ile Ala Thr Val Asn Gln	Val Lys Gln		
8045	8050	8055		
Thr Ala Thr Ser Leu Asp	Gln Ala Met Asp Gln Leu	Ser Gln Ala		
8060	8065	8070		
Ile Asn Asp Lys Ala Gln	Thr Leu Ala Asp Gly Asn	Tyr Leu Asn		
8075	8080	8085		
Ala Asp Pro Asp Lys Gln	Asn Ala Tyr Lys Gln Ala	Val Ala Lys		
8090	8095	8100		

-continued

Ala	Glu	Ala	Leu	Leu	Asn	Lys	Gln	Ser	Gly	Thr	Asn	Glu	Val	Gln
	8105					8110					8115			
Ala	Gln	Val	Glu	Ser	Ile	Thr	Asn	Glu	Val	Asn	Ala	Ala	Lys	Gln
	8120					8125					8130			
Ala	Leu	Asn	Gly	Asn	Asp	Asn	Leu	Ala	Asn	Ala	Lys	Gln	Gln	Ala
	8135					8140					8145			
Lys	Gln	Gln	Leu	Ala	Asn	Leu	Thr	His	Leu	Asn	Asp	Ala	Gln	Lys
	8150					8155					8160			
Gln	Ser	Phe	Glu	Ser	Gln	Ile	Thr	Gln	Ala	Pro	Leu	Val	Thr	Asp
	8165					8170					8175			
Val	Thr	Thr	Ile	Asn	Gln	Lys	Ala	Gln	Thr	Leu	Asp	His	Ala	Met
	8180					8185					8190			
Glu	Leu	Leu	Arg	Asn	Ser	Val	Ala	Asp	Asn	Gln	Thr	Thr	Leu	Ala
	8195					8200					8205			
Ser	Glu	Asp	Tyr	His	Asp	Ala	Thr	Ala	Gln	Arg	Gln	Asn	Asp	Tyr
	8210					8215					8220			
Asn	Gln	Ala	Val	Thr	Ala	Ala	Asn	Asn	Ile	Ile	Asn	Gln	Thr	Thr
	8225					8230					8235			
Ser	Pro	Thr	Met	Asn	Pro	Asp	Asp	Val	Asn	Gly	Ala	Thr	Thr	Gln
	8240					8245					8250			
Val	Asn	Asn	Thr	Lys	Val	Ala	Leu	Asp	Gly	Asp	Glu	Asn	Leu	Ala
	8255					8260					8265			
Ala	Ala	Lys	Gln	Gln	Ala	Asn	Asn	Arg	Leu	Asp	Gln	Leu	Asp	His
	8270					8275					8280			
Leu	Asn	Asn	Ala	Gln	Lys	Gln	Gln	Leu	Gln	Ser	Gln	Ile	Thr	Gln
	8285					8290					8295			
Ser	Ser	Asp	Ile	Ala	Ala	Val	Asn	Gly	His	Lys	Gln	Thr	Ala	Glu
	8300					8305					8310			
Ser	Leu	Asn	Thr	Ala	Met	Gly	Asn	Leu	Ile	Asn	Ala	Ile	Ala	Asp
	8315					8320					8325			
His	Gln	Ala	Val	Glu	Gln	Arg	Gly	Asn	Phe	Ile	Asn	Ala	Asp	Thr
	8330					8335					8340			
Asp	Lys	Gln	Thr	Ala	Tyr	Asn	Thr	Ala	Val	Asn	Glu	Ala	Ala	Ala
	8345					8350					8355			
Met	Ile	Asn	Lys	Gln	Thr	Gly	Gln	Asn	Ala	Asn	Gln	Thr	Glu	Val
	8360					8365					8370			
Glu	Gln	Ala	Ile	Thr	Lys	Val	Gln	Thr	Thr	Leu	Gln	Ala	Leu	Asn
	8375					8380					8385			
Gly	Asp	His	Asn	Leu	Gln	Val	Ala	Lys	Thr	Asn	Ala	Thr	Gln	Ala
	8390					8395					8400			
Ile	Asp	Ala	Leu	Thr	Ser	Leu	Asn	Asp	Pro	Gln	Lys	Thr	Ala	Leu
	8405					8410					8415			
Lys	Asp	Gln	Val	Thr	Ala	Ala	Thr	Leu	Val	Thr	Ala	Val	His	Gln
	8420					8425					8430			
Ile	Glu	Gln	Asn	Ala	Asn	Thr	Leu	Asn	Gln	Ala	Met	His	Gly	Leu
	8435					8440					8445			
Arg	Gln	Ser	Ile	Gln	Asp	Asn	Ala	Ala	Thr	Lys	Ala	Asn	Ser	Lys
	8450					8455					8460			
Tyr	Ile	Asn	Glu	Asp	Gln	Pro	Glu	Gln	Gln	Asn	Tyr	Asp	Gln	Ala
	8465					8470					8475			
Val	Gln	Ala	Ala	Asn	Asn	Ile	Ile	Asn	Glu	Gln	Thr	Ala	Thr	Leu
	8480					8485					8490			

-continued

---

Asp	Asn	Asn	Ala	Ile	Asn	Gln	Ala	Ala	Thr	Thr	Val	Asn	Thr	Thr
8495						8500						8505		
Lys	Ala	Ala	Leu	His	Gly	Asp	Val	Lys	Leu	Gln	Asn	Asp	Lys	Asp
8510						8515					8520			
His	Ala	Lys	Gln	Thr	Val	Ser	Gln	Leu	Ala	His	Leu	Asn	Asn	Ala
8525						8530					8535			
Gln	Lys	His	Met	Glu	Asp	Thr	Leu	Ile	Asp	Ser	Glu	Thr	Thr	Arg
8540						8545					8550			
Thr	Ala	Val	Lys	Gln	Asp	Leu	Thr	Glu	Ala	Gln	Ala	Leu	Asp	Gln
8555						8560					8565			
Leu	Met	Asp	Ala	Leu	Gln	Gln	Ser	Ile	Ala	Asp	Lys	Asp	Ala	Thr
8570						8575					8580			
Arg	Ala	Ser	Ser	Ala	Tyr	Val	Asn	Ala	Glu	Pro	Asn	Lys	Lys	Gln
8585						8590					8595			
Ser	Tyr	Asp	Glu	Ala	Val	Gln	Asn	Ala	Glu	Ser	Ile	Ile	Ala	Gly
8600						8605					8610			
Leu	Asn	Asn	Pro	Thr	Ile	Asn	Lys	Gly	Asn	Val	Ser	Ser	Ala	Thr
8615						8620					8625			
Gln	Ala	Val	Ile	Ser	Ser	Lys	Asn	Ala	Leu	Asp	Gly	Val	Glu	Arg
8630						8635					8640			
Leu	Ala	Gln	Asp	Lys	Gln	Thr	Ala	Gly	Asn	Ser	Leu	Asn	His	Leu
8645						8650					8655			
Asp	Gln	Leu	Thr	Pro	Ala	Gln	Gln	Gln	Ala	Leu	Glu	Asn	Gln	Ile
8660						8665					8670			
Asn	Asn	Ala	Thr	Thr	Arg	Gly	Glu	Val	Ala	Gln	Lys	Leu	Thr	Glu
8675						8680					8685			
Ala	Gln	Ala	Leu	Asn	Gln	Ala	Met	Glu	Ala	Leu	Arg	Asn	Ser	Ile
8690						8695					8700			
Gln	Asp	Gln	Gln	Gln	Thr	Glu	Ala	Gly	Ser	Lys	Phe	Ile	Asn	Glu
8705						8710					8715			
Asp	Lys	Pro	Gln	Lys	Asp	Ala	Tyr	Gln	Ala	Ala	Val	Gln	Asn	Ala
8720						8725					8730			
Lys	Asp	Leu	Ile	Asn	Gln	Thr	Asn	Asn	Pro	Thr	Leu	Asp	Lys	Ala
8735						8740					8745			
Gln	Val	Glu	Gln	Leu	Thr	Gln	Ala	Val	Asn	Gln	Ala	Lys	Asp	Asn
8750						8755					8760			
Leu	His	Gly	Asp	Gln	Lys	Leu	Ala	Asp	Asp	Lys	Gln	His	Ala	Val
8765						8770					8775			
Thr	Asp	Leu	Asn	Gln	Leu	Asn	Gly	Leu	Asn	Asn	Pro	Gln	Arg	Gln
8780						8785					8790			
Ala	Leu	Glu	Ser	Gln	Ile	Asn	Asn	Ala	Ala	Thr	Arg	Gly	Glu	Val
8795						8800					8805			
Ala	Gln	Lys	Leu	Ala	Glu	Ala	Lys	Ala	Leu	Asp	Gln	Ala	Met	Gln
8810						8815					8820			
Ala	Leu	Arg	Asn	Ser	Ile	Gln	Asp	Gln	Gln	Gln	Thr	Glu	Ser	Gly
8825						8830					8835			
Ser	Lys	Phe	Ile	Asn	Glu	Asp	Lys	Pro	Gln	Lys	Asp	Ala	Tyr	Gln
8840						8845					8850			
Ala	Ala	Val	Gln	Asn	Ala	Lys	Asp	Leu	Ile	Asn	Gln	Thr	Gly	Asn
8855						8860					8865			
Pro	Thr	Leu	Asp	Lys	Ser	Gln	Val	Glu	Gln	Leu	Thr	Gln	Ala	Val



-continued

8870	8875	8880
Thr Thr Ala Lys Asp Asn 8885	Leu His Gly Asp Gln 8890	Lys Leu Ala Arg 8895
Asp Gln Gln Gln Ala Val 8900	Thr Thr Val Asn Ala 8905	Leu Pro Asn Leu 8910
Asn His Ala Gln Gln Gln 8915	Ala Leu Thr Asp Ala 8920	Ile Asn Ala Ala 8925
Pro Thr Arg Thr Glu Val 8930	Ala Gln His Val Gln 8935	Thr Ala Thr Glu 8940
Leu Asp His Ala Met Glu 8945	Thr Leu Lys Asn Lys 8950	Val Asp Gln Val 8955
Asn Thr Asp Lys Ala Gln 8960	Pro Asn Tyr Thr Glu 8965	Ala Ser Thr Asp 8970
Lys Lys Glu Ala Val Asp 8975	Gln Ala Leu Gln Ala 8980	Ala Glu Ser Ile 8985
Thr Asp Pro Thr Asn Gly 8990	Ser Asn Ala Asn Lys 8995	Asp Ala Val Asp 9000
Gln Val Leu Thr Lys Leu 9005	Gln Glu Lys Glu Asn 9010	Glu Leu Asn Gly 9015
Asn Glu Arg Val Ala Glu 9020	Ala Lys Thr Gln Ala 9025	Lys Gln Thr Ile 9030
Asp Gln Leu Thr His Leu 9035	Asn Ala Asp Gln Ile 9040	Ala Thr Ala Lys 9045
Gln Asn Ile Asp Gln Ala 9050	Thr Lys Leu Gln Pro 9055	Ile Ala Glu Leu 9060
Val Asp Gln Ala Thr Gln 9065	Leu Asn Gln Ser Met 9070	Asp Gln Leu Gln 9075
Gln Ala Val Asn Glu His 9080	Ala Asn Val Glu Gln 9085	Thr Val Asp Tyr 9090
Thr Gln Ala Asp Ser Asp 9095	Lys Gln Asn Ala Tyr 9100	Lys Gln Ala Ile 9105
Ala Asp Ala Glu Asn Val 9110	Leu Lys Gln Asn Ala 9115	Asn Lys Gln Gln 9120
Val Asp Gln Ala Leu Gln 9125	Asn Ile Leu Asn Ala 9130	Lys Gln Ala Leu 9135
Asn Gly Asp Glu Arg Val 9140	Ala Leu Ala Lys Thr 9145	Asn Gly Lys His 9150
Asp Ile Asp Gln Leu Asn 9155	Ala Leu Asn Asn Ala 9160	Gln Gln Asp Gly 9165
Phe Lys Gly Arg Ile Asp 9170	Gln Ser Asn Asp Leu 9175	Asn Gln Ile Gln 9180
Gln Ile Val Asp Glu Ala 9185	Lys Ala Leu Asn Arg 9190	Ala Met Asp Gln 9195
Leu Ser Gln Glu Ile Thr 9200	Asp Asn Glu Gly Arg 9205	Thr Lys Gly Ser 9210
Thr Asn Tyr Val Asn Ala 9215	Asp Thr Gln Val Lys 9220	Gln Val Tyr Asp 9225
Glu Thr Val Asp Lys Ala 9230	Lys Gln Ala Leu Asp 9235	Lys Ser Thr Gly 9240
Gln Asn Leu Thr Ala Lys 9245	Gln Val Ile Lys Leu 9250	Asn Asp Ala Val 9255

-continued

Thr	Ala	Ala	Lys	Lys	Ala	Leu	Asn	Gly	Glu	Glu	Arg	Leu	Asn	Asn
9260						9265						9270		
Arg	Lys	Ala	Glu	Ala	Leu	Gln	Arg	Leu	Asp	Gln	Leu	Thr	His	Leu
9275						9280						9285		
Asn	Asn	Ala	Gln	Arg	Gln	Leu	Ala	Ile	Gln	Gln	Ile	Asn	Asn	Ala
9290						9295						9300		
Glu	Thr	Leu	Asn	Lys	Ala	Ser	Arg	Ala	Ile	Asn	Arg	Ala	Thr	Lys
9305						9310						9315		
Leu	Asp	Asn	Ala	Met	Gly	Ala	Val	Gln	Gln	Tyr	Ile	Asp	Glu	Gln
9320						9325						9330		
His	Leu	Gly	Val	Ile	Ser	Ser	Thr	Asn	Tyr	Ile	Asn	Ala	Asp	Asp
9335						9340						9345		
Asn	Leu	Lys	Ala	Asn	Tyr	Asp	Asn	Ala	Ile	Ala	Asn	Ala	Ala	His
9350						9355						9360		
Glu	Leu	Asp	Lys	Val	Gln	Gly	Asn	Ala	Ile	Ala	Lys	Ala	Glu	Ala
9365						9370						9375		
Glu	Gln	Leu	Lys	Gln	Asn	Ile	Ile	Asp	Ala	Gln	Asn	Ala	Leu	Asn
9380						9385						9390		
Gly	Asp	Gln	Asn	Leu	Ala	Asn	Ala	Lys	Asp	Lys	Ala	Asn	Ala	Phe
9395						9400						9405		
Val	Asn	Ser	Leu	Asn	Gly	Leu	Asn	Gln	Gln	Gln	Gln	Asp	Leu	Ala
9410						9415						9420		
His	Lys	Ala	Ile	Asn	Asn	Ala	Asp	Thr	Val	Ser	Asp	Val	Thr	Asp
9425						9430						9435		
Ile	Val	Asn	Asn	Gln	Ile	Asp	Leu	Asn	Asp	Ala	Met	Glu	Thr	Leu
9440						9445						9450		
Lys	His	Leu	Val	Asp	Asn	Glu	Ile	Pro	Asn	Ala	Glu	Gln	Thr	Val
9455						9460						9465		
Asn	Tyr	Gln	Asn	Ala	Asp	Asp	Asn	Ala	Lys	Thr	Asn	Phe	Asp	Asp
9470						9475						9480		
Ala	Lys	Arg	Leu	Ala	Asn	Thr	Leu	Leu	Asn	Ser	Asp	Asn	Thr	Asn
9485						9490						9495		
Val	Asn	Asp	Ile	Asn	Gly	Ala	Ile	Gln	Ala	Val	Asn	Asp	Ala	Ile
9500						9505						9510		
His	Asn	Leu	Asn	Gly	Asp	Gln	Arg	Leu	Gln	Asp	Ala	Lys	Asp	Lys
9515						9520						9525		
Ala	Ile	Gln	Ser	Ile	Asn	Gln	Ala	Leu	Ala	Asn	Lys	Leu	Lys	Glu
9530						9535						9540		
Ile	Glu	Ala	Ser	Asn	Ala	Thr	Asp	Gln	Asp	Lys	Leu	Ile	Ala	Lys
9545						9550						9555		
Asn	Lys	Ala	Glu	Glu	Leu	Ala	Asn	Ser	Ile	Ile	Asn	Asn	Ile	Asn
9560						9565						9570		
Lys	Ala	Thr	Ser	Asn	Gln	Ala	Val	Ser	Gln	Val	Gln	Thr	Ala	Gly
9575						9580						9585		
Asn	His	Ala	Ile	Glu	Gln	Val	His	Ala	Asn	Glu	Ile	Pro	Lys	Ala
9590						9595						9600		
Lys	Ile	Asp	Ala	Asn	Lys	Asp	Val	Asp	Lys	Gln	Val	Gln	Ala	Leu
9605						9610						9615		
Ile	Asp	Glu	Ile	Asp	Arg	Asn	Pro	Asn	Leu	Thr	Asp	Lys	Glu	Lys
9620						9625						9630		
Gln	Ala	Leu	Lys	Asp	Arg	Ile	Asn	Gln	Ile	Leu	Gln	Gln	Gly	His
9635						9640						9645		

-continued

---

Asn Gly 9650	Ile Asn Asn Ala 9655	Met Thr Lys Glu Glu 9660	Ile Glu Gln Ala
Lys Ala 9665	Gln Leu Ala Gln Ala 9670	Leu Gln Asp Ile Lys 9675	Asp Leu Val
Lys Ala 9680	Lys Glu Asp Ala Lys 9685	Gln Asp Val Asp Lys 9690	Gln Val Gln
Ala Leu 9695	Ile Asp Glu Ile Asp 9700	Gln Asn Pro Asn Leu 9705	Thr Asp Lys
Glu Lys 9710	Gln Ala Leu Lys Tyr 9715	Arg Ile Asn Gln Ile 9720	Leu Gln Gln
Gly His 9725	Asn Asp Ile Asn Asn 9730	Ala Leu Thr Lys Glu 9735	Glu Ile Glu
Gln Ala 9740	Lys Ala Gln Leu Ala 9745	Gln Ala Leu Gln Asp 9750	Ile Lys Asp
Leu Val 9755	Lys Ala Lys Glu Asp 9760	Ala Lys Asn Ala Ile 9765	Lys Ala Leu
Ala Asn 9770	Ala Lys Arg Asp Gln 9775	Ile Asn Ser Asn Pro 9780	Asp Leu Thr
Pro Glu 9785	Gln Lys Ala Lys Ala 9790	Leu Lys Glu Ile Asp 9795	Glu Ala Glu
Lys Arg 9800	Ala Leu Gln Asn Val 9805	Glu Asn Ala Gln Thr 9810	Ile Asp Gln
Leu Asn 9815	Arg Gly Leu Asn Leu 9820	Gly Leu Asp Asp Ile 9825	Arg Asn Thr
His Val 9830	Trp Glu Val Asp Glu 9835	Gln Pro Ala Val Asn 9840	Glu Ile Phe
Glu Ala 9845	Thr Pro Glu Gln Ile 9850	Leu Val Asn Gly Glu 9855	Leu Ile Val
His Arg 9860	Asp Asp Ile Ile Thr 9865	Glu Gln Asp Ile Leu 9870	Ala His Ile
Asn Leu 9875	Ile Asp Gln Leu Ser 9880	Ala Glu Val Ile Asp 9885	Thr Pro Ser
Thr Ala 9890	Thr Ile Ser Asp Ser 9895	Leu Thr Ala Lys Val 9900	Glu Val Thr
Leu Leu 9905	Asp Gly Ser Lys Val 9910	Ile Val Asn Val Pro 9915	Val Lys Val
Val Glu 9920	Lys Glu Leu Ser Val 9925	Val Lys Gln Gln Ala 9930	Ile Glu Ser
Ile Glu 9935	Asn Ala Ala Gln Gln 9940	Lys Ile Asn Glu Ile 9945	Asn Asn Ser
Val Thr 9950	Leu Thr Leu Glu Gln 9955	Lys Glu Ala Ala Ile 9960	Ala Glu Val
Asn Lys 9965	Leu Lys Gln Gln Ala 9970	Ile Asp His Val Asn 9975	Asn Ala Pro
Asp Val 9980	His Ser Val Glu Glu 9985	Ile Gln Gln Gln Glu 9990	Gln Ala His
Ile Glu 9995	Gln Phe Asn Pro Glu 10000	Gln Phe Thr Ile Glu 10005	Gln Ala Lys
Ser Asn 10010	Ala Ile Lys Ser Ile 10015	Glu Asp Ala Ile Gln 10020	His Met Ile
Asp Glu	Ile Lys Ala Arg Thr	Asp Leu Thr Asp Lys	Glu Lys Gln

-continued

10025	10030	10035
Glu Ala Ile Ala Lys Leu Asn 10040	Gln Leu Lys Glu Gln 10045	Ala Ile Gln 10050
Ala Ile Gln Arg Ala Gln Ser 10055	Ile Asp Glu Ile Ser 10060	Glu Gln Leu 10065
Glu Gln Phe Lys Ala Gln Met 10070	Lys Ala Ala Asn Pro 10075	Thr Ala Lys 10080
Glu Leu Ala Lys Arg Lys Gln 10085	Glu Ala Ile Ser Arg 10090	Ile Lys Asp 10095
Phe Ser Asn Glu Lys Ile Asn 10100	Ser Ile Arg Asn Ser 10105	Glu Ile Gly 10110
Thr Ala Asp Glu Lys Gln Ala 10115	Ala Met Asn Gln Ile 10120	Asn Glu Ile 10125
Val Leu Glu Thr Ile Arg Asp 10130	Ile Asn Asn Ala His 10135	Thr Leu Gln 10140
Gln Val Glu Ala Ala Leu Asn 10145	Asn Gly Ile Ala Arg 10150	Ile Ser Ala 10155
Val Gln Ile Val Thr Ser Asp 10160	Arg Ala Lys Gln Ser 10165	Ser Ser Thr 10170
Gly Asn Glu Ser Asn Ser His 10175	Leu Thr Ile Gly Tyr 10180	Gly Thr Ala 10185
Asn His Pro Phe Asn Ser Ser 10190	Thr Ile Gly His Lys 10195	Lys Lys Leu 10200
Asp Glu Asp Asp Asp Ile Asp 10205	Pro Leu His Met Arg 10210	His Phe Ser 10215
Asn Asn Phe Gly Asn Val Ile 10220	Lys Asn Ala Ile Gly 10225	Val Val Gly 10230
Ile Ser Gly Leu Leu Ala Ser 10235	Phe Trp Phe Phe Ile 10240	Ala Lys Arg 10245
Arg Arg Lys Glu Asp Glu Glu 10250	Glu Glu Leu Glu Ile 10255	Arg Asp Asn 10260
Asn Lys Asp Ser Ile Lys Glu 10265	Thr Leu Asp Asp Thr 10270	Lys His Leu 10275
Pro Leu Leu Phe Ala Lys Arg 10280	Arg Arg Lys Glu Asp 10285	Glu Glu Asp 10290
Val Thr Val Glu Glu Lys Asp 10295	Ser Leu Asn Asn Gly 10300	Glu Ser Leu 10305
Asp Lys Val Lys His Thr Pro 10310	Phe Phe Leu Pro Lys 10315	Arg Arg Arg 10320
Lys Glu Asp Glu Glu Asp Val 10325	Glu Val Thr Asn Glu 10330	Asn Thr Asp 10335
Glu Lys Val Leu Lys Asp Asn 10340	Glu His Ser Pro Leu 10345	Leu Phe Ala 10350
Lys Arg Arg Lys Asp Lys Glu 10355	Glu Asp Val Glu Thr 10360	Thr Thr Ser 10365
Ile Glu Ser Lys Asp Glu Asp 10370	Val Pro Leu Leu Leu 10375	Ala Lys Lys 10380
Lys Asn Gln Lys Asp Asn Gln 10385	Ser Lys Asp Lys Lys 10390	Ser Ala Ser 10395
Lys Asn Thr Ser Lys Lys Val 10400	Ala Ala Lys Lys Lys 10405	Lys Lys Lys 10410

-continued

Ala Lys Lys Asn Lys Lys  
10415

<210> SEQ ID NO 25

<211> LENGTH: 340

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 25

Met Lys Lys Lys Leu Leu Val Leu Thr Met Ser Thr Leu Phe Ala Thr  
1 5 10 15

Gln Ile Met Asn Ser Asn His Ala Lys Ala Ser Val Thr Glu Ser Val  
20 25 30

Asp Lys Lys Phe Val Val Pro Glu Ser Gly Ile Asn Lys Ile Ile Pro  
35 40 45

Ala Tyr Asp Glu Phe Lys Asn Ser Pro Lys Val Asn Val Ser Asn Leu  
50 55 60

Thr Asp Asn Lys Asn Phe Val Ala Ser Glu Asp Lys Leu Asn Lys Ile  
65 70 75 80

Ala Asp Ser Ser Ala Ala Ser Lys Ile Val Asp Lys Asn Phe Val Val  
85 90 95

Pro Glu Ser Lys Leu Gly Asn Ile Val Pro Glu Tyr Lys Glu Ile Asn  
100 105 110

Asn Arg Val Asn Val Ala Thr Asn Asn Pro Ala Ser Gln Gln Val Asp  
115 120 125

Lys His Phe Val Ala Lys Gly Pro Glu Val Asn Arg Phe Ile Thr Gln  
130 135 140

Asn Lys Val Asn His His Phe Ile Thr Thr Gln Thr His Tyr Lys Lys  
145 150 155 160

Val Ile Thr Ser Tyr Lys Ser Thr His Val His Lys His Val Asn His  
165 170 175

Ala Lys Asp Ser Ile Asn Lys His Phe Ile Val Lys Pro Ser Glu Ser  
180 185 190

Pro Arg Tyr Thr His Pro Ser Gln Ser Leu Ile Ile Lys His His Phe  
195 200 205

Ala Val Pro Gly Tyr His Ala His Lys Phe Val Thr Pro Gly His Ala  
210 215 220

Ser Ile Lys Ile Asn His Phe Cys Val Val Pro Gln Ile Asn Ser Phe  
225 230 235 240

Lys Val Ile Pro Pro Tyr Gly His Asn Ser His Arg Met His Val Pro  
245 250 255

Ser Phe Gln Asn Asn Thr Thr Ala Thr His Gln Asn Ala Lys Val Asn  
260 265 270

Lys Ala Tyr Asp Tyr Lys Tyr Phe Tyr Ser Tyr Lys Val Val Lys Gly  
275 280 285

Val Lys Lys Tyr Phe Ser Phe Ser Gln Ser Asn Gly Tyr Lys Ile Gly  
290 295 300

Lys Pro Ser Leu Asn Ile Lys Asn Val Asn Tyr Gln Tyr Ala Val Pro  
305 310 315 320

Ser Tyr Ser Pro Thr His Tyr Val Pro Glu Phe Lys Gly Ser Leu Pro  
325 330 335

Ala Pro Arg Val  
340

-continued

<210> SEQ ID NO 26  
 <211> LENGTH: 130  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 26

```

Met Asn Phe Asn Asp Ile Glu Thr Met Val Lys Ser Lys Phe Lys Asp
1          5          10          15
Ile Lys Lys His Ala Glu Glu Ile Ala His Glu Ile Glu Val Arg Ser
          20          25          30
Gly Tyr Leu Arg Lys Ala Glu Gln Tyr Lys Arg Leu Glu Phe Asn Leu
          35          40          45
Ser Phe Ala Leu Asp Asp Ile Glu Ser Thr Ala Lys Asp Val Gln Thr
          50          55          60
Ala Lys Ser Ser Ala Asn Lys Asp Ser Val Thr Val Lys Gly Lys Ala
65          70          75          80
Pro Asn Thr Leu Tyr Ile Glu Lys Arg Asn Leu Met Lys Gln Lys Leu
          85          90          95
Glu Met Leu Gly Glu Asp Ile Asp Lys Asn Lys Glu Ser Leu Gln Lys
          100          105          110
Ala Lys Glu Ile Ala Gly Glu Lys Ala Ser Glu Tyr Phe Asn Lys Ala
          115          120          125

Met Asn
          130
  
```

<210> SEQ ID NO 27  
 <211> LENGTH: 636  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 27

```

Met Lys Lys Gln Ile Ile Ser Leu Gly Ala Leu Ala Val Ala Ser Ser
1          5          10          15
Leu Phe Thr Trp Asp Asn Lys Ala Asp Ala Ile Val Thr Lys Asp Tyr
          20          25          30
Ser Gly Lys Ser Gln Val Asn Ala Gly Ser Lys Asn Gly Thr Leu Ile
          35          40          45
Asp Ser Arg Tyr Leu Asn Ser Ala Leu Tyr Tyr Leu Glu Asp Tyr Ile
          50          55          60
Ile Tyr Ala Ile Gly Leu Thr Asn Lys Tyr Glu Tyr Gly Asp Asn Ile
65          70          75          80
Tyr Lys Glu Ala Lys Asp Arg Leu Leu Glu Lys Val Leu Arg Glu Asp
          85          90          95
Gln Tyr Leu Leu Glu Arg Lys Lys Ser Gln Tyr Glu Asp Tyr Lys Gln
          100          105          110
Trp Tyr Ala Asn Tyr Lys Lys Glu Asn Pro Arg Thr Asp Leu Lys Met
          115          120          125
Ala Asn Phe His Lys Tyr Asn Leu Glu Glu Leu Ser Met Lys Glu Tyr
          130          135          140
Asn Glu Leu Gln Asp Ala Leu Lys Arg Ala Leu Asp Asp Phe His Arg
145          150          155          160
Glu Val Lys Asp Ile Lys Asp Lys Asn Ser Asp Leu Lys Thr Phe Asn
          165          170          175
Ala Ala Glu Glu Asp Lys Ala Thr Lys Glu Val Tyr Asp Leu Val Ser
  
```

-continued

180					185					190					
Glu	Ile	Asp	Thr	Leu	Val	Val	Ser	Tyr	Tyr	Gly	Asp	Lys	Asp	Tyr	Gly
		195					200					205			
Glu	His	Ala	Lys	Glu	Leu	Arg	Ala	Lys	Leu	Asp	Leu	Ile	Leu	Gly	Asp
		210					215					220			
Thr	Asp	Asn	Pro	His	Lys	Ile	Thr	Asn	Glu	Arg	Ile	Lys	Lys	Glu	Met
		225					230					235			
Ile	Asp	Asp	Leu	Asn	Ser	Ile	Ile	Asp	Asp	Phe	Phe	Met	Glu	Thr	Lys
				245								250			255
Gln	Asn	Arg	Pro	Lys	Ser	Ile	Thr	Lys	Tyr	Asn	Pro	Thr	Thr	His	Asn
				260								265			270
Tyr	Lys	Thr	Asn	Ser	Asp	Asn	Lys	Pro	Asn	Phe	Asp	Lys	Leu	Val	Glu
				275								280			285
Glu	Thr	Lys	Lys	Ala	Val	Lys	Glu	Ala	Asp	Asp	Ser	Trp	Lys	Lys	Lys
				290								295			300
Thr	Val	Lys	Lys	Tyr	Gly	Glu	Thr	Glu	Thr	Lys	Ser	Pro	Val	Val	Lys
				305								310			315
Glu	Glu	Lys	Lys	Val	Glu	Glu	Pro	Gln	Ala	Pro	Lys	Val	Asp	Asn	Gln
				325								330			335
Gln	Glu	Val	Lys	Thr	Thr	Ala	Gly	Lys	Ala	Glu	Glu	Thr	Thr	Gln	Pro
				340								345			350
Val	Ala	Gln	Pro	Leu	Val	Lys	Ile	Pro	Gln	Gly	Thr	Ile	Thr	Gly	Glu
				355								360			365
Ile	Val	Lys	Gly	Pro	Glu	Tyr	Pro	Thr	Met	Glu	Asn	Lys	Thr	Val	Gln
				370								375			380
Gly	Glu	Ile	Val	Gln	Gly	Pro	Asp	Phe	Leu	Thr	Met	Glu	Gln	Ser	Gly
				385								390			395
Pro	Ser	Leu	Ser	Asn	Asn	Tyr	Thr	Asn	Pro	Pro	Leu	Thr	Asn	Pro	Ile
				405								410			415
Leu	Glu	Gly	Leu	Glu	Gly	Ser	Ser	Ser	Lys	Leu	Glu	Ile	Lys	Pro	Gln
				420								425			430
Gly	Thr	Glu	Ser	Thr	Leu	Lys	Gly	Thr	Gln	Gly	Glu	Ser	Ser	Asp	Ile
				435								440			445
Glu	Val	Lys	Pro	Gln	Ala	Thr	Glu	Thr	Thr	Glu	Ala	Ser	Gln	Tyr	Gly
				450								455			460
Pro	Arg	Pro	Gln	Phe	Asn	Lys	Thr	Pro	Lys	Tyr	Val	Lys	Tyr	Arg	Asp
				465								470			475
Ala	Gly	Thr	Gly	Ile	Arg	Glu	Tyr	Asn	Asp	Gly	Thr	Phe	Gly	Tyr	Glu
				485								490			495
Ala	Arg	Pro	Arg	Phe	Asn	Lys	Pro	Ser	Glu	Thr	Asn	Ala	Tyr	Asn	Val
				500								505			510
Thr	Thr	His	Ala	Asn	Gly	Gln	Val	Ser	Tyr	Gly	Ala	Arg	Pro	Thr	Tyr
				515								520			525
Lys	Lys	Pro	Ser	Glu	Thr	Asn	Ala	Tyr	Asn	Val	Thr	Thr	His	Ala	Asn
				530								535			540
Gly	Gln	Val	Ser	Tyr	Gly	Ala	Arg	Pro	Thr	Gln	Asn	Lys	Pro	Ser	Lys
				545								550			555
Thr	Asn	Ala	Tyr	Asn	Val	Thr	Thr	His	Gly	Asn	Gly	Gln	Val	Ser	Tyr
				565								570			575
Gly	Ala	Arg	Pro	Thr	Gln	Asn	Lys	Pro	Ser	Lys	Thr	Asn	Ala	Tyr	Asn
				580								585			590

-continued

---

Val Thr Thr His Ala Asn Gly Gln Val Ser Tyr Gly Ala Arg Pro Thr  
 595 600 605

Tyr Lys Lys Pro Ser Lys Thr Asn Ala Tyr Asn Val Thr Thr His Ala  
 610 615 620

Asp Gly Thr Ala Thr Tyr Gly Pro Arg Val Thr Lys  
 625 630 635

<210> SEQ ID NO 28  
 <211> LENGTH: 745  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 28

Ala Glu Gln His Thr Pro Met Lys Ala His Ala Val Thr Thr Ile Asp  
 1 5 10 15

Lys Ala Thr Thr Asp Lys Gln Gln Val Pro Pro Thr Lys Glu Ala Ala  
 20 25 30

His His Ser Gly Lys Glu Ala Ala Thr Asn Val Ser Ala Ser Ala Gln  
 35 40 45

Gly Thr Ala Asp Asp Thr Asn Ser Lys Val Thr Ser Asn Ala Pro Ser  
 50 55 60

Asn Lys Pro Ser Thr Val Val Ser Thr Lys Val Asn Glu Thr Arg Asp  
 65 70 75 80

Val Asp Thr Gln Gln Ala Ser Thr Gln Lys Pro Thr His Thr Ala Thr  
 85 90 95

Phe Lys Leu Ser Asn Ala Lys Thr Ala Ser Leu Ser Pro Arg Met Phe  
 100 105 110

Ala Ala Asn Ala Pro Gln Thr Thr Thr His Lys Ile Leu His Thr Asn  
 115 120 125

Asp Ile His Gly Arg Leu Ala Glu Glu Lys Gly Arg Val Ile Gly Met  
 130 135 140

Ala Lys Leu Lys Thr Val Lys Glu Gln Glu Lys Pro Asp Leu Met Leu  
 145 150 155 160

Asp Ala Gly Asp Ala Phe Gln Gly Leu Pro Leu Ser Asn Gln Ser Lys  
 165 170 175

Gly Glu Glu Met Ala Lys Ala Met Asn Ala Val Gly Tyr Asp Ala Met  
 180 185 190

Ala Val Gly Asn His Glu Phe Asp Phe Gly Tyr Asp Gln Leu Lys Lys  
 195 200 205

Leu Glu Gly Met Leu Asp Phe Pro Met Leu Ser Thr Asn Val Tyr Lys  
 210 215 220

Asp Gly Lys Arg Ala Phe Lys Pro Ser Thr Ile Val Thr Lys Asn Gly  
 225 230 235 240

Ile Arg Tyr Gly Ile Ile Gly Val Thr Thr Pro Glu Thr Lys Thr Lys  
 245 250 255

Thr Arg Pro Glu Gly Ile Lys Gly Val Glu Phe Arg Asp Pro Leu Gln  
 260 265 270

Ser Val Thr Ala Glu Met Met Arg Ile Tyr Lys Asp Val Asp Thr Phe  
 275 280 285

Val Val Ile Ser His Leu Gly Ile Asp Pro Ser Thr Gln Glu Thr Trp  
 290 295 300

Arg Gly Asp Tyr Leu Val Lys Gln Leu Ser Gln Asn Pro Gln Leu Lys  
 305 310 315 320



-continued

---

Lys Arg Ile Thr Val Ile Asp Gly His Ser His Thr Val Leu Gln Asn  
 325 330 335

Gly Gln Ile Tyr Asn Asn Asp Ala Leu Ala Gln Thr Gly Thr Ala Leu  
 340 345 350

Ala Asn Ile Gly Lys Ile Thr Phe Asn Tyr Arg Asn Gly Glu Val Ser  
 355 360 365

Asn Ile Lys Pro Ser Leu Ile Asn Val Lys Asp Val Glu Asn Val Thr  
 370 375 380

Pro Asn Lys Ala Leu Ala Glu Gln Ile Asn Gln Ala Asp Gln Thr Phe  
 385 390 395 400

Arg Ala Gln Thr Ala Glu Val Ile Ile Pro Asn Asn Thr Ile Asp Phe  
 405 410 415

Lys Gly Glu Arg Asp Asp Val Arg Thr Arg Glu Thr Asn Leu Gly Asn  
 420 425 430

Ala Ile Ala Asp Ala Met Glu Ala Tyr Gly Val Lys Asn Phe Ser Lys  
 435 440 445

Lys Thr Asp Phe Ala Val Thr Asn Gly Gly Gly Ile Arg Ala Ser Ile  
 450 455 460

Ala Lys Gly Lys Val Thr Arg Tyr Asp Leu Ile Ser Val Leu Pro Phe  
 465 470 475 480

Gly Asn Thr Ile Ala Gln Ile Asp Val Lys Gly Ser Asp Val Trp Thr  
 485 490 495

Ala Phe Glu His Ser Leu Gly Ala Pro Thr Thr Gln Lys Asp Gly Lys  
 500 505 510

Thr Val Leu Thr Ala Asn Gly Gly Leu Leu His Ile Ser Asp Ser Ile  
 515 520 525

Arg Val Tyr Tyr Asp Ile Asn Lys Pro Ser Gly Lys Arg Ile Asn Ala  
 530 535 540

Ile Gln Ile Leu Asn Lys Glu Thr Gly Lys Phe Glu Asn Ile Asp Leu  
 545 550 555 560

Lys Arg Val Tyr His Val Thr Met Asn Asp Phe Thr Ala Ser Gly Gly  
 565 570 575

Asp Gly Tyr Ser Met Phe Gly Gly Pro Arg Glu Glu Gly Ile Ser Leu  
 580 585 590

Asp Gln Val Leu Ala Ser Tyr Leu Lys Thr Ala Asn Leu Ala Lys Tyr  
 595 600 605

Asp Thr Thr Glu Pro Gln Arg Met Leu Leu Gly Lys Pro Ala Val Ser  
 610 615 620

Glu Gln Pro Ala Lys Gly Gln Gln Gly Ser Lys Gly Ser Lys Ser Gly  
 625 630 635 640

Lys Asp Thr Gln Pro Ile Gly Asp Asp Lys Val Met Asp Pro Ala Lys  
 645 650 655

Lys Pro Ala Pro Gly Lys Val Val Leu Leu Leu Ala His Arg Gly Thr  
 660 665 670

Val Ser Ser Gly Thr Glu Gly Ser Gly Arg Thr Ile Glu Gly Ala Thr  
 675 680 685

Val Ser Ser Lys Ser Gly Lys Gln Leu Ala Arg Met Ser Val Pro Lys  
 690 695 700

Gly Ser Ala His Glu Lys Gln Leu Pro Lys Thr Gly Thr Asn Gln Ser  
 705 710 715 720

Ser Ser Pro Glu Ala Met Phe Val Leu Leu Ala Gly Ile Gly Leu Ile  
 725 730 735

-continued

Ala Thr Val Arg Arg Arg Lys Ala Ser  
740 745

<210> SEQ ID NO 29  
<211> LENGTH: 628  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 29

Met Ser Asp Arg Phe Ile Lys Phe Asn Asp Glu Gln Leu Asp Ala Lys  
1 5 10 15  
Gln Val Met Met Leu Gln Asp Leu Ala Arg Leu Leu Leu Lys Asn Glu  
20 25 30  
Gln Thr Gln Val Lys Ile Gln Lys Phe Pro Tyr Tyr Asn Pro Val Gln  
35 40 45  
Asn Val Leu Ile Thr Ser Trp Phe Trp Ser His Arg Pro Ser His Ile  
50 55 60  
Glu Met Ala Gly Leu Lys Thr Asp Val Met Leu Ala Ala Tyr Gly Tyr  
65 70 75 80  
His Met Met Asp Val Gln Ile Val Asn Glu Val Val Gln Asp Lys Thr  
85 90 95  
Phe Lys His Pro Lys Phe Tyr Gln Gln Leu Phe Lys Leu Leu Glu Asp  
100 105 110  
Met Arg Val Leu Asn Ser Ile Lys Val Glu Arg Pro Ser Thr Ala Lys  
115 120 125  
Leu Ile Asp Leu Arg Leu Asp Thr Arg Ile Ser Tyr Thr Glu Ser Gln  
130 135 140  
Ile Lys Val Tyr Arg Thr Lys Thr Gln Tyr Thr Asp Leu Leu Phe Leu  
145 150 155 160  
Tyr Leu Glu His Ala Phe Leu Ser Gln Asp Phe Phe Asp Ile Pro Ser  
165 170 175  
Ile His Ser Asp Leu Asp Asp Ile Leu Val Asn Met Phe Leu Tyr Leu  
180 185 190  
Pro Asn Phe Phe Gln Asn Gln Asn Ser Glu Asp Asn Met Tyr Leu Ala  
195 200 205  
Gln Arg Ile Met Tyr Gln Val Asp Asp Ile Leu Lys Glu Asp Met Leu  
210 215 220  
Asn Glu Tyr Tyr Tyr Leu Pro Lys Thr Leu Tyr Asn Thr Leu Ala Ser  
225 230 235 240  
Pro Glu Phe Asp Asp Leu Lys Arg Thr Asp Ala Ser Gln Val Asp Gly  
245 250 255  
Gln Asp Asp Thr Ser Glu Asp Asp Asp Asn Glu Ser Glu Lys Ala Asp  
260 265 270  
Ser Lys Ser Ala Asp Ser Glu Ser Lys Gly Gly Ala Tyr Leu Glu Met  
275 280 285  
Glu Leu His Glu Gly Gln Asn Ser Glu Thr Leu Gly Asn Asp Glu Ala  
290 295 300  
Arg Glu Gly Asp Ala Thr Asp Asp Met Thr Asp Met Met Thr Lys Lys  
305 310 315 320  
Gly Lys Gly Ser Asn Asp Thr Leu Asn Arg Glu Glu Gly Asp Ala Val  
325 330 335  
Gly Gln Ser Gln Ala Phe Gln Leu Asp Gly Val Asn Lys Asn Val Glu  
340 345 350

-continued

Ile Lys Trp Gln Ile Pro Glu Ile Glu Pro Gln Tyr Val Leu Glu Tyr  
 355 360 365  
 Gln Glu Ser Lys Gln Asp Val Gln Tyr Glu Ile Lys Asp Leu Ile Gln  
 370 375 380  
 Ile Ile Lys Lys Thr Ile Glu Arg Glu Gln Arg Asp Ala Arg Phe Asn  
 385 390 395 400  
 Leu Thr Lys Gly Arg Leu Gln Lys Asp Leu Ile Asn Trp Phe Ile Asp  
 405 410 415  
 Asp Gln Tyr Lys Leu Phe Tyr Lys Lys Gln Asp Leu Ser Lys Ser Phe  
 420 425 430  
 Asp Ala Thr Phe Thr Leu Leu Ile Asp Ala Ser Ala Ser Met His Asp  
 435 440 445  
 Lys Met Ala Glu Thr Lys Lys Gly Val Val Leu Phe His Glu Thr Leu  
 450 455 460  
 Lys Ala Leu Asn Ile Lys His Glu Ile Leu Ser Phe Ser Glu Asp Ala  
 465 470 475 480  
 Phe Asp Ser Asp Glu His Ala Gln Pro Asn Ile Ile Asn Glu Ile Ile  
 485 490 495  
 Asn Tyr Asp Tyr Ser Thr Phe Glu Lys Asp Gly Pro Arg Ile Met Ala  
 500 505 510  
 Leu Glu Pro Gln Asp Asp Asn Arg Asp Gly Val Ala Ile Arg Val Ala  
 515 520 525  
 Ser Glu Arg Leu Met Arg Arg Asn Gln His Gln Arg Phe Leu Ile Val  
 530 535 540  
 Phe Ser Asp Gly Glu Pro Ser Ala Phe Asn Tyr Ser Gln Asp Gly Ile  
 545 550 555 560  
 Ile Asp Thr Tyr Glu Ala Val Glu Met Ser Arg Lys Phe Gly Ile Glu  
 565 570 575  
 Val Phe Asn Val Phe Leu Ser Gln Asp Pro Ile Thr Glu Asp Val Glu  
 580 585 590  
 Gln Thr Ile His Asn Ile Tyr Gly Gln Tyr Ala Ile Phe Val Glu Gly  
 595 600 605  
 Val Ala His Leu Pro Gly His Leu Ser Pro Leu Leu Lys Lys Leu Leu  
 610 615 620  
 Leu Lys Ser Leu  
 625

&lt;210&gt; SEQ ID NO 30

&lt;211&gt; LENGTH: 154

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 30

Ala Glu Ile Asn Lys Gln Thr Thr Ser Gln Gly Val Thr Thr Glu Lys  
 1 5 10 15  
 Asn Asn Gly Ile Ala Val Leu Glu Gln Asp Val Ile Thr Pro Thr Val  
 20 25 30  
 Lys Pro Gln Ala Lys Gln Asp Ile Ile Gln Ala Val Thr Thr Arg Lys  
 35 40 45  
 Gln Gln Ile Lys Lys Ser Asn Ala Ser Leu Gln Asp Glu Lys Asp Val  
 50 55 60  
 Ala Asn Asp Lys Ile Gly Lys Ile Glu Thr Lys Ala Ile Lys Asp Ile  
 65 70 75 80

-continued

---

Asp Ala Ala Thr Thr Asn Ala Gln Val Glu Ala Ile Lys Thr Lys Ala  
                   85                                  90                                  95  
 Ile Asn Asp Ile Asn Gln Thr Thr Pro Ala Thr Thr Ala Lys Ala Ala  
                   100                                  105                                  110  
 Ala Leu Glu Glu Phe Asp Glu Val Val Gln Ala Gln Ile Asp Gln Ala  
                   115                                  120                                  125  
 Pro Leu Asn Pro Asp Thr Thr Asn Glu Glu Val Ala Glu Ala Ile Glu  
                   130                                  135                                  140  
 Arg Ile Asn Ala Ala Lys Val Ser Gly Val  
                   145                                  150

<210> SEQ ID NO 31  
 <211> LENGTH: 584  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 31

Met Lys Phe Lys Ser Leu Ile Thr Thr Thr Leu Ala Leu Gly Val Leu  
   1                  5                                  10                                  15  
 Ala Ser Thr Gly Ala Asn Phe Asn Asn Asn Glu Ala Ser Ala Ala Ala  
                   20                                  25                                  30  
 Lys Pro Leu Asp Lys Ser Ser Ser Ser Leu His His Gly Tyr Ser Lys  
                   35                                  40                                  45  
 Val His Val Pro Tyr Ala Ile Thr Val Asn Gly Thr Ser Gln Asn Ile  
                   50                                  55                                  60  
 Leu Ser Ser Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp  
   65                  70                                  75                                  80  
 Leu Glu Asp Arg Val Lys Ser Val Leu Lys Ser Asp Arg Gly Ile Ser  
                   85                                  90                                  95  
 Asp Ile Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Tyr Phe  
                   100                                  105                                  110  
 Lys Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ala Gly Ile Tyr Thr  
                   115                                  120                                  125  
 Ala Asp Leu Ile Asn Thr Ser Glu Ile Lys Ala Ile Asn Ile Asn Val  
                   130                                  135                                  140  
 Asp Thr Lys Lys Gln Val Glu Asp Lys Lys Lys Asp Lys Ala Asn Tyr  
   145                  150                                  155                                  160  
 Gln Val Pro Tyr Thr Ile Thr Val Asn Gly Thr Ser Gln Asn Ile Leu  
                   165                                  170                                  175  
 Ser Asn Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp Leu  
                   180                                  185                                  190  
 Glu Asp Lys Val Lys Ser Val Leu Glu Ser Asn Arg Gly Ile Thr Asp  
                   195                                  200                                  205  
 Val Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Asn Phe Lys  
                   210                                  215                                  220  
 Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ser Gly Ile Tyr Thr Ala  
   225                  230                                  235                                  240  
 Asn Leu Ile Asn Ser Ser Asp Ile Lys Ser Ile Asn Ile Asn Val Asp  
                   245                                  250                                  255  
 Thr Lys Lys His Ile Glu Asn Lys Ala Lys Arg Asn Tyr Gln Val Pro  
                   260                                  265                                  270  
 Tyr Ser Ile Asn Leu Asn Gly Thr Ser Thr Asn Ile Leu Ser Asn Leu  
                   275                                  280                                  285

-continued

---

Ser Phe Ser Asn Lys Pro Trp Thr Asn Tyr Lys Asn Leu Thr Ser Gln  
 290 295 300  
 Ile Lys Ser Val Leu Lys His Asp Arg Gly Ile Ser Glu Gln Asp Leu  
 305 310 315 320  
 Lys Tyr Ala Lys Lys Ala Tyr Tyr Thr Val Tyr Phe Lys Asn Gly Gly  
 325 330 335  
 Lys Arg Ile Leu Gln Leu Asn Ser Lys Asn Tyr Thr Ala Asn Leu Val  
 340 345 350  
 His Ala Lys Asp Val Lys Arg Ile Glu Ile Thr Val Lys Thr Gly Thr  
 355 360 365  
 Lys Ala Lys Ala Asp Arg Tyr Val Pro Tyr Thr Ile Ala Val Asn Gly  
 370 375 380  
 Thr Ser Thr Pro Ile Leu Ser Asp Leu Lys Phe Thr Gly Asp Pro Arg  
 385 390 395 400  
 Val Gly Tyr Lys Asp Ile Ser Lys Lys Val Lys Ser Val Leu Lys His  
 405 410 415  
 Asp Arg Gly Ile Gly Glu Arg Glu Leu Lys Tyr Ala Lys Lys Ala Thr  
 420 425 430  
 Tyr Thr Val His Phe Lys Asn Gly Thr Lys Lys Val Ile Asn Ile Asn  
 435 440 445  
 Ser Asn Ile Ser Gln Leu Asn Leu Leu Tyr Val Gln Asp Ile Lys Lys  
 450 455 460  
 Ile Asp Ile Asp Val Lys Thr Gly Thr Lys Ala Lys Ala Asp Ser Tyr  
 465 470 475 480  
 Val Pro Tyr Thr Ile Ala Val Asn Gly Thr Ser Thr Pro Ile Leu Ser  
 485 490 495  
 Lys Leu Lys Ile Ser Asn Lys Gln Leu Ile Ser Tyr Lys Tyr Leu Asn  
 500 505 510  
 Asp Lys Val Lys Ser Val Leu Lys Ser Glu Arg Gly Ile Ser Asp Leu  
 515 520 525  
 Asp Leu Lys Phe Ala Lys Gln Ala Lys Tyr Thr Val Tyr Phe Lys Asn  
 530 535 540  
 Gly Lys Lys Gln Val Val Asn Leu Lys Ser Asp Ile Phe Thr Pro Asn  
 545 550 555 560  
 Leu Phe Ser Ala Lys Asp Ile Lys Lys Ile Asp Ile Asp Val Lys Gln  
 565 570 575  
 Tyr Thr Lys Ser Lys Lys Asn Lys  
 580

&lt;210&gt; SEQ ID NO 32

&lt;211&gt; LENGTH: 508

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 32

Met Lys Asn Lys Leu Leu Val Leu Ser Leu Gly Ala Leu Cys Val Ser  
 1 5 10 15  
 Gln Ile Trp Glu Ser Asn Arg Ala Ser Ala Val Val Ser Gly Glu Lys  
 20 25 30  
 Asn Pro Tyr Val Ser Glu Ser Leu Lys Leu Thr Asn Asn Lys Asn Lys  
 35 40 45  
 Ser Arg Thr Val Glu Glu Tyr Lys Lys Ser Leu Asp Asp Leu Ile Trp  
 50 55 60

-continued

---

Ser Phe Pro Asn Leu Asp Asn Glu Arg Phe Asp Asn Pro Glu Tyr Lys  
 65 70 75 80  
 Glu Ala Met Lys Lys Tyr Gln Gln Arg Phe Met Ala Glu Asp Glu Ala  
 85 90 95  
 Leu Lys Lys Phe Phe Ser Glu Glu Lys Lys Ile Lys Asn Gly Asn Thr  
 100 105 110  
 Asp Asn Leu Asp Tyr Leu Gly Leu Ser His Glu Arg Tyr Glu Ser Val  
 115 120 125  
 Phe Asn Thr Leu Lys Lys Gln Ser Glu Glu Phe Leu Lys Glu Ile Glu  
 130 135 140  
 Asp Ile Lys Lys Asp Asn Pro Glu Leu Lys Asp Phe Asn Glu Glu Glu  
 145 150 155 160  
 Gln Leu Lys Cys Asp Leu Glu Leu Asn Lys Leu Glu Asn Gln Ile Leu  
 165 170 175  
 Met Leu Gly Lys Thr Phe Tyr Gln Asn Tyr Arg Asp Asp Val Glu Ser  
 180 185 190  
 Leu Tyr Ser Lys Leu Asp Leu Ile Met Gly Tyr Lys Asp Glu Glu Arg  
 195 200 205  
 Ala Asn Lys Lys Ala Val Asn Lys Arg Met Leu Glu Asn Lys Lys Glu  
 210 215 220  
 Asp Leu Glu Thr Ile Ile Asp Glu Phe Phe Ser Asp Ile Asp Lys Thr  
 225 230 235 240  
 Arg Pro Asn Asn Ile Pro Val Leu Glu Asp Glu Lys Gln Glu Glu Lys  
 245 250 255  
 Asn His Lys Asn Met Ala Gln Leu Lys Ser Asp Thr Glu Ala Ala Lys  
 260 265 270  
 Ser Asp Glu Ser Lys Arg Ser Lys Arg Ser Lys Arg Ser Leu Asn Thr  
 275 280 285  
 Gln Asn His Lys Pro Ala Ser Gln Glu Val Ser Glu Gln Gln Lys Ala  
 290 295 300  
 Glu Tyr Asp Lys Arg Ala Glu Glu Arg Lys Ala Arg Phe Leu Asp Asn  
 305 310 315 320  
 Gln Lys Ile Lys Lys Thr Pro Val Val Ser Leu Glu Tyr Asp Phe Glu  
 325 330 335  
 His Lys Gln Arg Ile Asp Asn Glu Asn Asp Lys Lys Leu Val Val Ser  
 340 345 350  
 Ala Pro Thr Lys Lys Pro Thr Ser Pro Thr Thr Tyr Thr Glu Thr Thr  
 355 360 365  
 Thr Gln Val Pro Met Pro Thr Val Glu Arg Gln Thr Gln Gln Gln Ile  
 370 375 380  
 Ile Tyr Asn Ala Pro Lys Gln Leu Ala Gly Leu Asn Gly Glu Ser His  
 385 390 395 400  
 Asp Phe Thr Thr Thr His Gln Ser Pro Thr Thr Ser Asn His Thr His  
 405 410 415  
 Asn Asn Val Val Glu Phe Glu Glu Thr Ser Ala Leu Pro Gly Arg Lys  
 420 425 430  
 Ser Gly Ser Leu Val Gly Ile Ser Gln Ile Asp Ser Ser His Leu Thr  
 435 440 445  
 Glu Arg Glu Lys Arg Val Ile Lys Arg Glu His Val Arg Glu Ala Gln  
 450 455 460  
 Lys Leu Val Asp Asn Tyr Lys Asp Thr His Ser Tyr Lys Asp Arg Ile

-continued

---

```

465                470                475                480
Asn Ala Gln Gln Lys Val Asn Thr Leu Ser Glu Gly His Gln Lys Arg
      485                490                495
Phe Asn Lys Gln Ile Asn Lys Val Tyr Asn Gly Lys
      500                505

<210> SEQ ID NO 33
<211> LENGTH: 520
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 33
Met Leu Thr Leu Gln Ile His Thr Gly Gly Ile Asn Leu Lys Lys Lys
 1                5                10                15
Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile Ala Ser Val Thr
 20                25                30
Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro Ala Ala Asn Ala
 35                40                45
Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr Gln Val Leu Asn
 50                55                60
Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu
 65                70                75                80
Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys
 85                90                95
Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln Gln Asn Asn Phe
 100               105               110
Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn
 115               120               125
Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp
 130               135               140
Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys Leu Asn Glu
 145               150               155               160
Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys Glu Gln Gln Asn
 165               170               175
Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn Glu Glu Gln Arg
 180               185               190
Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn
 195               200               205
Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala
 210               215               220
Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu
 225               230               235               240
His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser
 245               250               255
Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys
 260               265               270
Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys
 275               280               285
Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Thr
 290               295               300
Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser
 305               310               315               320
Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln

```

-continued

---

	325		330		335
Ala Pro Lys	Glu Asp Asn Asn	Lys Pro Gly Lys	Glu Asp Gly Asn		
	340	345	350		
Lys Pro Gly	Lys Glu Asp Asn Asn	Lys Pro Gly Lys	Glu Asp Asn Lys		
	355	360	365		
Lys Pro Gly	Lys Glu Asp Asn Asn	Lys Pro Gly Lys	Glu Asp Asn Asn		
	370	375	380		
Lys Pro Gly	Lys Glu Asp Gly Asn	Lys Pro Gly Lys	Glu Asp Asn Lys		
	385	390	395	400	
Lys Pro Gly	Lys Glu Asp Asn Asn	Lys Pro Gly Lys	Glu Asp Gly Asn		
	405	410	415		
Lys Pro Gly	Lys Glu Asp Gly Asn	Gly Val His Val	Val Lys Pro Gly		
	420	425	430		
Asp Thr Val	Asn Asp Ile Ala Lys	Ala Asn Gly Thr	Thr Ala Asp Lys		
	435	440	445		
Ile Ala Ala	Asp Asn Lys Leu Ala	Asp Lys Asn Met	Ile Lys Pro Gly		
	450	455	460		
Gln Glu Leu	Val Val Asp Lys Lys	Gln Pro Ala Asn	His Ala Asp Ala		
	465	470	475	480	
Asn Lys Ala	Gln Ala Leu Pro	Glu Thr Gly Glu	Glu Asn Pro Phe	Ile	
	485	490	495		
Gly Thr Thr	Val Phe Gly Gly	Leu Ser Leu Ala	Leu Gly Ala Ala	Leu	
	500	505	510		
Leu Ala Gly	Arg Arg Arg	Glu Leu			
	515	520			

&lt;210&gt; SEQ ID NO 34

&lt;211&gt; LENGTH: 291

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 34

Ala Gln His	Asp Glu Ala Lys Lys	Asn Ala Phe Tyr	Gln Val Leu Asn	
1	5	10	15	
Met Pro Asn	Leu Asn Ala Asp	Gln Arg Asn Gly	Phe Ile Gln Ser Leu	
	20	25	30	
Lys Ala Ala	Pro Ser Gln Ser	Ala Asn Val Leu	Gly Glu Ala Gln Lys	
	35	40	45	
Leu Asn Asp	Ser Gln Ala Pro	Lys Ala Asp Ala	Gln Gln Asn Asn Phe	
	50	55	60	
Asn Lys Asp	Lys Lys Ser Ala	Phe Tyr Glu Ile	Leu Asn Met Pro Asn	
	65	70	75	80
Leu Asn Glu	Ala Gln Arg Asn	Gly Phe Ile Gln	Ser Leu Lys Ala Ala	
	85	90	95	
Pro Ser Gln	Ser Thr Asn Val	Leu Gly Glu Ala	Lys Lys Leu Asn Glu	
	100	105	110	
Ser Gln Ala	Pro Lys Ala Asp	Asn Asn Phe Asn	Lys Glu Lys Lys Asn	
	115	120	125	
Ala Phe Tyr	Glu Ile Leu Asn	Met Pro Asn Leu	Asn Glu Glu Gln Arg	
	130	135	140	
Asn Gly Phe	Ile Gln Ser Leu	Lys Ala Ala Pro	Ser Gln Ser Ala Asn	
	145	150	155	160
Leu Leu Ser	Glu Ala Lys Lys	Leu Asn Glu Ser	Gln Ala Pro Lys Ala	



-continued

---

	165	170	175												
Asp	Asn	Lys	Phe	Asn	Lys	Glu	Lys	Lys	Asn	Ala	Phe	Tyr	Glu	Ile	Leu
	180				185								190		
His	Leu	Pro	Asn	Leu	Asn	Glu	Glu	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser
	195					200						205			
Leu	Lys	Ala	Ala	Pro	Ser	Gln	Ser	Ala	Asn	Leu	Leu	Ala	Glu	Ala	Lys
	210					215					220				
Lys	Leu	Asn	Asp	Ala	Gln	Ala	Pro	Lys	Ala	Asp	Asn	Lys	Phe	Asn	Lys
	225				230					235					240
Glu	Lys	Lys	Asn	Ala	Phe	Tyr	Glu	Ile	Leu	His	Leu	Pro	Asn	Leu	Thr
			245						250					255	
Glu	Glu	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	Ala	Ala	Pro	Ser
			260					265					270		
Val	Ser	Lys	Glu	Ile	Leu	Ala	Glu	Ala	Lys	Lys	Leu	Asn	Asp	Ala	Gln
	275						280					285			
Ala	Pro	Lys													
	290														

<210> SEQ ID NO 35  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 35  
 gctgcacata tggcgcaaca cgatgaagct caac 34

<210> SEQ ID NO 36  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 36  
 agtggatcct tatgctttgt tagcatctgc 30

<210> SEQ ID NO 37  
 <211> LENGTH: 19  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 37  
 Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro  
 1 5 10 15

Arg Gly Ser  
 <210> SEQ ID NO 38  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 38  
 aacatatggt caacaaagat caacaaagc 29

<210> SEQ ID NO 39  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 39

---

-continued

---

aaggatccag attcgtttaa ttttttagc 29

<210> SEQ ID NO 40  
<211> LENGTH: 43  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 40

cttcattcaa agtctttaaag cggccccaag ccaaagcact aac 43

<210> SEQ ID NO 41  
<211> LENGTH: 43  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 41

gtttagtgctt tggcttgggg cggtttaaag actttgaatg aag 43

<210> SEQ ID NO 42  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 42

catatgttca acaaagataa aaaaagcgcc ttctatgaaa tc 42

<210> SEQ ID NO 43  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 43

gatttcatag aaggcgcttt ttttatcttt gttgaacata tg 42

<210> SEQ ID NO 44  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 44

catatgttca acaaagatgg aggaagcgcc ttctatgaaa tc 42

<210> SEQ ID NO 45  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 45

gatttcatag aaggcgcttc ctccatcttt gttgaacata tg 42

<210> SEQ ID NO 46  
<211> LENGTH: 52  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 46

ggggacaagt ttgtacaaaa aagcaggctg atgactaagt tgaaaaaaga ag 52

<210> SEQ ID NO 47  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

---

-continued

---

<400> SEQUENCE: 47  
aaggatcccc tccaaatgt aattgccc 28

<210> SEQ ID NO 48  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 48  
aaggatccgt ttgtaactct atccaaagac 30

<210> SEQ ID NO 49  
<211> LENGTH: 49  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 49  
ggggaccact ttgtacaaga aagctgggtg acacctattg cagattcg 49

<210> SEQ ID NO 50  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 50  
ggggacaagt ttgtacaaaa aagcaggctc agatagcgat tcagattcag 50

<210> SEQ ID NO 51  
<211> LENGTH: 31  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 51  
aaggatccct gtattttctc cttaattttc c 31

<210> SEQ ID NO 52  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 52  
aaggatccca tggctgcaaa gcaaataatg 30

<210> SEQ ID NO 53  
<211> LENGTH: 51  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 53  
ggggaccact ttgtacaaga aagctgggtg ccctgggtgta acaaatttat g 51

<210> SEQ ID NO 54  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 54  
gaaggatccg tttattctag ttaatatata gttaatg 37

<210> SEQ ID NO 55

---

-continued

---

<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 55  
gaactgcagc tgtatgtctt tggatagagt tac 33

<210> SEQ ID NO 56  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 56  
gaaggatccg gtggcttttt tacttggatt ttc 33

<210> SEQ ID NO 57  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 57  
gaactgcagc gacaaactca ttatttgctt tgc 33

<210> SEQ ID NO 58  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 58  
gaactcgagt ctagcttatt tacatgg 27

<210> SEQ ID NO 59  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 59  
gaactcgaga tagaaggcag aatagtaaca aaggattata gtggg 45

<210> SEQ ID NO 60  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 60  
gtaggatcct gggatagagt tacaaac 27

<210> SEQ ID NO 61  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 61  
gaactcgagg cattatgtgt atcaciaaatt tggg 34

<210> SEQ ID NO 62  
<211> LENGTH: 43  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 62  
gaactcgaga tagaaggcag agtggtttct ggggagaaga atc 43

-continued

<210> SEQ ID NO 63  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 63

gaactcgagg cagccatgca ttaattattt gcc

33

<210> SEQ ID NO 64  
 <211> LENGTH: 940  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus aureus subst. Newman

<400> SEQUENCE: 64

Met Lys Ser Asn Leu Arg Tyr Gly Ile Arg Lys His Lys Leu Gly Ala  
 1 5 10 15  
 Ala Ser Val Phe Leu Gly Thr Met Ile Val Val Gly Met Gly Gln Glu  
 20 25 30  
 Lys Glu Ala Ala Ala Ser Glu Gln Asn Asn Thr Thr Val Glu Glu Ser  
 35 40 45  
 Gly Ser Ser Ala Thr Glu Ser Lys Ala Ser Glu Thr Gln Thr Thr Thr  
 50 55 60  
 Asn Asn Val Asn Thr Ile Asp Glu Thr Gln Ser Tyr Ser Ala Thr Ser  
 65 70 75 80  
 Thr Glu Gln Pro Ser Gln Ser Thr Gln Val Thr Thr Glu Glu Ala Pro  
 85 90 95  
 Lys Thr Val Gln Ala Pro Lys Val Glu Thr Ser Arg Val Asp Leu Pro  
 100 105 110  
 Ser Glu Lys Val Ala Asp Lys Glu Thr Thr Gly Thr Gln Val Asp Ile  
 115 120 125  
 Ala Gln Pro Ser Asn Val Ser Glu Ile Lys Pro Arg Met Lys Arg Ser  
 130 135 140  
 Thr Asp Val Thr Ala Val Ala Glu Lys Glu Val Val Glu Glu Thr Lys  
 145 150 155 160  
 Ala Thr Gly Thr Asp Val Thr Asn Lys Val Glu Val Glu Glu Gly Ser  
 165 170 175  
 Glu Ile Val Gly His Lys Gln Asp Thr Asn Val Val Asn Pro His Asn  
 180 185 190  
 Ala Glu Arg Val Thr Leu Lys Tyr Lys Trp Lys Phe Gly Glu Gly Ile  
 195 200 205  
 Lys Ala Gly Asp Tyr Phe Asp Phe Thr Leu Ser Asp Asn Val Glu Thr  
 210 215 220  
 His Gly Ile Ser Thr Leu Arg Lys Val Pro Glu Ile Lys Ser Thr Asp  
 225 230 235 240  
 Gly Gln Val Met Ala Thr Gly Glu Ile Ile Gly Glu Arg Lys Val Arg  
 245 250 255  
 Tyr Thr Phe Lys Glu Tyr Val Gln Glu Lys Lys Asp Leu Thr Ala Glu  
 260 265 270  
 Leu Ser Leu Asn Leu Phe Ile Asp Pro Thr Thr Val Thr Gln Lys Gly  
 275 280 285  
 Asn Gln Asn Val Glu Val Lys Leu Gly Glu Thr Thr Val Ser Lys Ile  
 290 295 300  
 Phe Asn Ile Gln Tyr Leu Gly Gly Val Arg Asp Asn Trp Gly Val Thr

-continued

---

305				310					315					320	
Ala	Asn	Gly	Arg	Ile	Asp	Thr	Leu	Asn	Lys	Val	Asp	Gly	Lys	Phe	Ser
				325					330					335	
His	Phe	Ala	Tyr	Met	Lys	Pro	Asn	Asn	Gln	Ser	Leu	Ser	Ser	Val	Thr
			340					345						350	
Val	Thr	Gly	Gln	Val	Thr	Lys	Gly	Asn	Lys	Pro	Gly	Val	Asn	Asn	Pro
		355					360					365			
Thr	Val	Lys	Val	Tyr	Lys	His	Ile	Gly	Ser	Asp	Asp	Leu	Ala	Glu	Ser
	370					375					380				
Val	Tyr	Ala	Lys	Leu	Asp	Asp	Val	Ser	Lys	Phe	Glu	Asp	Val	Thr	Asp
385					390					395					400
Asn	Met	Ser	Leu	Asp	Phe	Asp	Thr	Asn	Gly	Gly	Tyr	Ser	Leu	Asn	Phe
				405					410					415	
Asn	Asn	Leu	Asp	Gln	Ser	Lys	Asn	Tyr	Val	Ile	Lys	Tyr	Glu	Gly	Tyr
			420					425						430	
Tyr	Asp	Ser	Asn	Ala	Ser	Asn	Leu	Glu	Phe	Gln	Thr	His	Leu	Phe	Gly
	435						440					445			
Tyr	Tyr	Asn	Tyr	Tyr	Tyr	Thr	Ser	Asn	Leu	Thr	Trp	Lys	Asn	Gly	Val
	450					455					460				
Ala	Phe	Tyr	Ser	Asn	Asn	Ala	Gln	Gly	Asp	Gly	Lys	Asp	Lys	Leu	Lys
465						470				475					480
Glu	Pro	Ile	Ile	Glu	His	Ser	Thr	Pro	Ile	Glu	Leu	Glu	Phe	Lys	Ser
				485					490					495	
Glu	Pro	Pro	Val	Glu	Lys	His	Glu	Leu	Thr	Gly	Thr	Ile	Glu	Glu	Ser
			500					505						510	
Asn	Asp	Ser	Lys	Pro	Ile	Asp	Phe	Glu	Tyr	His	Thr	Ala	Val	Glu	Gly
		515					520					525			
Ala	Glu	Gly	His	Ala	Glu	Gly	Thr	Ile	Glu	Thr	Glu	Glu	Asp	Ser	Ile
	530					535					540				
His	Val	Asp	Phe	Glu	Glu	Ser	Thr	His	Glu	Asn	Ser	Lys	His	His	Ala
545						550				555					560
Asp	Val	Val	Glu	Tyr	Glu	Glu	Asp	Thr	Asn	Pro	Gly	Gly	Gly	Gln	Val
				565					570					575	
Thr	Thr	Glu	Ser	Asn	Leu	Val	Glu	Phe	Asp	Glu	Asp	Ser	Thr	Lys	Gly
		580						585						590	
Ile	Val	Thr	Gly	Ala	Val	Ser	Asp	His	Thr	Thr	Ile	Glu	Asp	Thr	Lys
		595					600					605			
Glu	Tyr	Thr	Thr	Glu	Ser	Asn	Leu	Ile	Glu	Leu	Val	Asp	Glu	Leu	Pro
	610						615					620			
Glu	Glu	His	Gly	Gln	Ala	Gln	Gly	Pro	Ile	Glu	Glu	Ile	Thr	Glu	Asn
625						630				635					640
Asn	His	His	Ile	Ser	His	Ser	Gly	Leu	Gly	Thr	Glu	Asn	Gly	His	Gly
				645					650					655	
Asn	Tyr	Gly	Val	Ile	Glu	Glu	Ile	Glu	Glu	Asn	Ser	His	Val	Asp	Ile
			660					665						670	
Lys	Ser	Glu	Leu	Gly	Tyr	Glu	Gly	Gly	Gln	Asn	Ser	Gly	Asn	Gln	Ser
		675					680						685		
Phe	Glu	Glu	Asp	Thr	Glu	Glu	Asp	Lys	Pro	Lys	Tyr	Glu	Gln	Gly	Gly
	690						695					700			
Asn	Ile	Val	Asp	Ile	Asp	Phe	Asp	Ser	Val	Pro	Gln	Ile	His	Gly	Gln
705					710						715				720

-continued

---

```

Asn Asn Gly Asn Gln Ser Phe Glu Glu Asp Thr Glu Lys Asp Lys Pro
      725                                730                    735

Lys Tyr Glu Gln Gly Gly Asn Ile Ile Asp Ile Asp Phe Asp Ser Val
      740                                745                    750

Pro His Ile His Gly Phe Asn Lys His Thr Glu Ile Ile Glu Glu Asp
      755                                760                    765

Thr Asn Lys Asp Lys Pro Asn Tyr Gln Phe Gly Gly His Asn Ser Val
      770                                775                    780

Asp Phe Glu Glu Asp Thr Leu Pro Gln Val Ser Gly His Asn Glu Gly
      785                                790                    795                    800

Gln Gln Thr Ile Glu Glu Asp Thr Thr Pro Pro Ile Val Pro Pro Thr
      805                                810                    815

Pro Pro Thr Pro Glu Val Pro Ser Glu Pro Glu Thr Pro Thr Pro Pro
      820                                825                    830

Thr Pro Glu Val Pro Ser Glu Pro Glu Thr Pro Thr Pro Pro Thr Pro
      835                                840                    845

Glu Val Pro Thr Glu Pro Gly Lys Pro Ile Pro Pro Ala Lys Glu Glu
      850                                855                    860

Pro Lys Lys Pro Ser Lys Pro Val Glu Gln Gly Lys Val Val Thr Pro
      865                                870                    875                    880

Val Ile Glu Ile Asn Glu Lys Val Lys Ala Val Val Pro Thr Lys Lys
      885                                890                    895

Ala Gln Ser Lys Lys Ser Glu Leu Pro Glu Thr Gly Gly Glu Glu Ser
      900                                905                    910

Thr Asn Asn Gly Met Leu Phe Gly Gly Leu Phe Ser Ile Leu Gly Leu
      915                                920                    925

Ala Leu Leu Arg Arg Asn Lys Lys Asn His Lys Ala
      930                                935                    940
    
```

```

<210> SEQ ID NO 65
<211> LENGTH: 1315
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus subst. Newman
    
```

```

<400> SEQUENCE: 65

Met Leu Asn Arg Glu Asn Lys Thr Ala Ile Thr Arg Lys Gly Met Val
 1      5      10

Ser Asn Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Thr Val Gly Thr
 20     25     30

Ala Ser Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Gly Asn Gln
 35     40     45

Glu Ala Lys Ala Ala Glu Ser Thr Asn Lys Glu Leu Asn Glu Ala Thr
 50     55     60

Thr Ser Ala Ser Asp Asn Gln Ser Ser Asp Lys Val Asp Met Gln Gln
 65     70     75     80

Leu Asn Gln Glu Asp Asn Thr Lys Asn Asp Asn Gln Lys Glu Met Val
 85     90     95

Ser Ser Gln Gly Asn Glu Thr Thr Ser Asn Gly Asn Lys Leu Ile Glu
100    105    110

Lys Glu Ser Val Gln Ser Thr Thr Gly Asn Lys Val Glu Val Ser Thr
115    120    125

Ala Lys Ser Asp Glu Gln Ala Ser Pro Lys Ser Thr Asn Glu Asp Leu
130    135    140
    
```





-continued

---

Gln Ser Gly Gly Ala Gly Gln Glu Val Tyr Lys Ile Gly Asn Tyr Val  
                   565                                  570                                  575  
 Trp Glu Asp Thr Asn Lys Asn Gly Val Gln Glu Leu Gly Glu Lys Gly  
                   580                                  585                                  590  
 Val Gly Asn Val Thr Val Thr Val Phe Asp Asn Asn Thr Asn Thr Lys  
                   595                                  600                                  605  
 Val Gly Glu Ala Val Thr Lys Glu Asp Gly Ser Tyr Leu Ile Pro Asn  
                   610                                  615                                  620  
 Leu Pro Asn Gly Asp Tyr Arg Val Glu Phe Ser Asn Leu Pro Lys Gly  
                   625                                  630                                  635                                  640  
 Tyr Glu Val Thr Pro Ser Lys Gln Gly Asn Asn Glu Glu Leu Asp Ser  
                   645                                  650                                  655  
 Asn Gly Leu Ser Ser Val Ile Thr Val Asn Gly Lys Asp Asn Leu Ser  
                   660                                  665                                  670  
 Ala Asp Leu Gly Ile Tyr Lys Pro Lys Tyr Asn Leu Gly Asp Tyr Val  
                   675                                  680                                  685  
 Trp Glu Asp Thr Asn Lys Asn Gly Ile Gln Asp Gln Asp Glu Lys Gly  
                   690                                  695                                  700  
 Ile Ser Gly Val Thr Val Thr Leu Lys Asp Glu Asn Gly Asn Val Leu  
                   705                                  710                                  715                                  720  
 Lys Thr Val Thr Thr Asp Ala Asp Gly Lys Tyr Lys Phe Thr Asp Leu  
                   725                                  730                                  735  
 Asp Asn Gly Asn Tyr Lys Val Glu Phe Thr Thr Pro Glu Gly Tyr Thr  
                   740                                  745                                  750  
 Pro Thr Thr Val Thr Ser Gly Ser Asp Ile Glu Lys Asp Ser Asn Gly  
                   755                                  760                                  765  
 Leu Thr Thr Thr Gly Val Ile Asn Gly Ala Asp Asn Met Thr Leu Asp  
                   770                                  775                                  780  
 Ser Gly Phe Tyr Lys Thr Pro Lys Tyr Asn Leu Gly Asn Tyr Val Trp  
                   785                                  790                                  795                                  800  
 Glu Asp Thr Asn Lys Asp Gly Lys Gln Asp Ser Thr Glu Lys Gly Ile  
                   805                                  810                                  815  
 Ser Gly Val Thr Val Thr Leu Lys Asn Glu Asn Gly Glu Val Leu Gln  
                   820                                  825                                  830  
 Thr Thr Lys Thr Asp Lys Asp Gly Lys Tyr Gln Phe Thr Gly Leu Glu  
                   835                                  840                                  845  
 Asn Gly Thr Tyr Lys Val Glu Phe Glu Thr Pro Ser Gly Tyr Thr Pro  
                   850                                  855                                  860  
 Thr Gln Val Gly Ser Gly Thr Asp Glu Gly Ile Asp Ser Asn Gly Thr  
                   865                                  870                                  875                                  880  
 Ser Thr Thr Gly Val Ile Lys Asp Lys Asp Asn Asp Thr Ile Asp Ser  
                   885                                  890                                  895  
 Gly Phe Tyr Lys Pro Thr Tyr Asn Leu Gly Asp Tyr Val Trp Glu Asp  
                   900                                  905                                  910  
 Thr Asn Lys Asn Gly Val Gln Asp Lys Asp Glu Lys Gly Ile Ser Gly  
                   915                                  920                                  925  
 Val Thr Val Thr Leu Lys Asp Glu Asn Asp Lys Val Leu Lys Thr Val  
                   930                                  935                                  940  
 Thr Thr Asp Glu Asn Gly Lys Tyr Gln Phe Thr Asp Leu Asn Asn Gly  
                   945                                  950                                  955                                  960  
 Thr Tyr Lys Val Glu Phe Glu Thr Pro Ser Gly Tyr Thr Pro Thr Ser

-continued

965					970					975					
Val	Thr	Ser	Gly	Asn	Asp	Thr	Glu	Lys	Asp	Ser	Asn	Gly	Leu	Thr	Thr
			980					985					990		
Thr	Gly	Val	Ile	Lys	Asp	Ala	Asp	Asn	Met	Thr	Leu	Asp	Ser	Gly	Phe
		995					1000					1005			
Tyr	Lys	Thr	Pro	Lys	Tyr	Ser	Leu	Gly	Asp	Tyr	Val	Trp	Tyr	Asp	
	1010					1015					1020				
Ser	Asn	Lys	Asp	Gly	Lys	Gln	Asp	Ser	Thr	Glu	Lys	Gly	Ile	Lys	
	1025					1030					1035				
Asp	Val	Lys	Val	Thr	Leu	Leu	Asn	Glu	Lys	Gly	Glu	Val	Ile	Gly	
	1040					1045					1050				
Thr	Thr	Lys	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	Cys	Phe	Asp	Asn	Leu	
	1055					1060					1065				
Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu	Lys	Pro	Ala	Gly	Leu	
	1070					1075					1080				
Thr	Gln	Thr	Gly	Thr	Asn	Thr	Thr	Glu	Asp	Asp	Lys	Asp	Ala	Asp	
	1085					1090					1095				
Gly	Gly	Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp	Asp	Phe	Thr	
	1100					1105					1110				
Leu	Asp	Asn	Gly	Tyr	Tyr	Glu	Glu	Glu	Thr	Ser	Asp	Ser	Asp	Ser	
	1115					1120					1125				
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Arg	Asp	Ser	Asp	Ser	Asp	
	1130					1135					1140				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
	1145					1150					1155				
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Arg	Asp	Ser	Asp	Ser	Asp	
	1160					1165					1170				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
	1175					1180					1185				
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
	1190					1195					1200				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
	1205					1210					1215				
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
	1220					1225					1230				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
	1235					1240					1245				
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ala	Gly	Lys	His	Thr	Pro	Val	Lys	
	1250					1255					1260				
Pro	Met	Ser	Thr	Thr	Lys	Asp	His	His	Asn	Lys	Ala	Lys	Ala	Leu	
	1265					1270					1275				
Pro	Glu	Thr	Gly	Asn	Glu	Asn	Ser	Gly	Ser	Asn	Asn	Ala	Thr	Leu	
	1280					1285					1290				
Phe	Gly	Gly	Leu	Phe	Ala	Ala	Leu	Gly	Ser	Leu	Leu	Leu	Phe	Gly	
	1295					1300					1305				
Arg	Arg	Lys	Lys	Gln	Asn	Lys									
	1310					1315									

&lt;210&gt; SEQ ID NO 66

&lt;211&gt; LENGTH: 933

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus aureus subst. Newman

-continued

&lt;400&gt; SEQUENCE: 66

Met Asn Met Lys Lys Lys Glu Lys His Ala Ile Arg Lys Lys Ser Ile  
 1 5 10 15  
 Gly Val Ala Ser Val Leu Val Gly Thr Leu Ile Gly Phe Gly Leu Leu  
 20 25 30  
 Ser Ser Lys Glu Ala Asp Ala Ser Glu Asn Ser Val Thr Gln Ser Asp  
 35 40 45  
 Ser Ala Ser Asn Glu Ser Lys Ser Asn Asp Ser Ser Val Ser Ala  
 50 55 60  
 Ala Pro Lys Thr Asp Asp Thr Asn Val Ser Asp Thr Lys Thr Ser Ser  
 65 70 75 80  
 Asn Thr Asn Asn Gly Glu Thr Ser Val Ala Gln Asn Pro Ala Gln Gln  
 85 90 95  
 Glu Thr Thr Gln Ser Ser Ser Thr Asn Ala Thr Thr Glu Glu Thr Pro  
 100 105 110  
 Val Thr Gly Glu Ala Thr Thr Thr Thr Asn Gln Ala Asn Thr Pro  
 115 120 125  
 Ala Thr Thr Gln Ser Ser Asn Thr Asn Ala Glu Glu Leu Val Asn Gln  
 130 135 140  
 Thr Ser Asn Glu Thr Thr Phe Asn Asp Thr Asn Thr Val Ser Ser Val  
 145 150 155 160  
 Asn Ser Pro Gln Asn Ser Thr Asn Ala Glu Asn Val Ser Thr Thr Gln  
 165 170 175  
 Asp Thr Ser Thr Glu Ala Thr Pro Ser Asn Asn Glu Ser Ala Pro Gln  
 180 185 190  
 Ser Thr Asp Ala Ser Asn Lys Asp Val Val Asn Gln Ala Val Asn Thr  
 195 200 205  
 Ser Ala Pro Arg Met Arg Ala Phe Ser Leu Ala Ala Val Ala Ala Asp  
 210 215 220  
 Ala Pro Ala Ala Gly Thr Asp Ile Thr Asn Gln Leu Thr Asn Val Thr  
 225 230 235 240  
 Val Gly Ile Asp Ser Gly Thr Thr Val Tyr Pro His Gln Ala Gly Tyr  
 245 250 255  
 Val Lys Leu Asn Tyr Gly Phe Ser Val Pro Asn Ser Ala Val Lys Gly  
 260 265 270  
 Asp Thr Phe Lys Ile Thr Val Pro Lys Glu Leu Asn Leu Asn Gly Val  
 275 280 285  
 Thr Ser Thr Ala Lys Val Pro Pro Ile Met Ala Gly Asp Gln Val Leu  
 290 295 300  
 Ala Asn Gly Val Ile Asp Ser Asp Gly Asn Val Ile Tyr Thr Phe Thr  
 305 310 315 320  
 Asp Tyr Val Asn Thr Lys Asp Asp Val Lys Ala Thr Leu Thr Met Pro  
 325 330 335  
 Ala Tyr Ile Asp Pro Glu Asn Val Lys Lys Thr Gly Asn Val Thr Leu  
 340 345 350  
 Ala Thr Gly Ile Gly Ser Thr Thr Ala Asn Lys Thr Val Leu Val Asp  
 355 360 365  
 Tyr Glu Lys Tyr Gly Lys Phe Tyr Asn Leu Ser Ile Lys Gly Thr Ile  
 370 375 380  
 Asp Gln Ile Asp Lys Thr Asn Asn Thr Tyr Arg Gln Thr Ile Tyr Val  
 385 390 395 400

-continued

---

Asn	Pro	Ser	Gly	Asp	Asn	Val	Ile	Ala	Pro	Val	Leu	Thr	Gly	Asn	Leu
				405					410					415	
Lys	Pro	Asn	Thr	Asp	Ser	Asn	Ala	Leu	Ile	Asp	Gln	Gln	Asn	Thr	Ser
			420					425					430		
Ile	Lys	Val	Tyr	Lys	Val	Asp	Asn	Ala	Ala	Asp	Leu	Ser	Glu	Ser	Tyr
		435					440					445			
Phe	Val	Asn	Pro	Glu	Asn	Phe	Glu	Asp	Val	Thr	Asn	Ser	Val	Asn	Ile
	450					455					460				
Thr	Phe	Pro	Asn	Pro	Asn	Gln	Tyr	Lys	Val	Glu	Phe	Asn	Thr	Pro	Asp
465					470					475					480
Asp	Gln	Ile	Thr	Thr	Pro	Tyr	Ile	Val	Val	Val	Asn	Gly	His	Ile	Asp
				485					490					495	
Pro	Asn	Ser	Lys	Gly	Asp	Leu	Ala	Leu	Arg	Ser	Thr	Leu	Tyr	Gly	Tyr
			500					505					510		
Asn	Ser	Asn	Ile	Ile	Trp	Arg	Ser	Met	Ser	Trp	Asp	Asn	Glu	Val	Ala
		515					520					525			
Phe	Asn	Asn	Gly	Ser	Gly	Ser	Gly	Asp	Gly	Ile	Asp	Lys	Pro	Val	Val
	530					535					540				
Pro	Glu	Gln	Pro	Asp	Glu	Pro	Gly	Glu	Ile	Glu	Pro	Ile	Pro	Glu	Asp
545					550					555					560
Ser	Asp	Ser	Asp	Pro	Gly	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asn	Ser	Asp
				565					570					575	
Ser	Gly	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Thr	Ser	Asp	Ser	Gly	Ser	Asp
			580					585					590		
Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp
		595					600						605		
Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Asp
	610						615						620		
Asn	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
625					630					635					640
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
				645						650				655	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			660							665				670	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		675											685		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		690					695						700		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
705					710					715					720
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
				725						730				735	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			740							745				750	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Ala
		755											765		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		770					775						780		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
785					790					795					800
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp
				805						810				815	

-continued

---

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp  
820 825 830

Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Gly Ser Asp Ser Asp  
835 840 845

Ser Ser Ser Asp Ser Asp Ser Glu Ser Asp Ser Asn Ser Asp Ser Glu  
850 855 860

Ser Gly Ser Asn Asn Asn Val Val Pro Pro Asn Ser Pro Lys Asn Gly  
865 870 875 880

Thr Asn Ala Ser Asn Lys Asn Glu Ala Lys Asp Ser Lys Glu Pro Leu  
885 890 895

Pro Asp Thr Gly Ser Glu Asp Glu Ala Asn Thr Ser Leu Ile Trp Gly  
900 905 910

Leu Leu Ala Ser Ile Gly Ser Leu Leu Leu Phe Arg Arg Lys Lys Glu  
915 920 925

Asn Lys Asp Lys Lys  
930

&lt;210&gt; SEQ ID NO 67

&lt;211&gt; LENGTH: 677

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus aureus subst. Newman

&lt;400&gt; SEQUENCE: 67

Met Lys Ser Asn Leu Arg Tyr Gly Ile Arg Lys His Lys Leu Gly Ala  
1 5 10 15

Ala Ser Val Phe Leu Gly Thr Met Ile Val Val Gly Met Gly Gln Glu  
20 25 30

Lys Glu Ala Ala Ala Ser Glu Gln Asn Asn Thr Thr Val Glu Glu Ser  
35 40 45

Gly Ser Ser Ala Thr Glu Ser Lys Ala Ser Glu Thr Gln Thr Thr Thr  
50 55 60

Asn Asn Val Asn Thr Ile Asp Glu Thr Gln Ser Tyr Ser Ala Thr Ser  
65 70 75 80

Thr Glu Gln Pro Ser Gln Ser Thr Gln Val Thr Thr Glu Glu Ala Pro  
85 90 95

Lys Thr Val Gln Ala Pro Lys Val Glu Thr Ser Arg Val Asp Leu Pro  
100 105 110

Ser Glu Lys Val Ala Asp Lys Glu Thr Thr Gly Thr Gln Val Asp Ile  
115 120 125

Ala Gln Pro Ser Asn Val Ser Glu Ile Lys Pro Arg Met Lys Arg Ser  
130 135 140

Thr Asp Val Thr Ala Val Ala Glu Lys Glu Val Val Glu Glu Thr Lys  
145 150 155 160

Ala Thr Gly Thr Asp Val Thr Asn Lys Val Glu Val Glu Glu Gly Ser  
165 170 175

Glu Ile Val Gly His Lys Gln Asp Thr Asn Val Val Asn Pro His Asn  
180 185 190

Ala Glu Arg Val Thr Leu Lys Tyr Lys Trp Lys Phe Gly Glu Gly Ile  
195 200 205

Lys Ala Gly Asp Tyr Phe Asp Phe Thr Leu Ser Asp Asn Val Glu Thr  
210 215 220

His Gly Ile Ser Thr Leu Arg Lys Val Pro Glu Ile Lys Ser Thr Asp  
225 230 235 240



-continued

645	650	655
Asn Tyr Gly Val Ile Glu Glu Ile Glu Glu Asn Ser His Val Asp Ile		
660	665	670
Lys Ser Glu Leu Gly		
675		

**1.** An immunogenic composition comprising an isolated peptide comprising a Protein A (SpA) variant having an amino acid substitution at amino acid positions 9, 10, 36, and 37 of SEQ ID NO:2 in combination with SdrD, ClfA, or FnbpB polypeptides.

**2.** The immunogenic composition of claim **1**, comprising isolated SdrD, ClfA, and FnbpB polypeptide.

**3.-8.** (canceled)

**9.** A method for treating a staphylococcal infection in a subject comprising providing to a subject having, suspected of having or at risk of developing a staphylococcal infection a composition comprising an effective amount of an isolated peptide comprising a Protein A (SpA) variant having an amino acid substitution at amino acid positions 9 and 10 of SEQ ID NO:2 in combination with isolated SdrD, ClfA, or FnbpB polypeptides.

**10.** The method of claim **9**, wherein the composition comprises isolated SdrD, ClfA, and FnbpB polypeptide.

**11.** The method of claim **9**, wherein the SdrD, ClfA, and FnbpB polypeptides are from *Staphylococcus aureus*.

**12.** The method of claim **10**, wherein the composition is essentially free of other staphylococcal polypeptides.

**13.** The method of claim **10**, wherein the composition is essentially free of other staphylococcal carbohydrates.

**14.** The method of claim **10**, wherein the staphylococcal polypeptides in the composition consist essentially of the SpA variant and the isolated SdrD, ClfA, and FnbpB polypeptides.

**15.** The method of claim **10**, wherein the composition consists essentially of the SpA variant, the isolated SdrD, ClfA, and FnbpB polypeptides and an adjuvant.

**16.** An immunogenic composition comprising isolated SdrD, ClfA, or FnbpB polypeptides.

**17.** The immunogenic composition of claim **16**, comprising isolated SdrD, ClfA, and FnbpB polypeptide.

**18.-23.** (canceled)

**24.** A method for treating a staphylococcal infection in a subject comprising providing to a subject having, suspected of having or at risk of developing a staphylococcal infection an effective amount of a peptide composition comprising isolated SdrD, ClfA, or FnbpB polypeptides.

**25.** The method of claim **24**, wherein the peptide composition comprises isolated SdrD, ClfA, and FnbpB polypeptide.

**26.** The method of claim **24**, wherein the SdrD, ClfA, and FnbpB polypeptides are from *Staphylococcus aureus*.

**27.** The method of claim **24**, wherein the peptide composition further comprising an adjuvant.

**28.** The method of claim **25**, wherein the peptide composition is essentially free of other staphylococcal polypeptides.

**29.** The method of claim **25**, wherein the peptide composition is essentially free of other staphylococcal carbohydrates.

**30.** The method of claim **25**, wherein the peptide composition comprises staphylococcal polypeptides consisting essentially of the isolated SdrD, ClfA, and FnbpB polypeptides.

**31.** The method of claim **25**, wherein the peptide composition consists essentially of the isolated SdrD, ClfA, and FnbpB polypeptides and an adjuvant.

\* \* \* \* \*