



(19) **United States**

(12) **Patent Application Publication**  
**Schneewind**

(10) **Pub. No.: US 2013/0171183 A1**

(43) **Pub. Date: Jul. 4, 2013**

(54) **COMPOSITIONS AND METHODS RELATED TO PROTEIN A (SPA) VARIANTS**

**Publication Classification**

(75) Inventor: **Olaf Schneewind**, Chicago, IL (US)

(51) **Int. Cl.**  
*A61K 39/085* (2006.01)  
*C07K 14/31* (2006.01)

(73) Assignee: **UNIVERSITY OF CHICAGO**,  
CHICAGO, IL (US)

(52) **U.S. Cl.**  
CPC ..... *A61K 39/085* (2013.01); *C07K 14/31* (2013.01)  
USPC ..... **424/190.1**; 530/324

(21) Appl. No.: **13/807,598**

(22) PCT Filed: **Jul. 1, 2011**

(86) PCT No.: **PCT/US11/42845**


(57) **ABSTRACT**

§ 371 (c)(1),  
(2), (4) Date: **Mar. 19, 2013**

The present invention concerns methods and compositions for treating or preventing a bacterial infection, particularly infection by a *Staphylococcus* bacterium. The invention provides methods and compositions for stimulating an immune response against the bacteria. In certain embodiments, the methods and compositions involve a non-toxicogenic Protein A (SpA) variant.

**Related U.S. Application Data**


(60) Provisional application No. 61/361,218, filed on Jul. 2, 2010, provisional application No. 61/370,725, filed on Aug. 4, 2010.



WO 2012/003474

PCT/US2011/042845

1/12



FIGs. 1A-1B

WO 2012/003474

PCT/US2011/042845




FIG. 2




FIG. 3

WO 2012/003474

PCT/US2011/042845




FIG. 4

WO 2012/003474

PCT/US2011/042845

5/12




FIG. 5

WO 2012/003474

PCT/US2011/042845

6/12




FIG. 6

WO 2012/003474

PCT/US2011/042845




FIG. 7



WO 2012/003474

PCT/US2011/042845




FIG. 8

WO 2012/003474

PCT/US2011/042845

9/12




FIG. 9

WO 2012/003474

PCT/US2011/042845

10/12




FIG. 10

WO 2012/003474

PCT/US2011/042845

11/12




FIG. 11

WO 2012/003474

PCT/US2011/042845

12/12




FIG. 12A-12C

## COMPOSITIONS AND METHODS RELATED TO PROTEIN A (SPA) VARIANTS

[0001] This application claims the benefit of priority to U.S. Provisional Patent Application Ser. No. 61/361,218 filed Jul. 2, 2010, and 61/370,725 filed Aug. 4, 2010, hereby incorporated by reference in their entirety.

[0002] This invention was made with government support under AI057153, AI052474, and GM007281 awarded by the National Institutes of Health. The government has certain rights in the invention.

### BACKGROUND OF THE INVENTION

[0003] I. Field of the Invention

[0004] The present invention relates generally to the fields of immunology, microbiology, and pathology. More particularly, it concerns methods and compositions involving bacterial Protein A variants, which can be used to invoke an immune response against the bacteria.

[0005] II. Background

[0006] The number of both community acquired and hospital acquired infections have increased over recent years with the increased use of intravascular devices. Hospital acquired (nosocomial) infections are a major cause of morbidity and mortality, more particularly in the United States, where it affects more than 2 million patients annually. The most frequent infections are urinary tract infections (33% of the infections), followed by pneumonia (15.5%), surgical site infections (14.8%) and primary bloodstream infections (13%) (Emori and Gaynes, 1993).

[0007] The major nosocomial pathogens include *Staphylococcus aureus*, coagulase-negative Staphylococci (mostly *Staphylococcus epidermidis*), *enterococcus* spp., *Escherichia coli* and *Pseudomonas aeruginosa*. Although these pathogens cause approximately the same number of infections, the severity of the disorders they can produce combined with the frequency of antibiotic resistant isolates balance this ranking towards *S. aureus* and *S. epidermidis* as being the most significant nosocomial pathogens.

[0008] Staphylococci can cause a wide variety of diseases in humans and other animals through either toxin production or invasion. Staphylococcal toxins are also a common cause of food poisoning, as the bacteria can grow in improperly stored food.

[0009] *Staphylococcus epidermidis* is a normal skin commensal which is also an important opportunistic pathogen responsible for infections of impaired medical devices and infections at sites of surgery. Medical devices infected by *S. epidermidis* include cardiac pacemakers, cerebrospinal fluid shunts, continuous ambulatory peritoneal dialysis catheters, orthopedic devices and prosthetic heart valves.

[0010] *Staphylococcus aureus* is the most common cause of nosocomial infections with a significant morbidity and mortality. It is the cause of some cases of osteomyelitis, endocarditis, septic arthritis, pneumonia, abscesses, and toxic shock syndrome. *S. aureus* can survive on dry surfaces, increasing the chance of transmission. Any *S. aureus* infection can cause the staphylococcal scalded skin syndrome, a cutaneous reaction to exotoxin absorbed into the bloodstream. It can also cause a type of septicemia called pyaemia that can be life-threatening. Problematically, Methicillin-resistant *Staphylococcus aureus* (MRSA) has become a major cause of hospital-acquired infections.

[0011] *S. aureus* and *S. epidermidis* infections are typically treated with antibiotics, with penicillin being the drug of choice, whereas vancomycin is used for methicillin resistant isolates. The percentage of staphylococcal strains exhibiting wide-spectrum resistance to antibiotics has become increasingly prevalent, posing a threat for effective antimicrobial therapy. In addition, the recent emergence of vancomycin resistant *S. aureus* strain has aroused fear that MRSA strains are emerging and spreading for which no effective therapy is available.

[0012] An alternative to antibiotic treatment for staphylococcal infections is under investigation that uses antibodies directed against staphylococcal antigens. This therapy involves administration of polyclonal antisera (WO00/15238, WO00/12132) or treatment with monoclonal antibodies against lipoteichoic acid (WO98/57994).

[0013] An alternative approach would be the use of active vaccination to generate an immune response against staphylococci. The *S. aureus* genome has been sequenced and many of the coding sequences have been identified (WO02/094868, EP0786519), which can lead to the identification of potential antigens. The same is true for *S. epidermidis* (WO01/34809). As a refinement of this approach, others have identified proteins that are recognized by hyperimmune sera from patients who have suffered staphylococcal infection (WO01/98499, WO02/059148).

[0014] *S. aureus* secretes a plethora of virulence factors into the extracellular milieu (Archer, 1998; Dinges et al., 2000; Foster, 2005; Shaw et al., 2004; Sibbald et al., 2006). Like most secreted proteins, these virulence factors are translocated by the Sec machinery across the plasma membrane. Proteins secreted by the Sec machinery bear an N-terminal leader peptide that is removed by leader peptidase once the pre-protein is engaged in the Sec translocon (Dalbey and Wickner, 1985; van Wely et al., 2001). Recent genome analysis suggests that Actinobacteria and members of the Firmicutes encode an additional secretion system that recognizes a subset of proteins in a Sec-independent manner (Pallen, 2002). ESAT-6 (early secreted antigen target 6 kDa) and CFP-10 (culture filtrate antigen 10 kDa) of *Mycobacterium tuberculosis* represent the first substrates of this novel secretion system termed ESX-1 or 5 nm in *M. tuberculosis* (Andersen et al., 1995; Hsu et al., 2003; Pym et al., 2003; Stanley et al., 2003). In *S. aureus*, two ESAT-6 like factors designated EsxA and EsxB are secreted by the Ess pathway (ESAT-6 secretion system) (Burts et al., 2005).

[0015] The first generation of vaccines targeted against *S. aureus* or against the exoproteins it produces have met with limited success (Lee, 1996). There remains a need to develop effective vaccines against staphylococcal infections. Additional compositions for treating staphylococcal infections are also needed.

### SUMMARY OF THE INVENTION

[0016] Protein A (SpA) (SEQ ID NO:33), a cell wall anchored surface protein of *Staphylococcus aureus*, provides for bacterial evasion from innate and adaptive immune responses. Protein A binds immunoglobulins at their Fc portion, interacts with the VH3 domain of B cell receptors inappropriately stimulating B cell proliferation and apoptosis, binds to von Willebrand factor A1 domains to activate intracellular clotting, and also binds to the TNF Receptor-1 to contribute to the pathogenesis of staphylococcal pneumonia. Due to the fact that Protein A captures immunoglobulin and

displays toxic attributes, the possibility that this surface molecule may function as a vaccine in humans has not been rigorously pursued. Here the inventors demonstrate that Protein A variants no longer able to bind to immunoglobulins, which are thereby removed of their toxigenic potential, i.e., are non-toxicogenic, stimulate humoral immune responses that protect against staphylococcal disease.

**[0017]** In certain embodiments the SpA variant is a full length SpA variant comprising a variant A, B, C, D, and/or E domain. In certain aspects, the SpA variant comprises or consists of the amino acid sequence that is 80, 90, 95, 98, 99, or 100% identical to the amino acid sequence of SEQ ID NO:34. In other embodiments the SpA variant comprises a segment of SpA. The SpA segment can comprise at least or at most 1, 2, 3, 4, 5 or more IgG binding domains. The IgG domains can be at least or at most 1, 2, 3, 4, 5 or more variant A, B, C, D, or E domains. In certain aspects the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant A domains. In a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant B domains. In still a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant C domains. In yet a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant D domains. In certain aspects the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant E domains. In a further aspect the SpA variant comprises a combination of A, B, C, D, and E domains in various combinations and permutations. The combinations can include all or part of a SpA signal peptide segment, a SpA region X segment, and/or a SpA sorting signal segment. In other aspects the SpA variant does not include a SpA signal peptide segment, a SpA region X segment, and/or a SpA sorting signal segment. In certain aspects a variant A domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:4. In another aspect a variant B domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:6. In still another aspect a variant C domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:5. In certain aspects a variant D domain comprises a substitution at position(s) 9, 10, 36, and/or 37 of SEQ ID NO:2. In a further aspect a variant E domain comprises a substitution at position(s) 6, 7, 33, and/or 34 of SEQ ID NO:3.

**[0018]** In certain aspects, an SpA domain D variant or its equivalent can comprise a mutation at position 9 and 36; 9 and 37; 9 and 10; 36 and 37; 10 and 36; 10 and 37; 9, 36, and 37; 10, 36, and 37, 9, 10 and 36; or 9, 10 and 37 of SEQ ID NO:2. In a further aspect, analogous mutations can be included in one or more of domains A, B, C, or E.

**[0019]** In further aspects, the amino acid glutamine (Q) at position 9 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an aspartic acid (D), a cysteine (C), a glutamic acid (E), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the glutamine at position 9 can be substituted with an arginine (R). In a further aspect, the glutamine at position 9 of SEQ ID NO:2, or its equivalent, can be substituted with a lysine or a glycine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

**[0020]** In another aspect, the amino acid glutamine (Q) at position 10 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an

asparagine (N), an aspartic acid (D), a cysteine (C), a glutamic acid (E), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the glutamine at position 10 can be substituted with an arginine (R). In a further aspect, the glutamine at position 10 of SEQ ID NO:2, or its equivalent, can be substituted with a lysine or a glycine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

**[0021]** In certain aspects, the aspartic acid (D) at position 36 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an arginine (R), a cysteine (C), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a glutamine (Q), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the aspartic acid at position 36 can be substituted with a glutamic acid (E). In certain aspects, an aspartic acid at position 36 of SEQ ID NO:2, or its equivalent, can be substituted with an alanine or a serine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

**[0022]** In another aspect, the aspartic acid (D) at position 37 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an arginine (R), a cysteine (C), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a glutamine (Q), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the aspartic acid at position 37 can be substituted with a glutamic acid (E). In certain aspects, an aspartic acid at position 37 of SEQ ID NO:2, or its equivalent, can be substituted with an alanine or a serine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

**[0023]** In a particular embodiment the amino at position 9 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 9 of SEQ ID NO:2 is replaced by a glycine. In a further aspect the amino acid at position 9 of SEQ ID NO:2 is replaced by a lysine.

**[0024]** In a particular embodiment the amino at position 10 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 10 of SEQ ID NO:2 is replaced by a glycine. In a further aspect the amino acid at position 10 of SEQ ID NO:2 is replaced by a lysine.

**[0025]** In a particular embodiment the amino at position 36 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 36 of SEQ ID NO:2 is replaced by a serine. In a further aspect the amino acid at position 36 of SEQ ID NO:2 is replaced by an alanine.

**[0026]** In a particular embodiment the amino at position 37 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a

valine (V). In certain aspects the amino acid at position 37 of SEQ ID NO:2 is replaced by a serine. In a further aspect the amino acid at position 37 of SEQ ID NO:2 is replaced by an alanine.

**[0027]** In certain aspects the SpA variant includes a substitution of (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a  $V_H3$  binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to  $V_H3$ . In still further aspects the amino acid sequence of a SpA variant comprises an amino acid sequence that is at least 50%, 60%, 70%, 80%, 90%, 95%, or 100% identical, including all values and ranges there between, to the amino acid sequence of SEQ ID NOs:2-6.

**[0028]** In a further aspect the SpA variant includes (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain D, or at a corresponding amino acid position in other IgG domains, that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a  $V_H3$  binding sub-domain of SpA domain D, or at a corresponding amino acid position in other IgG domains, that disrupts or decreases binding to  $V_H3$ . In certain aspects amino acid residue F5, Q9, Q10, S11, F13, Y14, L17, N28, I31, and/or K35 (SEQ ID NO:2, QQNNFNKDDQSSAFYEILNMPNLNEAQRNGFIQSLKDDPSQSTNVLGEAKKLNES) of the IgG Fc binding sub-domain of domain D are modified or substituted. In certain aspects amino acid residue Q26, G29, F30, S33, D36, D37, Q40, N43, and/or E47 (SEQ ID NO:2) of the  $V_H3$  binding sub-domain of domain D are modified or substituted such that binding to Fc or  $V_H3$  is attenuated. In further aspects corresponding modifications or substitutions can be engineered in corresponding positions of the domain A, B, C, and/or E. Corresponding positions are defined by alignment of the domain D amino acid sequence with one or more of the amino acid sequences from other IgG binding domains of SpA, for example see FIG. 2A. In certain aspects the amino acid substitution can be any of the other 20 amino acids. In a further aspect conservative amino acid substitutions can be specifically excluded from possible amino acid substitutions. In other aspects only non-conservative substitutions are included. In any event, any substitution or combination of substitutions that reduces the binding of the domain such that SpA toxicity is significantly reduced is contemplated. The significance of the reduction in binding refers to a variant that produces minimal to no toxicity when introduced into a subject and can be assessed using in vitro methods described herein.

**[0029]** In certain embodiments, a variant SpA comprises at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more variant SpA domain D peptides. In certain aspects 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 or more amino acid residues of the variant SpA are substituted or modified—including but not limited to amino acids F5, Q9, Q10, S11, F13, Y14, L17, N28, I31, and/or K35 (SEQ ID NO:2) of the IgG Fc binding sub-domain of domain D and amino acid residue Q26, G29, F30, S33, D36, D37, Q40, N43, and/or E47 (SEQ ID NO:2) of the  $V_H3$  binding sub-domain of domain D. In one aspect of the invention glutamine residues at position 9 and/or 10 of SEQ ID NO:2 (or corresponding positions in other domains) are mutated. In another aspect, aspartic acid residues 36 and/or 37 of SEQ ID NO:2 (or corresponding positions in other domains) are mutated. In a further aspect, glutamine 9 and 10, and aspartic acid residues 36 and 37 are

mutated. Purified non-toxicogenic SpA or SpA-D mutants/variants described herein are no longer able to significantly bind (i.e., demonstrate attenuated or disrupted binding affinity) Fcγ or F(ab)<sub>2</sub>  $V_H3$  and also do not stimulate B cell apoptosis. These non-toxicogenic Protein A variants can be used as subunit vaccines and raise humoral immune responses and confer protective immunity against *S. aureus* challenge. Compared to wild-type full-length Protein A or the wild-type SpA-domain D, immunization with SpA-D variants resulted in an increase in Protein A specific antibody. Using a mouse model of staphylococcal challenge and abscess formation, it was observed that immunization with the non-toxicogenic Protein A variants generated significant protection from staphylococcal infection and abscess formation. As virtually all *S. aureus* strains express Protein A, immunization of humans with the non-toxicogenic Protein A variants can neutralize this virulence factor and thereby establish protective immunity. In certain aspects the protective immunity protects or ameliorates infection by drug resistant strains of *Staphylococcus*, such as USA300 and other MRSA strains.

**[0030]** Embodiments include the use of Protein A variants in methods and compositions for the treatment of bacterial and/or staphylococcal infection. This application also provides an immunogenic composition comprising a Protein A variant or immunogenic fragment thereof. In certain aspects, the immunogenic fragment is a Protein A domain D segment. Furthermore, the present invention provides methods and compositions that can be used to treat (e.g., limiting staphylococcal abscess formation and/or persistence in a subject) or prevent bacterial infection. In some cases, methods for stimulating an immune response involve administering to the subject an effective amount of a composition including or encoding all or part of a Protein A variant polypeptide or antigen, and in certain aspects other bacterial proteins. Other bacterial proteins include, but are not limited to (i) a secreted virulence factor, and/or a cell surface protein or peptide, or (ii) a recombinant nucleic acid molecule encoding a secreted virulence factor, and/or a cell surface protein or peptide.

**[0031]** In other aspects, the subject can be administered all or part of a Protein A variant, such as a variant Protein A domain D segment. The polypeptide of the invention can be formulated in a pharmaceutically acceptable composition. The composition can further comprise one or more of at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 additional staphylococcal antigen or immunogenic fragment thereof (e.g., Eap, Ehb, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla (e.g., H35 mutants), IsdC, SasF, vWbp, or vWh). Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa (GenBank CAC80837), Aap (GenBank accession AJ249487), Ant (GenBank accession NP\_372518), autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins









TABLE 1-continued

SpA and staphylococcal antigen combinations.										
Hla <sub>H35.4</sub>						+	+	+	+	+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
ClfA	+	+	+	+	+	+	+	+	+	+
ClfB		+	+	+	+	+	+	+	+	+
Coa			+	+	+	+	+	+	+	+
Hla				+	+	+	+	+	+	+
Hla <sub>H35.4</sub>						+	+	+	+	+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
ClfB		+	+	+	+	+	+	+	+	+
Coa			+	+	+	+	+	+	+	+
Hla				+	+	+	+	+	+	+
Hla <sub>H35.4</sub>						+	+	+	+	+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
Coa		+	+	+	+	+	+	+	+	+
Hla			+	+	+	+	+	+	+	+
Hla <sub>H35.4</sub>				+	+	+	+	+	+	+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
Coa		+	+	+	+	+	+	+	+	+
Hla			+	+	+	+	+	+	+	+
Hla <sub>H35.4</sub>				+	+	+	+	+	+	+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
Hla		+	+	+	+	+	+	+	+	+
Hla <sub>H35.4</sub>				+	+	+	+	+	+	+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
Hla		+	+	+	+	+	+	+	+	+
Hla <sub>H35.4</sub>				+	+	+	+	+	+	+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
Hla		+	+	+	+	+	+	+	+	+
Hla <sub>H35.4</sub>				+	+	+	+	+	+	+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
IsdC							+	+	+	+
SasF								+	+	+
vWbp									+	+
vWh										+
SasF								+	+	+
vWbp									+	+
vWh										+
vWbp									+	+
vWh										+
vWh										+

**[0034]** In still further aspects, the isolated Protein A variant is multimerized, e.g., dimerized or a linear fusion of two or more polypeptides or peptide segments. In certain aspects of the invention, a composition comprises multimers or concatamers of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more isolated cell surface proteins or segments thereof. Concatamers are linear polypeptides having one or more repeating peptide units. SpA polypeptides or fragments can be consecutive or separated by a spacer or other peptide sequences, e.g., one or more additional bacterial peptide. In a further aspect, the other polypeptides or peptides contained in the multimer or concatamer can include, but are not limited to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 of Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh or immunogenic fragments thereof. Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin

binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg2+ transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein.

**[0035]** The term "Protein A variant" or "SpA variant" refers to polypeptides that include a SpA IgG domain having two or more amino acid substitutions that disrupt binding to Fc and

$V_{H3}$ . In certain aspect, a SpA variant includes a variant domain D peptide, as well as variants of SpA polypeptides and segments thereof that are non-toxicogenic and stimulate an immune response against *staphylococcus* bacteria Protein A and/or bacteria expressing such.

**[0036]** Embodiments of the present invention include methods for eliciting an immune response against a *staphylococcus* bacterium or staphylococci in a subject comprising providing to the subject an effective amount of a Protein A variant or a segment thereof. In certain aspects, the methods for eliciting an immune response against a *staphylococcus* bacterium or staphylococci in a subject comprising providing to the subject an effective amount of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or more secreted proteins and/or cell surface proteins or segments/fragments thereof. A secreted protein or cell surface protein includes, but is not limited to Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, and/or vWh proteins and immunogenic fragments thereof. Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg2+ transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein.

**[0037]** Embodiments of the invention include compositions that include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to Protein A, or a second protein or peptide that is a secreted bacterial protein or a bacterial cell surface protein. In a further embodiment of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Protein A domain D polypeptide (SEQ ID NO:2), domain E (SEQ ID NO:3), domain A (SEQ ID NO:4), domain C (SEQ ID NO:5), domain B (SEQ ID NO:6), or a nucleic acid sequence encoding a Protein A domain D, domain E, domain A, domain C, or domain B polypeptide. In certain aspects a Protein A polypeptide segment will have an amino acid sequence of SEQ ID NO:8. Similarity or identity, with identity being preferred, is known in the art and a number of different programs can be used to identify whether a protein (or nucleic acid) has sequence identity or similarity to a known sequence. Sequence identity and/or similarity is determined using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith & Waterman (1981), by the sequence identity alignment algorithm of Needleman & Wunsch (1970), by the search for similarity method of Pearson & Lipman (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software

Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux et al. (1984), preferably using the default settings, or by inspection. Preferably, percent identity is calculated by using alignment tools known to and readily ascertainable to those of skill in the art. Percent identity is essentially the number of identical amino acids divided by the total number of amino acids compared times one hundred.

**[0038]** Still further embodiments include methods for stimulating in a subject a protective or therapeutic immune response against a *staphylococcus* bacterium comprising administering to the subject an effective amount of a composition including (i) a SpA variant, e.g., a variant SpA domain D polypeptide or peptide thereof; or, (ii) a nucleic acid molecule encoding such a SpA variant polypeptide or peptide thereof, or (iii) administering a SpA variant domain D polypeptide with any combination or permutation of bacterial proteins described herein. In a preferred embodiment the composition is not a *staphylococcus* bacterium. In certain aspects the subject is a human or a cow. In a further aspect the composition is formulated in a pharmaceutically acceptable formulation. The staphylococci may be *Staphylococcus aureus*.

**[0039]** Yet still further embodiments include vaccines comprising a pharmaceutically acceptable composition having an isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein, wherein the composition is capable of stimulating an immune response against a *staphylococcus* bacterium. The vaccine may comprise an isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described. In certain aspects of the invention the isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described are multimerized, e.g., dimerized or concatamerized. In a further aspect, the vaccine composition is contaminated by less than about 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, 0.5, 0.25, 0.05% (or any range derivable therein) of other Staphylococcal proteins. A composition may further comprise an isolated non-SpA polypeptide. Typically the vaccine comprises an adjuvant. In certain aspects a protein or peptide of the invention is linked (covalently or non-covalently) to the adjuvant, preferably the adjuvant is chemically conjugated to the protein.

**[0040]** In still yet further embodiments, a vaccine composition is a pharmaceutically acceptable composition having a recombinant nucleic acid encoding all or part of a SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein, wherein the composition is capable of stimulating an immune response against a *staphylococcus* bacteria. The vaccine composition may comprise a recombinant nucleic acid encoding all or part of a SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein. In certain embodiments the recombinant nucleic acid contains a heterologous promoter. Preferably the recombinant nucleic acid is a vector. More preferably the vector is a plasmid or a viral vector. In some aspects the vaccine includes a recombinant, non-staphylococcus bacterium containing the nucleic acid. The recombinant non-staphylococci may be *Salmonella* or another gram-positive bacteria. The vaccine may comprise a pharmaceutically acceptable excipient, more preferably an adjuvant.

**[0041]** Still further embodiments include methods for stimulating in a subject a protective or therapeutic immune

response against a *staphylococcus* bacterium comprising administering to the subject an effective amount of a composition of a SpA variant polypeptide or segment/fragment thereof and further comprising one or more of a Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, or vWh protein or peptide thereof. In a preferred embodiment the composition comprises a non-staphylococcus bacterium. In a further aspect the composition is formulated in a pharmaceutically acceptable formulation. The staphylococci for which a subject is being treated may be *Staphylococcus aureus*. Methods of the invention also include SpA variant compositions that contain 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or more secreted virulence factors and/or cell surface proteins, such as Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, or vWh in various combinations. In certain aspects a vaccine formulation includes Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, and vWh. In certain aspects an antigen combination can include (1) a SpA variant and IsdA; (2) SpA variant and ClfB; (3) SpA variant and SdrD; (4) SpA variant and Hla or Hla variant; (5) SpA variant and ClfB, SdrD, and Hla or Hla variant; (6) SpA variant, IsdA, SdrD, and Hla or Hla variant; (7) SpA variant, IsdA, ClfB, and Hla or Hla variant; (8) SpA variant, IsdA, ClfB, and SdrD; (9) SpA variant, IsdA, ClfB, SdrD and Hla or Hla variant; (10) SpA variant, IsdA, ClfB, and SdrD; (11) SpA variant, IsdA, SdrD, and Hla or Hla variant; (12) SpA variant, IsdA, and Hla or Hla variant; (13) SpA variant, IsdA, ClfB, and Hla or Hla variant; (14) SpA variant, ClfB, and SdrD; (15) SpA variant, ClfB, and Hla or Hla variant; or (16) SpA variant, SdrD, and Hla or Hla variant.

**[0042]** In certain aspects, a bacterium delivering a composition of the invention will be limited or attenuated with respect to prolonged or persistent growth or abscess formation. In yet a further aspect, SpA variant(s) can be overexpressed in an attenuated bacterium to further enhance or supplement an immune response or vaccine formulation.

**[0043]** The term “EsxA protein” refers to a protein that includes isolated wild-type EsxA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsxA proteins.

**[0044]** The term “EsxB protein” refers to a protein that includes isolated wild-type EsxB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsxB proteins.

**[0045]** The term “SdrD protein” refers to a protein that includes isolated wild-type SdrD polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrD proteins.

**[0046]** The term “SdrE protein” refers to a protein that includes isolated wild-type SdrE polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrE proteins.

**[0047]** The term “IsdA protein” refers to a protein that includes isolated wild-type IsdA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdA proteins.

**[0048]** The term “IsdB protein” refers to a protein that includes isolated wild-type IsdB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdB proteins.

**[0049]** The term “Eap protein” refers to a protein that includes isolated wild-type Eap polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Eap proteins.

**[0050]** The term “Ebh protein” refers to a protein that includes isolated wild-type Ebh polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Ebh proteins.

**[0051]** The term “Emp protein” refers to a protein that includes isolated wild-type Emp polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Emp proteins.

**[0052]** The term “EsaB protein” refers to a protein that includes isolated wild-type EsaB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsaB proteins.

**[0053]** The term “EsaC protein” refers to a protein that includes isolated wild-type EsaC polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsaC proteins.

**[0054]** The term “SdrC protein” refers to a protein that includes isolated wild-type SdrC polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrC proteins.

**[0055]** The term “ClfA protein” refers to a protein that includes isolated wild-type ClfA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria ClfA proteins.

**[0056]** The term “ClfB protein” refers to a protein that includes isolated wild-type ClfB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria ClfB proteins.

**[0057]** The term “Coa protein” refers to a protein that includes isolated wild-type Coa polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Coa proteins.

**[0058]** The term “Hla protein” refers to a protein that includes isolated wild-type Hla polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Hla proteins.

**[0059]** The term “IsdC protein” refers to a protein that includes isolated wild-type IsdC polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdC proteins.

**[0060]** The term “SasF protein” refers to a protein that includes isolated wild-type SasF polypeptides from *staphy-*

*lococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SasF proteins.

**[0061]** The term “vWbp protein” refers to a protein that includes isolated wild-type vWbp (von Willebrand factor binding protein) polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria vWbp proteins.

**[0062]** The term “vWh protein” refers to a protein that includes isolated wild-type vWh (von Willebrand factor binding protein homolog) polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria vWh proteins.

**[0063]** An immune response refers to a humoral response, a cellular response, or both a humoral and cellular response in an organism. An immune response can be measured by assays that include, but are not limited to, assays measuring the presence or amount of antibodies that specifically recognize a protein or cell surface protein, assays measuring T-cell activation or proliferation, and/or assays that measure modulation in terms of activity or expression of one or more cytokines.

**[0064]** In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsxA protein. In certain aspects the EsxA protein will have all or part of the amino acid sequence of SEQ ID NO:11.

**[0065]** In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsxB protein. In certain aspects the EsxB protein will have all or part of the amino acid sequence of SEQ ID NO:12.

**[0066]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an SdrD protein. In certain aspects the SdrD protein will have all or part of the amino acid sequence of SEQ ID NO:13.

**[0067]** In further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an SdrE protein. In certain aspects the SdrE protein will have all or part of the amino acid sequence of SEQ ID NO:14.

**[0068]** In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdA protein. In certain aspects the IsdA protein will have all or part of the amino acid sequence of SEQ ID NO:15.

**[0069]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdB protein. In certain aspects the IsdB protein will have all or part of the amino acid sequence of SEQ ID NO:16.

**[0070]** Embodiments of the invention include compositions that include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%,

or 99% identical or similar to a EsaB protein. In certain aspects the EsaB protein will have all or part of the amino acid sequence of SEQ ID NO:17.

**[0071]** In a further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a ClfB protein. In certain aspects the ClfB protein will have all or part of the amino acid sequence of SEQ ID NO:18.

**[0072]** In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdC protein. In certain aspects the IsdC protein will have all or part of the amino acid sequence of SEQ ID NO:19.

**[0073]** In yet further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a SasF protein. In certain aspects the SasF protein will have all or part of the amino acid sequence of SEQ ID NO:20.

**[0074]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a SdrC protein. In certain aspects the SdrC protein will have all or part of the amino acid sequence of SEQ ID NO:21.

**[0075]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a ClfA protein. In certain aspects the ClfA protein will have all or part of the amino acid sequence of SEQ ID NO:22.

**[0076]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Eap protein. In certain aspects the Eap protein will have all or part of the amino acid sequence of SEQ ID NO:23.

**[0077]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Ebh protein. In certain aspects the Ebh protein will have all or part of the amino acid sequence of SEQ ID NO:24.

**[0078]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Emp protein. In certain aspects the Emp protein will have all or part of the amino acid sequence of SEQ ID NO:25.

**[0079]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsaC protein. In certain aspects the EsaC protein will have all or part of the amino acid sequence of SEQ ID NO:26. Sequence of EsaC polypeptides can be found in the protein databases and include, but are not limited to accession numbers ZP\_02760162 (GI:168727885), NP\_645081.1 (GI:21281993), and NP\_370813.1 (GI:15923279), each of which is incorporated herein by reference as of the priority date of this application.

**[0080]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Coa protein. In certain aspects the Coa protein will have all or part of the amino acid sequence of SEQ ID NO:27.

**[0081]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Hla protein. In certain aspects the Hla protein will have all or part of the amino acid sequence of SEQ ID NO:28.

**[0082]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a vWa protein. In certain aspects the vWa protein will have all or part of the amino acid sequence of SEQ ID NO:29.

**[0083]** In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a vWbp protein. In certain aspects the vWbp protein will have all or part of the amino acid sequence of SEQ ID NO:32.

**[0084]** In certain aspects, a polypeptide or segment/fragment can have a sequence that is at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% or more identical to the amino acid sequence of the reference polypeptide. The term "similarity" refers to a polypeptide that has a sequence that has a certain percentage of amino acids that are either identical with the reference polypeptide or constitute conservative substitutions with the reference polypeptides.

**[0085]** The polypeptides described herein may include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more variant amino acids within at least, or at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:2-30, or SEQ ID NO:32-34.

**[0086]** A polypeptide segment as described herein may include 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126,

127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:2-30, or SEQ ID NO:33-34.

**[0087]** The compositions may be formulated in a pharmaceutically acceptable composition. In certain aspects of the invention the *staphylococcus* bacterium is an *S. aureus* bacterium.

**[0088]** In further aspects, a composition may be administered more than one time to the subject, and may be administered 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20 or more times. The administration of the compositions include, but is not limited to oral, parenteral, subcutaneous, intramuscular, intravenous, or various combinations thereof, including inhalation or aspiration.

**[0089]** In still further embodiments, a composition comprises a recombinant nucleic acid molecule encoding a polypeptide described herein or segments/fragments thereof. Typically a recombinant nucleic acid molecule encoding a polypeptide described herein contains a heterologous promoter. In certain aspects, a recombinant nucleic acid molecule of the invention is a vector, in still other aspects the vector is a plasmid. In certain embodiments the vector is a viral vector. In certain aspects a composition includes a recombinant, non-*staphylococcus* bacterium containing or expressing a polypeptide described herein. In particular aspects the recombinant non-staphylococcus bacteria is *Salmonella* or another gram-positive bacteria. A composition is typically administered to mammals, such as human subjects, but administration to other animals that are capable of eliciting an immune response is contemplated. In further aspects the *staphylococcus* bacterium containing or expressing the polypeptide is *Staphylococcus aureus*. In further embodiments the immune response is a protective immune response.

**[0090]** In further embodiments a composition comprises a recombinant nucleic acid molecule encoding all or part of one or more of a Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, SpA, vWbp, or vWh protein or peptide or variant thereof. Additional staphylococcal antigens that can be used in combination with the polypeptides described herein include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg2+ transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/



saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein. In particular aspects, a bacteria is a recombinant non-staphylococcus bacteria, such as a *Salmonella* or other gram-positive bacteria.

**[0091]** Compositions of the invention are typically administered to human subjects, but administration to other animals that are capable of eliciting an immune response to a *staphylococcus* bacterium is contemplated, particularly cattle, horses, goats, sheep and other domestic animals, i.e., mammals.

**[0092]** In certain aspects the *staphylococcus* bacterium is a *Staphylococcus aureus*. In further embodiments the immune response is a protective immune response. In still further aspects, the methods and compositions of the invention can be used to prevent, ameliorate, reduce, or treat infection of tissues or glands, e.g., mammary glands, particularly mastitis and other infections. Other methods include, but are not limited to prophylactically reducing bacterial burden in a subject not exhibiting signs of infection, particularly those subjects suspected of or at risk of being colonized by a target bacteria, e.g., patients that are or will be at risk or susceptible to infection during a hospital stay, treatment, and/or recovery.

**[0093]** Any embodiment discussed with respect to one aspect of the invention applies to other aspects of the invention as well. In particular, any embodiment discussed in the context of a SpA variant polypeptide or peptide or nucleic acid may be implemented with respect to other antigens, such as Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein (or nucleic acids), and vice versa. It is also understood that any one or more of Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein can be specifically excluded from a claimed composition.

**[0094]** Embodiments of the invention include compositions that contain or do not contain a bacterium. A composition may or may not include an attenuated or viable or intact staphylococcal bacterium. In certain aspects, the composition comprises a bacterium that is not a staphylococcal bacterium or does not contain staphylococcal bacteria. In certain embodiments a bacterial composition comprises an isolated or recombinantly expressed staphylococcal Protein A variant or a nucleotide encoding the same. The composition may be or include a recombinantly engineered *staphylococcus* bacterium that has been altered in a way that comprises specifically altering the bacterium with respect to a secreted virulence factor or cell surface protein. For example, the bacteria may be recombinantly modified to express more of the virulence factor or cell surface protein than it would express if unmodified.

**[0095]** The term “isolated” can refer to a nucleic acid or polypeptide that is substantially free of cellular material, bacterial material, viral material, or culture medium (when produced by recombinant DNA techniques) of their source of origin, or chemical precursors or other chemicals (when chemically synthesized). Moreover, an isolated compound refers to one that can be administered to a subject as an isolated compound; in other words, the compound may not simply be considered “isolated” if it is adhered to a column or embedded in an agarose gel. Moreover, an “isolated nucleic acid fragment” or “isolated peptide” is a nucleic acid or protein fragment that is not naturally occurring as a fragment and/or is not typically in the functional state.

**[0096]** Moieties of the invention, such as polypeptides, peptides, antigens, or immunogens, may be conjugated or linked covalently or noncovalently to other moieties such as adjuvants, proteins, peptides, supports, fluorescence moieties, or labels. The term “conjugate” or “immunoconjugate” is broadly used to define the operative association of one moiety with another agent and is not intended to refer solely to any type of operative association, and is particularly not limited to chemical “conjugation.” Recombinant fusion proteins are particularly contemplated. Compositions of the invention may further comprise an adjuvant or a pharmaceutically acceptable excipient. An adjuvant may be covalently or non-covalently coupled to a polypeptide or peptide of the invention. In certain aspects, the adjuvant is chemically conjugated to a protein, polypeptide, or peptide.

**[0097]** The term “providing” is used according to its ordinary meaning to indicate “to supply or furnish for use.” In some embodiments, the protein is provided directly by administering the protein, while in other embodiments, the protein is effectively provided by administering a nucleic acid that encodes the protein. In certain aspects the invention contemplates compositions comprising various combinations of nucleic acid, antigens, peptides, and/or epitopes.

**[0098]** The subject will have (e.g., are diagnosed with a staphylococcal infection), will be suspected of having, or will be at risk of developing a staphylococcal infection. Compositions of the present invention include immunogenic compositions wherein the antigen(s) or epitope(s) are contained in an amount effective to achieve the intended purpose. More specifically, an effective amount means an amount of active ingredients necessary to stimulate or elicit an immune response, or provide resistance to, amelioration of, or mitigation of infection. In more specific aspects, an effective amount prevents, alleviates or ameliorates symptoms of disease or infection, or prolongs the survival of the subject being treated.

Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any preparation used in the methods of the invention, an effective amount or dose can be estimated initially from in vitro studies, cell culture, and/or animal model assays. For example, a dose can be formulated in animal models to achieve a desired immune response or circulating antibody concentration or titer. Such information can be used to more accurately determine useful doses in humans.

[0099] The embodiments in the Example section are understood to be embodiments of the invention that are applicable to all aspects of the invention.

[0100] The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” It is also contemplated that anything listed using the term “or” may also be specifically excluded.

[0101] Throughout this application, the term “about” is used to indicate that a value includes the standard deviation of error for the device or method being employed to determine the value.

[0102] Following long-standing patent law, the words “a” and “an,” when used in conjunction with the word “comprising” in the claims or specification, denotes one or more, unless specifically noted.

[0103] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

#### DESCRIPTION OF THE DRAWINGS

[0104] So that the matter in which the above-recited features, advantages and objects of the invention as well as others which will become clear are attained and can be understood in detail, more particular descriptions and certain embodiments of the invention briefly summarized above are illustrated in the appended drawings. These drawings form a part of the specification. It is to be noted, however, that the appended drawings illustrate certain embodiments of the invention and therefore are not to be considered limiting in their scope.

[0105] FIGS. 1A-1B. (FIG. 1A) Primary structure of the Protein A precursor with an N-terminal YSIRK motif signal peptide, five immunoglobulin binding domains as tandem repeats designated E, D, A, B, C, region X, and the LPXTG sorting signal. (FIG. 1B) Following synthesis of the Protein A precursor, staphylococci secrete this product via the Sec pathway, and sortase A cleaves the LPXTG sorting signal between the T and G residues. Nucleophilic attack of the amino group within lipid II at the sortase-Protein A thioester-linked intermediate forms the amide bond that links Protein A to the cell wall envelope and enables its display on the bacterial surface.

[0106] FIG. 2. Three dimensional model of the molecular interactions between the SpA-domain D of Protein A, the VH3 Fab domain of the B cell receptor, and of the Fc $\gamma$  domain of immunoglobulin. The model is derived from two crystal structures (Graillie et al., 2000 and Gouda et al., 1992) that revealed side chain residues involved in the formation of ionic

bonds that enable these complexes. Gln-9 and Gln-10 of SpA-D promote binding to Fc $\gamma$ , whereas Asp-36 and Asp-37 enable complex formation with VH3 Fab.

[0107] FIG. 3. Left panel—Coomassie Blue stained SDS-PAGE reveals the migrational position of purified His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub>, human IgG, and sortase A (SrtA), a control protein. Right panel—Coomassie Blue stained SDS-PAGE to reveal the elution of Protein A immunoglobulin complexes eluted following affinity chromatography of human IgG on Ni-NTA columns pre-charged with His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or SrtA.

[0108] FIG. 4. ELISA assays to quantify human immunoglobulin (hIgG), human F(ab)<sub>2</sub> IgG fragments and human Fc fragments of immunoglobulin (hFc). Plates were coated with equal amounts of His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or SrtA. hIgG-HRP, F(ab)<sub>2</sub>-HRP and hFc-HRP were added onto the plates and incubated for an hour. Absorbance at 450 nm was recorded and plotted to determine the half maximal titers.

[0109] FIG. 5. Purified SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or a PBS mock control were injected into the peritoneum of mice and analyzed for their ability to reduce the B cell population in the spleen of experimental BALB/c mice. Animals were killed 4 hours following injection, their spleen removed, tissue homogenized and stained with CD 19 antibodies directed against B cells. The number of B cells was quantified by FACS sorting.

[0110] FIG. 6 Generation of a non-toxicogenic protein A vaccine. a, Translational protein A (SpA) product of *S. aureus* Newman and USA300 LAC with an N-terminal signal peptide (white box), five immunoglobulin binding domains (IgBDs designated E, D, A, B and C), variable region X and C-terminal sorting signal (black box). b, Amino acid sequence of the five IgBDs as well as nontoxicogenic SpA-D<sub>KKAA</sub> with the positions of triple  $\alpha$ -helical bundles (H1, H2 and H3) as well as glutamine (Q) 9, 10 and aspartate (D) 36, 37 indicated. c, Coomassie Blue-stained SDS-PAGE of SpA, SpA-D, SpA-D<sub>KKAA</sub> or SrtA purified on Ni-NTA sepharose in the presence or absence of human immunoglobulin (hIgG). d, ELISA examining the association of immobilized SpA, SpA-D or SpA-D<sub>KKAA</sub> with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments and von Willebrand factor (vWF). e, CD19+ B lymphocytes in splenic tissue of BALB/c mice that had been mock immunized or treated with SpA-D or SpA-D<sub>KKAA</sub> were quantified by FACS.

[0111] FIG. 7 Non-toxicogenic protein A vaccine prevents abscess formation. Histopathology of renal tissue isolated during necropsy of BALB/c mice that had been mock immunized (PBS) or vaccinated with SpA, SpA-D as well as SpA-D<sub>KKAA</sub> and challenged with *S. aureus* Newman. Thin sectioned tissues were stained with hematoxylin-eosin. White arrows identify polymorphonuclear leukocyte (PMN) infiltrates. Dark arrows identify staphylococcal abscess communities.

[0112] FIG. 8 Antibodies raised by the non-toxicogenic protein A vaccine block the B cell superantigen function of SpA. a, Rabbit antibodies raised against SpA-D<sub>KKAA</sub> were purified on a matrix with immobilized antigen and analyzed by Coomassie Blue-stained SDS-PAGE. Antibodies were cleaved with pepsin and F(ab)<sub>2</sub> fragments were purified by a second round of affinity chromatography on SpA-D<sub>KKAA</sub> matrix. b, SpA-D<sub>KKAA</sub> specific F(ab)<sub>2</sub> interfere with the binding of SpA or SpA-D to human immunoglobulin (hIgG) or, c, to von Willebrand Factor (vWF).

**[0113]** FIG. 9 Full-length non-toxicigenic protein A generates improved immune responses. a, Full-length SpA<sub>KKAA</sub> was purified on Ni-NTA sepharose and analyzed by Coomassie-Blue stained SDS-PAGE. b, CD 19+ B lymphocytes in splenic tissue of BALB/c mice that had been mock immunized or treated with SpA or SpA<sub>KKAA</sub> were quantified by FACS. c, ELISA examining the association of immobilized SpA or SpA<sub>KKAA</sub> with human IgG as well as its Fc or F(ab)2 fragments or von Willebrand factor (vWF). d, Human or mouse serum antibody titers to diphtheria toxoid (CRM197) and non-toxicigenic SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub>. Human volunteers with a history of DTaP immunization and staphylococcal infection (n=16) as well as mice (n=20) that had been infected with *S. aureus* Newman or USA 300 LAC or immunized with SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub> were examined by quantitative dot blot.

**[0114]** FIG. 10 Staphylococcal infection does not generate protective immunity. BALB/c mice (n=20) were infected with *S. aureus* Newman or mock challenged (PBS) for thirty days and infection cleared with chloramphenicol treatment. Both cohorts of animals were then challenged with *S. aureus* Newman and bacterial load (CFU) in kidney tissue homogenate analyzed following necropsy on day 4.

**[0115]** FIG. 11 Comparison of abscess formation in mice treated with PBS, SpA, SpA-D, and SpA-D<sub>KKAA</sub>.

**[0116]** FIGS. 12A-12C (A) ELISA examining the association of immobilized SpA, SpA-D, SpA-DKKAA or SpA-DGGSS with human IgG as well as its Fc or F(ab)2 fragments and IgM. Statistical significance of SpA-DKKAA and SpA-DGGSS binding to each ligand was compared against SpA-D; SpA-D binding was compared against SpA (n=3); \* signifies P<0.05; \*\* signifies P<0.01. (B) ELISA examining the level of cross-reactive antibodies of hyper-immune sera samples collected from actively immunized mice (n=5) with SpA-D, SpA-DKKAA and SpA-DGGSS. (C) Abscess formation in mice treated with PBS, SpA-D, SpA-D<sub>KKAA</sub> and SpA-D<sub>GGSS</sub>.

#### DETAILED DESCRIPTION

**[0117]** *Staphylococcus aureus* is a commensal of the human skin and nares, and the leading cause of bloodstream, skin and soft tissue infections (Klevens et al., 2007). Recent dramatic increases in the mortality of staphylococcal diseases are attributed to the spread of methicillin-resistant *S. aureus* (MRSA) strains often not susceptible to antibiotics (Kennedy et al., 2008). In a large retrospective study, the incidence of MRSA infections was 4.6% of all hospital admissions in the United States (Klevens et al., 2007). The annual health care costs for 94,300 MRSA infected individuals in the United States exceed \$2.4 billion (Klevens et al., 2007). The current MRSA epidemic has precipitated a public health crisis that needs to be addressed by development of a preventive vaccine (Boucher and Corey, 2008). To date, an FDA licensed vaccine that prevents *S. aureus* diseases is not available.

**[0118]** The inventors describe here the use of Protein A, a cell wall anchored surface protein of staphylococci, for the generation of variants that can serve as subunit vaccines. The pathogenesis of staphylococcal infections is initiated as bacteria invade the skin or blood stream via trauma, surgical wounds, or medical devices (Lowy, 1998). Although the invading pathogen may be phagocytosed and killed, staphylococci can also escape innate immune defenses and seed infections in organ tissues, inducing inflammatory responses that attract macrophages, neutrophils, and other phagocytes

(Lowy, 1998). The responsive invasion of immune cells to the site of infection is accompanied by liquefaction necrosis as the host seeks to prevent staphylococcal spread and allow for removal of necrotic tissue debris (Lam et al., 1963). Such lesions can be observed by microscopy as hypercellular areas containing necrotic tissue, leukocytes, and a central nidus of bacteria (Lam et al., 1963). Unless staphylococcal abscesses are surgically drained and treated with antibiotics, disseminated infection and septicemia produce a lethal outcome (Sheagren, 1984).

#### II. STAPHYLOCOCCAL ANTIGENS

**[0119]** A. Staphylococcal Protein A (SpA)

**[0120]** All *Staphylococcus aureus* strains express the structural gene for Protein A (spa) (Jensen, 1958; Said-Salim et al., 2003), a well characterized virulence factor whose cell wall anchored surface protein product (SpA) encompasses five highly homologous immunoglobulin binding domains designated E, D, A, B, and C (Sjodahl, 1977). These domains display ~80% identity at the amino acid level, are 56 to 61 residues in length, and are organized as tandem repeats (Uhlen et al., 1984). SpA is synthesized as a precursor protein with an N-terminal YSIRK/GS signal peptide and a C-terminal LPXTG motif sorting signal (DeDent et al., 2008; Schneewind et al., 1992). Cell wall anchored Protein A is displayed in great abundance on the staphylococcal surface (DeDent et al., 2007; Sjoquist et al., 1972). Each of its immunoglobulin binding domains is composed of anti-parallel  $\alpha$ -helices that assemble into a three helix bundle and bind the Fc domain of immunoglobulin G (IgG) (Deisenhofer, 1981; Deisenhofer et al., 1978), the VH3 heavy chain (Fab) of IgM (i.e., the B cell receptor) (Graille et al., 2000), the von Willebrand factor at its A1 domain [vWF A1 is a ligand for platelets] (O'Seaghda et al., 2006) and the tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ) receptor I (TNFR1) (Gomez et al., 2006), which is displayed on surfaces of airway epithelia (Gomez et al., 2004; Gomez et al., 2007).

**[0121]** SpA impedes neutrophil phagocytosis of staphylococci through its attribute of binding the Fc component of IgG (Jensen, 1958; Uhlen et al., 1984). Moreover, SpA is able to activate intravascular clotting via its binding to von Willebrand factor A1 domains (Hartleib et al., 2000). Plasma proteins such as fibrinogen and fibronectin act as bridges between staphylococci (C1fA and C1fB) and the platelet integrin GPIIb/IIIa (O'Brien et al., 2002), an activity that is supplemented through Protein A association with vWF A1, which allows staphylococci to capture platelets via the GPIIb- $\alpha$  platelet receptor (Foster, 2005; O'Seaghda et al., 2006). SpA also binds TNFR1 and this interaction contributes to the pathogenesis of staphylococcal pneumonia (Gomez et al., 2004). SpA activates proinflammatory signaling through TNFR1 mediated activation of TRAF2, the p38/c-Jun kinase, mitogen activate protein kinase (MAPK) and the Rel-transcription factor NF-KB. SpA binding further induces TNFR1 shedding, an activity that appears to require the TNF-converting enzyme (TACE) (Gomez et al., 2007). All of the aforementioned SpA activities are mediated through its five IgG binding domains and can be perturbed by the same amino acid substitutions, initially defined by their requirement for the interaction between Protein A and human IgG1 (Cedergren et al., 1993).

**[0122]** SpA also functions as a B cell superantigen by capturing the Fab region of VH3 bearing IgM, the B cell receptor (Gomez et al., 2007; Goodyear et al., 2003; Goodyear and

Silverman, 2004; Roben et al., 1995). Following intravenous challenge, staphylococcal Protein A (SpA) mutations show a reduction in staphylococcal load in organ tissues and dramatically diminished ability to form abscesses (described herein). During infection with wildtype *S. aureus*, abscesses are formed within forty-eight hours and are detectable by light microscopy of hematoxylin-eosin stained, thin-sectioned kidney tissue, initially marked by an influx of polymorphonuclear leukocytes (PMNs). On day 5 of infection, abscesses increase in size and enclosed a central population of staphylococci, surrounded by a layer of eosinophilic, amorphous material and a large cuff of PMNs. Histopathology revealed massive necrosis of PMNs in proximity to the staphylococcal nidus at the center of abscess lesions as well as a mantle of healthy phagocytes. The inventors also observed a rim of necrotic PMNs at the periphery of abscess lesions, bordering the eosinophilic pseudocapsule that separated healthy renal tissue from the infectious lesion. Staphylococcal variants lacking Protein A are unable to establish the histopathology features of abscesses and are cleared during infection.

**[0123]** In previous studies, Cedergren et al. (1993) engineered five individual substitutions in the Fc fragment binding sub-domain of the B domain of SpA, L17D, N28A, I31A and K35A. These authors created these proteins to test data gathered from a three dimensional structure of a complex between one domain of SpA and Fc<sub>1</sub>. Cedergren et al. determined the effects of these mutations on stability and binding, but did not contemplate use of such substitutions for the production of a vaccine antigen.

**[0124]** Brown et al. (1998) describe studies designed to engineer new proteins based on SpA that allow the use of more favorable elution conditions when used as affinity ligands. The mutations studied included single mutations of Q13A, Q14H, N15A, N15H, F17H, Y18F, L21H, N32H, or K39H. Brown et al. report that Q13A, N15A, N15H, and N32H substitutions made little difference to the dissociation constant values and that the Y18F substitution resulted in a 2 fold decrease in binding affinity as compared to wild type SpA. Brown et al. also report that L21H and F17H substitutions decrease the binding affinity by five-fold and a hundred-fold respectively. The authors also studied analogous substitutions in two tandem domains. Thus, the Brown et al. studies were directed to generating a SpA with a more favorable elution profile, hence the use of His substitutions to provide a pH sensitive alteration in the binding affinity. Brown et al. is silent on the use of SpA as a vaccine antigen.

**[0125]** Graille et al. (2000) describe a crystal structure of domain D of SpA and the Fab fragment of a human IgM antibody. Graille et al. define by analysis of a crystal structure the D domain amino acid residues that interact with the Fab fragment as residues Q26, G29, F30, Q32, S33, D36, D37, Q40, N43, E47, or L51, as well as the amino acid residues that form the interface between the domain D sub-domains. Graille et al. define the molecular interactions of these two proteins, but is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

**[0126]** O'Seaghda et al. (2006) describe studies directed at elucidating which sub-domain of domain D binds vWF. The authors generated single mutations in either the Fc or VH3 binding sub-domains, i.e., amino acid residues F5A, Q9A, Q10A, F13A, Y14A, L17A, N28A, I31A, K35A, G29A, F30A, S33A, D36A, D37A, Q40A, E47A, or Q32A. The authors discovered that vWF binds the same sub-domain that binds Fc. O'Seaghda et al. define the sub-domain of

domain D responsible for binding vWF, but is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

**[0127]** Gomez et al. (2006) describe the identification of residues responsible for activation of the TNFR1 by using single mutations of F5A, F13A, Y14A, L17A, N21A, I31A, Q32A, and K35A. Gomez et al. is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

**[0128]** Recombinant affinity tagged Protein A, a polypeptide encompassing the five IgG domains (EDCAB) (Sjodahl, 1977) but lacking the C-terminal Region X (Guss et al., 1984), was purified from recombinant *E. coli* and used as a vaccine antigen (Stranger-Jones et al., 2006). Because of the attributes of SpA in binding the Fc portion of IgG, a specific humoral immune response to Protein A could not be measured (Stranger-Jones et al., 2006). The inventors have overcome this obstacle through the generation of SpA-DQ9,10K; D36,37A. BALB/c mice immunized with recombinant Protein A (SpA) displayed significant protection against intravenous challenge with *S. aureus* strains: a 2.951 log reduction in staphylococcal load as compared to the wild-type ( $P > 0.005$ ; Student's t-test) (Stranger-Jones et al., 2006). SpA specific antibodies may cause phagocytic clearance prior to abscess formation and/or impact the formation of the aforementioned eosinophilic barrier in abscesses that separate staphylococcal communities from immune cells since these do not form during infection with Protein A mutant strains. Each of the five SpA domains (i.e., domains formed from three helix bundles designated E, D, A, B, and C) exerts similar binding properties (Jansson et al., 1998). The solution and crystal structure of the domain D has been solved both with and without the Fc and VH3 (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille et al., 2000). Mutations in residues known to be involved in IgG binding (FS, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF AI and TNFR1 binding (Cedergren et al., 1993; Gomez et al., 2006; O'Seaghda et al., 2006), whereas residues important for the VH3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) appear to have no impact on the other binding activities (Graille et al., 2000; Jansson et al., 1998). SpA specifically targets a subset of B cells that express VH3 family related IgM on their surface, i.e., VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells proliferate and commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e., marginal zone B cells and follicular B2 cells) (Goodyear et al., 2003; Goodyear et al., 2004).

**[0129]** Molecular Basis of Protein a Surface Display and Function.

**[0130]** Protein A is synthesized as a precursor in the bacterial cytoplasm and secreted via its YSIRK signal peptide at the cross wall, i.e. the cell division septum of staphylococci (FIG. 1) (DeDent et al., 2007; DeDent et al., 2008). Following cleavage of the C-terminal LPXTG sorting signal, Protein A is anchored to bacterial peptidoglycan crossbridges by sortase A (Mazmanian et al., 1999; Schneewind et al., 1995; Mazmanian et al., 2000). Protein A is the most abundant surface protein of staphylococci; the molecule is expressed by virtually all *S. aureus* strains (Cespedes et al., 2005; Kennedy et al., 2008; Said-Salim et al., 2003). Staphylococci turn over 15-20% of their cell wall per division cycle (Navarre and Schneewind, 1999). Murine hydrolases cleave the glycan

strands and wall peptides of peptidoglycan, thereby releasing Protein A with its attached C-terminal cell wall disaccharide tetrapeptide into the extracellular medium (Ton-That et al., 1999). Thus, by physiological design, Protein A is both anchored to the cell wall and displayed on the bacterial surface but also released into surrounding tissues during host infection (Marraffini et al., 2006).

**[0131]** Protein A captures immunoglobulins on the bacterial surface and this biochemical activity enables staphylococcal escape from host innate and acquired immune responses (Jensen, 1958; Goodyear et al., 2004). Interestingly, region X of Protein A (Guss et al., 1984), a repeat domain that tethers the IgG binding domains to the LPXTG sorting signal/cell wall anchor, is perhaps the most variable portion of the staphylococcal genome (Said-Salim, 2003; Schneewind et al., 1992). Each of the five immunoglobulin binding domains of Protein A (SpA), formed from three helix bundles and designated E, D, A, B, and C, exerts similar structural and functional properties (Sjodahl, 1977; Jansson et al., 1998). The solution and crystal structure of the domain D has been solved both with and without the Fc and V<sub>H3</sub> (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille 2000).

**[0132]** In the crystal structure complex, the Fab interacts with helix II and helix III of domain D via a surface composed of four VH region  $\beta$ -strands (Graille 2000). The major axis of helix II of domain D is approximately 50° to the orientation of the strands, and the interhelical portion of domain D is most proximal to the C0 strand. The site of interaction on Fab is remote from the Ig light chain and the heavy chain constant region. The interaction involves the following domain D residues: Asp-36 of helix II, Asp-37 and Gln-40 in the loop between helix II and helix III and several other residues (Graille 2000). Both interacting surfaces are composed predominantly of polar side chains, with three negatively charged residues on domain D and two positively charged residues on the 2A2 Fab buried by the interaction, providing an overall electrostatic attraction between the two molecules. Of the five polar interactions identified between Fab and domain D, three are between side chains. A salt bridge is formed between Arg-H19 and Asp-36 and two hydrogen bonds are made between Tyr-H59 and Asp-37 and between Asn-H82a and Ser-33. Because of the conservation of Asp-36 and Asp-37 in all five IgG binding domains of Protein A, the inventors mutated these residues.

**[0133]** The SpA-D sites responsible for Fab binding are structurally separate from the domain surface that mediates Fc $\gamma$  binding. The interaction of Fc $\gamma$  with domain D primarily involves residues in helix I with lesser involvement of helix II (Gouda et al., 1992; Deisenhofer, 1981). With the exception of the Gln-32, a minor contact in both complexes, none of the residues that mediate the Fc $\gamma$  interaction are involved in Fab binding. To examine the spatial relationship between these different Ig-binding sites, the SpA domains in these complexes have been superimposed to construct a model of a complex between Fab, the SpA-domain D, and the Fc $\gamma$  molecule. In this ternary model, Fab and Fc $\gamma$  form a sandwich about opposite faces of the helix II without evidence of steric hindrance of either interaction. These findings illustrate how, despite its small size (i.e., 56-61 aa), an SpA domain can simultaneously display both activities, explaining experimental evidence that the interactions of Fab with an individual domain are noncompetitive. Residues for the interaction between SpA-D and Fc $\gamma$  are Gln-9 and Gln-10.

**[0134]** In contrast, occupancy of the Fc portion of IgG on the domain D blocks its interaction with vWF A1 and probably also TNFR1 (O'Seaghda et al., 2006). Mutations in residues essential for IgG Fc binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF A1 and TNFR1 binding (O'Seaghda et al., 2006; Cedergren et al., 1993; Gomez et al., 2006), whereas residues critical for the VH3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) have no impact on the binding activities of IgG Fc, vWF A1 or TNFR1 (Jansson et al., 1998; Graille et al., 2000). The Protein A immunoglobulin Fab binding activity targets a subset of B cells that express V<sub>H3</sub> family related IgM on their surface, i.e., these molecules function as VH3type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells rapidly proliferate and then commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e., marginal zone B cells and follicular B2 cells) (Goodyear and Silverman, 2004; Goodyear and Silverman, 2003). More than 40% of circulating B cells are targeted by the Protein A interaction and the V<sub>H3</sub> family represents the largest family of human B cell receptors to impart protective humoral responses against pathogens (Goodyear and Silverman, 2004; Goodyear and Silverman, 2003). Thus, Protein A functions analogously to staphylococcal superantigens (Roben et al., 1995), albeit that the latter class of molecules, for example SEB, TSST-1, TSST-2, form complexes with the T cell receptor to inappropriately stimulate host immune responses and thereby precipitating characteristic disease features of staphylococcal infections (Roben et al., 1995; Tiedemann et al., 1995). Together these findings document the contributions of Protein A in establishing staphylococcal infections and in modulating host immune responses.

**[0135]** In sum, Protein A domains can be viewed as displaying two different interfaces for binding with host molecules and any development of Protein A based vaccines must consider the generation of variants that do not perturb host cell signaling, platelet aggregation, sequestration of immunoglobulins or the induction of B cell proliferation and apoptosis. Such Protein A variants should also be useful in analyzing vaccines for the ability of raising antibodies that block the aforementioned SpA activities and occupy the five repeat domains at their dual binding interfaces. This goal is articulated and pursued here for the first time and methods are described in detail for the generation of Protein A variants that can be used as a safe vaccine for humans. To perturb IgG Fc $\gamma$ , vWF A1 and TNFR1 binding, glutamine (Q) 9 and 10 [numbering derived from the SpA domain D as described in Uhlen et al., 1984] were mutated, and generated lysine substitutions for both glutamines with the expectation that these abolish the ligand attributes at the first binding interface. To perturb IgM Fab VH3 binding, aspartate (D) 36 and 37 were mutated, each of which is required for the association with the B cell receptor. D36 and D37 were both substituted with alanine. Q9, I10K and D36,37A mutations are here combined in the recombinant molecule SpA-DQ9, I10K; D36,37A and tested for the binding attributes of Protein A. Further, SpA-D and SpA-DQ9, I10K; D36,37A are subjected to immunization studies in mice and rabbits and analyzed for [1] the production of specific antibodies (SpA-D Ab); [2] the ability of SpA-D Ab to block the association between Protein A and its four different ligands; and, [3] the attributes of SpA-D Ab to generate protective immunity against staphylococcal infections. (See Examples section below).

**[0136]** B. Staphylococcal Coagulases

**[0137]** Coagulases are enzymes produced by *Staphylococcus* bacteria that convert fibrinogen to fibrin. Coa and vW<sub>h</sub> activate prothrombin without proteolysis (Friedrich et al., 2003). The coagulase-prothrombin complex recognizes fibrinogen as a specific substrate, converting it directly into fibrin. The crystal structure of the active complex revealed binding of the D1 and D2 domains to prothrombin and insertion of its Ile<sup>1</sup>-Val<sup>2</sup> N-terminus into the Ile<sup>16</sup> pocket, inducing a functional active site in the zymogen through conformational change (Friedrich et al., 2003). Exosite I of α-thrombin, the fibrinogen recognition site, and proexosite I on prothrombin are blocked by the D2 of Coa (Friedrich et al., 2003). Nevertheless, association of the tetrameric (Coa-prothrombin)<sub>2</sub> complex binds fibrinogen at a new site with high affinity (Panizzi et al., 2006). This model explains the coagulant properties and efficient fibrinogen conversion by coagulase (Panizzi et al., 2006).

**[0138]** Fibrinogen is a large glycoprotein (Mr ~340,000), formed by three pairs of Aα-, Bβ-, and γ-chains covalently linked to form a “dimer of trimers,” where A and B designate the fibrinopeptides released by thrombin cleavage (Panizzi et al., 2006). The elongated molecule folds into three separate domains, a central fragment E that contains the N-termini of all six chains and two flanking fragments D formed mainly by the C-termini of the Bβ- and γ-chains. These globular domains are connected by long triple-helical structures. Coagulase-prothrombin complexes, which convert human fibrinogen to the self-polymerizing fibrin, are not targeted by circulating thrombin inhibitors (Panizzi et al., 2006). Thus, staphylococcal coagulases bypass the physiological blood coagulation pathway.

**[0139]** All *S. aureus* strains secrete coagulase and vWbp (Bjerketorp et al., 2004; Field and Smith, 1945). Although early work reported important contributions of coagulase to the pathogenesis of staphylococcal infections (Ekstedt and Yotis, 1960; Smith et al., 1947), more recent investigations with molecular genetics tools challenged this view by observing no virulence phenotypes with endocarditis, skin abscess and mastitis models in mice (Moreillon et al., 1995; Phonimdaeng et al., 1990). Generating isogenic variants of *S. aureus* Newman, a fully virulent clinical isolate (Duthie et al., 1952), it is described herein that coa mutants indeed display virulence defects in a lethal bacteremia and renal abscess model in mice. In the inventors experience, *S. aureus* 8325-4 is not fully virulent and it is presumed that mutational lesions in this strain may not be able to reveal virulence defects in vivo. Moreover, antibodies raised against Coa or vWbp perturb the pathogenesis of *S. aureus* Newman infections to a degree mirroring the impact of gene deletions. Coa and vWbp contribute to staphylococcal abscess formation and lethal bacteremia and may also function as protective antigens in subunit vaccines.

**[0140]** Biochemical studies document the biological value of antibodies against Coa and vWbp. By binding to antigen and blocking its association with clotting factors, the antibodies prevent the formation of Coa-prothrombin and vWbp-prothrombin complexes. Passive transfer studies revealed protection of experimental animals against staphylococcal abscess formation and lethal challenge by Coa and vWbp antibodies. Thus, Coa and vWbp neutralizing antibodies generate immune protection against staphylococcal disease.

**[0141]** Earlier studies revealed a requirement of coagulase for resisting phagocytosis in blood (Smith et al., 1947) and the

inventors observed a similar phenotype for Δcoa mutants in lepirudin-treated mouse blood (see Example 3 below). As vWbp displays higher affinity for human prothrombin than the mouse counterpart, it is suspected the same may be true for ΔvWbp variants in human blood. Further, expression of Coa and vWbp in abscess lesions as well as their striking distribution in the eosinophilic pseudocapsule surrounding (staphylococcal abscess communities (SACs) or the peripheral fibrin wall, suggest that secreted coagulases contribute to the establishment of these lesions. This hypothesis was tested and, indeed, Δcoa mutants were defective in the establishment of abscesses. A corresponding test, blocking Coa function with specific antibodies, produced the same effect. Consequently, it is proposed that the clotting of fibrin is a critical event in the establishment of staphylococcal abscesses that can be targeted for the development of protective vaccines. Due to their overlapping function on human prothrombin, both Coa and vWbp are considered excellent candidates for vaccine development.

**[0142]** C. Other Staphylococcal Antigens

**[0143]** Research over the past several decades identified *S. aureus* exotoxins, surface proteins and regulatory molecules as important virulence factors (Foster, 2005; Mazmanian et al., 2001; Novick, 2003). Much progress has been achieved regarding the regulation of these genes. For example, staphylococci perform a bacterial census via the secretion of auto-inducing peptides that bind to a cognate receptor at threshold concentration, thereby activating phospho-relay reactions and transcriptional activation of many of the exotoxin genes (Novick, 2003). The pathogenesis of staphylococcal infections relies on these virulence factors (secreted exotoxins, exopolysaccharides, and surface adhesins). The development of staphylococcal vaccines is hindered by the multifaceted nature of staphylococcal invasion mechanisms. It is well established that live attenuated micro-organisms are highly effective vaccines; immune responses elicited by such vaccines are often of greater magnitude and of longer duration than those produced by non-replicating immunogens. One explanation for this may be that live attenuated strains establish limited infections in the host and mimic the early stages of natural infection. Embodiments of the invention are directed to compositions and methods including variant SpA polypeptides and peptides, as well as other immunogenic extracellular proteins, polypeptides, and peptides (including both secreted and cell surface proteins or peptides) of gram positive bacteria for the use in mitigating or immunizing against infection. In particular embodiments the bacteria is a *staphylococcus* bacteria. Extracellular proteins, polypeptides, or peptides include, but are not limited to secreted and cell surface proteins of the targeted bacteria.

**[0144]** The human pathogen *S. aureus* secretes EsxA and EsxB, two ESAT-6 like proteins, across the bacterial envelope (Burts et al., 2005, which is incorporated herein by reference). Staphylococcal esxA and esxB are clustered with six other genes in the order of transcription: esxA esaA essA esaB essB essC esaC esxB. The acronyms esa, ess, and esx stand for ESAT-6 secretion accessory, system, and extracellular, respectively, depending whether the encoded proteins play an accessory (esa) or direct (ess) role for secretion, or are secreted (esx) in the extracellular milieu. The entire cluster of eight genes is herein referred to as the Ess cluster. EsxA, esxB, esaA, esaB, and esaC are all required for synthesis or secretion of EsxA and EsxB. Mutants that fail to produce EsxA, EsxB, and EssC display defects in the pathogenesis of

*S. aureus* murine abscesses, suggesting that this specialized secretion system may be a general strategy of human bacterial pathogenesis. Secretion of non-WXG100 substrates by the ESX-1 pathway has been reported for several antigens including EspA, EspB, Rv3483c, and Rv3615c (Fortune et al., 2005; MacGurn et al., 2005; McLaughlin et al., 2007; Xu et al., 2007). The alternate ESX-5 pathway has also been shown to secrete both WXG100 and non-WXG100 proteins in pathogenic mycobacteria (Abdallah et al., 2007; Abdallah et al., 2006).

**[0145]** The *Staphylococcus aureus* Ess pathway can be viewed as a secretion module equipped with specialized transport components (Ess), accessory factors (Esa) and cognate secretion substrates (Esx). EssA, EssB and EssC are required for EsxA and EsxB secretion. Because EssA, EssB and EssC are predicted to be transmembrane proteins, it is contemplated that these proteins form a secretion apparatus. Some of the proteins in the ess gene cluster may actively transport secreted substrates (acting as motor) while others may regulate transport (regulator). Regulation may be achieved, but need not be limited to, transcriptional or post-translational mechanisms for secreted polypeptides, sorting of specific substrates to defined locations (e.g., extracellular medium or host cells), or timing of secretion events during infection. At this point, it is unclear whether all secreted Esx proteins function as toxins or contribute indirectly to pathogenesis.

**[0146]** Staphylococci rely on surface protein mediated-adhesion to host cells or invasion of tissues as a strategy for escape from immune defenses. Furthermore, *S. aureus* utilize surface proteins to sequester iron from the host during infection. The majority of surface proteins involved in staphylococcal pathogenesis carry C-terminal sorting signals, i.e., they are covalently linked to the cell wall envelope by sortase. Further, staphylococcal strains lacking the genes required for surface protein anchoring, i.e., sortase A and B, display a dramatic defect in the virulence in several different mouse models of disease. Thus, surface protein antigens represent a validated vaccine target as the corresponding genes are essential for the development of staphylococcal disease and can be exploited in various embodiments of the invention. The sortase enzyme superfamily are Gram-positive transpeptidases responsible for anchoring surface protein virulence factors to the peptidoglycan cell wall layer. Two sortase isoforms have been identified in *Staphylococcus aureus*, SrtA and SrtB. These enzymes have been shown to recognize a LPXTG motif in substrate proteins. The SrtB isoform appears to be important in heme iron acquisition and iron homeostasis, whereas the SrtA isoform plays a critical role in the pathogenesis of Gram-positive bacteria by modulating the ability of the bacterium to adhere to host tissue via the covalent anchoring of adhesins and other proteins to the cell wall peptidoglycan. In certain embodiments the SpA variants described herein can be used in combination with other staphylococcal proteins such as Coa, Eap, Ebh, Emp, EsaC, EsaB, EsxA, EsxB, Hla, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, IsdC, SasF, vWbp, and/or vWh proteins.

**[0147]** Certain aspects of the invention include methods and compositions concerning proteinaceous compositions including polypeptides, peptides, or nucleic acid encoding SpA variant(s) and other staphylococcal antigens such as other proteins transported by the Ess pathway, or sortase substrates. These proteins may be modified by deletion, insertion, and/or substitution.

**[0148]** The Esx polypeptides include the amino acid sequence of Esx proteins from bacteria in the *Staphylococcus* genus. The Esx sequence may be from a particular *staphylococcus* species, such as *Staphylococcus aureus*, and may be from a particular strain, such as Newman. In certain embodiments, the EsxA sequence is SAV0282 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number Q99WU4 (gil68565539), which is hereby incorporated by reference. In other embodiments, the EsxB sequence is SAV0290 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number Q99WT7 (gil68565532), which is hereby incorporated by reference. In further embodiments, other polypeptides transported by the Ess pathway may be used, the sequences of which may be identified by one of skill in the art using databases and internet accessible resources.

**[0149]** The sortase substrate polypeptides include, but are not limited to the amino acid sequence of SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, IsdC or SasF proteins from bacteria in the *Staphylococcus* genus. The sortase substrate polypeptide sequence may be from a particular *staphylococcus* species, such as *Staphylococcus aureus*, and may be from a particular strain, such as Newman. In certain embodiments, the SdrD sequence is from strain N315 and can be accessed using Genbank Accession Number NP\_373773.1 (gil15926240), which is incorporated by reference. In other embodiments, the SdrE sequence is from strain N315 and can be accessed using Genbank Accession Number NP\_373774.1 (gil15926241), which is incorporated by reference. In other embodiments, the IsdA sequence is SAV1130 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number NP\_371654.1 (gil15924120), which is incorporated by reference. In other embodiments, the IsdB sequence is SAV1129 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number NP\_371653.1 (gil15924119), which is incorporated by reference. In further embodiments, other polypeptides transported by the Ess pathway or processed by sortase may be used, the sequences of which may be identified by one of skill in the art using databases and internet accessible resources.

**[0150]** Examples of various proteins that can be used in the context of the present invention can be identified by analysis of database submissions of bacterial genomes, including but not limited to accession numbers NC\_002951 (GI:57650036 and GenBank CP000046), NC\_002758 (GI:57634611 and GenBank BA000017), NC\_002745 (GI:29165615 and GenBank BA000018), NC\_003923 (GI:21281729 and GenBank BA000033), NC\_002952 (GI:49482253 and GenBank BX571856), NC\_002953 (GI:49484912 and GenBank BX571857), NC\_007793 (GI:87125858 and GenBank CP000255), NC\_007795 (GI:87201381 and GenBank CP000253) each of which are incorporated by reference.

**[0151]** As used herein, a "protein" or "polypeptide" refers to a molecule comprising at least ten amino acid residues. In some embodiments, a wild-type version of a protein or polypeptide are employed, however, in many embodiments of the invention, a modified protein or polypeptide is employed to generate an immune response. The terms described above may be used interchangeably. A "modified protein" or "modified polypeptide" or a "variant" refers to a protein or polypeptide whose chemical structure, particularly its amino acid

sequence, is altered with respect to the wild-type protein or polypeptide. In some embodiments, a modified/variant protein or polypeptide has at least one modified activity or function (recognizing that proteins or polypeptides may have multiple activities or functions). It is specifically contemplated that a modified/variant protein or polypeptide may be altered with respect to one activity or function yet retain a wild-type activity or function in other respects, such as immunogenicity.

**[0152]** In certain embodiments the size of a protein or polypeptide (wild-type or modified) may comprise, but is not limited to, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 875, 900, 925, 950, 975, 1000, 1100, 1200, 1300, 1400, 1500, 1750, 2000, 2250, 2500 amino molecules or greater, and any range derivable therein, or derivative of a corresponding amino sequence described or referenced herein. It is contemplated that polypeptides may be mutated by truncation, rendering them shorter than their corresponding wild-type form, but also they might be altered by fusing or conjugating a heterologous protein sequence with a particular function (e.g., for targeting or localization, for enhanced immunogenicity, for purification purposes, etc.).

**[0153]** As used herein, an “amino molecule” refers to any amino acid, amino acid derivative, or amino acid mimic known in the art. In certain embodiments, the residues of the proteinaceous molecule are sequential, without any non-amino molecule interrupting the sequence of amino molecule residues. In other embodiments, the sequence may comprise one or more non-amino molecule moieties. In particular embodiments, the sequence of residues of the proteinaceous molecule may be interrupted by one or more non-amino molecule moieties.

**[0154]** Accordingly, the term “proteinaceous composition” encompasses amino molecule sequences comprising at least one of the 20 common amino acids in naturally synthesized proteins, or at least one modified or unusual amino acid.

**[0155]** Proteinaceous compositions may be made by any technique known to those of skill in the art, including (i) the expression of proteins, polypeptides, or peptides through standard molecular biological techniques, (ii) the isolation of proteinaceous compounds from natural sources, or (iii) the chemical synthesis of proteinaceous materials. The nucleotide as well as the protein, polypeptide, and peptide sequences for various genes have been previously disclosed, and may be found in the recognized computerized databases. One such database is the National Center for Biotechnology Information’s Genbank and GenPept databases (on the World Wide Web at [ncbi.nlm.nih.gov/](http://ncbi.nlm.nih.gov/)). The coding regions for these

genes may be amplified and/or expressed using the techniques disclosed herein or as would be known to those of ordinary skill in the art.

**[0156]** Amino acid sequence variants of SpA, coagulases and other polypeptides of the invention can be substitutional, insertional, or deletion variants. A variation in a polypeptide of the invention may affect 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, or more non-contiguous or contiguous amino acids of the polypeptide, as compared to wild-type. A variant can comprise an amino acid sequence that is at least 50%, 60%, 70%, 80%, or 90%, including all values and ranges there between, identical to any sequence provided or referenced herein, e.g., SEQ ID NO:2-8 or SEQ ID NO:11-30. A variant can include 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more substitute amino acids. A polypeptide processed or secreted by the Ess pathway or other surface proteins (see Table 1) or sortase substrates from any *staphylococcus* species and strain are contemplated for use in compositions and methods described herein.

**[0157]** Deletion variants typically lack one or more residues of the native or wild-type protein. Individual residues can be deleted or a number of contiguous amino acids can be deleted. A stop codon may be introduced (by substitution or insertion) into an encoding nucleic acid sequence to generate a truncated protein. Insertional mutants typically involve the addition of material at a non-terminal point in the polypeptide. This may include the insertion of one or more residues. Terminal additions, called fusion proteins, may also be generated. These fusion proteins include multimers or concatamers of one or more peptide or polypeptide described or referenced herein.

**[0158]** Substitutional variants typically contain the exchange of one amino acid for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide, with or without the loss of other functions or properties. Substitutions may be conservative, that is, one amino acid is replaced with one of similar shape and charge. Conservative substitutions are well known in the art and include, for example, the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; and valine to isoleucine or leucine. Alternatively, substitutions may be non-conservative such that a function or activity of the polypeptide is affected. Non-conservative changes typically involve substituting a residue with one that is chemically dissimilar, such as a polar or charged amino acid for a nonpolar or uncharged amino acid, and vice versa.

TABLE 2

Exemplary surface proteins of <i>S. aureus</i> strains.								
SAV #	SA#	Surface	MW2	Mu50	N315	Newman	MRSA252*	MSSA476*
SAV0111	SA0107	Spa	492	450	450	520	516	492
SAV2503	SA2291	FnBPA	1015	1038	1038	741	—	1015



TABLE 2-continued

Exemplary surface proteins of <i>S. aureus</i> strains.								
SAV #	SA#	Surface	MW2	Mu50	N315	Newman	MRSA252*	MSSA476*
SAV2502	SA2290	FnBPB	943	961	961	677	965	957
SAV0811	SA0742	CIfA	946	935	989	933	1029	928
SAV2630	SA2423	ClfB	907	877	877	913	873	905
Np	Np	Cna	1183	—	—	—	1183	1183
SAV0561	SA0519	SdrC	955	953	953	947	906	957
SAV0562	SA0520	SdrD	1347	1385	1385	1315	—	1365
SAV0563	SA0521	SdrE	1141	1141	1141	1166	1137	1141
Np	Np	Pls	—	—	—	—	—	—
SAV2654	SA2447	SasA	2275	2271	2271	2271	1351	2275
SAV2160	SA1964	SasB	686	2481	2481	2481	2222	685
	SA1577	SasC	2186	213	2186	2186	2189	2186
SAV0134	SA0129	SasD	241	241	241	241	221	241
SAV1130	SA0977	SasE/IsdA	350	350	350	350	354	350
SAV2646	SA2439	SasF	635	635	635	635	627	635
SAV2496		SasG	1371	525	927	—	—	1371
SAV0023	SA0022	SasH	772	—	772	772	786	786
SAV1731	SA1552	SasI	895	891	891	891	534	895
SAV1129	SA0976	SasJ/IsdB	645	645	645	645	652	645
	SA2381	SasK	198	211	211	—	—	197
	Np	SasL	—	232	—	—	—	—
SAV1131	SA0978	IsdC	227	227	227	227	227	227

[0159] Proteins of the invention may be recombinant, or synthesized in vitro. Alternatively, a non-recombinant or recombinant protein may be isolated from bacteria. It is also contemplated that a bacteria containing such a variant may be implemented in compositions and methods of the invention. Consequently, a protein need not be isolated.

[0160] The term “functionally equivalent codon” is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine, and also refers to codons that encode biologically equivalent amino acids (see Table 2, below).

TABLE 3

Codon Table	
Amino Acids	Codons
Alanine	Ala A GCA GCC GCG GCU
Cysteine	Cys C UGC UGU
Aspartic acid	Asp D GAC GAU
Glutamic acid	Glu E GAA GAG
Phenylalanine	Phe F UUC UUU
Glycine	Gly G GGA GGC GGG GGU
Histidine	His H CAC CAU
Isoleucine	Ile I AUA AUC AUU
Lysine	Lys K AAA AAG
Leucine	Leu L UUA UUG CUA CUC CUG CUU
Methionine	Met M AUG
Asparagine	Asn N AAC AAU
Proline	Pro P CCA CCC CCG CCU
Glutamine	Gln Q CAA CAG

TABLE 3-continued

Codon Table	
Amino Acids	Codons
Arginine	Arg R AGA AGG CGA CGC CGG CGU
Serine	Ser S AGC AGU UCA UCC UCG UCU
Threonine	Thr T ACA ACC ACG ACU
Valine	Val V GUA GUC GUG GUU
Tryptophan	Trp W UGG
Tyrosine	Tyr Y UAC UAU

[0161] It also will be understood that amino acid and nucleic acid sequences may include additional residues, such as additional N- or C-terminal amino acids, or 5' or 3' sequences, respectively, and yet still be essentially as set forth in one of the sequences disclosed herein, so long as the sequence meets the criteria set forth above, including the maintenance of biological protein activity (e.g., immunogenicity) where protein expression is concerned. The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region.

[0162] The following is a discussion based upon changing of the amino acids of a protein to create a variant polypeptide or peptide. For example, certain amino acids may be substituted for other amino acids in a protein structure with or without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's functional activity, certain amino acid substitutions can be made in a protein sequence, and in its underlying DNA coding sequence, and nevertheless produce a protein

with a desirable property. It is thus contemplated by the inventors that various changes may be made in the DNA sequences of genes.

**[0163]** It is contemplated that in compositions of the invention, there is between about 0.001 mg and about 10 mg of total polypeptide, peptide, and/or protein per ml. The concentration of protein in a composition can be about, at least about or at most about 0.001, 0.010, 0.050, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5, 10.0 mg/ml or more (or any range derivable therein). Of this, about, at least about, or at most about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100% may be an SpA variant or a coagulase, and may be used in combination with other peptides or polypeptides, such as other bacterial peptides and/or antigens.

**[0164]** The present invention contemplates the administration of variant SpA polypeptides or peptides to effect a preventative therapy or therapeutic effect against the development of a disease or condition associated with infection by a *staphylococcus* pathogen.

**[0165]** In certain aspects, combinations of staphylococcal antigens are used in the production of an immunogenic composition that is effective at treating or preventing staphylococcal infection. Staphylococcal infections progress through several different stages. For example, the staphylococcal life cycle involves commensal colonization, initiation of infection by accessing adjoining tissues or the bloodstream, and/or anaerobic multiplication in the blood. The interplay between *S. aureus* virulence determinants and the host defense mechanisms can induce complications such as endocarditis, metastatic abscess formation, and sepsis syndrome. Different molecules on the surface of the bacterium are involved in different steps of the infection cycle. Combinations of certain antigens can elicit an immune response which protects against multiple stages of staphylococcal infection. The effectiveness of the immune response can be measured either in animal model assays and/or using an opsonophagocytic assay.

**[0166]** D. Polypeptides and Polypeptide Production

**[0167]** The present invention describes polypeptides, peptides, and proteins and immunogenic fragments thereof for use in various embodiments of the present invention. For example, specific polypeptides are assayed for or used to elicit an immune response. In specific embodiments, all or part of the proteins of the invention can also be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, (1984); Tam et al., (1983); Merrifield, (1986); and Barany and Merrifield (1979), each incorporated herein by reference.

**[0168]** Alternatively, recombinant DNA technology may be employed wherein a nucleotide sequence which encodes a peptide of the invention is inserted into an expression vector, transformed or transfected into an appropriate host cell and cultivated under conditions suitable for expression.

**[0169]** One embodiment of the invention includes the use of gene transfer to cells, including microorganisms, for the production and/or presentation of polypeptides or peptides. The gene for the polypeptide or peptide of interest may be

transferred into appropriate host cells followed by culture of cells under the appropriate conditions. The generation of recombinant expression vectors, and the elements included therein, are well known in the art and briefly discussed herein. Alternatively, the protein to be produced may be an endogenous protein normally synthesized by the cell that is isolated and purified.

**[0170]** Another embodiment of the present invention uses autologous B lymphocyte cell lines, which are transfected with a viral vector that expresses an immunogen product, and more specifically, a protein having immunogenic activity. Other examples of mammalian host cell lines include, but are not limited to Vero and HeLa cells, other B- and T-cell lines, such as CEM, 721.221, H9, Jurkat, Raji, as well as cell lines of Chinese hamster ovary, W138, BHK, COS-7, 293, HepG2, 3T3, RIN and MDCK cells. In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or that modifies and processes the gene product in the manner desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed.

**[0171]** A number of selection systems may be used including, but not limited to HSV thymidine kinase, hypoxanthine-guanine phosphoribosyltransferase, and adenine phosphoribosyltransferase genes, in tk-, hgprt- or apt- cells, respectively. Also, anti-metabolite resistance can be used as the basis of selection: for dhfr, which confers resistance to trimethoprim and methotrexate; gpt, which confers resistance to mycophenolic acid; neo, which confers resistance to the aminoglycoside G418; and hyg, which confers resistance to hygromycin.

**[0172]** Animal cells can be propagated in vitro in two modes: as non-anchorage-dependent cells growing in suspension throughout the bulk of the culture or as anchorage-dependent cells requiring attachment to a solid substrate for their propagation (i.e., a monolayer type of cell growth).

**[0173]** Non-anchorage dependent or suspension cultures from continuous established cell lines are the most widely used means of large scale production of cells and cell products. However, suspension cultured cells have limitations, such as tumorigenic potential and lower protein production than adherent cells.

**[0174]** Where a protein is specifically mentioned herein, it is preferably a reference to a native or recombinant protein or optionally a protein in which any signal sequence has been removed. The protein may be isolated directly from the staphylococcal strain or produced by recombinant DNA techniques. Immunogenic fragments of the protein may be incorporated into the immunogenic composition of the invention. These are fragments comprising at least 10 amino acids, 20 amino acids, 30 amino acids, 40 amino acids, 50 amino acids, or 100 amino acids, including all values and ranges there between, taken contiguously from the amino acid sequence of the protein. In addition, such immunogenic fragments are immunologically reactive with antibodies generated against the Staphylococcal proteins or with antibodies generated by infection of a mammalian host with Staphylococci. Immunogenic fragments also include fragments that when administered at an effective dose, (either alone or as a hapten bound to

a carrier), elicit a protective or therapeutic immune response against Staphylococcal infection, in certain aspects it is protective against *S. aureus* and/or *S. epidermidis* infection. Such an immunogenic fragment may include, for example, the protein lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment according to the invention comprises substantially all of the extracellular domain of a protein which has at least 80% identity, at least 85% identity, at least 90% identity, at least 95% identity, or at least 97-99% identity, including all values and ranges there between, to a sequence selected segment of a polypeptide described or referenced herein.

**[0175]** Also included in immunogenic compositions of the invention are fusion proteins composed of one or more Staphylococcal proteins, or immunogenic fragments of staphylococcal proteins. Such fusion proteins may be made recombinantly and may comprise one portion of at least 1, 2, 3, 4, 5, or 6 staphylococcal proteins or segments. Alternatively, a fusion protein may comprise multiple portions of at least 1, 2, 3, 4 or 5 staphylococcal proteins. These may combine different Staphylococcal proteins and/or multiples of the same protein or protein fragment, or immunogenic fragments in the same protein (forming a multimer or a concatamer). Alternatively, the invention also includes individual fusion proteins of Staphylococcal proteins or immunogenic fragments thereof, as a fusion protein with heterologous sequences such as a provider of T-cell epitopes or purification tags, for example:  $\beta$ -galactosidase, glutathione-S-transferase, green fluorescent proteins (GFP), epitope tags such as FLAG, myc tag, poly histidine, or viral surface proteins such as influenza virus haemagglutinin, or bacterial proteins such as tetanus toxoid, diphtheria toxoid, or CRM197.

## II. NUCLEIC ACIDS

**[0176]** In certain embodiments, the present invention concerns recombinant polynucleotides encoding the proteins, polypeptides, peptides of the invention. The nucleic acid sequences for SpA, coagulases and other bacterial proteins are included, all of which are incorporated by reference, and can be used to prepare peptides or polypeptides.

**[0177]** As used in this application, the term "polynucleotide" refers to a nucleic acid molecule that either is recombinant or has been isolated free of total genomic nucleic acid. Included within the term "polynucleotide" are oligonucleotides (nucleic acids of 100 residues or less in length), recombinant vectors, including, for example, plasmids, cosmids, phage, viruses, and the like. Polynucleotides include, in certain aspects, regulatory sequences, isolated substantially away from their naturally occurring genes or protein encoding sequences. Polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be RNA, DNA (genomic, cDNA or synthetic), analogs thereof, or a combination thereof. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide.

**[0178]** In this respect, the term "gene," "polynucleotide," or "nucleic acid" is used to refer to a nucleic acid that encodes a protein, polypeptide, or peptide (including any sequences required for proper transcription, post-translational modification, or localization). As will be understood by those in the art, this term encompasses genomic sequences, expression cassettes, cDNA sequences, and smaller engineered nucleic acid segments that express, or may be adapted to express, proteins, polypeptides, domains, peptides, fusion proteins,

and mutants. A nucleic acid encoding all or part of a polypeptide may contain a contiguous nucleic acid sequence of: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 441, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, 1000, 1010, 1020, 1030, 1040, 1050, 1060, 1070, 1080, 1090, 1095, 1100, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 9000, 10000, or more nucleotides, nucleosides, or base pairs, including all values and ranges therebetween, of a polynucleotide encoding one or more amino acid sequence described or referenced herein. It also is contemplated that a particular polypeptide may be encoded by nucleic acids containing variations having slightly different nucleic acid sequences but, nonetheless, encode the same or substantially similar protein (see Table 3 above).

**[0179]** In particular embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a variant SpA or coagulase. The term "recombinant" may be used in conjunction with a polynucleotide or polypeptide and generally refers to a polypeptide or polynucleotide produced and/or manipulated in vitro or that is a replication product of such a molecule.

**[0180]** In other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a variant SpA or coagulase polypeptide or peptide to generate an immune response in a subject. In various embodiments the nucleic acids of the invention may be used in genetic vaccines.

**[0181]** The nucleic acid segments used in the present invention can be combined with other nucleic acid sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant nucleic acid protocol. In some cases, a nucleic acid sequence may encode a polypeptide sequence with additional heterologous coding sequences, for example to allow for purification of the polypeptide, transport, secretion, post-translational modification, or for therapeutic benefits such as targeting or efficacy. As discussed above, a tag or other heterologous polypeptide may be added to the modified polypeptide-encoding sequence, wherein "heterologous" refers to a polypeptide that is not the same as the modified polypeptide.

**[0182]** In certain other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors that include within their sequence a contiguous nucleic acid sequence from SEQ ID NO:1 (SpA domain D) or SEQ ID NO:3 (SpA) or any other nucleic acid sequences encoding coagulases or other secreted virulence factors and/or surface proteins including proteins transported by the Ess pathway, processed by sortase, or proteins incorporated herein by reference.

**[0183]** In certain embodiments, the present invention provides polynucleotide variants having substantial identity to

the sequences disclosed herein; those comprising at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher sequence identity, including all values and ranges there between, compared to a polynucleotide sequence of this invention using the methods described herein (e.g., BLAST analysis using standard parameters).

**[0184]** The invention also contemplates the use of polynucleotides which are complementary to all the above described polynucleotides.

**[0185]** A. Vectors

**[0186]** Polypeptides of the invention may be encoded by a nucleic acid molecule comprised in a vector. The term “vector” is used to refer to a carrier nucleic acid molecule into which a heterologous nucleic acid sequence can be inserted for introduction into a cell where it can be replicated and expressed. A nucleic acid sequence can be “heterologous,” which means that it is in a context foreign to the cell in which the vector is being introduced or to the nucleic acid in which is incorporated, which includes a sequence homologous to a sequence in the cell or nucleic acid but in a position within the host cell or nucleic acid where it is ordinarily not found. Vectors include DNAs, RNAs, plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (e.g., YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (for example Sambrook et al., 2001; Ausubel et al., 1996, both incorporated herein by reference). In addition to encoding a variant SpA polypeptide the vector can encode other polypeptide sequences such as a one or more other bacterial peptide, a tag, or an immunogenicity enhancing peptide. Useful vectors encoding such fusion proteins include pIN vectors (Inouye et al., 1985), vectors encoding a stretch of histidines, and pGEX vectors, for use in generating glutathione S-transferase (GST) soluble fusion proteins for later purification and separation or cleavage.

**[0187]** The term “expression vector” refers to a vector containing a nucleic acid sequence coding for at least part of a gene product capable of being transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. Expression vectors can contain a variety of “control sequences,” which refer to nucleic acid sequences necessary for the transcription and possibly translation of an operably linked coding sequence in a particular host organism. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described herein.

**[0188]** 1. Promoters and Enhancers

**[0189]** A “promoter” is a control sequence. The promoter is typically a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind such as RNA polymerase and other transcription factors. The phrases “operatively positioned,” “operatively linked,” “under control,” and “under transcriptional control” mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and expression of that sequence. A promoter may or may not be used in conjunction with an “enhancer,” which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

**[0190]** Naturally, it may be important to employ a promoter and/or enhancer that effectively directs the expression of the

DNA segment in the cell type or organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression (see Sambrook et al., 2001, incorporated herein by reference). The promoters employed may be constitutive, tissue-specific, or inducible and in certain embodiments may direct high level expression of the introduced DNA segment under specified conditions, such as large-scale production of recombinant proteins or peptides.

**[0191]** Various elements/promoters may be employed in the context of the present invention to regulate the expression of a gene. Examples of such inducible elements, which are regions of a nucleic acid sequence that can be activated in response to a specific stimulus, include but are not limited to Immunoglobulin Heavy Chain (Banerji et al., 1983; Gilles et al., 1983; Grosschedl et al., 1985; Atchinson et al., 1986, 1987; Imler et al., 1987; Weinberger et al., 1984; Kiledjian et al., 1988; Porton et al.; 1990), Immunoglobulin Light Chain (Queen et al., 1983; Picard et al., 1984), T Cell Receptor (Luria et al., 1987; Winoto et al., 1989; Redondo et al.; 1990), HLA DQ  $\alpha$  and/or DQ  $\beta$  (Sullivan et al., 1987),  $\beta$  Interferon (Goodbourn et al., 1986; Fujita et al., 1987; Goodbourn et al., 1988), Interleukin-2 (Greene et al., 1989), Interleukin-2 Receptor (Greene et al., 1989; Lin et al., 1990), MHC Class II 5 (Koch et al., 1989), MHC Class II HLA-DR $\alpha$  (Sherman et al., 1989),  $\beta$ -Actin (Kawamoto et al., 1988; Ng et al.; 1989), Muscle Creatine Kinase (MCK) (Jaynes et al., 1988; Horlick et al., 1989; Johnson et al., 1989), Prealbumin (Transthyretin) (Costa et al., 1988), Elastase I (Ornitz et al., 1987), Metallothionein (MTII) (Karin et al., 1987; Culotta et al., 1989), Collagenase (Pinkert et al., 1987; Angel et al., 1987), Albumin (Pinkert et al., 1987; Tronche et al., 1989, 1990),  $\alpha$ -Fetoprotein (Godbout et al., 1988; Campere et al., 1989),  $\gamma$ -Globin (Bodine et al., 1987; Perez-Stable et al., 1990),  $\beta$ -Globin (Trudel et al., 1987), c-fos (Cohen et al., 1987), c-Ha-Ras (Triesman, 1986; Deschamps et al., 1985), Insulin (Edlund et al., 1985), Neural Cell Adhesion Molecule (NCAM) (Hirsh et al., 1990),  $\alpha$ 1-Antitrypsin (Latimer et al., 1990), H2B (TH2B) Histone (Hwang et al., 1990), Mouse and/or Type I Collagen (Ripe et al., 1989), Glucose-Regulated Proteins (GRP94 and GRP78) (Chang et al., 1989), Rat Growth Hormone (Larsen et al., 1986), Human Serum Amyloid A (SAA) (Edbrooke et al., 1989), Troponin I (TN I) (Yutzey et al., 1989), Platelet-Derived Growth Factor (PDGF) (Pech et al., 1989), Duchenne Muscular Dystrophy (Klamut et al., 1990), SV40 (Banerji et al., 1981; Moreau et al., 1981; Sleigh et al., 1985; Firak et al., 1986; Herr et al., 1986; Imbra et al., 1986; Kadesch et al., 1986; Wang et al., 1986; Ondek et al., 1987; Kuhl et al., 1987; Schaffner et al., 1988), Polyoma (Swartzendruber et al., 1975; Vasseur et al., 1980; Katinka et al., 1980, 1981; Tyndell et al., 1981; Dandolo et al., 1983; de Villiers et al., 1984; Hen et al., 1986; Satake et al., 1988; Campbell et al., 1988), Retroviruses (Kriegler et al., 1982, 1983; Levinson et al., 1982; Kriegler et al., 1983, 1984a, b, 1988; Bosze et al., 1986; Miksicek et al., 1986; Celander et al., 1987; Thiesen et al., 1988; Celander et al., 1988; Choi et al., 1988; Reisman et al., 1989), Papilloma Virus (Campo et al., 1983; Lusky et al., 1983; Spandidos and Wilkie, 1983; Spalholz et al., 1985; Lusky et al., 1986; Cripe et al., 1987; Gloss et al., 1987; Hirochika et al., 1987; Stephens et al., 1987), Hepatitis B Virus (Bulla et al., 1986; Jameel et al., 1986; Shaul et al., 1987; Spandau et al., 1988; Vannice et al., 1988), Human Immunodeficiency Virus (Muesing et al., 1987; Hauber et al., 1988; Jakobovits et al., 1988; Feng et al.,

1988; Takebe et al., 1988; Rosen et al., 1988; Berkhout et al., 1989; Laspia et al., 1989; Sharp et al., 1989; Braddock et al., 1989), Cytomegalovirus (CMV) IE (Weber et al., 1984; Boshart et al., 1985; Foecking et al., 1986), Gibbon Ape Leukemia Virus (Holbrook et al., 1987; Quinn et al., 1989).

**[0192]** Inducible elements include, but are not limited to MT II—Phorbol Ester (TPA)/Heavy metals (Palmiter et al., 1982; Haslinger et al., 1985; Searle et al., 1985; Stuart et al., 1985; Imagawa et al., 1987; Karin et al., 1987; Angel et al., 1987b; McNeall et al., 1989); MMTV (mouse mammary tumor virus)—Glucocorticoids (Huang et al., 1981; Lee et al., 1981; Majors et al., 1983; Chandler et al., 1983; Lee et al., 1984; Ponta et al., 1985; Sakai et al., 1988);  $\beta$ -Interferon—poly(rI)x/poly(rc) (Tavernier et al., 1983); Adenovirus 5 E2-E1A (Imperiale et al., 1984); Collagenase—Phorbol Ester (TPA) (Angel et al., 1987a); Stromelysin—Phorbol Ester (TPA) (Angel et al., 1987b); SV40—Phorbol Ester (TPA) (Angel et al., 1987b); Murine MX Gene—Interferon, Newcastle Disease Virus (Hug et al., 1988); GRP78 Gene—A23187 (Resendez et al., 1988);  $\alpha$ -2-Macroglobulin—IL-6 (Kunz et al., 1989); Vimentin—Serum (Rittling et al., 1989); MHC Class I Gene H-2kb—Interferon (Blonar et al., 1989); HSP70—E1A/SV40 Large T Antigen (Taylor et al., 1989, 1990a, 1990b); Proliferin—Phorbol Ester/TPA (Mordacq et al., 1989); Tumor Necrosis Factor—PMA (Hensel et al., 1989); and Thyroid Stimulating Hormone a Gene—Thyroid Hormone (Chatterjee et al., 1989).

**[0193]** The particular promoter that is employed to control the expression of peptide or protein encoding polynucleotide of the invention is not believed to be critical, so long as it is capable of expressing the polynucleotide in a targeted cell, preferably a bacterial cell. Where a human cell is targeted, it is preferable to position the polynucleotide coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a bacterial, human or viral promoter.

**[0194]** In embodiments in which a vector is administered to a subject for expression of the protein, it is contemplated that a desirable promoter for use with the vector is one that is not down-regulated by cytokines or one that is strong enough that even if down-regulated, it produces an effective amount of a variant SpA for eliciting an immune response. Non-limiting examples of these are CMV IE and RSV LTR. Tissue specific promoters can be used, particularly if expression is in cells in which expression of an antigen is desirable, such as dendritic cells or macrophages. The mammalian MHC I and MHC II promoters are examples of such tissue-specific promoters.

**[0195]** 2. Initiation Signals and Internal Ribosome Binding Sites (IRES)

**[0196]** A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals.

**[0197]** In certain embodiments of the invention, the use of internal ribosome entry sites (IRES) elements are used to create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, 1988; Macejak and Sarnow, 1991). IRES elements can be linked to heterologous

open reading frames. Multiple open reading frames can be transcribed together, each separated by an IRES, creating polycistronic messages. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message (see U.S. Pat. Nos. 5,925,565 and 5,935,819, herein incorporated by reference).

**[0198]** 3. Selectable and Screenable Markers

**[0199]** In certain embodiments of the invention, cells containing a nucleic acid construct of the present invention may be identified *in vitro* or *in vivo* by encoding a screenable or selectable marker in the expression vector. When transcribed and translated, a marker confers an identifiable change to the cell permitting easy identification of cells containing the expression vector. Generally, a selectable marker is one that confers a property that allows for selection. A positive selectable marker is one in which the presence of the marker allows for its selection, while a negative selectable marker is one in which its presence prevents its selection. An example of a positive selectable marker is a drug resistance marker.

**[0200]** B. Host Cells

**[0201]** As used herein, the terms “cell,” “cell line,” and “cell culture” may be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. In the context of expressing a heterologous nucleic acid sequence, “host cell” refers to a prokaryotic or eukaryotic cell, and it includes any transformable organism that is capable of replicating a vector or expressing a heterologous gene encoded by a vector. A host cell can, and has been, used as a recipient for vectors or viruses. A host cell may be “transfected” or “transformed,” which refers to a process by which exogenous nucleic acid, such as a recombinant protein-encoding sequence, is transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny.

**[0202]** Host cells may be derived from prokaryotes or eukaryotes, including bacteria, yeast cells, insect cells, and mammalian cells for replication of the vector or expression of part or all of the nucleic acid sequence(s). Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection (ATCC), which is an organization that serves as an archive for living cultures and genetic materials ([www.atcc.org](http://www.atcc.org)).

**[0203]** C. Expression Systems

**[0204]** Numerous expression systems exist that comprise at least a part or all of the compositions discussed above. Prokaryote- and/or eukaryote-based systems can be employed for use with the present invention to produce nucleic acid sequences, or their cognate polypeptides, proteins and peptides. Many such systems are commercially and widely available.

**[0205]** The insect cell/baculovirus system can produce a high level of protein expression of a heterologous nucleic acid segment, such as described in U.S. Pat. Nos. 5,871,986, 4,879,236, both herein incorporated by reference, and which can be bought, for example, under the name MAXBAC® 2.0 from INVITROGEN® and BACPACK™ BACULOVIRUS EXPRESSION SYSTEM FROM CLONTECH®.

**[0206]** In addition to the disclosed expression systems of the invention, other examples of expression systems include STRATAGENE®'s COMPLETE CONTROL™ Inducible Mammalian Expression System, which involves a synthetic ecdysone-inducible receptor, or its pET Expression System,

an *E. coli* expression system. Another example of an inducible expression system is available from INVITROGEN®, which carries the T-REX™ (tetracycline-regulated expression) System, an inducible mammalian expression system that uses the full-length CMV promoter. INVITROGEN® also provides a yeast expression system called the *Pichia methanolica* Expression System, which is designed for high-level production of recombinant proteins in the methylotrophic yeast *Pichia methanolica*. One of skill in the art would know how to express a vector, such as an expression construct, to produce a nucleic acid sequence or its cognate polypeptide, protein, or peptide.

### III. POLYSACCHARIDES

[0207] The immunogenic compositions of the invention may further comprise capsular polysaccharides including one or more of PIA (also known as PNAG) and/or *S. aureus* Type V and/or type VIII capsular polysaccharide and/or *S. epidermidis* Type I, and/or Type II and/or Type III capsular polysaccharide.

[0208] A. PIA (PNAG)

[0209] It is now clear that the various forms of staphylococcal surface polysaccharides identified as PS/A, PIA and SAA are the same chemical entity—PNAG (Maira-Litran et al., 2004). Therefore the term PIA or PNAG encompasses all these polysaccharides or oligosaccharides derived from them.

[0210] PIA is a polysaccharide intercellular adhesin and is composed of a polymer of  $\beta$ -(1 $\rightarrow$ 6)-linked glucosamine substituted with N-acetyl and O-succinyl constituents. This polysaccharide is present in both *S. aureus* and *S. epidermidis* and can be isolated from either source (Joyce et al., 2003; Maira-Litran et al., 2002). For example, PNAG may be isolated from *S. aureus* strain MN8m (WO04/43407). PIA isolated from *S. epidermidis* is an integral constituent of biofilm. It is responsible for mediating cell-cell adhesion and probably also functions to shield the growing colony from the host's immune response. The polysaccharide previously known as poly-N-succinyl- $\beta$ -(1 $\rightarrow$ 6)-glucosamine (PNSG) was recently shown not to have the expected structure since the identification of N-succinylation was incorrect (Maira-Litran et al., 2002). Therefore the polysaccharide formally known as PNSG and now found to be PNAG is also encompassed by the term PIA.

[0211] PIA (or PNAG) may be of different sizes varying from over 400 kDa to between 75 and 400 kDa to between 10 and 75 kDa to oligosaccharides composed of up to 30 repeat units (of  $\beta$ -(1 $\rightarrow$ 6)-linked glucosamine substituted with N-acetyl and O-succinyl constituents). Any size of PIA polysaccharide or oligosaccharide may be used in an immunogenic composition of the invention, in one aspect the polysaccharide is over 40 kDa. Sizing may be achieved by any method known in the art, for instance by microfluidization, ultrasonic irradiation or by chemical cleavage (WO 03/53462, EP497524, EP497525). In certain aspects PIA (PNAG) is at least or at most 40-400 kDa, 40-300 kDa, 50-350 kDa, 60-300 kDa, 50-250 kDa and 60-200 kDa.

[0212] PIA (PNAG) can have different degree of acetylation due to substitution on the amino groups by acetate. PIA produced in vitro is almost fully substituted on amino groups (95-100%). Alternatively, a deacetylated PIA (PNAG) can be used having less than 60%, 50%, 40%, 30%, 20%, 10% acetylation. Use of a deacetylated PIA (PNAG) is preferred since non-acetylated epitopes of PNAG are efficient at mediating opsonic killing of Gram positive bacteria, preferably *S.*

*aureus* and/or *S. epidermidis*. In certain aspects, the PIA (PNAG) has a size between 40 kDa and 300 kDa and is deacetylated so that less than 60%, 50%, 40%, 30% or 20% of amino groups are acetylated.

[0213] The term deacetylated PNAG (dPNAG) refers to a PNAG polysaccharide or oligosaccharide in which less than 60%, 50%, 40%, 30%, 20% or 10% of the amino groups are acetylated. In certain aspects, PNAG is deacetylated to form dPNAG by chemically treating the native polysaccharide. For example, the native PNAG is treated with a basic solution such that the pH rises to above 10. For instance the PNAG is treated with 0.1-5 M, 0.2-4 M, 0.3-3 M, 0.5-2 M, 0.75-1.5 M or 1 M NaOH, KOH or NH<sub>4</sub>OH. Treatment is for at least 10 to 30 minutes, or 1, 2, 3, 4, 5, 10, 15 or 20 hours at a temperature of 20-100, 25-80, 30-60 or 30-50 or 35-45° C. dPNAG may be prepared as described in WO 04/43405.

[0214] The polysaccharide(s) can be conjugated or unconjugated to a carrier protein.

[0215] B. Type 5 and Type 8 Polysaccharides from *S. aureus*

[0216] Most strains of *S. aureus* that cause infection in man contain either Type 5 or Type 8 polysaccharides. Approximately 60% of human strains are Type 8 and approximately 30% are Type 5. The structures of Type 5 and Type 8 capsular polysaccharide antigens are described in Moreau et al., (1990) and Fournier et al., (1984). Both have FucNAcp in their repeat unit as well as ManNAcA which can be used to introduce a sulfhydryl group. The structures are:

[0217] Type 5

[0218]  $\rightarrow$ 4)- $\beta$ -D-ManNAcA(3OAc)-(1 $\rightarrow$ 4)- $\alpha$ -L-FucNAc(1 $\rightarrow$ 3)- $\beta$ -D-FucNAc-(1 $\rightarrow$

[0219] Type 8

[0220]  $\rightarrow$ 3)- $\beta$ -D-ManNAcA(4OAc)-(1 $\rightarrow$ 3)- $\alpha$ -L-FucNAc(1 $\rightarrow$ 3)- $\beta$ -D-FucNAc-(1 $\rightarrow$

[0221] Recently (Jones, 2005) NMR spectroscopy revised the structures to:

[0222] Type 5

[0223]  $\rightarrow$ 4)- $\beta$ -D-ManNAcA-(1 $\rightarrow$ 4)- $\alpha$ -L-FucNAc(3OAc)-(1 $\rightarrow$ 3)- $\beta$ -D-FucNAc-(1 $\rightarrow$

[0224] Type 8

[0225]  $\rightarrow$ 3)- $\beta$ -D-ManNAcA(4OAc)-(1 $\rightarrow$ 3)- $\alpha$ -L-FucNAc(1 $\rightarrow$ 3)- $\alpha$ -D-FucNAc(1 $\rightarrow$

[0226] Polysaccharides may be extracted from the appropriate strain of *S. aureus* using method well known to of skill in the art, See U.S. Pat. No. 6,294,177. For example, ATCC 12902 is a Type 5 *S. aureus* strain and ATCC 12605 is a Type 8 *S. aureus* strain.

[0227] Polysaccharides are of native size or alternatively may be sized, for instance by microfluidisation, ultrasonic irradiation, or by chemical treatment. The invention also covers oligosaccharides derived from the type 5 and 8 polysaccharides from *S. aureus*. The type 5 and 8 polysaccharides included in the immunogenic composition of the invention are preferably conjugated to a carrier protein as described below or are alternatively unconjugated. The immunogenic compositions of the invention alternatively contains either type 5 or type 8 polysaccharide.

[0228] C. *S. aureus* 336 Antigen

[0229] In an embodiment, the immunogenic composition of the invention comprises the *S. aureus* 336 antigen described in U.S. Pat. No. 6,294,177. The 336 antigen comprises  $\beta$ -linked hexosamine, contains no O-acetyl groups, and specifically binds to antibodies to *S. aureus* Type 336 deposited under ATCC 55804. In an embodiment, the 336 antigen is

a polysaccharide which is of native size or alternatively may be sized, for instance by microfluidisation, ultrasonic irradiation, or by chemical treatment. The invention also covers oligosaccharides derived from the 336 antigen. The 336 antigen can be unconjugated or conjugated to a carrier protein.

**[0230]** D. Type I, II and III Polysaccharides from *S. epidermidis*

**[0231]** Amongst the problems associated with the use of polysaccharides in vaccination, is the fact that polysaccharides per se are poor immunogens. It is preferred that the polysaccharides utilized in the invention are linked to a protein carrier which provide bystander T-cell help to improve immunogenicity. Examples of such carriers which may be conjugated to polysaccharide immunogens include the Diphtheria and Tetanus toxoids (DT, DT CRM197 and TT respectively), Keyhole Limpet Haemocyanin (KLH), and the purified protein derivative of Tuberculin (PPD), *Pseudomonas aeruginosa* exoprotein A (rEPA), protein D from *Haemophilus influenzae*, pneumolysin or fragments of any of the above. Fragments suitable for use include fragments encompassing T-helper epitopes. In particular the protein D fragment from *H. influenzae* will preferably contain the N-terminal 1/3 of the protein. Protein D is an IgD-binding protein from *Haemophilus influenzae* (EP 0 594 610 B1) and is a potential immunogen. In addition, staphylococcal proteins may be used as a carrier protein in the polysaccharide conjugates of the invention.

**[0232]** A carrier protein that would be particularly advantageous to use in the context of a staphylococcal vaccine is staphylococcal alpha toxoid. The native form may be conjugated to a polysaccharide since the process of conjugation reduces toxicity. Preferably genetically detoxified alpha toxins such as the His35Leu or His35Arg variants are used as carriers since residual toxicity is lower. Alternatively the alpha toxin is chemically detoxified by treatment with a cross-linking reagent, formaldehyde or glutaraldehyde. A genetically detoxified alpha toxin is optionally chemically detoxified, preferably by treatment with a cross-linking reagent, formaldehyde or glutaraldehyde to further reduce toxicity.

**[0233]** The polysaccharides may be linked to the carrier protein(s) by any known method (for example those methods described in U.S. Pat. Nos. 4,372,945, 4,474,757, and 4,356,170). Preferably, CDAP conjugation chemistry is carried out (see WO95/08348). In CDAP, the cyanylating reagent 1-cyano-dimethylaminopyridinium tetrafluoroborate (CDAP) is preferably used for the synthesis of polysaccharide-protein conjugates. The cyanilation reaction can be performed under relatively mild conditions, which avoids hydrolysis of the alkaline sensitive polysaccharides. This synthesis allows direct coupling to a carrier protein.

**[0234]** Conjugation preferably involves producing a direct linkage between the carrier protein and polysaccharide. Optionally a spacer (such as adipic dihydride (ADH)) may be introduced between the carrier protein and the polysaccharide.

#### IV. IMMUNE RESPONSE AND ASSAYS

**[0235]** As discussed above, the invention concerns evoking or inducing an immune response in a subject against a variant SpA or coagulase peptide. In one embodiment, the immune response can protect against or treat a subject having, suspected of having, or at risk of developing an infection or related disease, particularly those related to staphylococci.

One use of the immunogenic compositions of the invention is to prevent nosocomial infections by inoculating a subject prior to undergoing procedures in a hospital or other environment having an increased risk of infection.

**[0236]** A. Immunoassays

**[0237]** The present invention includes the implementation of serological assays to evaluate whether and to what extent an immune response is induced or evoked by compositions of the invention. There are many types of immunoassays that can be implemented. Immunoassays encompassed by the present invention include, but are not limited to, those described in U.S. Pat. No. 4,367,110 (double monoclonal antibody sandwich assay) and U.S. Pat. No. 4,452,901 (western blot). Other assays include immunoprecipitation of labeled ligands and immunocytochemistry, both in vitro and in vivo.

**[0238]** Immunoassays generally are binding assays. Certain preferred immunoassays are the various types of enzyme linked immunosorbent assays (ELISAs) and radioimmunoassays (RIA) known in the art. Immunohistochemical detection using tissue sections is also particularly useful. In one example, antibodies or antigens are immobilized on a selected surface, such as a well in a polystyrene microtiter plate, dipstick, or column support. Then, a test composition suspected of containing the desired antigen or antibody, such as a clinical sample, is added to the wells. After binding and washing to remove non specifically bound immune complexes, the bound antigen or antibody may be detected. Detection is generally achieved by the addition of another antibody, specific for the desired antigen or antibody, that is linked to a detectable label. This type of ELISA is known as a "sandwich ELISA." Detection also may be achieved by the addition of a second antibody specific for the desired antigen, followed by the addition of a third antibody that has binding affinity for the second antibody, with the third antibody being linked to a detectable label.

**[0239]** Competition ELISAs are also possible implementations in which test samples compete for binding with known amounts of labeled antigens or antibodies. The amount of reactive species in the unknown sample is determined by mixing the sample with the known labeled species before or during incubation with coated wells. The presence of reactive species in the sample acts to reduce the amount of labeled species available for binding to the well and thus reduces the ultimate signal. Irrespective of the format employed, ELISAs have certain features in common, such as coating, incubating or binding, washing to remove non specifically bound species, and detecting the bound immune complexes.

**[0240]** Antigen or antibodies may also be linked to a solid support, such as in the form of plate, beads, dipstick, membrane, or column matrix, and the sample to be analyzed is applied to the immobilized antigen or antibody. In coating a plate with either antigen or antibody, one will generally incubate the wells of the plate with a solution of the antigen or antibody, either overnight or for a specified period. The wells of the plate will then be washed to remove incompletely-adsorbed material. Any remaining available surfaces of the wells are then "coated" with a nonspecific protein that is antigenically neutral with regard to the test antisera. These include bovine serum albumin (BSA), casein, and solutions of milk powder. The coating allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific binding of antisera onto the surface.

**[0241] B. Diagnosis of Bacterial Infection**

**[0242]** In addition to the use of proteins, polypeptides, and/or peptides, as well as antibodies binding these polypeptides, proteins, and/or peptides, to treat or prevent infection as described above, the present invention contemplates the use of these polypeptides, proteins, peptides, and/or antibodies in a variety of ways, including the detection of the presence of Staphylococci to diagnose an infection, whether in a patient or on medical equipment which may also become infected. In accordance with the invention, a preferred method of detecting the presence of infections involves the steps of obtaining a sample suspected of being infected by one or more staphylococcal bacteria species or strains, such as a sample taken from an individual, for example, from one's blood, saliva, tissues, bone, muscle, cartilage, or skin. Following isolation of the sample, diagnostic assays utilizing the polypeptides, proteins, peptides, and/or antibodies of the present invention may be carried out to detect the presence of staphylococci, and such assay techniques for determining such presence in a sample are well known to those skilled in the art and include methods such as radioimmunoassay, western blot analysis and ELISA assays. In general, in accordance with the invention, a method of diagnosing an infection is contemplated wherein a sample suspected of being infected with staphylococci has added to it the polypeptide, protein, peptide, antibody, or monoclonal antibody in accordance with the present invention, and staphylococci are indicated by antibody binding to the polypeptides, proteins, and/or peptides, or polypeptides, proteins, and/or peptides binding to the antibodies in the sample.

**[0243]** Accordingly, antibodies in accordance with the invention may be used for the prevention of infection from staphylococcal bacteria (i.e., passive immunization), for the treatment of an ongoing infection, or for use as research tools. The term "antibodies" as used herein includes monoclonal, polyclonal, chimeric, single chain, bispecific, simianized, and humanized or primatized antibodies as well as Fab fragments, such as those fragments which maintain the binding specificity of the antibodies, including the products of an Fab immunoglobulin expression library. Accordingly, the invention contemplates the use of single chains such as the variable heavy and light chains of the antibodies. Generation of any of these types of antibodies or antibody fragments is well known to those skilled in the art. Specific examples of the generation of an antibody to a bacterial protein can be found in U.S. Patent Application Pub. No. 20030153022, which is incorporated herein by reference in its entirety.

**[0244]** Any of the above described polypeptides, proteins, peptides, and/or antibodies may be labeled directly with a detectable label for identification and quantification of staphylococcal bacteria. Labels for use in immunoassays are generally known to those skilled in the art and include enzymes, radioisotopes, and fluorescent, luminescent and chromogenic substances, including colored particles such as colloidal gold or latex beads. Suitable immunoassays include enzyme-linked immunosorbent assays (ELISA).

**[0245] C. Protective Immunity**

**[0246]** In some embodiments of the invention, proteinaceous compositions confer protective immunity to a subject. Protective immunity refers to a body's ability to mount a specific immune response that protects the subject from developing a particular disease or condition that involves the

agent against which there is an immune response. An immunogenically effective amount is capable of conferring protective immunity to the subject.

**[0247]** As used herein in the specification and in the claims section that follows, the term polypeptide or peptide refer to a stretch of amino acids covalently linked there amongst via peptide bonds. Different polypeptides have different functionalities according to the present invention. While according to one aspect, a polypeptide is derived from an immunogen designed to induce an active immune response in a recipient, according to another aspect of the invention, a polypeptide is derived from an antibody which results following the elicitation of an active immune response in, for example, an animal, and which can serve to induce a passive immune response in the recipient. In both cases, however, the polypeptide is encoded by a polynucleotide according to any possible codon usage.

**[0248]** As used herein the phrase "immune response" or its equivalent "immunological response" refers to the development of a humoral (antibody mediated), cellular (mediated by antigen-specific T cells or their secretion products) or both humoral and cellular response directed against a protein, peptide, carbohydrate, or polypeptide of the invention in a recipient patient. Such a response can be an active response induced by administration of immunogen or a passive response induced by administration of antibody, antibody containing material, or primed T-cells. A cellular immune response is elicited by the presentation of polypeptide epitopes in association with Class I or Class II MHC molecules, to activate antigen-specific CD4 (+) T helper cells and/or CD8 (+) cytotoxic T cells. The response may also involve activation of monocytes, macrophages, NK cells, basophils, dendritic cells, astrocytes, microglia cells, eosinophils or other components of innate immunity. As used herein "active immunity" refers to any immunity conferred upon a subject by administration of an antigen.

**[0249]** As used herein "passive immunity" refers to any immunity conferred upon a subject without administration of an antigen to the subject. "Passive immunity" therefore includes, but is not limited to, administration of activated immune effectors including cellular mediators or protein mediators (e.g., monoclonal and/or polyclonal antibodies) of an immune response. A monoclonal or polyclonal antibody composition may be used in passive immunization for the prevention or treatment of infection by organisms that carry the antigen recognized by the antibody. An antibody composition may include antibodies that bind to a variety of antigens that may in turn be associated with various organisms. The antibody component can be a polyclonal antiserum. In certain aspects the antibody or antibodies are affinity purified from an animal or second subject that has been challenged with an antigen(s). Alternatively, an antibody mixture may be used, which is a mixture of monoclonal and/or polyclonal antibodies to antigens present in the same, related, or different microbes or organisms, such as gram-positive bacteria, gram-negative bacteria, including but not limited to *staphylococcus* bacteria.

**[0250]** Passive immunity may be imparted to a patient or subject by administering to the patient immunoglobulins (Ig) and/or other immune factors obtained from a donor or other non-patient source having a known immunoreactivity. In other aspects, an antigenic composition of the present invention can be administered to a subject who then acts as a source or donor for globulin, produced in response to challenge with



the antigenic composition (“hyperimmune globulin”), that contains antibodies directed against *Staphylococcus* or other organism. A subject thus treated would donate plasma from which hyperimmune globulin would then be obtained, via conventional plasma-fractionation methodology, and administered to another subject in order to impart resistance against or to treat *staphylococcus* infection. Hyperimmune globulins according to the invention are particularly useful for immune-compromised individuals, for individuals undergoing invasive procedures or where time does not permit the individual to produce their own antibodies in response to vaccination. See U.S. Pat. Nos. 6,936,258, 6,770,278, 6,756,361, 5,548,066, 5,512,282, 4,338,298, and 4,748,018, each of which is incorporated herein by reference in its entirety, for exemplary methods and compositions related to passive immunity.

**[0251]** For purposes of this specification and the accompanying claims the terms “epitope” and “antigenic determinant” are used interchangeably to refer to a site on an antigen to which B and/or T cells respond or recognize. B-cell epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, e.g., Epitope Mapping Protocols (1996). Antibodies that recognize the same epitope can be identified in a simple immunoassay showing the ability of one antibody to block the binding of another antibody to a target antigen. T-cells recognize continuous epitopes of about nine amino acids for CD8 cells or about 13-15 amino acids for CD4 cells. T cells that recognize the epitope can be identified by in vitro assays that measure antigen-dependent proliferation, as determined by <sup>3</sup>H-thymidine incorporation by primed T cells in response to an epitope (Burke et al., 1994), by antigen-dependent killing (cytotoxic T lymphocyte assay, Tigges et al., 1996) or by cytokine secretion.

**[0252]** The presence of a cell-mediated immunological response can be determined by proliferation assays (CD4 (+) T cells) or CTL (cytotoxic T lymphocyte) assays. The relative contributions of humoral and cellular responses to the protective or therapeutic effect of an immunogen can be distinguished by separately isolating IgG and T-cells from an immunized syngeneic animal and measuring protective or therapeutic effect in a second subject.

**[0253]** As used herein and in the claims, the terms “antibody” or “immunoglobulin” are used interchangeably and refer to any of several classes of structurally related proteins that function as part of the immune response of an animal or recipient, which proteins include IgG, IgD, IgE, IgA, IgM and related proteins.

**[0254]** Under normal physiological conditions antibodies are found in plasma and other body fluids and in the membrane of certain cells and are produced by lymphocytes of the type denoted B cells or their functional equivalent. Antibodies of the IgG class are made up of four polypeptide chains linked together by disulfide bonds. The four chains of intact IgG molecules are two identical heavy chains referred to as H-chains and two identical light chains referred to as L-chains.

**[0255]** In order to produce polyclonal antibodies, a host, such as a rabbit or goat, is immunized with the antigen or antigen fragment, generally with an adjuvant and, if necessary, coupled to a carrier. Antibodies to the antigen are subsequently collected from the sera of the host. The polyclonal antibody can be affinity purified against the antigen rendering it monospecific.

**[0256]** Monoclonal antibodies can be produced by hyperimmunization of an appropriate donor with the antigen or ex-vivo by use of primary cultures of splenic cells or cell lines derived from spleen (Anavi, 1998; Huston et al., 1991; Johnson et al., 1991; Mernaugh et al., 1995).

**[0257]** As used herein and in the claims, the phrase “an immunological portion of an antibody” includes a Fab fragment of an antibody, a Fv fragment of an antibody, a heavy chain of an antibody, a light chain of an antibody, a heterodimer consisting of a heavy chain and a light chain of an antibody, a variable fragment of a light chain of an antibody, a variable fragment of a heavy chain of an antibody, and a single chain variant of an antibody, which is also known as scFv. In addition, the term includes chimeric immunoglobulins which are the expression products of fused genes derived from different species, one of the species can be a human, in which case a chimeric immunoglobulin is said to be humanized. Typically, an immunological portion of an antibody competes with the intact antibody from which it was derived for specific binding to an antigen.

**[0258]** Optionally, an antibody or preferably an immunological portion of an antibody, can be chemically conjugated to, or expressed as, a fusion protein with other proteins. For purposes of this specification and the accompanying claims, all such fused proteins are included in the definition of antibodies or an immunological portion of an antibody.

**[0259]** As used herein the terms “immunogenic agent” or “immunogen” or “antigen” are used interchangeably to describe a molecule capable of inducing an immunological response against itself on administration to a recipient, either alone, in conjunction with an adjuvant, or presented on a display vehicle.

**[0260]** D. Treatment Methods

**[0261]** A method of the present invention includes treatment for a disease or condition caused by a *staphylococcus* pathogen. An immunogenic polypeptide of the invention can be given to induce an immune response in a person infected with *staphylococcus* or suspected of having been exposed to *staphylococcus*. Methods may be employed with respect to individuals who have tested positive for exposure to *staphylococcus* or who are deemed to be at risk for infection based on possible exposure.

**[0262]** In particular, the invention encompasses a method of treatment for staphylococcal infection, particularly hospital acquired nosocomial infections. The immunogenic compositions and vaccines of the invention are particularly advantageous to use in cases of elective surgery. Such patients will know the date of surgery in advance and could be inoculated in advance. The immunogenic compositions and vaccines of the invention are also advantageous to use to inoculate health care workers.

**[0263]** In some embodiments, the treatment is administered in the presence of adjuvants or carriers or other staphylococcal antigens. Furthermore, in some examples, treatment comprises administration of other agents commonly used against bacterial infection, such as one or more antibiotics.

[0264] The use of peptides for vaccination can require, but not necessarily, conjugation of the peptide to an immunogenic carrier protein, such as hepatitis B surface antigen, keyhole limpet hemocyanin, or bovine serum albumin. Methods for performing this conjugation are well known in the art.

#### V. VACCINE AND OTHER PHARMACEUTICAL COMPOSITIONS AND ADMINISTRATION

##### [0265] A. Vaccines

[0266] The present invention includes methods for preventing or ameliorating staphylococcal infections, particularly hospital acquired nosocomial infections. As such, the invention contemplates vaccines for use in both active and passive immunization embodiments. Immunogenic compositions, proposed to be suitable for use as a vaccine, may be prepared from immunogenic SpA polypeptide(s), such as a SpA domain D variant, or immunogenic coagulases. In other embodiments SpA or coagulases can be used in combination with other secreted virulence proteins, surface proteins or immunogenic fragments thereof. In certain aspects, antigenic material is extensively dialyzed to remove undesired small molecular weight molecules and/or lyophilized for more ready formulation into a desired vehicle.

[0267] Other options for a protein/peptide-based vaccine involve introducing nucleic acids encoding the antigen(s) as DNA vaccines. In this regard, recent reports described construction of recombinant vaccinia viruses expressing either 10 contiguous minimal CTL epitopes (Thomson, 1996) or a combination of B cell, cytotoxic T-lymphocyte (CTL), and T-helper (Th) epitopes from several microbes (An, 1997), and successful use of such constructs to immunize mice for priming protective immune responses. Thus, there is ample evidence in the literature for successful utilization of peptides, peptide-pulsed antigen presenting cells (APCs), and peptide-encoding constructs for efficient *in vivo* priming of protective immune responses. The use of nucleic acid sequences as vaccines is exemplified in U.S. Pat. Nos. 5,958,895 and 5,620,896.

[0268] The preparation of vaccines that contain polypeptide or peptide sequence(s) as active ingredients is generally well understood in the art, as exemplified by U.S. Pat. Nos. 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all of which are incorporated herein by reference. Typically, such vaccines are prepared as injectables either as liquid solutions or suspensions: solid forms suitable for solution in or suspension in liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients that are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants that enhance the effectiveness of the vaccines. In specific embodiments, vaccines are formulated with a combination of substances, as described in U.S. Pat. Nos. 6,793,923 and 6,733,754, which are incorporated herein by reference.

[0269] Vaccines may be conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, poly-

alkalene glycols or triglycerides: such suppositories may be formed from mixtures containing the active ingredient in the range of about 0.5% to about 10%, preferably about 1% to about 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 10% to about 95% of active ingredient, preferably about 25% to about 70%.

[0270] The polypeptides and polypeptide-encoding DNA constructs may be formulated into a vaccine as neutral or salt forms. Pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the peptide) and those that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like.

[0271] Typically, vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, including the capacity of the individual's immune system to synthesize antibodies and the degree of protection desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are of the order of several hundred micrograms of active ingredient per vaccination. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed by subsequent inoculations or other administrations.

[0272] The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These are believed to include oral application within a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection and the like. The dosage of the vaccine will depend on the route of administration and will vary according to the size and health of the subject.

[0273] In certain instances, it will be desirable to have multiple administrations of the vaccine, e.g., 2, 3, 4, 5, 6 or more administrations. The vaccinations can be at 1, 2, 3, 4, 5, 6, 7, 8, to 5, 6, 7, 8, 9, 10, 11, 12 twelve week intervals, including all ranges there between. Periodic boosters at intervals of 1-5 years will be desirable to maintain protective levels of the antibodies. The course of the immunization may be followed by assays for antibodies against the antigens, as described in U.S. Pat. Nos. 3,791,932; 4,174,384 and 3,949,064.

##### [0274] 1. Carriers

[0275] A given composition may vary in its immunogenicity. It is often necessary therefore to boost the host immune system, as may be achieved by coupling a peptide or polypeptide to a carrier. Exemplary and preferred carriers are keyhole limpet hemocyanin (KLH) and bovine serum albumin (BSA). Other albumins such as ovalbumin, mouse serum albumin, or rabbit serum albumin can also be used as carriers. Means for conjugating a polypeptide to a carrier protein are well known in the art and include glutaraldehyde, m-maleimidobencoyl-N-hydroxysuccinimide ester, carbodiimide, and bis-biazo-tized benzidine.

**[0276]** 2. Adjuvants

**[0277]** The immunogenicity of polypeptide or peptide compositions can be enhanced by the use of non-specific stimulators of the immune response, known as adjuvants. Suitable adjuvants include all acceptable immunostimulatory compounds, such as cytokines, toxins, or synthetic compositions. A number of adjuvants can be used to enhance an antibody response against a variant SpA polypeptide or coagulase, or any other bacterial protein or combination contemplated herein. Adjuvants can (1) trap the antigen in the body to cause a slow release; (2) attract cells involved in the immune response to the site of administration; (3) induce proliferation or activation of immune system cells; or (4) improve the spread of the antigen throughout the subject's body.

**[0278]** Adjuvants include, but are not limited to, oil-in-water emulsions, water-in-oil emulsions, mineral salts, polynucleotides, and natural substances. Specific adjuvants that may be used include IL-1, IL-2, IL-4, IL-7, IL-12,  $\gamma$ -interferon, GMCSF, BCG, aluminum salts, such as aluminum hydroxide or other aluminum compound, MDP compounds, such as thur-MDP and nor-MDP, CGP (MTP-PE), lipid A, and monophosphoryl lipid A (MPL). RIBI, which contains three components extracted from bacteria, MPL, trehalose dimycolate (TDM), and cell wall skeleton (CWS) in a 2% squalene/Tween 80 emulsion. MHC antigens may even be used. Others adjuvants or methods are exemplified in U.S. Pat. Nos. 6,814,971, 5,084,269, 6,656,462, each of which is incorporated herein by reference).

**[0279]** Various methods of achieving adjuvant affect for the vaccine includes use of agents such as aluminum hydroxide or phosphate (alum), commonly used as about 0.05 to about 0.1% solution in phosphate buffered saline, admixture with synthetic polymers of sugars (Carbopol®) used as an about 0.25% solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between about 70° to about 101° C. for a 30-second to 2-minute period, respectively. Aggregation by reactivating with pepsin-treated (Fab) antibodies to albumin; mixture with bacterial cells (e.g., *C. parvum*), endotoxins or lipopolysaccharide components of Gram-negative bacteria; emulsion in physiologically acceptable oil vehicles (e.g., mannide mono-oleate (Aracel A)); or emulsion with a 20% solution of a perfluorocarbon (Fluosol-DA®) used as a block substitute may also be employed to produce an adjuvant effect.

**[0280]** Examples of and often preferred adjuvants include complete Freund's adjuvant (a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*), incomplete Freund's adjuvants, and aluminum hydroxide.

**[0281]** In some aspects, it is preferred that the adjuvant be selected to be a preferential inducer of either a Th1 or a Th2 type of response. High levels of Th1-type cytokines tend to favor the induction of cell mediated immune responses to a given antigen, while high levels of Th2-type cytokines tend to favor the induction of humoral immune responses to the antigen.

**[0282]** The distinction of Th1 and Th2-type immune response is not absolute. In reality an individual will support an immune response which is described as being predominantly Th1 or predominantly Th2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4+ T cell clones by Mosmann and Coffman (Mosmann, and Coffman, 1989). Traditionally,

Th1-type responses are associated with the production of the INF- $\gamma$  and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of Th1-type immune responses are not produced by T-cells, such as IL-12. In contrast, Th2-type responses are associated with the secretion of IL-4, IL-5, IL-6, IL-10.

**[0283]** In addition to adjuvants, it may be desirable to co-administer biologic response modifiers (BRM) to enhance immune responses. BRMs have been shown to upregulate T cell immunity or downregulate suppressor cell activity. Such BRMs include, but are not limited to, Cimetidine (CIM; 1200 mg/d) (Smith/Kline, PA); or low-dose Cyclophosphamide (CYP; 300 mg/m<sup>2</sup>) (Johnson/Mead, NJ) and cytokines such as  $\gamma$ -interferon, IL-2, or IL-12 or genes encoding proteins involved in immune helper functions, such as B-7.

**[0284]** B. Lipid Components and Moieties

**[0285]** In certain embodiments, the present invention concerns compositions comprising one or more lipids associated with a nucleic acid or a polypeptide/peptide. A lipid is a substance that is insoluble in water and extractable with an organic solvent. Compounds other than those specifically described herein are understood by one of skill in the art as lipids, and are encompassed by the compositions and methods of the present invention. A lipid component and a non-lipid may be attached to one another, either covalently or non-covalently.

**[0286]** A lipid may be a naturally occurring lipid or a synthetic lipid. However, a lipid is usually a biological substance. Biological lipids are well known in the art, and include for example, neutral fats, phospholipids, phosphoglycerides, steroids, terpenes, lysolipids, glycosphingolipids, glucolipids, sulphatides, lipids with ether and ester-linked fatty acids and polymerizable lipids, and combinations thereof.

**[0287]** A nucleic acid molecule or a polypeptide/peptide, associated with a lipid may be dispersed in a solution containing a lipid, dissolved with a lipid, emulsified with a lipid, mixed with a lipid, combined with a lipid, covalently bonded to a lipid, contained as a suspension in a lipid or otherwise associated with a lipid. A lipid or lipid-poxvirus-associated composition of the present invention is not limited to any particular structure. For example, they may also simply be interspersed in a solution, possibly forming aggregates which are not uniform in either size or shape. In another example, they may be present in a bilayer structure, as micelles, or with a "collapsed" structure. In another non-limiting example, a lipofectamine (Gibco BRL)-poxvirus or Superfect (Qiagen)-poxvirus complex is also contemplated.

**[0288]** In certain embodiments, a composition may comprise about 1%, about 2%, about 3%, about 4%, about 5%, about 6%, about 7%, about 8%, about 9%, about 10%, about 11%, about 12%, about 13%, about 14%, about 15%, about 16%, about 17%, about 18%, about 19%, about 20%, about 21%, about 22%, about 23%, about 24%, about 25%, about 26%, about 27%, about 28%, about 29%, about 30%, about 31%, about 32%, about 33%, about 34%, about 35%, about 36%, about 37%, about 38%, about 39%, about 40%, about 41%, about 42%, about 43%, about 44%, about 45%, about 46%, about 47%, about 48%, about 49%, about 50%, about 51%, about 52%, about 53%, about 54%, about 55%, about 56%, about 57%, about 58%, about 59%, about 60%, about 61%, about 62%, about 63%, about 64%, about 65%, about 66%, about 67%, about 68%, about 69%, about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about

81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or any range therebetween, of a particular lipid, lipid type, or non-lipid component such as an adjuvant, antigen, peptide, polypeptide, sugar, nucleic acid or other material disclosed herein or as would be known to one of skill in the art. In a non-limiting example, a composition may comprise about 10% to about 20% neutral lipids, and about 33% to about 34% of a cerebroside, and about 1% cholesterol. In another non-limiting example, a liposome may comprise about 4% to about 12% terpenes, wherein about 1% of the micelle is specifically lycopene, leaving about 3% to about 11% of the liposome as comprising other terpenes; and about 10% to about 35% phosphatidyl choline, and about 1% of a non-lipid component. Thus, it is contemplated that compositions of the present invention may comprise any of the lipids, lipid types or other components in any combination or percentage range.

#### [0289] C. Combination Therapy

[0290] The compositions and related methods of the present invention, particularly administration of a secreted virulence factor or surface protein, including a variant SpA polypeptide or peptide, and/or other bacterial peptides or proteins to a patient/subject, may also be used in combination with the administration of traditional therapies. These include, but are not limited to, the administration of antibiotics such as streptomycin, ciprofloxacin, doxycycline, gentamycin, chloramphenicol, trimethoprim, sulfamethoxazole, ampicillin, tetracycline or various combinations of antibiotics.

[0291] In one aspect, it is contemplated that a polypeptide vaccine and/or therapy is used in conjunction with antibacterial treatment. Alternatively, the therapy may precede or follow the other agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agents and/or a proteins or polynucleotides are administered separately, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and antigenic composition would still be able to exert an advantageously combined effect on the subject. In such instances, it is contemplated that one may administer both modalities within about 12-24 h of each other or within about 6-12 h of each other. In some situations, it may be desirable to extend the time period for administration significantly, where several days (2, 3, 4, 5, 6 or 7) to several weeks (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

[0292] Various combinations may be employed, for example antibiotic therapy is "A" and the immunogenic molecule given as part of an immune therapy regime, such as an antigen, is "B":

[0293] A/B/A B/A/B B/B/A A/A/B A/B/B B/A/A A/B/B B/A/B/B

[0294] B/B/B/A B/B/A/B A/A/B/B A/B/A/B A/B/B/A B/B/A/A

[0295] B/A/B/A B/A/A/B A/A/A/B B/A/A/A A/B/A/A A/A/B/A

[0296] Administration of the immunogenic compositions of the present invention to a patient/subject will follow general protocols for the administration of such compounds, taking into account the toxicity, if any, of the SpA composition, or other compositions described herein. It is expected that the treatment cycles would be repeated as necessary. It

also is contemplated that various standard therapies, such as hydration, may be applied in combination with the described therapy.

#### [0297] D. General Pharmaceutical Compositions

[0298] In some embodiments, pharmaceutical compositions are administered to a subject. Different aspects of the present invention involve administering an effective amount of a composition to a subject. In some embodiments of the present invention, staphylococcal antigens, members of the Ess pathway, including polypeptides or peptides of the Esa or Esx class, and/or members of sortase substrates may be administered to the patient to protect against infection by one or more *staphylococcus* pathogens. Alternatively, an expression vector encoding one or more such polypeptides or peptides may be given to a patient as a preventative treatment. Additionally, such compounds can be administered in combination with an antibiotic or an antibacterial. Such compositions will generally be dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium.

[0299] In addition to the compounds formulated for parenteral administration, such as those for intravenous or intramuscular injection, other pharmaceutically acceptable forms include, e.g., tablets or other solids for oral administration; time release capsules; and any other form currently used, including creams, lotions, mouthwashes, inhalants and the like.

[0300] The active compounds of the present invention can be formulated for parenteral administration, e.g., formulated for injection via the intravenous, intramuscular, sub-cutaneous, or even intraperitoneal routes. The preparation of an aqueous composition that contains a compound or compounds that increase the expression of an MHC class I molecule will be known to those of skill in the art in light of the present disclosure. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for use to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and, the preparations can also be emulsified.

[0301] Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[0302] The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil, or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that it may be easily injected. It also should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

[0303] The proteinaceous compositions may be formulated into a neutral or salt form. Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example,

sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

**[0304]** The carrier also can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion, and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

**[0305]** Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques, which yield a powder of the active ingredient, plus any additional desired ingredient from a previously sterile-filtered solution thereof.

**[0306]** Administration of the compositions according to the present invention will typically be via any common route. This includes, but is not limited to oral, nasal, or buccal administration. Alternatively, administration may be by orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal, intranasal, or intravenous injection. In certain embodiments, a vaccine composition may be inhaled (e.g., U.S. Pat. No. 6,651,655, which is specifically incorporated by reference). Such compositions would normally be administered as pharmaceutically acceptable compositions that include physiologically acceptable carriers, buffers or other excipients. As used herein, the term "pharmaceutically acceptable" refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem complications commensurate with a reasonable benefit/risk ratio. The term "pharmaceutically acceptable carrier," means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting a chemical agent.

**[0307]** For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous, and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in

isotonic NaCl solution and either added to hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, Remington's Pharmaceutical Sciences, 1990). Some variation in dosage will necessarily occur depending on the condition of the subject. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

**[0308]** An effective amount of therapeutic or prophylactic composition is determined based on the intended goal. The term "unit dose" or "dosage" refers to physically discrete units suitable for use in a subject, each unit containing a predetermined quantity of the composition calculated to produce the desired responses discussed above in association with its administration, i.e., the appropriate route and regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the protection desired.

**[0309]** Precise amounts of the composition also depend on the judgment of the practitioner and are peculiar to each individual. Factors affecting dose include physical and clinical state of the subject, route of administration, intended goal of treatment (alleviation of symptoms versus cure), and potency, stability, and toxicity of the particular composition.

**[0310]** Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically or prophylactically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above.

**[0311]** E. In Vitro, Ex Vivo, or In Vivo Administration

**[0312]** As used herein, the term in vitro administration refers to manipulations performed on cells removed from or outside of a subject, including, but not limited to cells in culture. The term ex vivo administration refers to cells which have been manipulated in vitro, and are subsequently administered to a subject. The term in vivo administration includes all manipulations performed within a subject.

**[0313]** In certain aspects of the present invention, the compositions may be administered either in vitro, ex vivo, or in vivo. In certain in vitro embodiments, autologous B-lymphocyte cell lines are incubated with a virus vector of the instant invention for 24 to 48 hours or with a variant SpA and/or cogulase and/or any other composition described herein for two hours. The transduced cells can then be used for in vitro analysis, or alternatively for ex vivo administration. U.S. Pat. Nos. 4,690,915 and 5,199,942, both incorporated herein by reference, disclose methods for ex vivo manipulation of blood mononuclear cells and bone marrow cells for use in therapeutic applications.

**[0314]** F. Antibodies and Passive Immunization

**[0315]** Another aspect of the invention is a method of preparing an immunoglobulin for use in prevention or treatment of staphylococcal infection comprising the steps of immunizing a recipient or donor with the vaccine of the invention and isolating immunoglobulin from the recipient or donor. An immunoglobulin prepared by this method is a further aspect of the invention. A pharmaceutical composition comprising the immunoglobulin of the invention and a pharmaceutically acceptable carrier is a further aspect of the invention which could be used in the manufacture of a medicament for the treatment or prevention of staphylococcal disease. A method for treatment or prevention of staphylococcal infection com-

prising a step of administering to a patient an effective amount of the pharmaceutical preparation of the invention is a further aspect of the invention.

[0316] Inocula for polyclonal antibody production are typically prepared by dispersing the antigenic composition in a physiologically tolerable diluent such as saline or other adjuvants suitable for human use to form an aqueous composition. An immunostimulatory amount of inoculum is administered to a mammal and the inoculated mammal is then maintained for a time sufficient for the antigenic composition to induce protective antibodies.

[0317] The antibodies can be isolated to the extent desired by well known techniques such as affinity chromatography (Harlow and Lane, 1988). Antibodies can include antiserum preparations from a variety of commonly used animals, e.g. goats, primates, donkeys, swine, horses, guinea pigs, rats or man.

[0318] An immunoglobulin produced in accordance with the present invention can include whole antibodies, antibody fragments or subfragments. Antibodies can be whole immunoglobulins of any class (e.g., IgG, IgM, IgA, IgD or IgE), chimeric antibodies or hybrid antibodies with dual specificity to two or more antigens of the invention. They may also be fragments (e.g., F(ab')<sub>2</sub>, Fab', Fab, Fv and the like) including hybrid fragments. An immunoglobulin also includes natural, synthetic, or genetically engineered proteins that act like an antibody by binding to specific antigens to form a complex.

[0319] A vaccine of the present invention can be administered to a recipient who then acts as a source of immunoglobulin, produced in response to challenge from the specific vaccine. A subject thus treated would donate plasma from which hyperimmune globulin would be obtained via conventional plasma fractionation methodology. The hyperimmune globulin would be administered to another subject in order to impart resistance against or treat staphylococcal infection. Hyperimmune globulins of the invention are particularly useful for treatment or prevention of staphylococcal disease in infants, immune compromised individuals, or where treatment is required and there is no time for the individual to produce antibodies in response to vaccination.

[0320] An additional aspect of the invention is a pharmaceutical composition comprising two or more monoclonal antibodies (or fragments thereof; preferably human or humanised) reactive against at least two constituents of the immunogenic composition of the invention, which could be used to treat or prevent infection by Gram positive bacteria, preferably staphylococci, more preferably *S. aureus* or *S. epidermidis*. Such pharmaceutical compositions comprise monoclonal antibodies that can be whole immunoglobulins of any class, chimeric antibodies, or hybrid antibodies with specificity to two or more antigens of the invention. They may also be fragments (e.g., F(ab')<sub>2</sub>, Fab', Fab, Fv and the like) including hybrid fragments.

[0321] Methods of making monoclonal antibodies are well known in the art and can include the fusion of splenocytes with myeloma cells (Kohler and Milstein, 1975; Harlow and Lane, 1988). Alternatively, monoclonal Fv fragments can be obtained by screening a suitable phage display library (Vaughan et al., 1998). Monoclonal antibodies may be humanized or part humanized by known methods.

## VI. EXAMPLES

[0322] The following examples are given for the purpose of illustrating various embodiments of the invention and are not

meant to limit the present invention in any fashion. One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the methods described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses which are encompassed within the spirit of the invention as defined by the scope of the claims will occur to those skilled in the art.

### Example 1

#### Non-Toxicigenic Protein A Variants as Subunit Vaccines to Prevent *Staphylococcus aureus* Infections

##### [0323] A. Results

[0324] An animal model for *S. aureus* infection BALB/c mice were infected by intravenous injection with  $1 \times 10^7$  CFU of the human clinical isolate *S. aureus* Newman (Baba et al., 2007). Within 6 hours following infection, 99.999% of staphylococci disappeared from the blood stream and were distributed via the vasculature. Staphylococcal dissemination to peripheral tissues occurred rapidly, as the bacterial load in kidney and other peripheral organ tissues reached  $1 \times 10^5$  CFU  $g^{-1}$  within the first three hours. The staphylococcal load in kidney tissues increased by 1.5 log CFU within twenty-four hours. Forty-eight hours following infection, mice developed disseminated abscesses in multiple organs, detectable by light microscopy of hematoxylin-eosin stained, thin-sectioned kidney tissue. The initial abscess diameter was  $524 \mu\text{M}$  ( $\pm 65 \mu\text{M}$ ); lesions were initially marked by an influx of polymorphonuclear leukocytes (PMNs) and harbored no discernable organization of staphylococci, most of which appeared to reside within PMNs. On day 5 of infection, abscesses increased in size and enclosed a central population of staphylococci, surrounded by a layer of eosinophilic, amorphous material and a large cuff of PMNs. Histopathology revealed massive necrosis of PMNs in proximity to the staphylococcal nidus at the center of abscess lesions as well as a mantle of healthy phagocytes. A rim of necrotic PMNs were observed at the periphery of abscess lesions, bordering eosinophilic, amorphous material that separates healthy renal tissue from lesions. Abscesses eventually reached a diameter of  $\geq 1,524 \mu\text{M}$  on day 15 or 36. At later time intervals, the staphylococcal load was increased to  $10^4$ - $10^6$  CFU  $g^{-1}$  and growing abscess lesions migrated towards the organ capsule. Peripheral lesions were prone to rupture, thereby releasing necrotic material and staphylococci into the peritoneal cavity or the retroperitoneal space. These events resulted in bacteremia as well as a secondary wave of abscesses, eventually precipitating a lethal outcome.

[0325] To enumerate staphylococcal load in renal tissue, animals were killed, their kidneys excised and tissue homogenate spread on agar media for colony formation. On day 5 of infection, a mean of  $1 \times 10^6$  CFU  $g^{-1}$  renal tissue for *S. aureus* Newman was observed. To quantify abscess formation, kidneys were visually inspected, and each individual organ was given a score of one or zero. The final sum was divided by the total number of kidneys to calculate percent surface abscesses (Table 4). In addition, randomly chosen kidneys were fixed in formalin, embedded, thin sectioned, and stained with hematoxylin-eosin. For each kidney, four

sagittal sections at 200  $\mu\text{M}$  intervals were viewed by microscopy. The numbers of lesions were counted for each section and averaged to quantify the number of abscesses within the kidneys. *S. aureus* Newman caused  $4.364 \pm 0.889$  abscesses per kidney, and surface abscesses were observed on 14 out of 20 kidneys (70%) (Table 4).

**[0326]** When examined by scanning electron microscopy, *S. aureus* Newman was located in tightly associated lawns at the center of abscesses. Staphylococci were contained by an amorphous pseudocapsule that separated bacteria from the cuff of abscesses leukocytes. No immune cells were observed in these central nests of staphylococci, however occasional red blood cells were located among the bacteria. Bacterial populations at the abscess center, designated staphylococcal abscess communities (SAC), appeared homogenous and coated by an electron-dense, granular material. The kinetics of the appearance of infectious lesions and the morphological attributes of abscesses formed by *S. aureus* Newman were similar to those observed following mouse infection with *S. aureus* USA300 (LAC), the current epidemic community-acquired methicillin-resistant *S. aureus* (CA-MRSA) clone in the United States (Diep et al., 2006).

were cleared from renal tissues, a  $\geq 3.5 \log_{10} \text{CFU g}^{-1}$  reduction compared to the wild-type (Table 3). Thus, sortase A anchored surface proteins enable the formation of abscess lesions and the persistence of bacteria in host tissues, wherein staphylococci replicate as communities embedded in an extracellular matrix and shielded from surrounding leukocytes by an amorphous pseudocapsule.

**[0329]** Sortase A anchors a large spectrum of proteins with LPXTG motif sorting signals to the cell wall envelope, thereby providing for the surface display of many virulence factors (Mazmanian et al., 2002). To identify surface proteins required for staphylococcal abscess formation, bursa aurealis insertions were introduced in 5' coding sequences of genes that encode polypeptides with LPXTG motif proteins (Bae et al., 2004) and these mutations were transduced into *S. aureus* Newman. Mutations in the structural gene for Protein A (*spa*) reduced the staphylococcal load in infected mouse kidney tissues by  $1.004 \log_{10}$  ( $P=0.0144$ ). When analyzed for their ability to form abscesses in kidney tissues by histopathology, we observed that the *spa* mutants were unable to form abscesses as compared with the wild-type parent strain *S. aureus* Newman (wild-type *S. aureus* Newman  $4.364 \pm 0.889$

TABLE 4

Genotype	Staphylococcal load in kidney tissue			Abscess formation in kidney tissue		
	<sup>a</sup> $\log_{10} \text{CFU g}^{-1}$ tissue	<sup>b</sup> Significance (P-value)	<sup>c</sup> Reduction ( $\log_{10} \text{CFU g}^{-1}$ )	<sup>d</sup> Surface abscesses (%)	<sup>e</sup> Number of abscesses per kidney	<sup>f</sup> Significance (P-value)
	wild-type	$6.141 \pm 0.192$	—	—	70	$4.364 \pm 0.889$
$\Delta\text{srtA}$	$4.095 \pm 0.347$	$6.7 \times 10^{-6}$	2.046	0	$0.000 \pm 0.000$	0.0216
<i>spa</i>	$5.137 \pm 0.374$	0.0144	1.004	13	$0.375 \pm 0.374$	0.0356

<sup>a</sup>Means of staphylococcal load calculated as  $\log_{10} \text{CFU g}^{-1}$  in homogenized renal tissues 5 days following infection in cohorts of fifteen BALB/c mice per challenge strain. Standard error of the means ( $\pm\text{SEM}$ ) is indicated.

<sup>b</sup>Statistical significance was calculated with the Student's t-test and P-values recorded; P-values  $< 0.05$  were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as  $\log_{10} \text{CFU g}^{-1}$ .

<sup>d</sup>Abscess formation in kidney tissues five days following infection was measured by macroscopic inspection (% positive)

<sup>e</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys from eight to ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean ( $\pm\text{SEM}$ ).

<sup>f</sup>Statistical significance was calculated with the Student's t-test and P-values recorded; P-values  $< 0.05$  were deemed significant.

**[0327]** *S. aureus* Protein A (Spa) Mutants are Avirulent and Cannot Form Abscesses

**[0328]** Sortase A is a transpeptidase that immobilizes nineteen surface proteins in the envelope of *S. aureus* strain Newman (Mazmanian et al., 1999; Mazmanian et al., 2000). Earlier work identified sortase A as a virulence factor in multiple animal model systems, however the contributions of this enzyme and its anchored surface proteins to abscess formation or persistence have not yet been revealed (Jonsson et al., 2002; Weiss et al., 2004). Compared to the wild-type parent (Baba et al., 2007), an isogenic *srtA* variant ( $\Delta\text{srtA}$ ) failed to form abscess lesions on either macroscopic or histopathology examination on days 2, 5, or 15. In mice infected with the *strA* mutant, only  $1 \times 10^4 \text{CFU g}^{-1}$  was recovered from kidney tissue on day 5 of infection, which is a  $2.046 \log_{10} \text{CFU g}^{-1}$  reduction compared to the wild-type parent strain ( $P=6.73 \times 10^{-6}$ ). A similar defect was observed for the *srtA* mutant of MRSA strain USA300 (data not shown). Scanning electron microscopy showed that *srtA* mutants were highly dispersed and often associated with leukocytes in otherwise healthy renal tissue. On day fifteen following infection, *srtA* mutants

abscesses per kidney vs. the isogenic *spa* mutant with  $0.375 \pm 0.374$  lesions;  $P=0.0356$ ).

**[0330]** Protein A Blocks Innate and Adaptive Immune Responses.

**[0331]** Studies identified Protein A as a critical virulence factor during the pathogenesis of *S. aureus* infections. Earlier work demonstrated that Protein A impedes phagocytosis of staphylococci by binding the Fc component of immunoglobulin (Jensen 1958; Uhlén et al., 1984), activates platelet aggregation via the von Willebrand factor (Hartleib et al., 2000), functions as a B cell superantigen by capturing the F(ab)<sub>2</sub> region of VH3 bearing IgM (Roben et al., 1995), and, through its activation of TNFR1, can initiate staphylococcal pneumonia (Gomez et al., 2004). Due to the fact that Protein A captures immunoglobulin and displays toxic attributes, the possibility that this surface molecule may function as a vaccine in humans has not been rigorously pursued. The inventors demonstrate for the first time that Protein A variants no longer able to bind to immunoglobulins, vWF and TNFR-1 are removed of their toxigenic potential and are able to stimulate humoral immune responses that protect against staphylococcal disease.

**[0332]** Molecular Basis of Protein A Surface Display and Function.

**[0333]** Protein A is synthesized as a precursor in the bacterial cytoplasm and secreted via its YSIRK signal peptide at the cross wall, i.e., the cell division septum of staphylococci (FIG. 1). (DeDent et al., 2007; DeDent et al., 2008). Following cleavage of the C-terminal LPXTG sorting signal, Protein A is anchored to bacterial peptidoglycan crossbridges by sortase A (Schneewind et al., 1995; Mazmanian et al., 1999; Mazmanian et al., 2000). Protein A is the most abundant surface protein of staphylococci; the molecule is expressed by virtually all *S. aureus* strains (Said-Salim et al., 2003; Cespedes et al., 2005; Kennedy et al., 2008). Staphylococci turn over 15-20% of their cell wall per division cycle (Navarre and Schneewind 1999). Murine hydrolases cleave the glycan strands and wall peptides of peptidoglycan, thereby releasing Protein A with its attached C-terminal cell wall disaccharide tetrapeptide into the extracellular medium (Ton-That et al., 1999). Thus, by physiological design, Protein A is both anchored to the cell wall and displayed on the bacterial surface but also released into surrounding tissues during host infection (Marraffini et al., 2006).

**[0334]** Protein A captures immunoglobulins on the bacterial surface and this biochemical activity enables staphylococcal escape from host innate and acquired immune responses (Jensen 1958; Goodyear and Silverman 2004). Interestingly, region X of Protein A (Guss et al., 1984), a repeat domain that tethers the IgG binding domains to the LPXTG sorting signal/cell wall anchor, is perhaps the most variable portion of the staphylococcal genome (Schneewind et al., 1992; Said-Salim et al., 2003). Each of the five immunoglobulin binding domains of Protein A (SpA), formed from three helix bundles and designated E, D, A, B, and C, exerts similar structural and functional properties (Sjödahl 1977; Jansson et al., 1998). The solution and crystal structure of domain D has been solved both with and without the Fc and V<sub>H3</sub> (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille et al., 2000).

**[0335]** In the crystal structure complex, the Fab interacts with helix II and helix III of domain D via a surface composed of four VH region  $\beta$ -strands (Graille et al., 2000). The major axis of helix II of domain D is approximately 50° to the orientation of the strands, and the interhelical portion of domain D is most proximal to the C0 strand. The site of interaction on Fab is remote from the Ig light chain and the heavy chain constant region. The interaction involves the following domain D residues: Asp-36 of helix II as well as Asp-37 and Gln-40 in the loop between helix II and helix III, in addition to several other residues with SpA-D (Graille et al., 2000). Both interacting surfaces are composed predominantly of polar side chains, with three negatively charged residues on domain D and two positively charged residues on the 2A2 Fab buried by the interaction, providing an overall electrostatic attraction between the two molecules. Of the five polar interactions identified between Fab and domain D, three are between side chains. A salt bridge is formed between Arg-H19 and Asp-36 and two hydrogen bonds are made between Tyr-H59 and Asp-37 and between Asn-H82a and Ser-33. Because of the conservation of Asp-36 and Asp-37 in all five IgG binding domains of Protein A, these residues were selected for mutagenesis.

**[0336]** The SpA-D sites responsible for Fab binding are structurally separate from the domain surface that mediates Fc $\gamma$  binding. The interaction of Fc $\gamma$  with domain B primarily

involves residues in helix I with lesser involvement of helix II (Deisenhofer 1981; Gouda et al., 1992). With the exception of the Gln-32, a minor contact in both complexes, none of the residues that mediate the Fc $\gamma$  interaction are involved in Fab binding. To examine the spatial relationship between these different Ig-binding sites, the SpA domains in these complexes have been superimposed to construct a model of a complex between Fab, the SpA-domain D, and the Fc $\gamma$  molecule. In this ternary model, Fab and Fc $\gamma$  form a sandwich about opposite faces of the helix II without evidence of steric hindrance of either interaction. These findings illustrate how, despite its small size (i.e., 56-61 aa), a SpA domain can simultaneously display both activities, explaining experimental evidence that the interactions of Fab with an individual domain are noncompetitive. Residues for the interaction between SpA-D and Fc $\gamma$  are Gln-9 and Gln-10.

**[0337]** In contrast, occupancy of the Fc portion of IgG on the domain D blocks its interaction with vWF A1 and probably also TNFR1 (O'Seaghdha et al., 2006). Mutations in residues essential for IgG Fc binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF A1 and TNFR1 binding (Cedergren et al., 1993; Gomez et al., 2006; O'Seaghdha et al. 2006), whereas residues critical for the V<sub>H3</sub> interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) have no impact on the binding activities of IgG Fc, vWF A1 or TNFR1 (Jansson et al., 1998; Graille et al., 2000). The Protein A immunoglobulin Fab binding activity targets a subset of B cells that express VH3 family related IgM on their surface, i.e. these molecules function as VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells rapidly proliferate and then commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e. marginal zone B cells and follicular B2 cells) (Goodyear and Silverman 2003; Goodyear and Silverman 2004). It is important to note that more than 40% of circulating B cells are targeted by the Protein A interaction and the VH3 family represents the largest family of human B cell receptors to impart protective humoral responses against pathogens (Goodyear and Silverman 2003; Goodyear and Silverman 2004). Thus, Protein A functions analogously to staphylococcal superantigens (Roben et al., 1995), albeit that the latter class of molecules, for example SEB, TSST-1, TSST-2, form complexes with the T cell receptor to inappropriately stimulate host immune responses and thereby precipitating characteristic disease features of staphylococcal infections (Roben et al., 1995; Tiedemann et al., 1995). Together these findings document the contributions of Protein A in establishing staphylococcal infections and in modulating host immune responses.

**[0338]** Non-Toxicogenic Variant of Protein A.

**[0339]** The inventors have developed a non-toxicogenic variant of staphylococcal Protein A and, with this reagent in hand, aimed for the first time to measure the immune response of animals to Protein A immunization. Further, the inventors address whether immunization of animals with a non-toxicogenic variant of Protein A could generate immune responses that raise protective immunity against staphylococcal infection.

**[0340]** To perturb the IgG Fc, vWF A1 and TNFR1 binding activities of Protein A, glutamine (Q) residues 9 and 10 [the numbering here is derived from that established for the SpA domain D] were modified generating lysine or glycine substitutions for both glutamines with the expectation that these substitutions abolish the ion bonds formed between wild-type



Protein A and its ligands. The added effect of the dual lysine substitutions may be that these positively charged residues institute a repellent charge for immunoglobulins. To perturb IgM Fab VH3 binding, the inventors selected the aspartate (D) residues 36 and 37 of SpA-D, each of which is required for the association of Protein A with the B cell receptor. D36 and D37 were both substituted with alanine. The Q9,10K and D36,37A mutations were combined in the recombinant molecule SpA-D<sub>Q9,10K;D36,37A</sub> and examined for the binding attributes of Protein A.

**[0341]** In brief, the Protein A (spa) genomic sequence of *Staphylococcus aureus* N315 was PCR amplified with the primers (GCTGCACATATGGCGCAACACGATGAAGCTCAAC [5' primer](SEQ ID NO:35) and AGTGATCCTTATGCTTTGTTAGCATCTGC [3' primer] (SEQ ID NO:36)), cloned into the pET15b vector (pYJS1, codons 48-486) (Stranger-Jones, et al., 2006) and recombinant plasmid transformed into *E. coli* BL21(DE3) (Studier et al., 1990). The Protein A product derived from pYJS1 harbors SpA residues 36-265 fused to the N-terminal His tag (MGSSHHHHHSSGLVPRGS (SEQ ID NO:37)). Following IPTG inducible expression, recombinant N-terminal His<sub>6</sub>-tagged SpA was purified by affinity chromatography on Ni-NTA resin (Stranger-Jones et al., 2006). The domain D of SpA (SpA-D) was PCR amplified with a pair of specific primers (AACATATGTTCAACAAAGATCAACAAAGC [5' primer](SEQ ID NO:38) and AAGGATCCAGATTCGTTAATTTTTAGC [3' primer] (SEQ ID NO:39)), sub-cloned into the pET15b vector (pHAN1, spa codons 212-261) and recombinant plasmid transformed into *E. coli* BL21(DE3) to express and purify recombinant N-terminal His<sub>6</sub>-tagged protein. To generate mutations in the SpA-D coding sequence, sets of two pairs of primers were synthesized (for D to A substitutions: CTTCAATCAAAGTCTTAAAGCCGC-CCCAAGCCAAAGCACTAAC [5' primer] (SEQ ID NO:40) and GTIAGTGCTTTGGCTTGGGGCGGCTT-TAAGACTTTGAATGAAG [3' primer] (SEQ ID NO:41); for Q to K substitutions CATATGTTCAACAAA-GATAAAAAAAGCGCCTTCTATGAAATC [5' primer] (SEQ ID NO:42) and GATTCATAGAAGGCGCTTTTTT-TATCTTTGTTGAACATATG [3' primer] (SEQ ID NO:43); for Q to G substitutions CATATGTTCAACAAAGATG-GAGGAAGCGCCTTCTATGAAATC [5' primer] (SEQ ID NO:44) and GATTCATAGAAGGCGCTTCTC-CATCTTTGTTGAACATATG [3' primer] (SEQ ID NO:45). Primers were used for quick-change mutagenesis protocols. Following mutagenesis, DNA sequences were confirmed for each of the recombinant proteins: SpA, SpA-D and SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10K;D36,37A</sub>. All proteins were purified from lysates of recombinant *E. coli* using Ni-NTA chromatography and subsequently dialyzed against PBS and stored at 4° C.

**[0342]** To measure binding of immunoglobulin to Protein A and its variants, 200 µg of purified protein was diluted into a 1 ml volume using column buffer (50 mM Tris-HCl, 150 mM NaCl, pH7.5) and then loaded onto a pre-equilibrated Ni-NTA column (1 ml bed volume). Columns were washed with 10 ml of column buffer. 200 µg of purified human IgG was diluted in a total volume of 1 ml column buffer and then applied to each of the columns charged with Protein A and its variants. The columns were subsequently washed with 5 ml wash buffer (10 mM imidazole in column buffer) and 5 ml column buffer. Protein samples were eluted with 2 ml elution buffer (500 mM imidazole in column buffer), fractions col-

lected and aliquots subjected to SDS-PAGE gel electrophoresis, followed by Coomassie-Blue staining. As shown in FIG. 3, wild-type Protein A (SpA) and its SpA-domain D both retained immunoglobulin during chromatography. In contrast, the SpA-D<sub>Q9,10K;D36,37A</sub> variant did not bind to immunoglobulin.

**[0343]** To quantify the binding of Protein A and its variants to the Fc portion of immunoglobulin and the VH3 domain of Fab, HRP conjugated human immunoglobulin G [hIgG], the Fc portion of human IgG [hFc] and the F(ab)<sub>2</sub> portion of human IgG [hF(ab)<sub>2</sub>] as well as ELISA assays were used to quantify the relative amount binding to Protein A and its variants. The data in FIG. 4 demonstrate the binding of SpA and SpA-D to hIgG and hFc, whereas SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10K;D36,37A</sub> displayed only background binding activities. SpA bound similar amounts of hFc and hF(ab)<sub>2</sub>, however the binding of SpA-D to hF(ab)<sub>2</sub> was reduced compared to full length SpA. This result suggests that the presence of multiple IgG binding domains may cooperatively increase the ability of Protein A to bind to the B cell receptor. When compared with the reduced binding power of SpA-D for hF(ab)<sub>2</sub>, of the two variants only SpA-D<sub>Q9,10K;D36,37A</sub> displayed a significant reduction in the ability to bind the VH3 domain of immunoglobulin. To examine the toxicogenic attributes of SpA-D and its variants, purified proteins were injected into mice, which were sacrificed after 4 hours to remove their spleens. Organ tissue was homogenized, capsular material removed and B cells stained with fluorescent CD19 antibodies. Following FACS analysis to quantify the abundance of B cells in splenic tissues, it was observed that SpA-D caused a 5% drop in the B cell count compared to a mock (PBS) control (FIG. 5). In contrast, SpA-D<sub>Q9,10K;D36,37A</sub> did not cause a reduction in B-cell counts, indicating that the mutant molecule had lost its toxicogenic attributes of stimulating B cell proliferation and death (FIG. 5). In summary, amino acid substitutions in the SpA-D residues Q9, Q10, D36, and D37 abolished the ability of Protein A domains to bind immunoglobulins or exert toxicogenic functions in human and animal tissues.

**[0344]** Non-Toxicogenic Protein A Variants Elicit Vaccine Protection.

**[0345]** To test whether or not Protein A and its variants can function as vaccine antigens, SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub>, and SpA-D<sub>Q9,10K;D36,37A</sub> were emulsified with complete or incomplete Freund's adjuvant and immunized 4 week old BALB/c mice on day 1 and day 11 with 50 µg of purified protein. Cohort of animals (n=5) were analyzed for humoral immune responses to immunization by bleeding the animals before (day 0) and after the immunization schedule (day 21). Table 5 indicates that immunized mice generated only a modest humoral immune response directed at wild-type Protein A or its SpA-D module, whereas the amount of antibody raised following immunization with SpA-D<sub>Q9,10K;D36,37A</sub> or SpA-D<sub>Q9,10K;D36,37A</sub> was increased four to five fold. Following intravenous challenge with 1×10<sup>7</sup> CFU *S. aureus* Newman, animals were killed on day 4, their kidneys removed and either analyzed for staphylococcal load (by plating tissue homogenate on agar plates and enumerating colony forming units, CFU) or histopathology. As expected, mock (PBS) immunized mice (n=19) harbored 6.46 log<sub>10</sub> (±0.25) CFU in kidney tissue and infectious lesions were organized into 3.7 (±1.2) abscesses per organ (n=10) (Table 5). Immunization of animals with SpA led to a 2.51 log<sub>10</sub> CFU reduction on day 5 (P=0.0003) with 2.1 (±1.2) abscesses per organ. The latter

data indicate that there was no significant reduction in abscess formation ( $P=0.35$ ). Immunization with SpA-D generated similar results: a  $2.03 \log_{10}$  CFU reduction on day 5 ( $P=0.0001$ ) with  $1.5 (\pm 0.8)$  abscesses per organ ( $P=0.15$ ). In contrast, immunization with SpA-D<sub>Q9,10K;D36,37A</sub> or SpA-D<sub>Q9,10G;D36,37A</sub> created increased protection, with  $3.07 \log_{10}$  and  $3.03 \log_{10}$  CFU reduction on day 4, respectively (statistical significance  $P<0.0001$  for both observations). Further, immunization with both SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10G;D36,37A</sub> generated significant protection from staphylococcal abscess formation, as only  $0.5 (\pm 0.4)$  and  $0.8 (\pm 0.5)$  infectious lesions per organ ( $P=0.02$  and  $P=0.04$ ) were identified. Thus, immunization with non-toxicogenic Protein A variants generates increased humoral immune responses for Protein A and provides protective immunity against staphylococcal challenge. These data indicate that Protein A is an ideal candidate for a human vaccine that prevents *S. aureus* disease.

**[0346]** These exciting results have several implications for the design of a human vaccine. First, the generation of substitution mutations that affect the ability of the immunoglobulin binding domains of Protein A, either alone or in combination of two or more domains, can generate non-toxicogenic variants suitable for vaccine development. It seems likely that a combination of mutant IgG binding domains closely resembling the structure of Protein A can generate even better humoral immune responses as is reported here for the SpA-domain D alone. Further, a likely attribute of Protein A specific antibodies may be that the interaction of antigen binding sites with the microbial surface can neutralize the ability of staphylococci to capture immunoglobulins via their Fc portion or to stimulate the B cell receptor via the VH3 binding activities.

**[0349]** Murine Abscess—

**[0350]** BALB/c mice (24-day-old female, 8-10 mice per group, Charles River Laboratories, Wilmington, Mass.) are immunized by intramuscular injection into the hind leg with purified protein (Chang et al., 2003; Schneewind et al., 1992). Purified SpA, SpA-D or SpA-DQ9,10K;D36,37A (50  $\mu$ g protein) is administered on days 0 (emulsified 1:1 with complete Freund's adjuvant) and 11 (emulsified 1:1 with incomplete Freund's adjuvant). Blood samples are drawn by retroorbital bleeding on days 0, 11, and 20. Sera are examined by ELISA for IgG titers for specific SpA-D and SpA-DQ9,10K;D36,37A binding activity. Immunized animals are challenged on day 21 by retroorbital injection of 100  $\mu$ l of *S. aureus* Newman or *S. aureus* USA300 suspension ( $1 \times 10^7$  cfu). For this, overnight cultures of *S. aureus* Newman are diluted 1:100 into fresh tryptic soy broth and grown for 3 h at 37° C. Staphylococci are centrifuged, washed twice, and diluted in PBS to yield an A<sub>600</sub> of 0.4 ( $1 \times 10^8$  cfu per ml). Dilutions are verified experimentally by agar plating and colony formation. Mice are anesthetized by intraperitoneal injection of 80-120 mg of ketamine and 3-6 mg of xylazine per kilogram of body weight and infected by retroorbital injection. On day 5 or 15 following challenge, mice are euthanized by compressed CO<sub>2</sub> inhalation. Kidneys are removed and homogenized in 1% Triton X-100. Aliquots are diluted and plated on agar medium for triplicate determination of cfu. For histology, kidney tissue is incubated at room temperature in 10% formalin for 24 h. Tissues are embedded in paraffin, thin-sectioned, stained with hematoxylin/eosin, and examined by microscopy.

**[0351]** Murine Lethal Infection—

**[0352]** BALB/c mice (24-day-old female, 8-10 mice per group, Charles River Laboratories, Wilmington, Mass.) are immunized by intramuscular injection into the hind leg with

TABLE 5

Antigen	Bacterial load in kidney (n = number of mice)			IgG titer	Abscess formation in mice (n = number of mice)				
	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> Reduction	<sup>c</sup> P value		<sup>d</sup> Surface abscess	Reduction	<sup>e</sup> Histopathology	Reduction	<sup>f</sup> p value
Mock	6.46 ± 0.25 (n = 19)	—	—	<100	14/19 (70%)	—	3.7 ± 1.2 (n = 10)	—	—
SpA	3.95 ± 0.56 (n = 20)	2.51	0.0003	1706 ± 370	10/20 (50%)	32%	2.1 ± 1.2 (n = 10)	2.2	0.35
SpA-D	4.43 ± 0.41 (n = 18)	2.03	0.0001	381 ± 27	10/18 (55%)	25%	1.5 ± 0.8 (n = 10)	2.2	0.15
SpA-D1	3.39 ± 0.50 (n = 19)	3.07	<0.0001	5600 ± 801	6/20 (30%)	59%	0.5 ± 0.4 (n = 10)	3.2	0.02
SpA-D2	3.43 ± 0.46 (n = 19)	3.03	<0.0001	3980 ± 676	6/19 (32%)	57%	0.8 ± 0.5 (n = 10)	2.9	0.04

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of 18 to 20 BALB/c mice. Standard error of the means ( $\pm$ SEM) is indicated.

<sup>c</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>b</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Abscess formation in kidney tissues four days following infection was measured by macroscopic inspection (% positive)

<sup>e</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys from ten animals; the number of abscesses per kidney was recorded and averaged for the final mean ( $\pm$ SEM).

<sup>f</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

SpA-D1 and SpA-D2 represent SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10G;D36,37A</sub> respectively.

**[0347]** Vaccine Protection in Murine Abscess, Murine Lethal Infection, and Murine Pneumonia Models.

**[0348]** Three animal models have been established for the study of *S. aureus* infectious disease. These models are used here to examine the level of protective immunity provided via the generation of Protein A specific antibodies.

purified SpA, SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> (50  $\mu$ g protein). Vaccine is administered on days 0 (emulsified 1:1 with complete Freund's adjuvant) and 11 (emulsified 1:1 with incomplete Freund's adjuvant). Blood samples are drawn by retroorbital bleeding on days 0, 11, and 20. Sera are examined by ELISA for IgG titers with specific SpA-D and SpA-D<sub>Q9,10K;D36,37A</sub> binding activity. Immunized animals are challenged

on day 21 by retroorbital injection of 100  $\mu$ l of *S. aureus* Newman or *S. aureus* USA300 suspension ( $15 \times 10^7$  cfu) (34). For this, overnight cultures of *S. aureus* Newman are diluted 1:100 into fresh tryptic soy broth and grown for 3 h at 37° C. Staphylococci are centrifuged, washed twice, diluted in PBS to yield an  $A_{600}$  of 0.4 ( $1 \times 10^8$  cfu per ml) and concentrated. Dilutions are verified experimentally by agar plating and colony formation. Mice are anesthetized by intraperitoneal injection of 80-120 mg of ketamine and 3-6 mg of xylazine per kilogram of body weight. Immunized animals are challenged on day 21 by intraperitoneal inject with  $2 \times 10^{10}$  cfu of *S. aureus* Newman or  $3-10 \times 10^9$  cfu of clinical *S. aureus* isolates. Animals are monitored for 14 days, and lethal disease is recorded.

#### [0353] Murine Pneumonia Model—

[0354] *S. aureus* strains Newman or USA300 (LAC) are grown at 37° C. in tryptic soy broth/agar to OD<sub>660</sub> 0.5. 50-ml culture aliquots are centrifuged, washed in PBS, and suspended in 750  $\mu$ l PBS for mortality studies ( $3-4 \times 10^8$  CFU per 30- $\mu$ l volume), or 1,250  $\mu$ l PBS ( $2 \times 10^8$  CFU per 30  $\mu$ l volume) for bacterial load and histopathology experiments (2, 3). For lung infection, 7-wk-old C57BL/6J mice (The Jackson Laboratory) are anesthetized before inoculation of 30  $\mu$ l of *S. aureus* suspension into the left nare. Animals are placed into the cage in a supine position for recovery and observed for 14 days. For active immunization, 4-wk-old mice receive 20  $\mu$ g SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> in CFA on day 0 via the i.m. route, followed by a boost with 20  $\mu$ g SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> in incomplete Freund's adjuvant (IFA) on day 10. Animals are challenged with *S. aureus* on day 21. Sera are collected before immunization and on day 20 to assess specific antibody production. For passive immunization studies, 7-wk-old mice receive 100  $\mu$ l of either NRS (normal rabbit serum) or SpA-D-specific rabbit antisera via i.p. injection 24 h before challenge. To assess the pathological correlates of pneumonia, infected animals are killed via forced CO<sub>2</sub> inhalation before removal of both lungs. The right lung is homogenized for enumeration of lung bacterial load. The left lung is placed in 1% formalin and paraffin embedded, thin sectioned, stained with hematoxylin-eosin, and analyzed by microscopy.

#### [0355] Rabbit Antibodies—

[0356] Purified 200  $\mu$ g SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> is used as an immunogen for the production of rabbit antisera. 200  $\mu$ g protein is emulsified with CFA for injection at day 0, followed by booster injections with 200  $\mu$ g protein emulsified with IFA on days 21 and 42. Rabbit antibody titers are determined by ELISA. Purified antibodies are obtained by affinity chromatography of rabbit serum on SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> sepharose. The concentration of eluted antibodies is measured by absorbance at A<sub>280</sub> and specific antibody titers are determined by ELISA.

#### [0357] Active Immunization with SpA-Domain D Variants.—

[0358] To determine vaccine efficacy, animals are actively immunized with purified SpA-D or SpAD<sub>Q9,10K;D36,37A</sub>. As a control, animals are immunized with adjuvant alone. Antibody titers against Protein A preparations are determined using SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> as antigens; note that the SpA-D<sub>Q9,10K;D36,37A</sub> variant cannot bind the Fc or Fab portion of IgG. Using infectious disease models described above, any reduction in bacterial load (murine abscess and pneumonia), histopathology evidence of staphylococcal dis-

ease (murine abscess and pneumonia) and protection from lethal disease (murine lethal challenge and pneumonia) is measured.

#### [0359] Passive Immunization with Affinity Purified Rabbit Polyclonal Antibodies Generated Against SpA-Domain D Variants.

[0360] To determine protective immunity of Protein A specific rabbit antibodies, mice are passively immunized with 5 mg/kg of purified SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> derived rabbit antibodies. Both of these antibody preparations are purified by affinity chromatography using immobilized SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub>. As a control, animals are passively immunized with rV10 antibodies (a plague protective antigen that has no impact on the outcome of staphylococcal infections). Antibody titers against all Protein A preparations are determined using SpA-D<sub>Q9,10K;D36,37A</sub> as an antigen, as this variant cannot bind the Fc or Fab portion of IgG. Using the infectious disease models described above, the reduction in bacterial load (murine abscess and pneumonia), histopathology evidence of staphylococcal disease (murine abscess and pneumonia), and the protection from lethal disease (murine lethal challenge and pneumonia) is measured.

### Example 2

#### Non-Toxigenic Protein A Vaccine for Methicillin-Resistant *Staphylococcus aureus* Infection

[0361] Clinical isolates of *S. aureus* express protein A (Shopsin et al., 1999, whose primary translational product is comprised of an N-terminal signal peptide (DeDent et al., 2008), five Ig-BDs (designated E, D, A, B and C) (Sjodahl, 1977), region X with variable repeats of an eight residue peptide (Guss et al., 1984), and C-terminal sorting signal for the cell wall anchoring of SpA (Schneewind et al., 1992; Schneewind et al., 1995) (FIG. 6). Guided by amino acid homology (Uhlen et al., 1984), the triple  $\alpha$ -helical bundle structure of IgBDs (Deisenhofer et al., 1978; Deisenhofer et al., 1981) and their atomic interactions with Fab V<sub>H</sub>3 (Graille et al., 2000) or Fc $\gamma$  (Gouda et al., 1998), glutamine 9 and 10 were selected as well as aspartate 36 and 37 as critical for the association of SpA with antibodies or B cell receptor, respectively. Substitutions Gln9Lys, Gln10Lys, Asp36Ala and Asp37Ala were introduced into the D domain to generate SpA-D<sub>KKAA</sub> (FIG. 6). The ability of isolated SpA-D or SpA-D<sub>KKAA</sub> to bind human IgG was analyzed by affinity chromatography (FIG. 6). Polyhistidine tagged SpA-D as well as full-length SpA retained human IgG on Ni-NTA, whereas SpA-D<sub>KKAA</sub> and a negative control (SrtA) did not (FIG. 6). A similar result was observed with von Willebrand factor (Hartleib et al., 2000), which, along with tumor necrosis factor receptor 1 (TNFR1) (Gomez et al., 2004), can also bind protein A via glutamine 9 and 10 (FIG. 6). Human immunoglobulin encompasses 60-70% V<sub>H</sub>3-type IgG. The inventors distinguish between Fc domain and B cell receptor activation of Igs and measured association of human Fc $\gamma$  and F(ab)<sub>2</sub> fragments, both of which bound to full-length SpA or SpA-D, but not to SpA-D<sub>KKAA</sub> (FIG. 6). Injection of SpA-D into the peritoneal cavity of mice resulted in B cell expansion followed by apoptotic collapse of CD19+ lymphocytes in spleen tissue of BALB/c mice (Goodyear and Silverman, 2003) (FIG. 6). B cell superantigen activity was not observed following injection with SpA-D<sub>KKAA</sub> and TUNEL-staining of

splenic tissue failed to detect the increase in apoptotic cells that follows injection of SpA or SpA-D (FIG. 6).

**[0362]** Naive six week old BALB/c mice were injected with 50 µg each of purified SpA, SpA-D or SpA-D<sub>KKAA</sub> emulsified in CFA and boosted with the same antigen emulsified in IFA. In agreement with the hypothesis that SpA-D promotes the apoptotic collapse of activated clonal B cell populations, the inventors observed a ten-fold higher titer of SpA-D<sub>KKAA</sub> specific antibodies following immunization of mice with the non-toxicogenic variant as compared to the B cell superantigen (SpA-D vs. SpA-D<sub>KKAA</sub> P<0.0001, Table 6). Antibody titers raised by immunization with full-length SpA were higher than those elicited by SpA-D (P=0.0022), which is likely due to the larger size and reiterative domain structure of this antigen (Table 6). Nevertheless, even SpA elicited lower antibody titers than SpA-D<sub>KKAA</sub> (P=0.0003), which encompasses only 50 amino acids of protein A (520 residues, SEQ ID NO:33). Immunized mice were challenged by intravenous inoculation with *S. aureus* Newman and the ability of staphylococci to seed abscesses in renal tissues was examined by necropsy four days after challenge. In homogenized renal tissue of mock (PBS/adjuvant) immunized mice, an average staphylococcal load of 6.46 log<sub>10</sub> CFU g<sup>-1</sup> was enumerated (Table 6). Immunization of mice with SpA or SpA-D led to a reduction in staphylococcal load, however SpA-D<sub>KKAA</sub> vaccinated animals displayed an even greater, 3.07 log<sub>10</sub> CFU g<sup>-1</sup> reduction of *S. aureus* Newman in renal tissues (P<0.0001, Table 6). Abscess formation in kidneys was analyzed by histopathology (FIG. 7). Mock immunized animals harbored an average of 3.7 (±1.2) abscesses per kidney (Table 6). Vaccination with SpA-D<sub>KKAA</sub> reduced the average number of abscesses to 0.5 (±0.4) (P=0.0204), whereas immunization with SpA or SpA-D did not cause a significant reduction in the number of abscess lesions (Table 6). Lesions from SpA-D<sub>KKAA</sub> vaccinated animals were smaller in size, with fewer infiltrating PMNs and characteristically lacked staphylococcal abscess communities (Cheng et al., 2009) (FIG. 7). Abscesses in animals that had been immunized with SpA or SpA-D displayed the same overall structure of lesions in mock immunized animals (FIG. 7).

**[0363]** The inventors examined whether SpA-D<sub>KKAA</sub> immunization can protect mice against MRSA strains and selected the USA300 LAC isolate for animal challenge (Diep et al., 2006). This highly virulent CA-MRSA strain spread rapidly throughout the United States, causing significant human morbidity and mortality (Kennedy et al., 2008). Compared to adjuvant control mice, SpA-D<sub>KKAA</sub> immunized animals harbored a 1.07 log<sub>10</sub> CFU g<sup>-1</sup> reduction in bacterial load of infected kidney tissues. Histopathology examination of renal tissue following *S. aureus* USA300 challenge

revealed that the average number of abscesses was reduced from 4.04 (±0.8) to 1.6 (±0.6) (P=0.02774). In contrast, SpA or SpA-D immunization did not cause a significant reduction in bacterial load or abscess formation (Table 6).

**[0364]** Rabbits were immunized with SpA-D<sub>KKAA</sub> and specific antibodies were purified on SpA-D<sub>KKAA</sub> affinity column followed by SDS-PAGE (FIG. 8). SpA-D<sub>KKAA</sub> specific IgG was cleaved with pepsin to generate Fcγ and F(ab)<sub>2</sub> fragments, the latter of which were purified by chromatography on SpA-D<sub>KKAA</sub> column (FIG. 8). Binding of human IgG or vWF to SpA or SpA-D was perturbed by SpA-D<sub>KKAA</sub> specific F(ab)<sub>2</sub>, indicating that SpA-D<sub>KKAA</sub> derived antibodies neutralize the B cell superantigen function of protein A as well as its interactions with Ig (FIG. 8).

**[0365]** To further improve the vaccine properties for non-toxicogenic protein A, the inventors generated SpA<sub>KKAA</sub> which includes all five IgBDs with four amino acid substitutions—substitutions corresponding to Gln9Lys, Gln10Lys, Asp36Ala and Asp37Ala of domain D—in each of its five domains (E, D, A, B and C). Polyhistidine tagged SpA<sub>KKAA</sub> was purified by affinity chromatography and analyzed by Coomassie Blue-stained SDS-PAGE (FIG. 9). Unlike full-length SpA, SpA<sub>KKAA</sub> did not bind human IgG, Fc and F(ab)<sub>2</sub> or vWF (FIG. 9). SpA<sub>KKAA</sub> failed to display B cell superantigen activity, as injection of the variant into BALB/c mice did not cause a depletion of CD19+ B cells in splenic tissue (FIG. 9). SpA<sub>KKAA</sub> vaccination generated higher specific antibody titers than SpA-D<sub>KKAA</sub> immunization and provided mice with elevated protection against *S. aureus* USA300 challenge (Table 6). Four days following challenge, SpA<sub>KKAA</sub> vaccinated animals harbored 3.54 log<sub>10</sub> CFU g<sup>-1</sup> fewer staphylococci in renal tissues (P=0.0001) and also caused a greater reduction in the number of abscess lesions (P=0.0109) (Table 6).

**[0366]** SpA<sub>KKAA</sub> was used to immunize rabbits. Rabbit antibodies specific for SpA-D<sub>KKAA</sub> or SpA<sub>KKAA</sub> were affinity purified on matrices with immobilized cognate antigen and injected at a concentration of 5 mg kg<sup>-1</sup> body weight into the peritoneal cavity of BALB/c mice (Table 7). Twenty-four hours later, specific antibody titers were determined in serum and animals challenged by intravenous inoculation with *S. aureus* Newman. Passive transfer reduced the staphylococcal load in kidney tissues for SpA-D<sub>KKAA</sub> (P=0.0016) or SpA<sub>KKAA</sub> (P=0.0005) specific antibodies. On histopathology examination, both antibodies reduced the abundance of abscess lesions in the kidneys of mice challenged with *S. aureus* Newman (Table 7). Together these data reveal that vaccine protection following immunization with SpA-D<sub>KKAA</sub> or SpA<sub>KKAA</sub> is conferred by antibodies that neutralize protein A.

TABLE 6

Immunization of mice with protein A vaccines.						
Staphylococcal load and abscess formation in renal tissue						
Antigen	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> P-value	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>d</sup> IgG Titer	<sup>e</sup> Number of abscesses	<sup>f</sup> P-value
<i>S. aureus</i> Newman challenge						
Mock	6.46 ± 0.25	—	—	<100	3.7 ± 1.2	—
SpA	3.95 ± 0.56	0.0003	2.51	1706 ± 370	2.1 ± 1.2	0.3581
SpA-D	4.43 ± 0.41	0.0001	2.03	381 ± 27	1.5 ± 0.8	0.1480
SpA-D <sub>KKAA</sub>	3.39 ± 0.50	<0.0001	3.07	5600 ± 801	0.5 ± 0.4	0.0204

TABLE 6-continued

Immunization of mice with protein A vaccines.						
Staphylococcal load and abscess formation in renal tissue						
Antigen	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> P-value	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>d</sup> IgG Titer	<sup>e</sup> Number of abscesses	<sup>f</sup> P-value
<i>S. aureus</i> USA300 (LAC) challenge						
Mock	7.20 ± 0.24	—	—	<100	4.0 ± 0.8	—
SpA	6.81 ± 0.26	0.2819	0.39	476 ± 60	3.3 ± 1.0	0.5969
SpA-D	6.34 ± 0.52	0.1249	0.86	358 ± 19	2.2 ± 0.6	0.0912
SpA-D <sub>KKAA</sub>	6.00 ± 0.42	0.0189	1.20	3710 ± 1147	1.6 ± 0.6	0.0277
SpA <sub>KKAA</sub>	3.66 ± 0.76	0.0001	3.54	10200 ± 2476	1.2 ± 0.5	0.0109

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of fifteen to twenty BALB/c mice per immunization. Representative of two independent and reproducible animal experiments is shown. Standard error of the means (±SEM) is indicated.

<sup>b</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Means of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA.

<sup>e</sup>Histopathology of hematoxyline-eosin stained, thin sectioned kidneys from ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

TABLE 7

Passive immunization of mice with antibodies against protein A.						
Staphylococcal load and abscess formation in renal tissue						
<sup>a</sup> Antibody	<sup>b</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>c</sup> P-value	<sup>d</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>e</sup> IgG Titer	<sup>f</sup> Number of abscesses	<sup>g</sup> P-value
Mock	7.10 ± 0.14	—	—	<100	4.5 ± 0.8	—
α-SpA-D <sub>KKAA</sub>	5.53 ± 0.43	0.0016	1.57	466 ± 114	1.9 ± 0.7	0.0235
α-SpA <sub>KKAA</sub>	5.69 ± 0.34	0.0005	1.41	1575 ± 152	1.6 ± 0.5	0.0062

<sup>a</sup>Affinity purified antibodies were injected into the peritoneal cavity of BALB/c mice at a concentration of 5 mg · kg<sup>-1</sup> twenty-four hours prior to intravenous challenge with 1 × 10<sup>7</sup> CFU *S. aureus* Newman.

<sup>b</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of fifteen BALB/c mice per immunization. Representative of two independent and reproducible animal experiments is shown. Standard error of the means (±SEM) is indicated.

<sup>c</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>d</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>e</sup>Means of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA.

<sup>f</sup>Histopathology of hematoxyline-eosin stained, thin sectioned kidneys from ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

**[0367]** Following infection with virulent *S. aureus*, mice do not develop protective immunity against subsequent infection with the same strain (Burts et al., 2008) (FIG. 10). The average abundance of SpA-D<sub>KKAA</sub> specific IgG in these animals was determined by dot blot as 0.20 μg ml<sup>-1</sup> (±0.04) and 0.14 μg ml<sup>-1</sup> (±0.01) for strains Newman and USA300 LAC, respectively (FIG. 9). The minimal concentration of protein A-specific IgG required for disease protection in SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub> vaccinated animals (P .05 log<sub>10</sub> reduction in staphylococcal CFU g<sup>-1</sup> renal tissue) was calculated as 4.05 μg ml<sup>-1</sup> (±0.88). Average serum concentration of SpA-specific IgG in adult healthy human volunteers (n=16) was 0.21 μg ml<sup>-1</sup> (±0.02). Thus, *S. aureus* infections in mice or humans are not associated with immune responses that raise significant levels of neutralizing antibodies directed against protein A, which is likely due to the B cell superantigen attributes of this molecule. In contrast, the average serum concentration of IgG specific for diphtheria toxin in human volunteers, 0.068 μg ml<sup>-1</sup> (±0.20), was within range for protective immunity against diphtheria (Behring, 1890; Lagergard et al., 1992).

**[0368]** Clinical *S. aureus* isolates express protein A, an essential virulence factor whose B cell superantigen activity and evasive attributes towards opsono-phagocytic clearance are absolutely required for staphylococcal abscess formation

(Palmqvist et al., 2005; Cheng et al., 2009; Silverman and Goodyear, 2006). Protein A can thus be thought of as a toxin, essential for pathogenesis, whose molecular attributes must be neutralized in order to achieve protective immunity. By generating non-toxigenic variants unable to bind Igs via Fcγ or VH<sub>3</sub>-Fab domains, the inventors measure here for the first time protein A neutralizing immune responses as a correlate for protective immunity against *S. aureus* infection. In contrast to many methicillin-sensitive strains, CA-MRSA isolate USA300 LAC is significantly more virulent (Cheng et al., 2009). For example, immunization of experimental animals with the surface protein IsdB (Kuklin et al., 2006; Stranger-Jones et al., 2006) raises antibodies that confer protection against *S. aureus* Newman (Stranger-Jones et al., 2009) but not against USA300 challenge.

**[0369]** The methods utilized include:

**[0370]** Bacterial Strains and Growth.

**[0371]** *Staphylococcus aureus* strains Newman and USA300 were grown in tryptic soy broth (TSB) at 37° C. *Escherichia coli* strains DH5α and BL21 (DE3) were grown in Luria-Bertani (LB) broth with 100 μg ml<sup>-1</sup> ampicillin at 37° C.

**[0372]** Rabbit Antibodies.

**[0373]** The coding sequence for SpA was PCR-amplified with two primers, gctgcacatattggcgcgaacacgatgaagctcaac (SEQ ID NO:35) and agtggatccttatgcttgagctttgttagcctgc (SEQ ID NO:36) using *S. aureus* Newman template DNA. SpA-D was PCR-amplified with two primers, aacatattgcaacaaagatcaacaaagc (SEQ ID NO:38) and aaggatccagatgctt-taatttttagc (SEQ ID NO:39). The sequence for SpA-D<sub>KKAA</sub> was mutagenized with two sets of primers catatgttcaacaaagataaaaaagcgccttctatgaaac (SEQ ID NO:42) and gatttcata-gaagcgcctttttatctttgttgaacatag (SEQ ID NO:43) for Q9K, Q10K as well as cttcattcaagctttaaagccgc-cccaagcacaagcactaac (SEQ ID NO:40) and gttagtgtcttgct-tggggcgcgttgaagcttgaatgaag (SEQ ID NO:41) for D36A, D37A. The sequence of SpA<sub>KKAA</sub> was synthesized by Integrated DNA Technologies, Inc. PCR products were cloned into pET-15b generating N-terminal His<sub>6</sub> tagged recombinant protein. Plasmids were transformed into BL21 (DE3). Overnight cultures of transformants were diluted 1:100 into fresh media and grown at 37° C. to an OD<sub>600</sub> 0.5, at which point cultures were induced with 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) and grown for an additional three hours. Bacterial cells were sedimented by centrifugation, suspended in column buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl) and disrupted with a French pressure cell at 14,000 psi. Lysates were cleared of membrane and insoluble components by ultracentrifugation at 40,000×g. Proteins in the soluble lysate were subjected to nickel-nitrilotriacetic acid (Ni-NTA, Qiagen) affinity chromatography. Proteins were eluted in column buffer containing successively higher concentrations of imidazole (100-500 mM). Protein concentrations were determined by bicinchonic acid (BCA) assay (Thermo Scientific). For antibody generation, rabbits (6 month old New-Zealand white, female, Charles River Laboratories) were immunized with 500 μg protein emulsified in Complete Freund's Adjuvant (Difco) by subscapular injection. For booster immunizations, proteins emulsified in Incomplete Freund's Adjuvant and injected 24 or 48 days following the initial immunization. On day 60, rabbits were bled and serum recovered.

**[0374]** Purified antigen (5 mg protein) was covalently linked to HiTrap NHS-activated HP columns (GE Healthcare). Antigen-matrix was used for affinity chromatography of 10-20 ml of rabbit serum at 4° C. Charged matrix was washed with 50 column volumes of PBS, antibodies eluted with elution buffer (1 M glycine, pH 2.5, 0.5 M NaCl) and immediately neutralized with 1M Tris-HCl, pH 8.5. Purified antibodies were dialyzed overnight against PBS at 4° C.

**[0375]** F(ab)<sub>2</sub> Fragments.

**[0376]** Affinity purified antibodies were mixed with 3 mg of pepsin at 37° C. for 30 minutes. The reaction was quenched with 1 M Tris-HCl, pH 8.5 and F(ab)<sub>2</sub> fragments were affinity purified with specific antigen-conjugated HiTrap NHS-activated HP columns. Purified antibodies were dialyzed overnight against PBS at 4° C., loaded onto SDS-PAGE gel and visualized with Coomassie Blue staining.

**[0377]** Active and Passive Immunization.

**[0378]** BALB/c mice (3 week old, female, Charles River Laboratories) were immunized with 50 μg protein emulsified in Complete Freund's Adjuvant (Difco) by intramuscular injection. For booster immunizations, proteins were emulsified in Incomplete Freund's Adjuvant and injected 11 days following the initial immunization. On day 20 following

immunization, 5 mice were bled to obtain sera for specific antibody titers by enzyme-linked immunosorbent assay (ELISA).

**[0379]** Affinity purified antibodies in PBS were injected at a concentration 5 mg kg<sup>-1</sup> of experimental animal weight into the peritoneal cavity of BALB/c mice (6 week old, female, Charles River Laboratories) 24 hours prior to challenge with *S. aureus*. Animal blood was collected via periorbital vein puncture. Blood cells were removed with heparinized micro-hematocrit capillary tubes (Fisher) and Z-gel serum separation micro tubes (Sarstedt) were used to collect and measure antigen specific antibody titers by ELISA.

**[0380]** Mouse Renal Abscess.

**[0381]** Overnight cultures of *S. aureus* Newman or USA300 (LAC) were diluted 1:100 into fresh TSB and grown for 2 hours at 37° C. Staphylococci were sedimented, washed and suspended PBS at OD<sub>600</sub> of 0.4 (~1×10<sup>8</sup> CFU ml<sup>-1</sup>). Inocula were quantified by spreading sample aliquots on TSA and enumerating colonies formed. BALB/c mice (6 week old, female, Charles River Laboratories) were anesthetized via intraperitoneal injection with 100 mg ml<sup>-1</sup> ketamine and 20 mg ml<sup>-1</sup> xylazine per kilogram of body weight. Mice were infected by retro-orbital injection with 1×10<sup>7</sup> CFU of *S. aureus* Newman or 5×10<sup>6</sup> CFU of *S. aureus* USA300. On day 4 following challenge, mice were killed by CO<sub>2</sub> inhalation. Both kidneys were removed, and the staphylococcal load in one organ was analyzed by homogenizing renal tissue with PBS, 1% Triton X-100. Serial dilutions of homogenate were spread on TSA and incubated for colony formation. The remaining organ was examined by histopathology. Briefly, kidneys were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with hematoxylin-eosin, and inspected by light microscopy to enumerate abscess lesions. All mouse experiments were performed in accordance with the institutional guidelines following experimental protocol review and approval by the Institutional Biosafety Committee (IBC) and the Institutional Animal Care and Use Committee (IACUC) at the University of Chicago.

**[0382]** Protein A Binding.

**[0383]** For human IgG binding, Ni-NTA affinity columns were pre-charged with 200 μg of purified proteins (SpA, SpA-D, SpA-D<sub>KKAA</sub>, and SrtA) in column buffer. After washing, 200 μg of human IgG (Sigma) was loaded onto the column. Protein samples were collected from washes and elutions and subjected to SDS-PAGE gel electrophoresis, followed by Coomassie Blue staining. Purified proteins (SpA, SpA<sub>KKAA</sub>, SpA-D and SpA-D<sub>KKAA</sub>) were coated onto MaxiSorp ELISA plates (NUNC) in 0.1M carbonate buffer (pH 9.5) at 1 μg ml<sup>-1</sup> concentration overnight at 4° C. Plates were next blocked with 5% whole milk followed by incubation with serial dilutions of peroxidase-conjugated human IgG, Fc or F(ab)<sub>2</sub> fragments for one hour. Plates were washed and developed using OptEIA ELISA reagents (BD). Reactions were quenched with 1 M phosphoric acid and A<sub>450</sub> readings were used to calculate half maximal titer and percent binding.

**[0384]** von Willebrand Factor (vWF) Binding Assays.

**[0385]** Purified proteins (SpA, SpA<sub>KKAA</sub>, SpA D and SpA-D<sub>KKAA</sub>) were coated and blocked as described above. Plates were incubated with human vWF at 1 μg ml<sup>-1</sup> concentration for two hours, then washed and blocked with human IgG for another hour. After washing, plates were incubated with serial dilution of peroxidase-conjugated antibody directed against

human vWF for one hour. Plates were washed and developed using OptEIA ELISA reagents (BD). Reactions were quenched with 1 M phosphoric acid and  $A_{450}$  readings were used to calculate half maximal titer and percent binding. For inhibition assays, plates were incubated with affinity purified F(ab)<sub>2</sub> fragments specific for SpA-D<sub>KKAA</sub> at 10  $\mu\text{ml}^{-1}$  concentration for one hour prior to ligand binding assays.

**[0386]** Splenocyte Apoptosis.

**[0387]** Affinity purified proteins (150  $\mu\text{g}$  of SpA, SpA-D, SpA<sub>KKAA</sub>, and SpA-D<sub>KKAA</sub>) were injected into the peritoneal cavity of BALB/c mice (6 week old, female, Charles River Laboratories). Four hours following injection, animals were killed by CO<sub>2</sub> inhalation. Their spleens were removed and homogenized. Cell debris were removed using cell strainer and suspended cells were transferred to ACK lysis buffer (0.15 M NH<sub>4</sub>Cl, 10 mM KHCO<sub>3</sub>, 0.1 mM EDTA) to lyse red blood cells. White blood cells were sedimented by centrifugation, suspended in PBS and stained with 1:250 diluted R-PE conjugated anti-CD19 monoclonal antibody (Invitrogen) on ice and in the dark for one hour. Cells were washed with 1% FBS and fixed with 4% formalin overnight at 4° C. The following day, cells were diluted in PBS and analyzed by flow cytometry. The remaining organ was examined for histopathology. Briefly, spleens were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with the Apoptosis detection kit (Millipore), and inspected by light microscopy.

**[0388]** Antibody Quantification.

**[0389]** Sera were collected from healthy human volunteers or BALB/c mice that had been either infected with *S. aureus* Newman or USA300 for 30 days or that had been immunized with SpA-D<sub>KKAA</sub>/SpA<sub>KKAA</sub> as described above. Human/mouse IgG (Jackson Immunology Laboratory), SpA<sub>KKAA</sub>, and CRM<sub>197</sub> were blotted onto nitrocellulose membrane. Membranes were blocked with 5% whole milk, followed by incubation with either human or mouse sera. IRDye 700DX conjugated affinity purified anti-human/mouse IgG (Rockland) was used to quantify signal intensities using the Odyssey™ infrared imaging system (Li-cor). Experiments with blood from human volunteers involved protocols that were reviewed, approved and performed under regulatory supervision of The University of Chicago's Institutional Review Board (IRB).

**[0390]** Statistical Analysis.

**[0391]** Two tailed Student's t tests were performed to analyze the statistical significance of renal abscess, ELISA, and B cell superantigen data.

#### REFERENCES

**[0392]** The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

**[0393]** U.S. Pat. No. 3,791,932

**[0394]** U.S. Pat. No. 3,949,064

**[0395]** U.S. Pat. No. 4,174,384

**[0396]** U.S. Pat. No. 4,338,298

**[0397]** U.S. Pat. No. 4,356,170

**[0398]** U.S. Pat. No. 4,367,110

**[0399]** U.S. Pat. No. 4,372,945

**[0400]** U.S. Pat. No. 4,452,901

**[0401]** U.S. Pat. No. 4,474,757

**[0402]** U.S. Pat. No. 4,554,101

**[0403]** U.S. Pat. No. 4,578,770

**[0404]** U.S. Pat. No. 4,596,792

**[0405]** U.S. Pat. No. 4,599,230

**[0406]** U.S. Pat. No. 4,599,231

**[0407]** U.S. Pat. No. 4,601,903

**[0408]** U.S. Pat. No. 4,608,251

**[0409]** U.S. Pat. No. 4,683,195

**[0410]** U.S. Pat. No. 4,683,202

**[0411]** U.S. Pat. No. 4,684,611

**[0412]** U.S. Pat. No. 4,690,915

**[0413]** U.S. Pat. No. 4,690,915

**[0414]** U.S. Pat. No. 4,748,018

**[0415]** U.S. Pat. No. 4,800,159

**[0416]** U.S. Pat. No. 4,879,236

**[0417]** U.S. Pat. No. 4,952,500

**[0418]** U.S. Pat. No. 5,084,269

**[0419]** U.S. Pat. No. 5,199,942

**[0420]** U.S. Pat. No. 5,221,605

**[0421]** U.S. Pat. No. 5,238,808

**[0422]** U.S. Pat. No. 5,302,523

**[0423]** U.S. Pat. No. 5,310,687

**[0424]** U.S. Pat. No. 5,322,783

**[0425]** U.S. Pat. No. 5,384,253

**[0426]** U.S. Pat. No. 5,464,765

**[0427]** U.S. Pat. No. 5,512,282

**[0428]** U.S. Pat. No. 5,512,282

**[0429]** U.S. Pat. No. 5,538,877

**[0430]** U.S. Pat. No. 5,538,880

**[0431]** U.S. Pat. No. 5,548,066

**[0432]** U.S. Pat. No. 5,550,318

**[0433]** U.S. Pat. No. 5,563,055

**[0434]** U.S. Pat. No. 5,580,859

**[0435]** U.S. Pat. No. 5,589,466

**[0436]** U.S. Pat. No. 5,591,616

**[0437]** U.S. Pat. No. 5,610,042

**[0438]** U.S. Pat. No. 5,620,896

**[0439]** U.S. Pat. No. 5,648,240

**[0440]** U.S. Pat. No. 5,656,610

**[0441]** U.S. Pat. No. 5,702,932

**[0442]** U.S. Pat. No. 5,736,524

**[0443]** U.S. Pat. No. 5,780,448

**[0444]** U.S. Pat. No. 5,789,215

**[0445]** U.S. Pat. No. 5,801,234

**[0446]** U.S. Pat. No. 5,840,846

**[0447]** U.S. Pat. No. 5,843,650

**[0448]** U.S. Pat. No. 5,846,709

**[0449]** U.S. Pat. No. 5,846,783

**[0450]** U.S. Pat. No. 5,849,497

**[0451]** U.S. Pat. No. 5,849,546

**[0452]** U.S. Pat. No. 5,849,547

**[0453]** U.S. Pat. No. 5,858,652

**[0454]** U.S. Pat. No. 5,866,366

**[0455]** U.S. Pat. No. 5,871,986

**[0456]** U.S. Pat. No. 5,916,776

**[0457]** U.S. Pat. No. 5,922,574

**[0458]** U.S. Pat. No. 5,925,565

**[0459]** U.S. Pat. No. 5,925,565

**[0460]** U.S. Pat. No. 5,928,905

**[0461]** U.S. Pat. No. 5,928,906

**[0462]** U.S. Pat. No. 5,932,451

**[0463]** U.S. Pat. No. 5,935,819

**[0464]** U.S. Pat. No. 5,935,825

**[0465]** U.S. Pat. No. 5,939,291

**[0466]** U.S. Pat. No. 5,942,391

**[0467]** U.S. Pat. No. 5,945,100

- [0468] U.S. Pat. No. 5,958,895  
 [0469] U.S. Pat. No. 5,981,274  
 [0470] U.S. Pat. No. 5,994,624  
 [0471] U.S. Pat. No. 6,008,341  
 [0472] U.S. Pat. No. 6,288,214  
 [0473] U.S. Pat. No. 6,294,177  
 [0474] U.S. Pat. No. 6,651,655  
 [0475] U.S. Pat. No. 6,656,462  
 [0476] U.S. Pat. No. 6,733,754  
 [0477] U.S. Pat. No. 6,756,361  
 [0478] U.S. Pat. No. 6,770,278  
 [0479] U.S. Pat. No. 6,793,923  
 [0480] U.S. Pat. No. 6,814,971  
 [0481] U.S. Pat. No. 6,936,258  
 [0482] U.S. Patent Appln. 2002/0169288  
 [0483] U.S. Patent Appln. 2003/0153022  
 [0484] Abdallah et al., *Mol. Microbiol.*, 62, 667-679, 2006.  
 [0485] Abdallah et al., *Nat. Rev. Microbiol.*, 5, 883-891, 2007.  
 [0486] Albus et al., *Infect. Immun.*, 59:1008-1014, 1991.  
 [0487] An, *J. Virol.*, 71(3):2292-302, 1997.  
 [0488] Anavi, Sc. thesis from the department of Molecular Microbiology and Biotechnology of the Tel-Aviv University, Israel, 1998.  
 [0489] Andersen et al., *J. Immunol.*, 154, 3359-3372, 1995.  
 [0490] Angel et al., *Cell*, 49:729, 1987b.  
 [0491] Angel et al., *Mol. Cell. Biol.*, 7:2256, 1987a.  
 [0492] Archer, *Clin. Infect. Dis.*, 26, 1179-1181, 1998.  
 [0493] Atchison and Perry, *Cell*, 46:253, 1986.  
 [0494] Atchison and Perry, *Cell*, 48:121, 1987.  
 [0495] Ausubel et al., *In: Current Protocols in Molecular Biology*, John, Wiley & Sons, Inc, New York, 1996.  
 [0496] Baba et al., *J. Bacteriol.* 190:300-310, 2007.  
 [0497] Bae and Schneewind, *Plasmid*, 55:58-63, 2006.  
 [0498] Bae et al., *Proc. Natl. Acad. Sci. USA*, 101, 12312-12317, 2004.  
 [0499] Banerji et al., *Cell*, 27(2 Pt 1):299-308, 1981.  
 [0500] Banerji et al., *Cell*, 33(3):729-740, 1983.  
 [0501] Barany and Merrifield, *In: The Peptides*, Gross and Meienhofer (Eds.), Academic Press, NY, 1-284, 1979.  
 [0502] Behring E A. Über das Zustandekommen der Diphtherie—Immunität bei Thieren. Deutsche Medizinische Wochenschrift, 16:1145-8, 1890.  
 [0503] Bellus, *J. Macromol. Sci. Pure Appl. Chem.*, A31(1): 1355-1376, 1994.  
 [0504] Berkhout et al., *Cell*, 59:273-282, 1989.  
 [0505] Birch-Hirschfeld, L. 1934. Über die Agglutination von Staphylokokken durch Bestandteile des Säugetierblutplasmas. Klinische Wochenschrift 13:331.  
 [0506] Bjerketorp et al., *FEMS Microbiol. Lett.*, 234:309-314, 2004.  
 [0507] Blonar et al., *EMBO J.*, 8:1139, 1989.  
 [0508] Bodine and Ley, *EMBO J.*, 6:2997, 1987.  
 [0509] Borrebaeck, *In: Antibody Engineering—A Practical Guide*, W. H. Freeman and Co., 1992.  
 [0510] Boshart et al., *Cell*, 41:521, 1985.  
 [0511] Bosze et al., *EMBO* 1, 5(7):1615-1623, 1986.  
 [0512] Boucher and Corey. *Clin. Infect. Dis.* 46:S334-S349, 2008.  
 [0513] Braddock et al., *Cell*, 58:269, 1989.  
 [0514] Brown et al., *Biochemistry*, 37:4397-4406, 1998.  
 [0515] Bubeck Wardenburg and Schneewind. *J. Exp. Med.* 205:287-294, 2008.  
 [0516] Bubeck-Wardenburg et al., *Infect. Immun.* 74:1040-1044, 2007.  
 [0517] Bubeck-Wardenburg et al., *Proc. Natl. Acad. Sci. USA*, 103:13831-13836, 2006.  
 [0518] Bulla and Siddiqui, *J. Virol.*, 62:1437, 1986.  
 [0519] Burke et al., *J. Inf. Dis.*, 170:1110-1119, 1994.  
 [0520] Burlak et al., *Cell Microbiol.*, 9:1172-1190, 2007.  
 [0521] Burts and Missiakas, *Mol. Microbiol.*, 69:736-46, 2008.  
 [0522] Burts et al., *Proc. Natl. Acad. Sci. USA*, 102:1169-1174, 2005.  
 [0523] Campbell and Villarreal, *Mol. Cell. Biol.*, 8:1993, 1988.  
 [0524] Campere and Tilghman, *Genes and Dev.*, 3:537, 1989.  
 [0525] Campo et al., *Nature*, 303:77, 1983.  
 [0526] Carbonelli et al., *FEMS Microbiol. Lett.*, 177(1):75-82, 1999.  
 [0527] Cedergren et al., *Protein Eng.*, 6:441-448, 1993.  
 [0528] Celander and Haseltine, *J. Virology*, 61:269, 1987.  
 [0529] Celander et al., *J. Virology*, 62:1314, 1988.  
 [0530] Cespedes et al., *J. Infect. Dis.*, 191(3):444-52, 2005.  
 [0531] Champion et al., *Science*, 313:1632-1636, 2006.  
 [0532] Chandler et al., *Cell*, 33:489, 1983.  
 [0533] Chandler et al., *Proc. Natl. Acad. Sci. USA*, 94(8): 3596-601, 1997.  
 [0534] Chang et al., *Lancet.*, 362(9381):362-369, 2003.  
 [0535] Chang et al., *Mol. Cell. Biol.*, 9:2153, 1989.  
 [0536] Chatterjee et al., *Proc. Natl. Acad. Sci. USA*, 86:9114, 1989.  
 [0537] Chen and Okayama, *Mol. Cell Biol.*, 7(8):2745-2752, 1987.  
 [0538] Cheng et al., *FASEB J.*, 23:1-12, 2009.  
 [0539] Choi et al., *Cell*, 53:519, 1988.  
 [0540] Cocea, *Biotechniques*, 23(5):814-816, 1997.  
 [0541] Cohen et al., *J. Cell. Physiol.*, 5:75, 1987.  
 [0542] Cosgrove et al., *Infect. Control Hosp. Epidemiol.* 26:166-174, 2005.  
 [0543] Costa et al., *Mol. Cell. Biol.*, 8:81, 1988.  
 [0544] Cripe et al., *EMBO J.*, 6:3745, 1987.  
 [0545] Culotta and Hamer, *Mol. Cell. Biol.*, 9:1376, 1989.  
 [0546] Dalbey and Widmer, *J. Biol. Chem.*, 260:15925-15931, 1985.  
 [0547] Dandolo et al., *J. Virology*, 47:55-64, 1983.  
 [0548] De Villiers et al., *Nature*, 312(5991):242-246, 1984.  
 [0549] DeBord et al., *Infect. Immun.*, 74:4910-4914, 2006.  
 [0550] DeDent et al., *EMBO J.* 27:2656-2668, 2008.  
 [0551] DeDent et al., *J. Bacteriol.* 189:4473-4484, 2007.  
 [0552] Deisenhofer et al., *Hoppe-Seyh Zeitsch. Physiol. Chem.* 359:975-985, 1978.  
 [0553] Deisenhofer, *Biochemistry* 20:2361-2370, 1981.  
 [0554] Deschamps et al., *Science*, 230:1174-1177, 1985.  
 [0555] Devereux et al., *Nucl. Acid Res.*, 12:387-395, 1984.  
 [0556] Diep et al., *J. Infect. Dis.*, 193:1495-1503, 2006a.  
 [0557] Diep et al., *Lancet.*, 367:731-739, 2006b.  
 [0558] Dinges et al., *Clin. Microbiol. Rev.*, 13:16-34, 2000.  
 [0559] Duthie and Lorenz, *J. Gen. Microbiol.*, 6:95-107, 1952.  
 [0560] Edbrooke et al., *Mol. Cell. Biol.*, 9:1908, 1989.  
 [0561] Edlund et al., *Science*, 230:912-916, 1985.  
 [0562] Ekstedt and Yotis, *Ann. N.Y. Acad. Sci.*, 80:496-500, 1960.  
 [0563] Emori and Gaynes, *Clin. Microbiol. Rev.*, 6:428-442, 1993.



- [0564] EP 0786519  
 [0565] EP 497524  
 [0566] EP 497525  
 [0567] Epitope Mapping Protocols In: *Methods in Molecular Biology*, Vol. 66, Morris (Ed.), 1996.  
 [0568] Fehchheimer, et al., *Proc Natl. Acad. Sci. USA*, 84:8463-8467, 1987.  
 [0569] Feng and Holland, *Nature*, 334:6178, 1988.  
 [0570] Field and Smith, *J. Comp. Pathol.*, 55:63, 1945.  
 [0571] Firak and Subramanian, *Mol. Cell. Biol.*, 6:3667, 1986.  
 [0572] Foecking and Hofstetter, *Gene*, 45(1):101-105, 1986.  
 [0573] Fortune et al., *Proc Natl. Acad. Sci. USA*, 102:10676-10681, 2005.  
 [0574] Foster, *Nat. Rev. Microbiol.*, 3:948-958, 2005.  
 [0575] Fournier et al., *Infect. Immun.*, 45:87-93, 1984.  
 [0576] Fraley et al., *Proc. Natl. Acad. Sci. USA*, 76:3348-3352, 1979.  
 [0577] Friedrich et al., *Nature*, 425:535-539, 2003.  
 [0578] Fujita et al., *Cell*, 49:357, 1987.  
 [0579] GB Appln. 2 202 328  
 [0580] Gilles et al., *Cell*, 33:717, 1983.  
 [0581] Gloss et al., *EMBO J.*, 6:3735, 1987.  
 [0582] Godbout et al., *Mol. Cell. Biol.*, 8:1169, 1988.  
 [0583] Gomez et al., *EMBO J.* 26:701-709, 2007.  
 [0584] Gomez et al., *J. Biol. Chem.* 281:20190-20196, 2006.  
 [0585] Gomez et al., *Nature Med.* 10:842-8, 2004.  
 [0586] Goodbourn and Maniatis, *Proc. Natl. Acad. Sci. USA*, 85:1447, 1988.  
 [0587] Goodbourn et al., *Cell*, 45:601, 1986.  
 [0588] Goodyear and Silverman, *J. Exp. Med.*, 197:1125-1139, 2003.  
 [0589] Goodyear and Silverman, *Proc. Nat. Acad. Sci. USA*, 101:11392-11397, 2004.  
 [0590] Gopal, *Mol. Cell Biol.*, 5:1188-1190, 1985.  
 [0591] Gouda et al., *Biochemistry*, 31(40):9665-72, 1992.  
 [0592] Gouda et al., *Biochemistry*, 37:129-36, 1998.  
 [0593] Graham and Van Der Eb, *Virology*, 52:456-467, 1973.  
 [0594] Graille et al., *Proc. Nat. Acad. Sci. USA* 97:5399-5404, 2000.  
 [0595] Greene et al., *Immunology Today*, 10:272, 1989  
 [0596] Grosschedl and Baltimore, *Cell*, 41:885, 1985.  
 [0597] Guinn et al., *Mol. Microbiol.*, 51:359-370, 2004.  
 [0598] Guss et al., *Eur. J. Biochem.* 138:413-420, 1984.  
 [0599] Harland and Weintraub, *J. Cell Biol.*, 101(3):1094-1099, 1985.  
 [0600] Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., Chapter 8, 1988.  
 [0601] Hartleib et al., *Blood* 96:2149-2156, 2000.  
 [0602] Harvey et al., *Proc. Natl. Acad. Sci. USA*, 83:1084-1088, 1986.  
 [0603] Haslinger and Karin, *Proc. Natl. Acad. Sci. USA*, 82:8572, 1985.  
 [0604] Hauber and Cullen, *J Virology*, 62:673, 1988.  
 [0605] Hen et al., *Nature*, 321:249, 1986.  
 [0606] Hensel et al., *Lymphokine Res.*, 8:347, 1989.  
 [0607] Herr and Clarke, *Cell*, 45:461, 1986.  
 [0608] Hirochika et al., *J. Virol.*, 61:2599, 1987.  
 [0609] Hirsch et al., *Mol. Cell. Biol.*, 10:1959, 1990.  
 [0610] Holbrook et al., *Virology*, 157:211, 1987.  
 [0611] Horlick and Benfield, *Mol. Cell. Biol.*, 9:2396, 1989.  
 [0612] Hsu et al., *Proc. Natl. Acad. Sci. USA*, 100:12420-12425, 2003.  
 [0613] Huang et al., *Cell*, 27:245, 1981.  
 [0614] Hug et al., *Mol. Cell. Biol.*, 8:3065, 1988.  
 [0615] Huston et al., In: *Methods in Enzymology*, Langone (Ed.), Academic Press, NY, 203:46-88, 1991.  
 [0616] Hwang et al., *Mol. Cell. Biol.*, 10:585, 1990.  
 [0617] Imagawa et al., *Cell*, 51:251, 1987.  
 [0618] Imbra and Karin, *Nature*, 323:555, 1986.  
 [0619] Imler et al., *Mol. Cell. Biol.*, 7:2558, 1987.  
 [0620] Imperiale and Nevins, *Mol. Cell. Biol.*, 4:875, 1984.  
 [0621] Innis et al., *Proc Natl Acad Sci USA*, 85(24):9436-9440, 1988.  
 [0622] Inouye and Inouye, *Nucleic Acids Res.*, 13: 3101-3109, 1985.  
 [0623] Jakobovits et al., *Mol. Cell. Biol.*, 8:2555, 1988.  
 [0624] Jameel and Siddiqui, *Mol. Cell. Biol.*, 6:710, 1986.  
 [0625] Jansson et al., *FEMS Immunol. Med. Microbiol.* 20:69-78 1998.  
 [0626] Jaynes et al., *Mol. Cell. Biol.*, 8:62, 1988.  
 [0627] Jensen, *Acta Path. Microbiol. Scand.* 44:421-428, 1958.  
 [0628] Johnson et al., *Methods in Enzymol.*, 203:88-99, 1991.  
 [0629] Johnson et al., *Mol. Cell. Biol.*, 9:3393, 1989.  
 [0630] Jones, *Carb. Research*, 340:1097-1106, 2005.  
 [0631] Jonsson et al., *Oral Dis.*, 8(3):130-140, 2002.  
 [0632] Joyce et al., *Carbohydrate Research* 338:903-922 (2003)  
 [0633] Kadesch and Berg, *Mol. Cell. Biol.*, 6:2593, 1986.  
 [0634] Kaeppler et al., *Plant Cell Rep.*, 8:415-418, 1990.  
 [0635] Kaneda et al., *Science*, 243:375-378, 1989.  
 [0636] Karin et al., *Mol. Cell. Biol.*, 7:606, 1987.  
 [0637] Katinka et al., *Cell*, 20:393, 1980.  
 [0638] Kato et al., *J. Biol. Chem.*, 266:3361-3364, 1991.  
 [0639] Kawamoto et al., *Mol. Cell. Biol.*, 8:267, 1988.  
 [0640] Kennedy et al., *Proc. Natl. Acad. Sci. USA* 105:1327-1332, 2008.  
 [0641] Kiledjian et al., *Mol. Cell. Biol.*, 8:145, 1988.  
 [0642] Kinoshita, M., N. Kobayashi, S. Nagashima, M. Ishino, S. Otokozawa, K. Mise, A. Sumi, H. Tsutsumi, N. Uehara, N. Watanabe, and M. Endo. 2008. Diversity of staphylocoagulase and identification of novel variants of staphylocoagulase gene in *Staphylococcus aureus*. *Microbiol. Immunol.* 52:334-348.  
 [0643] Klamut et al., *Mol. Cell. Biol.*, 10:193, 1990.  
 [0644] Klevens et al., *Clin. Infect. Dis.*, 2008; 47:927-30, 2008.  
 [0645] Klevens et al., *JAMA*, 298:1763-1771, 2007.  
 [0646] Koch et al., *Mol. Cell. Biol.*, 9:303, 1989.  
 [0647] Kohler and Milstein, *Nature* 256:495-497 (1975)  
 [0648] Kriegler and Botchan, In: *Eukaryotic Viral Vectors*, Gluzman (Ed.), Cold Spring Harbor: Cold Spring Harbor Laboratory, NY, 1982.  
 [0649] Kriegler and Botchan, *Mol. Cell. Biol.*, 3:325, 1983.  
 [0650] Kriegler et al., *Cell*, 38:483, 1984a.  
 [0651] Kriegler et al., *Cell*, 53:45, 1988.  
 [0652] Kriegler et al., In: *Cancer Cells 2/Oncogenes and Viral Genes*, Van de Woude et al. eds, Cold Spring Harbor, Cold Spring Harbor Laboratory, 1984b.  
 [0653] Kroh et al., *Proc. Natl. Acad. Sci. USA*, 106:7786-7791, 2009.

- [0654] Kuhl et al., *Cell*, 50:1057, 1987.
- [0655] Kuklin et al., *Infect. Immun.*, 74:2215-23, 2006.
- [0656] Kunz et al., *Nucl. Acids Res.*, 17:1121, 1989.
- [0657] Kuroda et al., *Lancet*, 357:1225-1240, 2001.
- [0658] Kyte and Doolittle, *J. Mol. Biol.*, 157(1):105-132, 1982.
- [0659] Lagergard et al., *Eur. J. Clin. Microbiol. Infect. Dis.*, 11:341-5, 1992.
- [0660] Lam et al., *J. Bacteriol.*, 86:87-91, 1963.
- [0661] Larsen et al., *Proc Natl. Acad. Sci. USA*, 83:8283, 1986, 1963.
- [0662] Laspia et al., *Cell*, 59:283, 1989.
- [0663] Latimer et al., *Mol. Cell. Biol.*, 10:760, 1990.
- [0664] Lee et al., *Nature*, 294:228, 1981.
- [0665] Lee et al., *Nucleic Acids Res.*, 12:4191-206, 1984.
- [0666] Lee, *Trends Microbiol.* 4(4):162-166, 1996.
- [0667] Levenson et al., *Hum. Gene Ther.*, 9(8):1233-1236, 1998.
- [0668] Levinson et al., *Nature*, 295:79, 1982.
- [0669] Lin et al., *Mol. Cell. Biol.*, 10:850, 1990.
- [0670] Lowy, *New Engl. J. Med.*, 339:520-532, 1998.
- [0671] Luria et al., *EMBO J.*, 6:3307, 1987.
- [0672] Lusky and Botchan, *Proc. Natl. Acad. Sci. USA*, 83:3609, 1986.
- [0673] Lusky et al., *Mol. Cell. Biol.*, 3:1108, 1983.
- [0674] Macejak and Sarnow, *Nature*, 353:90-94, 1991.
- [0675] MacGurn et al., *Mol. Microbiol.*, 57:1653-1663, 2005.
- [0676] Maira-Litran et al., *Infect. Immun.*, 70:4433-4440, 2002.
- [0677] Maira-Litran et al., *Vaccine*, 22:872-879, 2004.
- [0678] Majors and Varmus, *Proc. Natl. Acad. Sci. USA*, 80:5866, 1983.
- [0679] Markwardt, *Untersuchungen über Hirudin. Naturwissenschaften*, 41:537-538, 1955.
- [0680] Mazmanian et al., *Mol. Microbiol.* 40, 1049-1057, 2001.
- [0681] Mazmanian et al., *Mol. Microbiol.*, 40(5):1049-1057, 2001.
- [0682] Mazmanian et al., *Proc. Natl. Acad. Sci. USA*, 97:5510-5515, 2000.
- [0683] Mazmanian et al., *Science*, 285(5428):760-3, 1999.
- [0684] McLaughlin et al., *PLoS Pathog.*, 3:e105, 2007.
- [0685] McNeall et al., *Gene*, 76:81, 1989.
- [0686] Mernaugh et al., In: *Molecular Methods in Plant Pathology*, Singh et al. (Eds.), CRC Press Inc., Boca Raton, Fla., 359-365, 1995.
- [0687] Merrifield, *Science*, 232(4748):341-347, 1986.
- [0688] Miksicek et al., *Cell*, 46:203, 1986.
- [0689] Mordacq and Linzer, *Genes and Dev.*, 3:760, 1989.
- [0690] Moreau et al., *Carbohydrate Res.*, 201:285-297, 1990.
- [0691] Moreau et al., *Nucl. Acids Res.*, 9:6047, 1981.
- [0692] Moreillon et al., *Infect. Immun.*, 63:4738-4743, 1995.
- [0693] Moreillon et al., *Infect. Immun.*, 63:4738-4743, 1995.
- [0694] Mosmann and Coffman, *Ann. Rev. Immunol.*, 7:145-173, 1989.
- [0695] Muesing et al., *Cell*, 48:691, 1987.
- [0696] Musher et al., *Medicine (Baltimore)*, 73:186-208, 1994.
- [0697] Navarre and Schneewind, *J. Biol. Chem.*, 274:15847-15856, 1999.
- [0698] Needleman & Wunsch, *J. Mol. Biol.*, 48:443, 1970.
- [0699] Ng et al., *Nuc. Acids Res.*, 17:601, 1989.
- [0700] Nicolau and Sene, *Biochim. Biophys. Acta*, 721:185-190, 1982.
- [0701] Nicolau et al., *Methods Enzymol.*, 149:157-176, 1987.
- [0702] Novick, *Mol. Microbiol.*, 48:1429-1449, 2003.
- [0703] O'Brien et al., *Mol. Microbiol.* 44:1033-1044, 2002.
- [0704] O'Seaghdha et al., *FEBS J.* 273:4831-4841, 2006.
- [0705] Omirulleh et al., *Plant Mol. Biol.*, 21(3):415-28, 1993.
- [0706] Ondek et al., *EMBO J.*, 6:1017, 1987.
- [0707] Ornitz et al., *Mol. Cell. Biol.*, 7:3466, 1987.
- [0708] Pallen, *Trends Microbiol.*, 10:209-212, 2002.
- [0709] Palmiter et al., *Nature*, 300:611, 1982.
- [0710] Palmqvist et al., *Microbes. Infect.*, 7:1501-11, 2005.
- [0711] Panizzi et al., *J. Biol. Chem.*, 281:1179-1187, 2006.
- [0712] PCT Appln. PCT/US89/01025
- [0713] PCT Appln. WO 00/02523
- [0714] PCT Appln. WO 00/12132
- [0715] PCT Appln. WO 00/12689
- [0716] PCT Appln. WO 00/15238
- [0717] PCT Appln. WO 01/34809
- [0718] PCT Appln. WO 01/60852
- [0719] PCT Appln. WO 01/98499
- [0720] PCT Appln. WO 02/059148
- [0721] PCT Appln. WO 02/094868
- [0722] PCT Appln. WO 03/53462
- [0723] PCT Appln. WO 04/43407
- [0724] PCT Appln. WO 06/032472
- [0725] PCT Appln. WO 06/032475
- [0726] PCT Appln. WO 06/032500
- [0727] PCT Appln. WO 07/113,222
- [0728] PCT Appln. WO 07/113,223
- [0729] PCT Appln. WO 94/09699
- [0730] PCT Appln. WO 95/06128
- [0731] PCT Appln. WO 95/08348
- [0732] PCT Appln. WO 98/57994
- [0733] Pearson & Lipman, *Proc. Natl. Acad. Sci. USA*, 85:2444, 1988.
- [0734] Pech et al., *Mol. Cell. Biol.*, 9:396, 1989.
- [0735] Pelletier and Sonenberg, *Nature*, 334(6180):320-325, 1988.
- [0736] Perez-Stable and Constantini, *Mol. Cell. Biol.*, 10:1116, 1990.
- [0737] Phonimdaeng et al., *Mol. Microbiol.*, 4:393-404, 1990.
- [0738] Picard and Schaffner, *Nature*, 307:83, 1984.
- [0739] Pinkert et al., *Genes and Dev.*, 1:268, 1987.
- [0740] Ponta et al., *Proc. Natl. Acad. Sci. USA*, 82:1020, 1985.
- [0741] Porton et al., *Mol. Cell. Biol.*, 10:1076, 1990.
- [0742] Potrykus et al., *Mol. Gen. Genet.*, 199(2):169-177, 1985.
- [0743] Pugsley, *Microbiol. Rev.*, 57:50-108, 1993.
- [0744] Pym et al., *Mol. Microbiol.*, 46:709-717, 2002.
- [0745] Pym et al., *Nat. Med.*, 9:533-539, 2003.
- [0746] Queen and Baltimore, *Cell*, 35:741, 1983.
- [0747] Quinn et al., *Mol. Cell. Biol.*, 9:4713, 1989.
- [0748] Redondo et al., *Science*, 247:1225, 1990.
- [0749] Reisman and Rotter, *Mol. Cell. Biol.*, 9:3571, 1989.
- [0750] Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1289-1329, 1990.
- [0751] Resendez Jr. et al., *Mol. Cell. Biol.*, 8:4579, 1988.

- [0752] Ripe et al., *Mol. Cell. Biol.*, 9:2224, 1989.  
 [0753] Rippe, et al., *Mol. Cell. Biol.*, 10:689-695, 1990.  
 [0754] Rittling et al., *Nuc. Acids Res.*, 17:1619, 1989.  
 [0755] Roben et al., *J. Immunol.* 154:6437-6445, 1995.  
 [0756] Rosen et al., *Cell*, 41:813, 1988.  
 [0757] Sakai et al., *Genes and Dev.*, 2:1144, 1988.  
 [0758] Salid-Salim et al., *Infect. Control Hosp. Epidemiol.* 24:451-455, 2003.  
 [0759] Sambrook et al., *In: Molecular cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001.  
 [0760] Schaffner et al., *J. Mol. Biol.*, 201:81, 1988.  
 [0761] Schneewind et al., *Cell* 70:267-281, 1992.  
 [0762] Schneewind et al., *EMBO*, 12:4803-4811, 1993.  
 [0763] Schneewind et al., *Science*, 268:103-6, 1995.  
 [0764] Searle et al., *Mol. Cell. Biol.*, 5:1480, 1985.  
 [0765] Sharp and Marciniak, *Cell*, 59:229, 1989.  
 [0766] Shaul and Ben-Levy, *EMBO J.*, 6:1913, 1987.  
 [0767] Shaw et al., *Microbiology*, 150:217-228, 2004.  
 [0768] Sheagren, *N. Engl. J. Med.* 310:1368-1373, 1984.  
 [0769] Sherman et al., *Mol. Cell. Biol.*, 9:50, 1989.  
 [0770] Shopsin et al., *J. Clin. Microbiol.*, 37:3556-63, 1999.  
 [0771] Sibbald et al., *Microbiol. Mol. Biol. Rev.*, 70:755-788, 2006.  
 [0772] Silverman and Goodyear. *Nat. Rev. Immunol.*, 6:465-75, 2006.  
 [0773] Sjudahl, *Eur. J. Biochem.* 73:343-351, 1977.  
 [0774] Sjoquist et al., *Eur. J. Biochem.* 30:190-194, 1972.  
 [0775] Sleigh and Lockett, *J. EMBO*, 4:3831, 1985.  
 [0776] Smith & Waterman, *Adv. Appl. Math.*, 2:482, 1981.  
 [0777] Smith et al., *Brit. J. Exp. Pathol.*, 28:57, 1947.  
 [0778] Sorensen et al., *Infect. Immun.*, 63:1710-1717, 1995.  
 [0779] Spalholz et al., *Cell*, 42:183, 1985.  
 [0780] Spandau and Lee, *J. Virology*, 62:427, 1988.  
 [0781] Spandidos and Wilkie, *EMBO J.*, 2:1193, 1983.  
 [0782] Stanley et al., *Proc. Natl. Acad. Sci. USA*, 100:13001-13006, 2003.  
 [0783] Stephens and Hentschel, *Biochem. J.*, 248:1, 1987.  
 [0784] Stewart and Young, In: *Solid Phase Peptide Synthesis*, 2d. ed., Pierce Chemical Co., 1984.  
 [0785] Stranger-Jones et al., *Proc. Nat. Acad. Sci. USA*, 103:16942-16947, 2006.  
 [0786] Stuart et al., *Nature*, 317:828, 1985.  
 [0787] Studier et al., *Methods Enzymol.* 185:60-89 1990.  
 [0788] Sullivan and Peterlin, *Mol. Cell. Biol.*, 7:3315, 1987.  
 [0789] Swartzendruber and Lehman, *J. Cell. Physiology*, 85:179, 1975.  
 [0790] Takebe et al., *Mol. Cell. Biol.*, 8:466, 1988.  
 [0791] Tam et al., *J. Am. Chem. Soc.*, 105:6442, 1983.  
 [0792] Tavernier et al., *Nature*, 301:634, 1983.  
 [0793] Taylor and Kingston, *Mol. Cell. Biol.*, 10:165, 1990a.  
 [0794] Taylor and Kingston, *Mol. Cell. Biol.*, 10:176, 1990b.  
 [0795] Taylor et al., *J. Biol. Chem.*, 264:15160, 1989.  
 [0796] Thiesen et al., *J. Virology*, 62:614, 1988.  
 [0797] Thomson et al., *J. Immunol.*, 157(2):822-826, 1996.  
 [0798] Tigges et al., *J. Immunol.*, 156(10):3901-3910, 1996.  
 [0799] Tigges et al., *J. Immunol.*, 156(10):3901-3910, 1996.  
 [0800] Ton-That et al., *Proc. Natl. Acad. Sci. USA*, 96(22):12424-9, 1999.  
 [0801] Treisman, *Cell*, 42:889, 1985.  
 [0802] Tronche et al., *Mol. Biol. Med.*, 7:173, 1990.  
 [0803] Trudel and Constantini, *Genes and Dev.*, 6:954, 1987.  
 [0804] Tyndell et al., *Nuc. Acids Res.*, 9:6231, 1981.  
 [0805] Uhlen et al., *J. Biol. Chem.* 259:1695-1702 and 13628 (Corr.) 1984.  
 [0806] van den Ent and Lowe, *FEBS Lett.*, 579:3837-3841, 2005.  
 [0807] van Wely et al., *FEMS Microbiol. Rev.*, 25:437-454, 2001.  
 [0808] Vannice and Levinson, *J. Virology*, 62:1305, 1988.  
 [0809] Vasseur et al., *Proc Natl. Acad. Sci. USA*, 77:1068, 1980.  
 [0810] Vaughan, et al., *Nat. Biotech.* 16; 535-539, 1998.  
 [0811] Wang and Calame, *Cell*, 47:241, 1986.  
 [0812] Weber et al., *Cell*, 36:983, 1984.  
 [0813] Weinberger et al. *Mol. Cell. Biol.*, 8:988, 1984.  
 [0814] Weiss et al., *J. Antimicrob. Chemother.*, 53(3):480-6, 2004.  
 [0815] Winoto and Baltimore, *Cell*, 59:649, 1989.  
 [0816] Wong et al., *Gene*, 10:87-94, 1980.  
 [0817] Xu et al., *J. Infect. Dis.*, 189:2323-2333, 2004.  
 [0818] Xu et al., *Mol. Microbiol.*, 66(3):787-800, 2007.  
 [0819] Yutzey et al. *Mol. Cell. Biol.*, 9:1397, 1989.

## SEQUENCE LISTING

&lt;160&gt; NUMBER OF SEQ ID NOS: 64

&lt;210&gt; SEQ ID NO 1

&lt;211&gt; LENGTH: 150

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 1

ttcaacaaag atcaacaaag cgccttctat gaaatcttga acatgcctaa cttaaacgaa 60

gcgcaacgta acgggttcat tcaaagtctt aaagacgacc caagccaaag cactaatgtt 120

ttaggtgaag ctaaaaaatt aaacgaatct 150

&lt;210&gt; SEQ ID NO 2

-continued

---

```

<211> LENGTH: 54
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 2

Gln Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile
1          5          10          15
Leu Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln
20          25          30
Ser Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala
35          40          45
Lys Lys Leu Asn Glu Ser
50

```

```

<210> SEQ ID NO 3
<211> LENGTH: 51
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 3

Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr Gln Val Leu Asn Met
1          5          10          15
Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys
20          25          30
Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys Leu
35          40          45
Asn Asp Ser
50

```

```

<210> SEQ ID NO 4
<211> LENGTH: 52
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 4

Asn Asn Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn
1          5          10          15
Met Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
20          25          30
Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys
35          40          45
Leu Asn Glu Ser
50

```

```

<210> SEQ ID NO 5
<211> LENGTH: 52
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 5

Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His
1          5          10          15
Leu Pro Asn Leu Thr Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
20          25          30
Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys
35          40          45
Leu Asn Asp Ala
50

```

-continued

<210> SEQ ID NO 6  
 <211> LENGTH: 52  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 6

```

Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His
1           5           10           15
Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
          20           25           30
Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys
          35           40           45
Leu Asn Asp Ala
          50
  
```

<210> SEQ ID NO 7  
 <211> LENGTH: 52  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (7)..(8)  
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (34)..(35)  
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 7

```

Asn Asn Phe Asn Lys Asp Xaa Xaa Ser Ala Phe Tyr Glu Ile Leu Asn
1           5           10           15
Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu
          20           25           30
Lys Xaa Xaa Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys
          35           40           45
Leu Asn Glu Ser
          50
  
```

<210> SEQ ID NO 8  
 <211> LENGTH: 52  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (7)..(8)  
 <223> OTHER INFORMATION: where X is any amino acid other than Q  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (12)..(35)  
 <223> OTHER INFORMATION: where Y is any amion acid other than D

<400> SEQUENCE: 8

```

Asn Asn Phe Asn Lys Asp Xaa Xaa Ser Ala Phe Tyr Glu Ile Leu Asn
1           5           10           15
Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu
          20           25           30
Lys Tyr Tyr Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys
          35           40           45
Leu Asn Glu Ser
          50
  
```

-continued

---

```

<210> SEQ ID NO 9
<211> LENGTH: 450
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 9

Met Lys Lys Lys Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile
1      5      10     15
Ala Ser Val Thr Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro
20     25     30
Ala Ala Asn Ala Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr
35     40     45
Gln Val Leu Asn Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe
50     55     60
Ile Gln Ser Leu Lys Asp Pro Ser Gln Ser Ala Asn Val Leu Gly
65     70     75     80
Glu Ala Gln Lys Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln
85     90     95
Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu
100    105    110
Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser
115    120    125
Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys
130    135    140
Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys
145    150    155    160
Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn
165    170    175
Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser
180    185    190
Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln
195    200    205
Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe
210    215    220
Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly
225    230    235    240
Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu
245    250    255
Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Glu Glu Asp
260    265    270
Asn Lys Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp
275    280    285
Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys Lys Pro Gly Lys Glu Asp
290    295    300
Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp
305    310    315    320
Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp
325    330    335
Gly Asn Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp
340    345    350
Gly Asn Gly Val His Val Val Lys Pro Gly Asp Thr Val Asn Asp Ile
355    360    365

```

-continued

---

Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys Ile Ala Ala Asp Asn Lys  
 370 375 380

Leu Ala Asp Lys Asn Met Ile Lys Pro Gly Gln Glu Leu Val Val Asp  
 385 390 395 400

Lys Lys Gln Pro Ala Asn His Ala Asp Ala Asn Lys Ala Gln Ala Leu  
 405 410 415

Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile Gly Thr Thr Val Phe Gly  
 420 425 430

Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu Leu Ala Gly Arg Arg Arg  
 435 440 445

Glu Leu  
 450

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 450

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 10

Met Lys Lys Lys Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile  
 1 5 10 15

Ala Ser Val Thr Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro  
 20 25 30

Ala Ala Asn Ala Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr  
 35 40 45

Gln Val Leu Asn Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe  
 50 55 60

Ile Gln Ser Leu Lys Asp Pro Ser Gln Ser Ala Asn Val Leu Gly  
 65 70 75 80

Glu Ala Gln Lys Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln  
 85 90 95

Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu  
 100 105 110

Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser  
 115 120 125

Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys  
 130 135 140

Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys  
 145 150 155 160

Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn  
 165 170 175

Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser  
 180 185 190

Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln  
 195 200 205

Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe  
 210 215 220

Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly  
 225 230 235 240

Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu  
 245 250 255

Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Glu Glu Asp  
 260 265 270

-continued

---

Asn Lys Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp  
 275 280 285  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys Lys Pro Gly Lys Glu Asp  
 290 295 300  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp  
 305 310 315 320  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp  
 325 330 335  
 Gly Asn Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp  
 340 345 350  
 Gly Asn Gly Val His Val Val Lys Pro Gly Asp Thr Val Asn Asp Ile  
 355 360 365  
 Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys Ile Ala Ala Asp Asn Lys  
 370 375 380  
 Leu Ala Asp Lys Asn Met Ile Lys Pro Gly Gln Glu Leu Val Val Asp  
 385 390 395 400  
 Lys Lys Gln Pro Ala Asn His Ala Asp Ala Asn Lys Ala Gln Ala Leu  
 405 410 415  
 Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile Gly Thr Thr Val Phe Gly  
 420 425 430  
 Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu Leu Ala Gly Arg Arg Arg  
 435 440 445  
 Glu Leu  
 450

<210> SEQ ID NO 11  
 <211> LENGTH: 97  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 11

Met Ala Met Ile Lys Met Ser Pro Glu Glu Ile Arg Ala Lys Ser Gln  
 1 5 10 15  
 Ser Tyr Gly Gln Gly Ser Asp Gln Ile Arg Gln Ile Leu Ser Asp Leu  
 20 25 30  
 Thr Arg Ala Gln Gly Glu Ile Ala Ala Asn Trp Glu Gly Gln Ala Phe  
 35 40 45  
 Ser Arg Phe Glu Glu Gln Phe Gln Gln Leu Ser Pro Lys Val Glu Lys  
 50 55 60  
 Phe Ala Gln Leu Leu Glu Glu Ile Lys Gln Gln Leu Asn Ser Thr Ala  
 65 70 75 80  
 Asp Ala Val Gln Glu Gln Asp Gln Gln Leu Ser Asn Asn Phe Gly Leu  
 85 90 95  
 Gln

<210> SEQ ID NO 12  
 <211> LENGTH: 102  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 12

Met Gly Gly Tyr Lys Gly Ile Lys Ala Asp Gly Gly Lys Val Asn Gln  
 1 5 10 15  
 Ala Lys Gln Leu Ala Ala Lys Ile Ala Lys Asp Ile Glu Ala Cys Gln  
 20 25 30



-continued

---

Lys Gln Thr Gln Gln Leu Ala Glu Tyr Ile Glu Gly Ser Asp Trp Glu  
 35 40 45

Gly Gln Phe Ala Asn Lys Val Lys Asp Val Leu Leu Ile Met Ala Lys  
 50 55 60

Phe Gln Glu Glu Leu Val Gln Pro Met Ala Asp His Gln Lys Ala Ile  
 65 70 75 80

Asp Asn Leu Ser Gln Asn Leu Ala Lys Tyr Asp Thr Leu Ser Ile Lys  
 85 90 95

Gln Gly Leu Asp Arg Val  
 100

<210> SEQ ID NO 13  
 <211> LENGTH: 1385  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 13

Met Leu Asn Arg Glu Asn Lys Thr Ala Ile Thr Arg Lys Gly Met Val  
 1 5 10 15

Ser Asn Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Thr Val Gly Thr  
 20 25 30

Ala Ser Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Gly Asn Gln  
 35 40 45

Glu Ala Lys Ala Ala Glu Ser Thr Asn Lys Glu Leu Asn Glu Ala Thr  
 50 55 60

Thr Ser Ala Ser Asp Asn Gln Ser Ser Asp Lys Val Asp Met Gln Gln  
 65 70 75 80

Leu Asn Gln Glu Asp Asn Thr Lys Asn Asp Asn Gln Lys Glu Met Val  
 85 90 95

Ser Ser Gln Gly Asn Glu Thr Thr Ser Asn Gly Asn Lys Ser Ile Glu  
 100 105 110

Lys Glu Ser Val Gln Ser Thr Thr Gly Asn Lys Val Glu Val Ser Thr  
 115 120 125

Ala Lys Ser Asp Glu Gln Ala Ser Pro Lys Ser Thr Asn Glu Asp Leu  
 130 135 140

Asn Thr Lys Gln Thr Ile Ser Asn Gln Glu Gly Leu Gln Pro Asp Leu  
 145 150 155 160

Leu Glu Asn Lys Ser Val Val Asn Val Gln Pro Thr Asn Glu Glu Asn  
 165 170 175

Lys Lys Val Asp Ala Lys Thr Glu Ser Thr Thr Leu Asn Val Lys Ser  
 180 185 190

Asp Ala Ile Lys Ser Asn Ala Glu Thr Leu Val Asp Asn Asn Ser Asn  
 195 200 205

Ser Asn Asn Glu Asn Asn Ala Asp Ile Ile Leu Pro Lys Ser Thr Ala  
 210 215 220

Pro Lys Ser Leu Asn Thr Arg Met Arg Met Ala Ala Ile Gln Pro Asn  
 225 230 235 240

Ser Thr Asp Ser Lys Asn Val Asn Asp Leu Ile Thr Ser Asn Thr Thr  
 245 250 255

Leu Thr Val Val Asp Ala Asp Asn Ser Lys Thr Ile Val Pro Ala Gln  
 260 265 270

Asp Tyr Leu Ser Leu Lys Ser Gln Ile Thr Val Asp Asp Lys Val Lys  
 275 280 285

-continued

---

Ser Gly Asp Tyr Phe Thr Ile Lys Tyr Ser Asp Thr Val Gln Val Tyr  
 290 295 300  
 Gly Leu Asn Pro Glu Asp Ile Lys Asn Ile Gly Asp Ile Lys Asp Pro  
 305 310 315 320  
 Asn Asn Gly Glu Thr Ile Ala Thr Ala Lys His Asp Thr Ala Asn Asn  
 325 330 335  
 Leu Ile Thr Tyr Thr Phe Thr Asp Tyr Val Asp Arg Phe Asn Ser Val  
 340 345 350  
 Lys Met Gly Ile Asn Tyr Ser Ile Tyr Met Asp Ala Asp Thr Ile Pro  
 355 360 365  
 Val Asp Lys Lys Asp Val Pro Phe Ser Val Thr Ile Gly Asn Gln Ile  
 370 375 380  
 Thr Thr Thr Thr Ala Asp Ile Thr Tyr Pro Ala Tyr Lys Glu Ala Asp  
 385 390 395 400  
 Asn Asn Ser Ile Gly Ser Ala Phe Thr Glu Thr Val Ser His Val Gly  
 405 410 415  
 Asn Val Glu Asp Pro Gly Tyr Tyr Asn Gln Val Val Tyr Val Asn Pro  
 420 425 430  
 Met Asp Lys Asp Leu Lys Gly Ala Lys Leu Lys Val Glu Ala Tyr His  
 435 440 445  
 Pro Lys Tyr Pro Thr Asn Ile Gly Gln Ile Asn Gln Asn Val Thr Asn  
 450 455 460  
 Ile Lys Ile Tyr Arg Val Pro Glu Gly Tyr Thr Leu Asn Lys Gly Tyr  
 465 470 475 480  
 Asp Val Asn Thr Asn Asp Leu Val Asp Val Thr Asp Glu Phe Lys Asn  
 485 490 495  
 Lys Met Thr Tyr Gly Ser Asn Gln Ser Val Asn Leu Asp Phe Gly Asp  
 500 505 510  
 Ile Thr Ser Ala Tyr Val Val Met Val Asn Thr Lys Phe Gln Tyr Thr  
 515 520 525  
 Asn Ser Glu Ser Pro Thr Leu Val Gln Met Ala Thr Leu Ser Ser Thr  
 530 535 540  
 Gly Asn Lys Ser Val Ser Thr Gly Asn Ala Leu Gly Phe Thr Asn Asn  
 545 550 555 560  
 Gln Ser Gly Gly Ala Gly Gln Glu Val Tyr Lys Ile Gly Asn Tyr Val  
 565 570 575  
 Trp Glu Asp Thr Asn Lys Asn Gly Val Gln Glu Leu Gly Glu Lys Gly  
 580 585 590  
 Val Gly Asn Val Thr Val Thr Val Phe Asp Asn Asn Thr Asn Thr Lys  
 595 600 605  
 Val Gly Glu Ala Val Thr Lys Glu Asp Gly Ser Tyr Leu Ile Pro Asn  
 610 615 620  
 Leu Pro Asn Gly Asp Tyr Arg Val Glu Phe Ser Asn Leu Pro Lys Gly  
 625 630 635 640  
 Tyr Glu Val Thr Pro Ser Lys Gln Gly Asn Asn Glu Glu Leu Asp Ser  
 645 650 655  
 Asn Gly Leu Ser Ser Val Ile Thr Val Asn Gly Lys Asp Asn Leu Ser  
 660 665 670  
 Ala Asp Leu Gly Ile Tyr Lys Pro Lys Tyr Asn Leu Gly Asp Tyr Val  
 675 680 685  
 Trp Glu Asp Thr Asn Lys Asn Gly Ile Gln Asp Gln Asp Glu Lys Gly

-continued

690					695					700					
Ile	Ser	Gly	Val	Thr	Val	Thr	Leu	Lys	Asp	Glu	Asn	Gly	Asn	Val	Leu
705					710					715					720
Lys	Thr	Val	Thr	Thr	Asp	Ala	Asp	Gly	Lys	Tyr	Lys	Phe	Thr	Asp	Leu
					725					730					735
Asp	Asn	Gly	Asn	Tyr	Lys	Val	Glu	Phe	Thr	Thr	Pro	Glu	Gly	Tyr	Thr
					740					745					750
Pro	Thr	Thr	Val	Thr	Ser	Gly	Ser	Asp	Ile	Glu	Lys	Asp	Ser	Asn	Gly
					755					760					765
Leu	Thr	Thr	Thr	Gly	Val	Ile	Asn	Gly	Ala	Asp	Asn	Met	Thr	Leu	Asp
					770					775					780
Ser	Gly	Phe	Tyr	Lys	Thr	Pro	Lys	Tyr	Asn	Leu	Gly	Asn	Tyr	Val	Trp
					785					790					800
Glu	Asp	Thr	Asn	Lys	Asp	Gly	Lys	Gln	Asp	Ser	Thr	Glu	Lys	Gly	Ile
					805					810					815
Ser	Gly	Val	Thr	Val	Thr	Leu	Lys	Asn	Glu	Asn	Gly	Glu	Val	Leu	Gln
					820					825					830
Thr	Thr	Lys	Thr	Asp	Lys	Asp	Gly	Lys	Tyr	Gln	Phe	Thr	Gly	Leu	Glu
					835					840					845
Asn	Gly	Thr	Tyr	Lys	Val	Glu	Phe	Glu	Thr	Pro	Ser	Gly	Tyr	Thr	Pro
					850					855					860
Thr	Gln	Val	Gly	Ser	Gly	Thr	Asp	Glu	Gly	Ile	Asp	Ser	Asn	Gly	Thr
					865					870					880
Ser	Thr	Thr	Gly	Val	Ile	Lys	Asp	Lys	Asp	Asn	Asp	Thr	Ile	Asp	Ser
					885					890					895
Gly	Phe	Tyr	Lys	Pro	Thr	Tyr	Asn	Leu	Gly	Asp	Tyr	Val	Trp	Glu	Asp
					900					905					910
Thr	Asn	Lys	Asn	Gly	Val	Gln	Asp	Lys	Asp	Glu	Lys	Gly	Ile	Ser	Gly
					915					920					925
Val	Thr	Val	Thr	Leu	Lys	Asp	Glu	Asn	Asp	Lys	Val	Leu	Lys	Thr	Val
					930					935					940
Thr	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	Gln	Phe	Thr	Asp	Leu	Asn	Asn	Gly
					945					950					960
Thr	Tyr	Lys	Val	Glu	Phe	Glu	Thr	Pro	Ser	Gly	Tyr	Thr	Pro	Thr	Ser
					965					970					975
Val	Thr	Ser	Gly	Asn	Asp	Thr	Glu	Lys	Asp	Ser	Asn	Gly	Leu	Thr	Thr
					980					985					990
Thr	Gly	Val	Ile	Lys	Asp	Ala	Asp	Asn	Met	Thr	Leu	Asp	Ser	Gly	Phe
					995					1000					1005
Tyr	Lys	Thr	Pro	Lys	Tyr	Ser	Leu	Gly	Asp	Tyr	Val	Trp	Tyr	Asp	
					1010					1015					1020
Ser	Asn	Lys	Asp	Gly	Lys	Gln	Asp	Ser	Thr	Glu	Lys	Gly	Ile	Lys	
					1025					1030					1035
Asp	Val	Lys	Val	Ile	Leu	Leu	Asn	Glu	Lys	Gly	Glu	Val	Ile	Gly	
					1040					1045					1050
Thr	Thr	Lys	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	Arg	Phe	Asp	Asn	Leu	
					1055					1060					1065
Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu	Lys	Pro	Thr	Gly	Leu	
					1070					1075					1080
Thr	Gln	Thr	Gly	Thr	Asn	Thr	Thr	Glu	Asp	Asp	Lys	Asp	Ala	Asp	
					1085					1090					1095

-continued

---

Gly	Gly	Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp	Asp	Phe	Thr
	1100					1105						1110		
Leu	Asp	Asn	Gly	Tyr	Tyr	Glu	Glu	Glu	Thr	Ser	Asp	Ser	Asp	Ser
	1115					1120						1125		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1130					1135						1140		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1145					1150						1155		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1160					1165						1170		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1175					1180						1185		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1190					1195						1200		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1205					1210						1215		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1220					1225						1230		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1235					1240						1245		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1250					1255						1260		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1265					1270						1275		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1280					1285						1290		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	1295					1300						1305		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	1310					1315						1320		
Ser	Asp	Ala	Gly	Lys	His	Thr	Pro	Val	Lys	Pro	Met	Ser	Thr	Thr
	1325					1330						1335		
Lys	Asp	His	His	Asn	Lys	Ala	Lys	Ala	Leu	Pro	Glu	Thr	Gly	Asn
	1340					1345						1350		
Glu	Asn	Ser	Gly	Ser	Asn	Asn	Ala	Thr	Leu	Phe	Gly	Gly	Leu	Phe
	1355					1360						1365		
Ala	Ala	Leu	Gly	Ser	Leu	Leu	Leu	Phe	Gly	Arg	Arg	Lys	Lys	Gln
	1370					1375						1380		
Asn	Lys													
	1385													

<210> SEQ ID NO 14  
 <211> LENGTH: 1141  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 14

Met	Ile	Asn	Arg	Asp	Asn	Lys	Lys	Ala	Ile	Thr	Lys	Lys	Gly	Met	Ile
1			5						10					15	
Ser	Asn	Arg	Leu	Asn	Lys	Phe	Ser	Ile	Arg	Lys	Tyr	Thr	Val	Gly	Thr
			20				25						30		
Ala	Ser	Ile	Leu	Val	Gly	Thr	Thr	Leu	Ile	Phe	Gly	Leu	Gly	Asn	Gln
			35				40					45			

-continued

---

Glu	Ala	Lys	Ala	Ala	Glu	Asn	Thr	Ser	Thr	Glu	Asn	Ala	Lys	Gln	Asp	
50						55					60					
Asp	Ala	Thr	Thr	Ser	Asp	Asn	Lys	Glu	Val	Val	Ser	Glu	Thr	Glu	Asn	
65					70					75					80	
Asn	Ser	Thr	Thr	Glu	Asn	Asp	Ser	Thr	Asn	Pro	Ile	Lys	Lys	Glu	Thr	
				85					90					95		
Asn	Thr	Asp	Ser	Gln	Pro	Glu	Ala	Lys	Glu	Glu	Ser	Thr	Thr	Ser	Ser	
			100					105					110			
Thr	Gln	Gln	Gln	Gln	Asn	Asn	Val	Thr	Ala	Thr	Thr	Glu	Thr	Lys	Pro	
		115					120					125				
Gln	Asn	Ile	Glu	Lys	Glu	Asn	Val	Lys	Pro	Ser	Thr	Asp	Lys	Thr	Ala	
130						135					140					
Thr	Glu	Asp	Thr	Ser	Val	Ile	Leu	Glu	Glu	Lys	Lys	Ala	Pro	Asn	Tyr	
145					150					155					160	
Thr	Asn	Asn	Asp	Val	Thr	Thr	Lys	Pro	Ser	Thr	Ser	Glu	Ile	Gln	Thr	
				165					170					175		
Lys	Pro	Thr	Thr	Pro	Gln	Glu	Ser	Thr	Asn	Ile	Glu	Asn	Ser	Gln	Pro	
			180					185						190		
Gln	Pro	Thr	Pro	Ser	Lys	Val	Asp	Asn	Gln	Val	Thr	Asp	Ala	Thr	Asn	
		195					200					205				
Pro	Lys	Glu	Pro	Val	Asn	Val	Ser	Lys	Glu	Glu	Leu	Lys	Asn	Asn	Pro	
	210					215					220					
Glu	Lys	Leu	Lys	Glu	Leu	Val	Arg	Asn	Asp	Asn	Asn	Thr	Asp	Arg	Ser	
225					230					235					240	
Thr	Lys	Pro	Val	Ala	Thr	Ala	Pro	Thr	Ser	Val	Ala	Pro	Lys	Arg	Leu	
				245					250					255		
Asn	Ala	Lys	Met	Arg	Phe	Ala	Val	Ala	Gln	Pro	Ala	Ala	Val	Ala	Ser	
			260						265				270			
Asn	Asn	Val	Asn	Asp	Leu	Ile	Thr	Val	Thr	Lys	Gln	Thr	Ile	Lys	Val	
		275					280						285			
Gly	Asp	Gly	Lys	Asp	Asn	Val	Ala	Ala	Ala	His	Asp	Gly	Lys	Asp	Ile	
	290					295					300					
Glu	Tyr	Asp	Thr	Glu	Phe	Thr	Ile	Asp	Asn	Lys	Val	Lys	Lys	Gly	Asp	
305					310					315					320	
Thr	Met	Thr	Ile	Asn	Tyr	Asp	Lys	Asn	Val	Ile	Pro	Ser	Asp	Leu	Thr	
				325					330					335		
Asp	Lys	Asn	Asp	Pro	Ile	Asp	Ile	Thr	Asp	Pro	Ser	Gly	Glu	Val	Ile	
			340					345					350			
Ala	Lys	Gly	Thr	Phe	Asp	Lys	Ala	Thr	Lys	Gln	Ile	Thr	Tyr	Thr	Phe	
		355					360					365				
Thr	Asp	Tyr	Val	Asp	Lys	Tyr	Glu	Asp	Ile	Lys	Ala	Arg	Leu	Thr	Leu	
	370					375					380					
Tyr	Ser	Tyr	Ile	Asp	Lys	Gln	Ala	Val	Pro	Asn	Glu	Thr	Ser	Leu	Asn	
385					390					395					400	
Leu	Thr	Phe	Ala	Thr	Ala	Gly	Lys	Glu	Thr	Ser	Gln	Asn	Val	Ser	Val	
				405					410					415		
Asp	Tyr	Gln	Asp	Pro	Met	Val	His	Gly	Asp	Ser	Asn	Ile	Gln	Ser	Ile	
		420						425					430			
Phe	Thr	Lys	Leu	Asp	Glu	Asn	Lys	Gln	Thr	Ile	Glu	Gln	Gln	Ile	Tyr	
		435					440					445				
Val	Asn	Pro	Leu	Lys	Lys	Thr	Ala	Thr	Asn	Thr	Lys	Val	Asp	Ile	Ala	
	450					455					460					

-continued

---

Gly Ser Gln Val Asp Asp Tyr Gly Asn Ile Lys Leu Gly Asn Gly Ser  
 465 470 475 480  
 Thr Ile Ile Asp Gln Asn Thr Glu Ile Lys Val Tyr Lys Val Asn Pro  
 485 490 495  
 Asn Gln Gln Leu Pro Gln Ser Asn Arg Ile Tyr Asp Phe Ser Gln Tyr  
 500 505 510  
 Glu Asp Val Thr Ser Gln Phe Asp Asn Lys Lys Ser Phe Ser Asn Asn  
 515 520 525  
 Val Ala Thr Leu Asp Phe Gly Asp Ile Asn Ser Ala Tyr Ile Ile Lys  
 530 535 540  
 Val Val Ser Lys Tyr Thr Pro Thr Ser Asp Gly Glu Leu Asp Ile Ala  
 545 550 555 560  
 Gln Gly Thr Ser Met Arg Thr Thr Asp Lys Tyr Gly Tyr Tyr Asn Tyr  
 565 570 575  
 Ala Gly Tyr Ser Asn Phe Ile Val Thr Ser Asn Asp Thr Gly Gly Gly  
 580 585 590  
 Asp Gly Thr Val Lys Pro Glu Glu Lys Leu Tyr Lys Ile Gly Asp Tyr  
 595 600 605  
 Val Trp Glu Asp Val Asp Lys Asp Gly Val Gln Gly Thr Asp Ser Lys  
 610 615 620  
 Glu Lys Pro Met Ala Asn Val Leu Val Thr Leu Thr Tyr Pro Asp Gly  
 625 630 635 640  
 Thr Thr Lys Ser Val Arg Thr Asp Ala Asn Gly His Tyr Glu Phe Gly  
 645 650 655  
 Gly Leu Lys Asp Gly Glu Thr Tyr Thr Val Lys Phe Glu Thr Pro Ala  
 660 665 670  
 Gly Tyr Leu Pro Thr Lys Val Asn Gly Thr Thr Asp Gly Glu Lys Asp  
 675 680 685  
 Ser Asn Gly Ser Ser Ile Thr Val Lys Ile Asn Gly Lys Asp Asp Met  
 690 695 700  
 Ser Leu Asp Thr Gly Phe Tyr Lys Glu Pro Lys Tyr Asn Leu Gly Asp  
 705 710 715 720  
 Tyr Val Trp Glu Asp Thr Asn Lys Asp Gly Ile Gln Asp Ala Asn Glu  
 725 730 735  
 Pro Gly Ile Lys Asp Val Lys Val Thr Leu Lys Asp Ser Thr Gly Lys  
 740 745 750  
 Val Ile Gly Thr Thr Thr Thr Asp Ala Ser Gly Lys Tyr Lys Phe Thr  
 755 760 765  
 Asp Leu Asp Asn Gly Asn Tyr Thr Val Glu Phe Glu Thr Pro Ala Gly  
 770 775 780  
 Tyr Thr Pro Thr Val Lys Asn Thr Thr Ala Glu Asp Lys Asp Ser Asn  
 785 790 795 800  
 Gly Leu Thr Thr Thr Gly Val Ile Lys Asp Ala Asp Asn Met Thr Leu  
 805 810 815  
 Asp Ser Gly Phe Tyr Lys Thr Pro Lys Tyr Ser Leu Gly Asp Tyr Val  
 820 825 830  
 Trp Tyr Asp Ser Asn Lys Asp Gly Lys Gln Asp Ser Thr Glu Lys Gly  
 835 840 845  
 Ile Lys Asp Val Lys Val Thr Leu Leu Asn Glu Lys Gly Glu Val Ile  
 850 855 860  
 Gly Thr Thr Lys Thr Asp Glu Asn Gly Lys Tyr Arg Phe Asp Asn Leu



-continued

---

		85						90						95	
Gln	Asn	Asn	Lys	Tyr	Tyr	Phe	Gln	Thr	Val	Leu	Asn	Asn	Ala	Ser	Phe
		100						105					110		
Trp	Lys	Glu	Tyr	Lys	Phe	Tyr	Asn	Ala	Asn	Asn	Gln	Glu	Leu	Ala	Thr
		115					120				125				
Thr	Val	Val	Asn	Asp	Asn	Lys	Lys	Ala	Asp	Thr	Arg	Thr	Ile	Asn	Val
		130				135					140				
Ala	Val	Glu	Pro	Gly	Tyr	Lys	Ser	Leu	Thr	Thr	Lys	Val	His	Ile	Val
		145			150						155			160	
Val	Pro	Gln	Ile	Asn	Tyr	Asn	His	Arg	Tyr	Thr	Thr	His	Leu	Glu	Phe
				165					170					175	
Glu	Lys	Ala	Ile	Pro	Thr	Leu	Ala	Asp	Ala	Ala	Lys	Pro	Asn	Asn	Val
		180						185					190		
Lys	Pro	Val	Gln	Pro	Lys	Pro	Ala	Gln	Pro	Lys	Thr	Pro	Thr	Glu	Gln
		195					200					205			
Thr	Lys	Pro	Val	Gln	Pro	Lys	Val	Glu	Lys	Val	Lys	Pro	Thr	Val	Thr
		210				215					220				
Thr	Thr	Ser	Lys	Val	Glu	Asp	Asn	His	Ser	Thr	Lys	Val	Val	Ser	Thr
		225			230					235					240
Asp	Thr	Thr	Lys	Asp	Gln	Thr	Lys	Thr	Gln	Thr	Ala	His	Thr	Val	Lys
				245					250					255	
Thr	Ala	Gln	Thr	Ala	Gln	Glu	Gln	Asn	Lys	Val	Gln	Thr	Pro	Val	Lys
			260					265					270		
Asp	Val	Ala	Thr	Ala	Lys	Ser	Glu	Ser	Asn	Asn	Gln	Ala	Val	Ser	Asp
		275					280					285			
Asn	Lys	Ser	Gln	Gln	Thr	Asn	Lys	Val	Thr	Lys	His	Asn	Glu	Thr	Pro
		290				295					300				
Lys	Gln	Ala	Ser	Lys	Ala	Lys	Glu	Leu	Pro	Lys	Thr	Gly	Leu	Thr	Ser
		305			310					315					320
Val	Asp	Asn	Phe	Ile	Ser	Thr	Val	Ala	Phe	Ala	Thr	Leu	Ala	Leu	Leu
				325					330					335	
Gly	Ser	Leu	Ser	Leu	Leu	Leu	Phe	Lys	Arg	Lys	Glu	Ser	Lys		
			340					345					350		

<210> SEQ ID NO 16  
 <211> LENGTH: 645  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 16

Met	Asn	Lys	Gln	Gln	Lys	Glu	Phe	Lys	Ser	Phe	Tyr	Ser	Ile	Arg	Lys
1			5						10					15	
Ser	Ser	Leu	Gly	Val	Ala	Ser	Val	Ala	Ile	Ser	Thr	Leu	Leu	Leu	Leu
		20						25					30		
Met	Ser	Asn	Gly	Glu	Ala	Gln	Ala	Ala	Ala	Glu	Glu	Thr	Gly	Gly	Thr
		35					40					45			
Asn	Thr	Glu	Ala	Gln	Pro	Lys	Thr	Glu	Ala	Val	Ala	Ser	Pro	Thr	Thr
		50				55						60			
Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser
		65			70					75				80	
Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala
				85					90					95	
Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Ala	Val	Lys



-continued

100					105					110					
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu
	115						120					125			
Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Asp	His	Ser
	130					135					140				
Ala	Pro	Asn	Ser	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Glu	Asn	Gly
	145					150					155				160
Glu	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Lys	Pro	Ala	Arg	Val
				165					170					175	
Ile	Phe	Thr	Asp	Ser	Lys	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly
			180					185					190		
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro
		195					200					205			
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg
	210					215					220				
Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Ala	Val	Lys	Ile	Val	Ser	Ser	Thr
	225					230					235				240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe
				245					250					255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp
			260					265					270		
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu
		275					280					285			
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu
	290					295					300				
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala
	305					310					315				320
Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln
				325					330					335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val
			340					345					350		
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys
		355					360					365			
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met
	370					375					380				
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln
	385					390					395				400
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile
				405					410					415	
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys
			420					425					430		
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile
		435					440					445			
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys
		450				455					460				
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr
	465					470					475				480
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln
				485					490					495	
Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu
			500					505						510	

-continued

---

```

Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
   515                               520                    525

Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
   530                               535                               540

Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
   545                               550                               555                    560

Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
   565                               570                               575

Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
   580                               585                               590

His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
   595                               600                               605

Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
   610                               615                               620

Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
   625                               630                               635                    640

Arg Lys Arg Lys Asn
   645

```

```

<210> SEQ ID NO 17
<211> LENGTH: 80
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

```

```

<400> SEQUENCE: 17

```

```

Met Asn Gln His Val Lys Val Thr Phe Asp Phe Thr Asn Tyr Asn Tyr
 1                               5                               10                    15

Gly Thr Tyr Asp Leu Ala Val Pro Ala Tyr Leu Pro Ile Lys Asn Leu
   20                               25                               30

Ile Ala Leu Val Leu Asp Ser Leu Asp Ile Ser Ile Phe Asp Val Asn
   35                               40                               45

Thr Gln Ile Lys Val Met Thr Lys Gly Gln Leu Leu Val Glu Asn Asp
   50                               55                               60

Arg Leu Ile Asp Tyr Gln Ile Ala Asp Gly Asp Ile Leu Lys Leu Leu
   65                               70                               75                    80

```

```

<210> SEQ ID NO 18
<211> LENGTH: 877
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

```

```

<400> SEQUENCE: 18

```

```

Met Lys Lys Arg Ile Asp Tyr Leu Ser Asn Lys Gln Asn Lys Tyr Ser
 1                               5                               10                    15

Ile Arg Arg Phe Thr Val Gly Thr Thr Ser Val Ile Val Gly Ala Thr
   20                               25                               30

Ile Leu Phe Gly Ile Gly Asn His Gln Ala Gln Ala Ser Glu Gln Ser
   35                               40                               45

Asn Asp Thr Thr Gln Ser Ser Lys Asn Asn Ala Ser Ala Asp Ser Glu
   50                               55                               60

Lys Asn Asn Met Ile Glu Thr Pro Gln Leu Asn Thr Thr Ala Asn Asp
   65                               70                               75                    80

Thr Ser Asp Ile Ser Ala Asn Thr Asn Ser Ala Asn Val Asp Ser Thr
   85                               90                               95

Thr Lys Pro Met Ser Thr Gln Thr Ser Asn Thr Thr Thr Thr Glu Pro

```

-continued

100					105					110					
Ala	Ser	Thr	Asn	Glu	Thr	Pro	Gln	Pro	Thr	Ala	Ile	Lys	Asn	Gln	Ala
			115				120					125			
Thr	Ala	Ala	Lys	Met	Gln	Asp	Gln	Thr	Val	Pro	Gln	Glu	Ala	Asn	Ser
			130				135					140			
Gln	Val	Asp	Asn	Lys	Thr	Thr	Asn	Asp	Ala	Asn	Ser	Ile	Ala	Thr	Asn
			145				150					155			
Ser	Glu	Leu	Lys	Asn	Ser	Gln	Thr	Leu	Asp	Leu	Pro	Gln	Ser	Ser	Pro
				165					170					175	
Gln	Thr	Ile	Ser	Asn	Ala	Gln	Gly	Thr	Ser	Lys	Pro	Ser	Val	Arg	Thr
				180					185					190	
Arg	Ala	Val	Arg	Ser	Leu	Ala	Val	Ala	Glu	Pro	Val	Val	Asn	Ala	Ala
				195					200					205	
Asp	Ala	Lys	Gly	Thr	Asn	Val	Asn	Asp	Lys	Val	Thr	Ala	Ser	Asn	Phe
				210					215					220	
Lys	Leu	Glu	Lys	Thr	Thr	Phe	Asp	Pro	Asn	Gln	Ser	Gly	Asn	Thr	Phe
				225					230					235	
Met	Ala	Ala	Asn	Phe	Thr	Val	Thr	Asp	Lys	Val	Lys	Ser	Gly	Asp	Tyr
				245					250					255	
Phe	Thr	Ala	Lys	Leu	Pro	Asp	Ser	Leu	Thr	Gly	Asn	Gly	Asp	Val	Asp
				260					265					270	
Tyr	Ser	Asn	Ser	Asn	Asn	Thr	Met	Pro	Ile	Ala	Asp	Ile	Lys	Ser	Thr
				275					280					285	
Asn	Gly	Asp	Val	Val	Ala	Lys	Ala	Thr	Tyr	Asp	Ile	Leu	Thr	Lys	Thr
				290					295					300	
Tyr	Thr	Phe	Val	Phe	Thr	Asp	Tyr	Val	Asn	Asn	Lys	Glu	Asn	Ile	Asn
				305					310					315	
Gly	Gln	Phe	Ser	Leu	Pro	Leu	Phe	Thr	Asp	Arg	Ala	Lys	Ala	Pro	Lys
				325					330					335	
Ser	Gly	Thr	Tyr	Asp	Ala	Asn	Ile	Asn	Ile	Ala	Asp	Glu	Met	Phe	Asn
				340					345					350	
Asn	Lys	Ile	Thr	Tyr	Asn	Tyr	Ser	Ser	Pro	Ile	Ala	Gly	Ile	Asp	Lys
				355					360					365	
Pro	Asn	Gly	Ala	Asn	Ile	Ser	Ser	Gln	Ile	Ile	Gly	Val	Asp	Thr	Ala
				370					375					380	
Ser	Gly	Gln	Asn	Thr	Tyr	Lys	Gln	Thr	Val	Phe	Val	Asn	Pro	Lys	Gln
				385					390					395	
Arg	Val	Leu	Gly	Asn	Thr	Trp	Val	Tyr	Ile	Lys	Gly	Tyr	Gln	Asp	Lys
				405					410					415	
Ile	Glu	Glu	Ser	Ser	Gly	Lys	Val	Ser	Ala	Thr	Asp	Thr	Lys	Leu	Arg
				420					425					430	
Ile	Phe	Glu	Val	Asn	Asp	Thr	Ser	Lys	Leu	Ser	Asp	Ser	Tyr	Tyr	Ala
				435					440					445	
Asp	Pro	Asn	Asp	Ser	Asn	Leu	Lys	Glu	Val	Thr	Asp	Gln	Phe	Lys	Asn
				450					455					460	
Arg	Ile	Tyr	Tyr	Glu	His	Pro	Asn	Val	Ala	Ser	Ile	Lys	Phe	Gly	Asp
				465					470					475	
Ile	Thr	Lys	Thr	Tyr	Val	Val	Leu	Val	Glu	Gly	His	Tyr	Asp	Asn	Thr
				485					490					495	
Gly	Lys	Asn	Leu	Lys	Thr	Gln	Val	Ile	Gln	Glu	Asn	Val	Asp	Pro	Val
				500					505					510	

-continued

---

```

Thr Asn Arg Asp Tyr Ser Ile Phe Gly Trp Asn Asn Glu Asn Val Val
   515                               520                               525

Arg Tyr Gly Gly Gly Ser Ala Asp Gly Asp Ser Ala Val Asn Pro Lys
   530                               535                               540

Asp Pro Thr Pro Gly Pro Pro Val Asp Pro Glu Pro Ser Pro Asp Pro
545                               550                               555                               560

Glu Pro Glu Pro Thr Pro Asp Pro Glu Pro Ser Pro Asp Pro Glu Pro
   565                               570                               575

Glu Pro Ser Pro Asp Pro Asp Pro Asp Ser Asp Ser Asp Ser Asp Ser
   580                               585                               590

Gly Ser Asp Ser Asp Ser Gly Ser Asp Ser Asp Ser Glu Ser Asp Ser
   595                               600                               605

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Glu Ser
   610                               615                               620

Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
625                               630                               635                               640

Asp Ser Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser
   645                               650                               655

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser
   660                               665                               670

Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
   675                               680                               685

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
   690                               695                               700

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser
705                               710                               715                               720

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
   725                               730                               735

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
   740                               745                               750

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
   755                               760                               765

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
   770                               775                               780

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
785                               790                               795                               800

Asp Ser Asp Ser Arg Val Thr Pro Pro Asn Asn Glu Gln Lys Ala Pro
   805                               810                               815

Ser Asn Pro Lys Gly Glu Val Asn His Ser Asn Lys Val Ser Lys Gln
   820                               825                               830

His Lys Thr Asp Ala Leu Pro Glu Thr Gly Asp Lys Ser Glu Asn Thr
   835                               840                               845

Asn Ala Thr Leu Phe Gly Ala Met Met Ala Leu Leu Gly Ser Leu Leu
   850                               855                               860

Leu Phe Arg Lys Arg Lys Gln Asp His Lys Glu Lys Ala
865                               870                               875

```

&lt;210&gt; SEQ ID NO 19

&lt;211&gt; LENGTH: 227

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 19

-continued

---

```

Met Lys Asn Ile Leu Lys Val Phe Asn Thr Thr Ile Leu Ala Leu Ile
1      5      10      15

Ile Ile Ile Ala Thr Phe Ser Asn Ser Ala Asn Ala Ala Asp Ser Gly
      20      25      30

Thr Leu Asn Tyr Glu Val Tyr Lys Tyr Asn Thr Asn Asp Thr Ser Ile
      35      40      45

Ala Asn Asp Tyr Phe Asn Lys Pro Ala Lys Tyr Ile Lys Lys Asn Gly
      50      55      60

Lys Leu Tyr Val Gln Ile Thr Val Asn His Ser His Trp Ile Thr Gly
65      70      75      80

Met Ser Ile Glu Gly His Lys Glu Asn Ile Ile Ser Lys Asn Thr Ala
      85      90      95

Lys Asp Glu Arg Thr Ser Glu Phe Glu Val Ser Lys Leu Asn Gly Lys
      100      105      110

Ile Asp Gly Lys Ile Asp Val Tyr Ile Asp Glu Lys Val Asn Gly Lys
      115      120      125

Pro Phe Lys Tyr Asp His His Tyr Asn Ile Thr Tyr Lys Phe Asn Gly
      130      135      140

Pro Thr Asp Val Ala Gly Ala Asn Ala Pro Gly Lys Asp Asp Lys Asn
145      150      155      160

Ser Ala Ser Gly Ser Asp Lys Gly Ser Asp Gly Thr Thr Thr Gly Gln
      165      170      175

Ser Glu Ser Asn Ser Ser Asn Lys Asp Lys Val Glu Asn Pro Gln Thr
      180      185      190

Asn Ala Gly Thr Pro Ala Tyr Ile Tyr Ala Ile Pro Val Ala Ser Leu
      195      200      205

Ala Leu Leu Ile Ala Ile Thr Leu Phe Val Arg Lys Lys Ser Lys Gly
210      215      220

Asn Val Glu
225

```

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 635

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 20

```

Met Ala Lys Tyr Arg Gly Lys Pro Phe Gln Leu Tyr Val Lys Leu Ser
1      5      10      15

Cys Ser Thr Met Met Ala Ser Ser Ile Ile Leu Thr Asn Ile Leu Pro
      20      25      30

Tyr Asp Ala Gln Ala Ala Ser Glu Lys Asp Thr Glu Ile Ser Lys Glu
      35      40      45

Ile Leu Ser Lys Gln Asp Leu Leu Asp Lys Val Asp Lys Ala Ile Arg
      50      55      60

Gln Ile Glu Gln Leu Lys Gln Leu Ser Ala Ser Ser Lys Ala His Tyr
65      70      75      80

Lys Ala Gln Leu Asn Glu Ala Lys Thr Ala Ser Gln Ile Asp Glu Ile
      85      90      95

Ile Lys Arg Ala Asn Glu Leu Asp Ser Lys Glu Asn Lys Ser Ser His
      100      105      110

Thr Glu Met Asn Gly Gln Ser Asp Ile Asp Ser Lys Leu Asp Gln Leu
115      120      125

```

-continued

---

Leu	Lys	Asp	Leu	Asn	Glu	Val	Ser	Ser	Asn	Val	Asp	Arg	Gly	Gln	Gln
130						135					140				
Ser	Gly	Glu	Asp	Asp	Leu	Asn	Ala	Met	Lys	Asn	Asp	Met	Ser	Gln	Thr
145					150					155					160
Ala	Thr	Thr	Lys	Tyr	Gly	Glu	Lys	Asp	Asp	Lys	Asn	Asp	Glu	Ala	Met
				165					170					175	
Val	Asn	Lys	Ala	Leu	Glu	Asp	Leu	Asp	His	Leu	Asn	Gln	Gln	Ile	His
			180					185					190		
Lys	Ser	Lys	Asp	Ala	Leu	Lys	Asp	Ala	Ser	Lys	Asp	Pro	Ala	Val	Ser
		195					200					205			
Thr	Thr	Asp	Ser	Asn	His	Glu	Val	Ala	Lys	Thr	Pro	Asn	Asn	Asp	Gly
	210					215					220				
Ser	Gly	His	Val	Val	Leu	Asn	Lys	Phe	Leu	Ser	Asn	Glu	Glu	Asn	Gln
225					230					235					240
Ser	His	Ser	Asn	Gln	Leu	Thr	Asp	Lys	Leu	Gln	Gly	Ser	Asp	Lys	Ile
				245					250					255	
Asn	His	Ala	Met	Ile	Glu	Lys	Leu	Ala	Lys	Ser	Asn	Ala	Ser	Thr	Gln
			260					265						270	
His	Tyr	Thr	Tyr	His	Lys	Leu	Asn	Thr	Leu	Gln	Ser	Leu	Asp	Gln	Arg
		275					280						285		
Ile	Ala	Asn	Thr	Gln	Leu	Pro	Lys	Asn	Gln	Lys	Ser	Asp	Leu	Met	Ser
	290					295					300				
Glu	Val	Asn	Lys	Thr	Lys	Glu	Arg	Ile	Lys	Ser	Gln	Arg	Asn	Ile	Ile
305					310					315					320
Leu	Glu	Glu	Leu	Ala	Arg	Thr	Asp	Asp	Lys	Lys	Tyr	Ala	Thr	Gln	Ser
				325					330					335	
Ile	Leu	Glu	Ser	Ile	Phe	Asn	Lys	Asp	Glu	Ala	Asp	Lys	Ile	Leu	Lys
			340					345					350		
Asp	Ile	Arg	Val	Asp	Gly	Lys	Thr	Asp	Gln	Gln	Ile	Ala	Asp	Gln	Ile
		355					360						365		
Thr	Arg	His	Ile	Asp	Gln	Leu	Ser	Leu	Thr	Thr	Ser	Asp	Asp	Leu	Leu
	370					375					380				
Thr	Ser	Leu	Ile	Asp	Gln	Ser	Gln	Asp	Lys	Ser	Leu	Leu	Ile	Ser	Gln
385					390					395					400
Ile	Leu	Gln	Thr	Lys	Leu	Gly	Lys	Ala	Glu	Ala	Asp	Lys	Leu	Ala	Lys
				405					410					415	
Asp	Trp	Thr	Asn	Lys	Gly	Leu	Ser	Asn	Arg	Gln	Ile	Val	Asp	Gln	Leu
			420					425					430		
Lys	Lys	His	Phe	Ala	Ser	Thr	Gly	Asp	Thr	Ser	Ser	Asp	Asp	Ile	Leu
		435					440						445		
Lys	Ala	Ile	Leu	Asn	Asn	Ala	Lys	Asp	Lys	Lys	Gln	Ala	Ile	Glu	Thr
	450					455					460				
Ile	Leu	Ala	Thr	Arg	Ile	Glu	Arg	Gln	Lys	Ala	Lys	Leu	Leu	Ala	Asp
465					470					475					480
Leu	Ile	Thr	Lys	Ile	Glu	Thr	Asp	Gln	Asn	Lys	Ile	Phe	Asn	Leu	Val
				485					490					495	
Lys	Ser	Ala	Leu	Asn	Gly	Lys	Ala	Asp	Asp	Leu	Leu	Asn	Leu	Gln	Lys
			500					505					510		
Arg	Leu	Asn	Gln	Thr	Lys	Lys	Asp	Ile	Asp	Tyr	Ile	Leu	Ser	Pro	Ile
		515					520						525		
Val	Asn	Arg	Pro	Ser	Leu	Leu	Asp	Arg	Leu	Asn	Lys	Asn	Gly	Lys	Thr
	530						535					540			

-continued

Thr Asp Leu Asn Lys Leu Ala Asn Leu Met Asn Gln Gly Ser Asn Leu  
 545 550 555 560  
 Leu Asp Ser Ile Pro Asp Ile Pro Thr Pro Lys Pro Glu Lys Thr Leu  
 565 570 575  
 Thr Leu Gly Lys Gly Asn Gly Leu Leu Ser Gly Leu Leu Asn Ala Asp  
 580 585 590  
 Gly Asn Val Ser Leu Pro Lys Ala Gly Glu Thr Ile Lys Glu His Trp  
 595 600 605  
 Leu Pro Ile Ser Val Ile Val Gly Ala Met Gly Val Leu Met Ile Trp  
 610 615 620  
 Leu Ser Arg Arg Asn Lys Leu Lys Asn Lys Ala  
 625 630 635

<210> SEQ ID NO 21  
 <211> LENGTH: 953  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.  
 <400> SEQUENCE: 21

Met Asn Asn Lys Lys Thr Ala Thr Asn Arg Lys Gly Met Ile Pro Asn  
 1 5 10 15  
 Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Ser Val Gly Thr Ala Ser  
 20 25 30  
 Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Ser Gly His Glu Ala  
 35 40 45  
 Lys Ala Ala Glu His Thr Asn Gly Glu Leu Asn Gln Ser Lys Asn Glu  
 50 55 60  
 Thr Thr Ala Pro Ser Glu Asn Lys Thr Thr Glu Lys Val Asp Ser Arg  
 65 70 75 80  
 Gln Leu Lys Asp Asn Thr Gln Thr Ala Thr Ala Asp Gln Pro Lys Val  
 85 90 95  
 Thr Met Ser Asp Ser Ala Thr Val Lys Glu Thr Ser Ser Asn Met Gln  
 100 105 110  
 Ser Pro Gln Asn Ala Thr Ala Ser Gln Ser Thr Thr Gln Thr Ser Asn  
 115 120 125  
 Val Thr Thr Asn Asp Lys Ser Ser Thr Thr Tyr Ser Asn Glu Thr Asp  
 130 135 140  
 Lys Ser Asn Leu Thr Gln Ala Lys Asn Val Ser Thr Thr Pro Lys Thr  
 145 150 155 160  
 Thr Thr Ile Lys Gln Arg Ala Leu Asn Arg Met Ala Val Asn Thr Val  
 165 170 175  
 Ala Ala Pro Gln Gln Gly Thr Asn Val Asn Asp Lys Val His Phe Thr  
 180 185 190  
 Asn Ile Asp Ile Ala Ile Asp Lys Gly His Val Asn Lys Thr Thr Gly  
 195 200 205  
 Asn Thr Glu Phe Trp Ala Thr Ser Ser Asp Val Leu Lys Leu Lys Ala  
 210 215 220  
 Asn Tyr Thr Ile Asp Asp Ser Val Lys Glu Gly Asp Thr Phe Thr Phe  
 225 230 235 240  
 Lys Tyr Gly Gln Tyr Phe Arg Pro Gly Ser Val Arg Leu Pro Ser Gln  
 245 250 255  
 Thr Gln Asn Leu Tyr Asn Ala Gln Gly Asn Ile Ile Ala Lys Gly Ile  
 260 265 270

-continued

---

Tyr Asp Ser Lys Thr Asn Thr Thr Thr Tyr Thr Phe Thr Asn Tyr Val  
 275 280 285  
 Asp Gln Tyr Thr Asn Val Ser Gly Ser Phe Glu Gln Val Ala Phe Ala  
 290 295 300  
 Lys Arg Glu Asn Ala Thr Thr Asp Lys Thr Ala Tyr Lys Met Glu Val  
 305 310 315 320  
 Thr Leu Gly Asn Asp Thr Tyr Ser Lys Asp Val Ile Val Asp Tyr Gly  
 325 330 335  
 Asn Gln Lys Gly Gln Gln Leu Ile Ser Ser Thr Asn Tyr Ile Asn Asn  
 340 345 350  
 Glu Asp Leu Ser Arg Asn Met Thr Val Tyr Val Asn Gln Pro Lys Lys  
 355 360 365  
 Thr Tyr Thr Lys Glu Thr Phe Val Thr Asn Leu Thr Gly Tyr Lys Phe  
 370 375 380  
 Asn Pro Asp Ala Lys Asn Phe Lys Ile Tyr Glu Val Thr Asp Gln Asn  
 385 390 395 400  
 Gln Phe Val Asp Ser Phe Thr Pro Asp Thr Ser Lys Leu Lys Asp Val  
 405 410 415  
 Thr Gly Gln Phe Asp Val Ile Tyr Ser Asn Asp Asn Lys Thr Ala Thr  
 420 425 430  
 Val Asp Leu Leu Asn Gly Gln Ser Ser Ser Asp Lys Gln Tyr Ile Ile  
 435 440 445  
 Gln Gln Val Ala Tyr Pro Asp Asn Ser Ser Thr Asp Asn Gly Lys Ile  
 450 455 460  
 Asp Tyr Thr Leu Glu Thr Gln Asn Gly Lys Ser Ser Trp Ser Asn Ser  
 465 470 475 480  
 Tyr Ser Asn Val Asn Gly Ser Ser Thr Ala Asn Gly Asp Gln Lys Lys  
 485 490 495  
 Tyr Asn Leu Gly Asp Tyr Val Trp Glu Asp Thr Asn Lys Asp Gly Lys  
 500 505 510  
 Gln Asp Ala Asn Glu Lys Gly Ile Lys Gly Val Tyr Val Ile Leu Lys  
 515 520 525  
 Asp Ser Asn Gly Lys Glu Leu Asp Arg Thr Thr Thr Asp Glu Asn Gly  
 530 535 540  
 Lys Tyr Gln Phe Thr Gly Leu Ser Asn Gly Thr Tyr Ser Val Glu Phe  
 545 550 555 560  
 Ser Thr Pro Ala Gly Tyr Thr Pro Thr Thr Ala Asn Ala Gly Thr Asp  
 565 570 575  
 Asp Ala Val Asp Ser Asp Gly Leu Thr Thr Thr Gly Val Ile Lys Asp  
 580 585 590  
 Ala Asp Asn Met Thr Leu Asp Ser Gly Phe Tyr Lys Thr Pro Lys Tyr  
 595 600 605  
 Ser Leu Gly Asp Tyr Val Trp Tyr Asp Ser Asn Lys Asp Gly Lys Gln  
 610 615 620  
 Asp Ser Thr Glu Lys Gly Ile Lys Gly Val Lys Val Thr Leu Gln Asn  
 625 630 635 640  
 Glu Lys Gly Glu Val Ile Gly Thr Thr Glu Thr Asp Glu Asn Gly Lys  
 645 650 655  
 Tyr Arg Phe Asp Asn Leu Asp Ser Gly Lys Tyr Lys Val Ile Phe Glu  
 660 665 670  
 Lys Pro Ala Gly Leu Thr Gln Thr Gly Thr Asn Thr Thr Glu Asp Asp



-continued

---

```

        675             680             685
Lys Asp Ala Asp Gly Gly Glu Val Asp Val Thr Ile Thr Asp His Asp
 690          695
Asp Phe Thr Leu Asp Asn Gly Tyr Tyr Glu Glu Thr Ser Asp Ser
 705          710          715          720
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 725          730          735
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 740          745          750
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 755          760          765
Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser
 770          775          780
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 785          790          795          800
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 805          810          815
Asp Ser Asp Ser Asp Ser Asp Asn Asp Ser Asp Ser Asp Ser Asp Ser
 820          825          830
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 835          840          845
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 850          855          860
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 865          870          875          880
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ala Gly Lys
 885          890          895
His Thr Pro Thr Lys Pro Met Ser Thr Val Lys Asp Gln His Lys Thr
 900          905          910
Ala Lys Ala Leu Pro Glu Thr Gly Ser Glu Asn Asn Asn Ser Asn Asn
 915          920          925
Gly Thr Leu Phe Gly Gly Leu Phe Ala Ala Leu Gly Ser Leu Leu Leu
 930          935          940
Phe Gly Arg Arg Lys Lys Gln Asn Lys
 945          950
<210> SEQ ID NO 22
<211> LENGTH: 989
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.
<400> SEQUENCE: 22
Met Asn Met Lys Lys Lys Glu Lys His Ala Ile Arg Lys Lys Ser Ile
 1          5          10          15
Gly Val Ala Ser Val Leu Val Gly Thr Leu Ile Gly Phe Gly Leu Leu
 20          25          30
Ser Ser Lys Glu Ala Asp Ala Ser Glu Asn Ser Val Thr Gln Ser Asp
 35          40          45
Ser Ala Ser Asn Glu Ser Lys Ser Asn Asp Ser Ser Ser Val Ser Ala
 50          55          60
Ala Pro Lys Thr Asp Asp Thr Asn Val Ser Asp Thr Lys Thr Ser Ser
 65          70          75          80
Asn Thr Asn Asn Gly Glu Thr Ser Val Ala Gln Asn Pro Ala Gln Gln

```

-continued

85				90				95							
Glu	Thr	Thr	Gln	Ser	Ser	Ser	Thr	Asn	Ala	Thr	Thr	Glu	Glu	Thr	Pro
			100					105				110			
Val	Thr	Gly	Glu	Ala	Thr	Thr	Thr	Thr	Thr	Asn	Gln	Ala	Asn	Thr	Pro
		115					120					125			
Ala	Thr	Thr	Gln	Ser	Ser	Asn	Thr	Asn	Ala	Glu	Glu	Leu	Val	Asn	Gln
		130				135					140				
Thr	Ser	Asn	Glu	Thr	Thr	Ser	Asn	Asp	Thr	Asn	Thr	Val	Ser	Ser	Val
				145		150				155					160
Asn	Ser	Pro	Gln	Asn	Ser	Thr	Asn	Ala	Glu	Asn	Val	Ser	Thr	Thr	Gln
			165					170						175	
Asp	Thr	Ser	Thr	Glu	Ala	Thr	Pro	Ser	Asn	Asn	Glu	Ser	Ala	Pro	Gln
			180					185					190		
Asn	Thr	Asp	Ala	Ser	Asn	Lys	Asp	Val	Val	Ser	Gln	Ala	Val	Asn	Pro
		195					200					205			
Ser	Thr	Pro	Arg	Met	Arg	Ala	Phe	Ser	Leu	Ala	Ala	Val	Ala	Ala	Asp
		210				215				220					
Ala	Pro	Ala	Ala	Gly	Thr	Asp	Ile	Thr	Asn	Gln	Leu	Thr	Asp	Val	Lys
		225			230					235					240
Val	Thr	Ile	Asp	Ser	Gly	Thr	Thr	Val	Tyr	Pro	His	Gln	Ala	Gly	Tyr
			245					250						255	
Val	Lys	Leu	Asn	Tyr	Gly	Phe	Ser	Val	Pro	Asn	Ser	Ala	Val	Lys	Gly
		260						265					270		
Asp	Thr	Phe	Lys	Ile	Thr	Val	Pro	Lys	Glu	Leu	Asn	Leu	Asn	Gly	Val
		275				280					285				
Thr	Ser	Thr	Ala	Lys	Val	Pro	Pro	Ile	Met	Ala	Gly	Asp	Gln	Val	Leu
		290				295					300				
Ala	Asn	Gly	Val	Ile	Asp	Ser	Asp	Gly	Asn	Val	Ile	Tyr	Thr	Phe	Thr
		305			310					315					320
Asp	Tyr	Val	Asp	Asn	Lys	Glu	Asn	Val	Thr	Ala	Asn	Ile	Thr	Met	Pro
			325					330						335	
Ala	Tyr	Ile	Asp	Pro	Glu	Asn	Val	Thr	Lys	Thr	Gly	Asn	Val	Thr	Leu
			340					345					350		
Thr	Thr	Gly	Ile	Gly	Thr	Asn	Thr	Ala	Ser	Lys	Thr	Val	Leu	Ile	Asp
		355					360					365			
Tyr	Glu	Lys	Tyr	Gly	Gln	Phe	His	Asn	Leu	Ser	Ile	Lys	Gly	Thr	Ile
		370				375					380				
Asp	Gln	Ile	Asp	Lys	Thr	Asn	Asn	Thr	Tyr	Arg	Gln	Thr	Ile	Tyr	Val
		385			390					395					400
Asn	Pro	Ser	Gly	Asp	Asn	Val	Val	Leu	Pro	Ala	Leu	Thr	Gly	Asn	Leu
			405					410						415	
Ile	Pro	Asn	Thr	Lys	Ser	Asn	Ala	Leu	Ile	Asp	Ala	Lys	Asn	Thr	Asp
			420					425					430		
Ile	Lys	Val	Tyr	Arg	Val	Asp	Asn	Ala	Asn	Asp	Leu	Ser	Glu	Ser	Tyr
		435					440					445			
Tyr	Val	Asn	Pro	Ser	Asp	Phe	Glu	Asp	Val	Thr	Asn	Gln	Val	Arg	Ile
		450				455					460				
Ser	Phe	Pro	Asn	Ala	Asn	Gln	Tyr	Lys	Val	Glu	Phe	Pro	Thr	Asp	Asp
		465			470					475					480
Asp	Gln	Ile	Thr	Thr	Pro	Tyr	Ile	Val	Val	Val	Asn	Gly	His	Ile	Asp
			485					490						495	

-continued

---

Pro	Ala	Ser	Thr	Gly	Asp	Leu	Ala	Leu	Arg	Ser	Thr	Phe	Tyr	Gly	Tyr
			500					505					510		
Asp	Ser	Asn	Phe	Ile	Trp	Arg	Ser	Met	Ser	Trp	Asp	Asn	Glu	Val	Ala
		515					520					525			
Phe	Asn	Asn	Gly	Ser	Gly	Ser	Gly	Asp	Gly	Ile	Asp	Lys	Pro	Val	Val
	530					535					540				
Pro	Glu	Gln	Pro	Asp	Glu	Pro	Gly	Glu	Ile	Glu	Pro	Ile	Pro	Glu	Asp
545					550					555					560
Ser	Asp	Ser	Asp	Pro	Gly	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asn	Ser	Asp
				565					570					575	
Ser	Gly	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Thr	Ser	Asp	Ser	Gly	Ser	Asp
			580					585					590		
Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp
		595					600					605			
Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala
	610					615					620				
Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp
625					630					635					640
Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp
				645					650					655	
Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Asp
			660					665					670		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		675				680						685			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	690					695					700				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
705					710					715					720
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
				725					730					735	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			740					745					750		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		755					760					765			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	770					775					780				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
785					790					795					800
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
				805					810					815	
Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu
				820					825					830	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		835					840					845			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	850						855					860			
Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
865					870					875					880
Ser	Ala	Ser	Asp	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asp	Ser	Ser	Ser	Asp
				885					890						895
Ser	Asp	Ser	Asp	Ser	Thr	Ser	Asp	Thr	Gly	Ser	Asp	Asn	Asp	Ser	Asp
			900					905						910	

-continued

---

Ser Asp Ser Asn Ser Asp Ser Glu Ser Gly Ser Asn Asn Asn Val Val  
 915 920 925

Pro Pro Asn Ser Pro Lys Asn Gly Thr Asn Ala Ser Asn Lys Asn Glu  
 930 935 940

Ala Lys Asp Ser Lys Glu Pro Leu Pro Asp Thr Gly Ser Glu Asp Glu  
 945 950 955 960

Ala Asn Thr Ser Leu Ile Trp Gly Leu Leu Ala Ser Leu Gly Ser Leu  
 965 970 975

Leu Leu Phe Arg Arg Lys Lys Glu Asn Lys Asp Lys Lys  
 980 985

<210> SEQ ID NO 23  
 <211> LENGTH: 584  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 23

Met Lys Phe Lys Ser Leu Ile Thr Thr Thr Leu Ala Leu Gly Val Leu  
 1 5 10 15

Ala Ser Thr Gly Ala Asn Phe Asn Asn Asn Glu Ala Ser Ala Ala Ala  
 20 25 30

Lys Pro Leu Asp Lys Ser Ser Ser Ser Leu His His Gly Tyr Ser Lys  
 35 40 45

Val His Val Pro Tyr Ala Ile Thr Val Asn Gly Thr Ser Gln Asn Ile  
 50 55 60

Leu Ser Ser Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp  
 65 70 75 80

Leu Glu Asp Arg Val Lys Ser Val Leu Lys Ser Asp Arg Gly Ile Ser  
 85 90 95

Asp Ile Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Tyr Phe  
 100 105 110

Lys Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ala Gly Ile Tyr Thr  
 115 120 125

Ala Asp Leu Ile Asn Thr Ser Glu Ile Lys Ala Ile Asn Ile Asn Val  
 130 135 140

Asp Thr Lys Lys Gln Val Glu Asp Lys Lys Lys Asp Lys Ala Asn Tyr  
 145 150 155 160

Gln Val Pro Tyr Thr Ile Thr Val Asn Gly Thr Ser Gln Asn Ile Leu  
 165 170 175

Ser Asn Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp Leu  
 180 185 190

Glu Asp Lys Val Lys Ser Val Leu Glu Ser Asn Arg Gly Ile Thr Asp  
 195 200 205

Val Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Asn Phe Lys  
 210 215 220

Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ser Gly Ile Tyr Thr Ala  
 225 230 235 240

Asn Leu Ile Asn Ser Ser Asp Ile Lys Ser Ile Asn Ile Asn Val Asp  
 245 250 255

Thr Lys Lys His Ile Glu Asn Lys Ala Lys Arg Asn Tyr Gln Val Pro  
 260 265 270

Tyr Ser Ile Asn Leu Asn Gly Thr Ser Thr Asn Ile Leu Ser Asn Leu  
 275 280 285

-continued

---

Ser Phe Ser Asn Lys Pro Trp Thr Asn Tyr Lys Asn Leu Thr Ser Gln  
 290 295 300  
 Ile Lys Ser Val Leu Lys His Asp Arg Gly Ile Ser Glu Gln Asp Leu  
 305 310 315 320  
 Lys Tyr Ala Lys Lys Ala Tyr Tyr Thr Val Tyr Phe Lys Asn Gly Gly  
 325 330 335  
 Lys Arg Ile Leu Gln Leu Asn Ser Lys Asn Tyr Thr Ala Asn Leu Val  
 340 345 350  
 His Ala Lys Asp Val Lys Arg Ile Glu Ile Thr Val Lys Thr Gly Thr  
 355 360 365  
 Lys Ala Lys Ala Asp Arg Tyr Val Pro Tyr Thr Ile Ala Val Asn Gly  
 370 375 380  
 Thr Ser Thr Pro Ile Leu Ser Asp Leu Lys Phe Thr Gly Asp Pro Arg  
 385 390 395 400  
 Val Gly Tyr Lys Asp Ile Ser Lys Lys Val Lys Ser Val Leu Lys His  
 405 410 415  
 Asp Arg Gly Ile Gly Glu Arg Glu Leu Lys Tyr Ala Lys Lys Ala Thr  
 420 425 430  
 Tyr Thr Val His Phe Lys Asn Gly Thr Lys Lys Val Ile Asn Ile Asn  
 435 440 445  
 Ser Asn Ile Ser Gln Leu Asn Leu Leu Tyr Val Gln Asp Ile Lys Lys  
 450 455 460  
 Ile Asp Ile Asp Val Lys Thr Gly Thr Lys Ala Lys Ala Asp Ser Tyr  
 465 470 475 480  
 Val Pro Tyr Thr Ile Ala Val Asn Gly Thr Ser Thr Pro Ile Leu Ser  
 485 490 495  
 Lys Leu Lys Ile Ser Asn Lys Gln Leu Ile Ser Tyr Lys Tyr Leu Asn  
 500 505 510  
 Asp Lys Val Lys Ser Val Leu Lys Ser Glu Arg Gly Ile Ser Asp Leu  
 515 520 525  
 Asp Leu Lys Phe Ala Lys Gln Ala Lys Tyr Thr Val Tyr Phe Lys Asn  
 530 535 540  
 Gly Lys Lys Gln Val Val Asn Leu Lys Ser Asp Ile Phe Thr Pro Asn  
 545 550 555 560  
 Leu Phe Ser Ala Lys Asp Ile Lys Lys Ile Asp Ile Asp Val Lys Gln  
 565 570 575  
 Tyr Thr Lys Ser Lys Lys Asn Lys  
 580

&lt;210&gt; SEQ ID NO 24

&lt;211&gt; LENGTH: 10419

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 24

Met Asn Tyr Arg Asp Lys Ile Gln Lys Phe Ser Ile Arg Lys Tyr Thr  
 1 5 10 15  
 Val Gly Thr Phe Ser Thr Val Ile Ala Thr Leu Val Phe Leu Gly Phe  
 20 25 30  
 Asn Thr Ser Gln Ala His Ala Ala Glu Thr Asn Gln Pro Ala Ser Val  
 35 40 45  
 Val Lys Gln Lys Gln Gln Ser Asn Asn Glu Gln Thr Glu Asn Arg Glu  
 50 55 60

-continued

---

Ser Gln Val Gln Asn Ser Gln Asn Ser Gln Asn Gly Gln Ser Leu Ser  
 65 70 75 80  
 Ala Thr His Glu Asn Glu Gln Pro Asn Ile Ser Gln Ala Asn Leu Val  
 85 90 95  
 Asp Gln Lys Val Ala Gln Ser Ser Thr Thr Asn Asp Glu Gln Pro Ala  
 100 105 110  
 Ser Gln Asn Val Asn Thr Lys Lys Asp Ser Ala Thr Ala Ala Thr Thr  
 115 120 125  
 Gln Pro Asp Lys Glu Gln Ser Lys His Lys Gln Asn Glu Ser Gln Ser  
 130 135 140  
 Ala Asn Lys Asn Gly Asn Asp Asn Arg Ala Ala His Val Glu Asn His  
 145 150 155 160  
 Glu Ala Asn Val Val Thr Ala Ser Asp Ser Ser Asp Asn Gly Asn Val  
 165 170 175  
 Gln His Asp Arg Asn Glu Leu Gln Ala Phe Phe Asp Ala Asn Tyr His  
 180 185 190  
 Asp Tyr Arg Phe Ile Asp Arg Glu Asn Ala Asp Ser Gly Thr Phe Asn  
 195 200 205  
 Tyr Val Lys Gly Ile Phe Asp Lys Ile Asn Thr Leu Leu Gly Ser Asn  
 210 215 220  
 Asp Pro Ile Asn Asn Lys Asp Leu Gln Leu Ala Tyr Lys Glu Leu Glu  
 225 230 235 240  
 Gln Ala Val Ala Leu Ile Arg Thr Met Pro Gln Arg Gln Gln Thr Ser  
 245 250 255  
 Arg Arg Ser Asn Arg Ile Gln Thr Arg Ser Val Glu Ser Arg Ala Ala  
 260 265 270  
 Glu Pro Arg Ser Val Ser Asp Tyr Gln Asn Ala Asn Ser Ser Tyr Tyr  
 275 280 285  
 Val Glu Asn Ala Asn Asp Gly Ser Gly Tyr Pro Val Gly Thr Tyr Ile  
 290 295 300  
 Asn Ala Ser Ser Lys Gly Ala Pro Tyr Asn Leu Pro Thr Thr Pro Trp  
 305 310 315 320  
 Asn Thr Leu Lys Ala Ser Asp Ser Lys Glu Ile Ala Leu Met Thr Ala  
 325 330 335  
 Lys Gln Thr Gly Asp Gly Tyr Gln Trp Val Ile Lys Phe Asn Lys Gly  
 340 345 350  
 His Ala Pro His Gln Asn Met Ile Phe Trp Phe Ala Leu Pro Ala Asp  
 355 360 365  
 Gln Val Pro Val Gly Arg Thr Asp Phe Val Thr Val Asn Ser Asp Gly  
 370 375 380  
 Thr Asn Val Gln Trp Ser His Gly Ala Gly Ala Gly Ala Asn Lys Pro  
 385 390 395 400  
 Leu Gln Gln Met Trp Glu Tyr Gly Val Asn Asp Pro His Arg Ser His  
 405 410 415  
 Asp Phe Lys Ile Arg Asn Arg Ser Gly Gln Val Ile Tyr Asp Trp Pro  
 420 425 430  
 Thr Val His Ile Tyr Ser Leu Glu Asp Leu Ser Arg Ala Ser Asp Tyr  
 435 440 445  
 Phe Ser Glu Ala Gly Ala Thr Pro Ala Thr Lys Ala Phe Gly Arg Gln  
 450 455 460  
 Asn Phe Glu Tyr Ile Asn Gly Gln Lys Pro Ala Glu Ser Pro Gly Val

-continued

---

465		470		475		480									
Pro	Lys	Val	Tyr	Thr	Phe	Ile	Gly	Gln	Gly	Asp	Ala	Ser	Tyr	Thr	Ile
			485						490					495	
Ser	Phe	Lys	Thr	Gln	Gly	Pro	Thr	Val	Asn	Lys	Leu	Tyr	Tyr	Ala	Ala
		500						505					510		
Gly	Gly	Arg	Ala	Leu	Glu	Tyr	Asn	Gln	Leu	Phe	Met	Tyr	Ser	Gln	Leu
		515					520					525			
Tyr	Val	Glu	Ser	Thr	Gln	Asp	His	Gln	Gln	Arg	Leu	Asn	Gly	Leu	Arg
	530					535					540				
Gln	Val	Val	Asn	Arg	Thr	Tyr	Arg	Ile	Gly	Thr	Thr	Lys	Arg	Val	Glu
545					550					555					560
Val	Ser	Gln	Gly	Asn	Val	Gln	Thr	Lys	Lys	Val	Leu	Glu	Ser	Thr	Asn
				565					570					575	
Leu	Asn	Ile	Asp	Asp	Phe	Val	Asp	Asp	Pro	Leu	Ser	Tyr	Val	Lys	Thr
			580					585					590		
Pro	Ser	Asn	Lys	Val	Leu	Gly	Phe	Tyr	Ser	Asn	Asn	Ala	Asn	Thr	Asn
		595					600					605			
Ala	Phe	Arg	Pro	Gly	Gly	Ala	Gln	Gln	Leu	Asn	Glu	Tyr	Gln	Leu	Ser
	610					615					620				
Gln	Leu	Phe	Thr	Asp	Gln	Lys	Leu	Gln	Glu	Ala	Ala	Arg	Thr	Arg	Asn
625					630					635					640
Pro	Ile	Arg	Leu	Met	Ile	Gly	Phe	Asp	Tyr	Pro	Asp	Ala	Tyr	Gly	Asn
			645						650					655	
Ser	Glu	Thr	Leu	Val	Pro	Val	Asn	Leu	Thr	Val	Leu	Pro	Glu	Ile	Gln
			660					665					670		
His	Asn	Ile	Lys	Phe	Phe	Lys	Asn	Asp	Asp	Thr	Gln	Asn	Ile	Ala	Glu
	675						680					685			
Lys	Pro	Phe	Ser	Lys	Gln	Ala	Gly	His	Pro	Val	Phe	Tyr	Val	Tyr	Ala
	690					695					700				
Gly	Asn	Gln	Gly	Asn	Ala	Ser	Val	Asn	Leu	Gly	Gly	Ser	Val	Thr	Ser
705					710					715					720
Ile	Gln	Pro	Leu	Arg	Ile	Asn	Leu	Thr	Ser	Asn	Glu	Asn	Phe	Thr	Asp
			725						730					735	
Lys	Asp	Trp	Gln	Ile	Thr	Gly	Ile	Pro	Arg	Thr	Leu	His	Ile	Glu	Asn
		740						745					750		
Ser	Thr	Asn	Arg	Pro	Asn	Asn	Ala	Arg	Glu	Arg	Asn	Ile	Glu	Leu	Val
		755					760					765			
Gly	Asn	Leu	Leu	Pro	Gly	Asp	Tyr	Phe	Gly	Thr	Ile	Arg	Phe	Gly	Arg
	770					775					780				
Lys	Glu	Gln	Leu	Phe	Glu	Ile	Arg	Val	Lys	Pro	His	Thr	Pro	Thr	Ile
785					790					795					800
Thr	Thr	Thr	Ala	Glu	Gln	Leu	Arg	Gly	Thr	Ala	Leu	Gln	Lys	Val	Pro
			805						810					815	
Val	Asn	Ile	Ser	Gly	Ile	Pro	Leu	Asp	Pro	Ser	Ala	Leu	Val	Tyr	Leu
			820					825					830		
Val	Ala	Pro	Thr	Asn	Gln	Thr	Thr	Asn	Gly	Gly	Ser	Glu	Ala	Asp	Gln
		835					840					845			
Ile	Pro	Ser	Gly	Tyr	Thr	Ile	Leu	Ala	Thr	Gly	Thr	Pro	Asp	Gly	Val
	850					855						860			
His	Asn	Thr	Ile	Thr	Ile	Arg	Pro	Gln	Asp	Tyr	Val	Val	Phe	Ile	Pro
865					870					875					880

-continued

---

Pro	Val	Gly	Lys	Gln	Ile	Arg	Ala	Val	Val	Tyr	Tyr	Asn	Lys	Val	Val
				885					890					895	
Ala	Ser	Asn	Met	Ser	Asn	Ala	Val	Thr	Ile	Leu	Pro	Asp	Asp	Ile	Pro
			900					905					910		
Pro	Thr	Ile	Asn	Asn	Pro	Val	Gly	Ile	Asn	Ala	Lys	Tyr	Tyr	Arg	Gly
		915					920					925			
Asp	Glu	Val	Asn	Phe	Thr	Met	Gly	Val	Ser	Asp	Arg	His	Ser	Gly	Ile
	930					935					940				
Lys	Asn	Thr	Thr	Ile	Thr	Thr	Leu	Pro	Asn	Gly	Trp	Thr	Ser	Asn	Leu
945					950					955					960
Thr	Lys	Ala	Asp	Lys	Asn	Asn	Gly	Ser	Leu	Ser	Ile	Thr	Gly	Arg	Val
				965					970					975	
Ser	Met	Asn	Gln	Ala	Phe	Asn	Ser	Asp	Ile	Thr	Phe	Lys	Val	Ser	Ala
			980					985					990		
Thr	Asp	Asn	Val	Asn	Asn	Thr	Thr	Asn	Asp	Ser	Gln	Ser	Lys	His	Val
		995					1000					1005			
Ser	Ile	His	Val	Gly	Lys	Ile	Ser	Glu	Asp	Ala	His	Pro	Ile	Val	
	1010					1015					1020				
Leu	Gly	Asn	Thr	Glu	Lys	Val	Val	Val	Val	Asn	Pro	Thr	Ala	Val	
	1025					1030					1035				
Ser	Asn	Asp	Glu	Lys	Gln	Ser	Ile	Ile	Thr	Ala	Phe	Met	Asn	Lys	
	1040					1045					1050				
Asn	Gln	Asn	Ile	Arg	Gly	Tyr	Leu	Ala	Ser	Thr	Asp	Pro	Val	Thr	
	1055					1060					1065				
Val	Asp	Asn	Asn	Gly	Asn	Val	Thr	Leu	His	Tyr	Arg	Asp	Gly	Ser	
	1070					1075					1080				
Ser	Thr	Thr	Leu	Asp	Ala	Thr	Asn	Val	Met	Thr	Tyr	Glu	Pro	Val	
	1085					1090					1095				
Val	Lys	Pro	Glu	Tyr	Gln	Thr	Val	Asn	Ala	Ala	Lys	Thr	Ala	Thr	
	1100					1105					1110				
Val	Thr	Ile	Ala	Lys	Gly	Gln	Ser	Phe	Ser	Ile	Gly	Asp	Ile	Lys	
	1115					1120					1125				
Gln	Tyr	Phe	Thr	Leu	Ser	Asn	Gly	Gln	Pro	Ile	Pro	Ser	Gly	Thr	
	1130					1135					1140				
Phe	Thr	Asn	Ile	Thr	Ser	Asp	Arg	Thr	Ile	Pro	Thr	Ala	Gln	Glu	
	1145					1150					1155				
Val	Ser	Gln	Met	Asn	Ala	Gly	Thr	Gln	Leu	Tyr	His	Ile	Thr	Ala	
	1160					1165					1170				
Thr	Asn	Ala	Tyr	His	Lys	Asp	Ser	Glu	Asp	Phe	Tyr	Ile	Ser	Leu	
	1175					1180					1185				
Lys	Ile	Ile	Asp	Val	Lys	Gln	Pro	Glu	Gly	Asp	Gln	Arg	Val	Tyr	
	1190					1195					1200				
Arg	Thr	Ser	Thr	Tyr	Asp	Leu	Thr	Thr	Asp	Glu	Ile	Ser	Lys	Val	
	1205					1210					1215				
Lys	Gln	Ala	Phe	Ile	Asn	Ala	Asn	Arg	Asp	Val	Ile	Thr	Leu	Ala	
	1220					1225					1230				
Glu	Gly	Asp	Ile	Ser	Val	Thr	Asn	Thr	Pro	Asn	Gly	Ala	Asn	Val	
	1235					1240					1245				
Ser	Thr	Ile	Thr	Val	Asn	Ile	Asn	Lys	Gly	Arg	Leu	Thr	Lys	Ser	
	1250					1255					1260				
Phe	Ala	Ser	Asn	Leu	Ala	Asn	Met	Asn	Phe	Leu	Arg	Trp	Val	Asn	
	1265					1270					1275				



-continued

---

Phe	Pro	Gln	Asp	Tyr	Thr	Val	Thr	Trp	Thr	Asn	Ala	Lys	Ile	Ala
1280						1285					1290			
Asn	Arg	Pro	Thr	Asp	Gly	Gly	Leu	Ser	Trp	Ser	Asp	Asp	His	Lys
1295						1300					1305			
Ser	Leu	Ile	Tyr	Arg	Tyr	Asp	Ala	Thr	Leu	Gly	Thr	Gln	Ile	Thr
1310						1315					1320			
Thr	Asn	Asp	Ile	Leu	Thr	Met	Leu	Lys	Ala	Thr	Thr	Thr	Val	Pro
1325						1330					1335			
Gly	Leu	Arg	Asn	Asn	Ile	Thr	Gly	Asn	Glu	Lys	Ser	Gln	Ala	Glu
1340						1345					1350			
Ala	Gly	Gly	Arg	Pro	Asn	Phe	Arg	Thr	Thr	Gly	Tyr	Ser	Gln	Ser
1355						1360					1365			
Asn	Ala	Thr	Thr	Asp	Gly	Gln	Arg	Gln	Phe	Thr	Leu	Asn	Gly	Gln
1370						1375					1380			
Val	Ile	Gln	Val	Leu	Asp	Ile	Ile	Asn	Pro	Ser	Asn	Gly	Tyr	Gly
1385						1390					1395			
Gly	Gln	Pro	Val	Thr	Asn	Ser	Asn	Thr	Arg	Ala	Asn	His	Ser	Asn
1400						1405					1410			
Ser	Thr	Val	Val	Asn	Val	Asn	Glu	Pro	Ala	Ala	Asn	Gly	Ala	Gly
1415						1420					1425			
Ala	Phe	Thr	Ile	Asp	His	Val	Val	Lys	Ser	Asn	Ser	Thr	His	Asn
1430						1435					1440			
Ala	Ser	Asp	Ala	Val	Tyr	Lys	Ala	Gln	Leu	Tyr	Leu	Thr	Pro	Tyr
1445						1450					1455			
Gly	Pro	Lys	Gln	Tyr	Val	Glu	His	Leu	Asn	Gln	Asn	Thr	Gly	Asn
1460						1465					1470			
Thr	Thr	Asp	Ala	Ile	Asn	Ile	Tyr	Phe	Val	Pro	Ser	Asp	Leu	Val
1475						1480					1485			
Asn	Pro	Thr	Ile	Ser	Val	Gly	Asn	Tyr	Thr	Asn	His	Gln	Val	Phe
1490						1495					1500			
Ser	Gly	Glu	Thr	Phe	Thr	Asn	Thr	Ile	Thr	Ala	Asn	Asp	Asn	Phe
1505						1510					1515			
Gly	Val	Gln	Ser	Val	Thr	Val	Pro	Asn	Thr	Ser	Gln	Ile	Thr	Gly
1520						1525					1530			
Thr	Val	Asp	Asn	Asn	His	Gln	His	Val	Ser	Ala	Thr	Ala	Pro	Asn
1535						1540					1545			
Val	Thr	Ser	Ala	Thr	Asn	Lys	Thr	Ile	Asn	Leu	Leu	Ala	Thr	Asp
1550						1555					1560			
Thr	Ser	Gly	Asn	Thr	Ala	Thr	Thr	Ser	Phe	Asn	Val	Thr	Val	Lys
1565						1570					1575			
Pro	Leu	Arg	Asp	Lys	Tyr	Arg	Val	Gly	Thr	Ser	Ser	Thr	Ala	Ala
1580						1585					1590			
Asn	Pro	Val	Arg	Ile	Ala	Asn	Ile	Ser	Asn	Asn	Ala	Thr	Val	Ser
1595						1600					1605			
Gln	Ala	Asp	Gln	Thr	Thr	Ile	Ile	Asn	Ser	Leu	Thr	Phe	Thr	Glu
1610						1615					1620			
Thr	Val	Pro	Asn	Arg	Ser	Tyr	Ala	Arg	Ala	Ser	Ala	Asn	Glu	Ile
1625						1630					1635			
Thr	Ser	Lys	Thr	Val	Ser	Asn	Val	Ser	Arg	Thr	Gly	Asn	Asn	Ala
1640						1645					1650			
Asn	Val	Thr	Val	Thr	Val	Thr	Tyr	Gln	Asp	Gly	Thr	Thr	Ser	Thr

-continued

1655	1660	1665
Val Thr Val Pro Val Lys His Val Ile Pro Glu Ile Val Ala His 1670 1675 1680		
Ser His Tyr Thr Val Gln Gly Gln Asp Phe Pro Ala Gly Asn Gly 1685 1690 1695		
Ser Ser Ala Ser Asp Tyr Phe Lys Leu Ser Asn Gly Ser Asp Ile 1700 1705 1710		
Ala Asp Ala Thr Ile Thr Trp Val Ser Gly Gln Ala Pro Asn Lys 1715 1720 1725		
Asp Asn Thr Arg Ile Gly Glu Asp Ile Thr Val Thr Ala His Ile 1730 1735 1740		
Leu Ile Asp Gly Glu Thr Thr Pro Ile Thr Lys Thr Ala Thr Tyr 1745 1750 1755		
Lys Val Val Arg Thr Val Pro Lys His Val Phe Glu Thr Ala Arg 1760 1765 1770		
Gly Val Leu Tyr Pro Gly Val Ser Asp Met Tyr Asp Ala Lys Gln 1775 1780 1785		
Tyr Val Lys Pro Val Asn Asn Ser Trp Ser Thr Asn Ala Gln His 1790 1795 1800		
Met Asn Phe Gln Phe Val Gly Thr Tyr Gly Pro Asn Lys Asp Val 1805 1810 1815		
Val Gly Ile Ser Thr Arg Leu Ile Arg Val Thr Tyr Asp Asn Arg 1820 1825 1830		
Gln Thr Glu Asp Leu Thr Ile Leu Ser Lys Val Lys Pro Asp Pro 1835 1840 1845		
Pro Arg Ile Asp Ala Asn Ser Val Thr Tyr Lys Ala Gly Leu Thr 1850 1855 1860		
Asn Gln Glu Ile Lys Val Asn Asn Val Leu Asn Asn Ser Ser Val 1865 1870 1875		
Lys Leu Phe Lys Ala Asp Asn Thr Pro Leu Asn Val Thr Asn Ile 1880 1885 1890		
Thr His Gly Ser Gly Phe Ser Ser Val Val Thr Val Ser Asp Ala 1895 1900 1905		
Leu Pro Asn Gly Gly Ile Lys Ala Lys Ser Ser Ile Ser Met Asn 1910 1915 1920		
Asn Val Thr Tyr Thr Thr Gln Asp Glu His Gly Gln Val Val Thr 1925 1930 1935		
Val Thr Arg Asn Glu Ser Val Asp Ser Asn Asp Ser Ala Thr Val 1940 1945 1950		
Thr Val Thr Pro Gln Leu Gln Ala Thr Thr Glu Gly Ala Val Phe 1955 1960 1965		
Ile Lys Gly Gly Asp Gly Phe Asp Phe Gly His Val Glu Arg Phe 1970 1975 1980		
Ile Gln Asn Pro Pro His Gly Ala Thr Val Ala Trp His Asp Ser 1985 1990 1995		
Pro Asp Thr Trp Lys Asn Thr Val Gly Asn Thr His Lys Thr Ala 2000 2005 2010		
Val Val Thr Leu Pro Asn Gly Gln Gly Thr Arg Asn Val Glu Val 2015 2020 2025		
Pro Val Lys Val Tyr Pro Val Ala Asn Ala Lys Ala Pro Ser Arg 2030 2035 2040		

-continued

Asp Val	Lys Gly	Gln Asn	Leu Thr	Asn Gly	Thr Asp	Ala Met	Asn	
2045			2050			2055		
Tyr Ile	Thr Phe	Asp Pro	Asn Thr	Asn Thr	Asn Gly	Ile Thr	Ala	
2060			2065			2070		
Ala Trp	Ala Asn	Arg Gln	Gln Pro	Asn Asn	Gln Gln	Ala Gly	Val	
2075			2080			2085		
Gln His	Leu Asn	Val Asp	Val Thr	Tyr Pro	Gly Ile	Ser Ala	Ala	
2090			2095			2100		
Lys Arg	Val Pro	Val Thr	Val Asn	Val Tyr	Gln Phe	Glu Phe	Pro	
2105			2110			2115		
Gln Thr	Thr Tyr	Thr Thr	Thr Val	Gly Gly	Thr Leu	Ala Ser	Gly	
2120			2125			2130		
Thr Gln	Ala Ser	Gly Tyr	Ala His	Met Gln	Asn Ala	Thr Gly	Leu	
2135			2140			2145		
Pro Thr	Asp Gly	Phe Thr	Tyr Lys	Trp Asn	Arg Asp	Thr Thr	Gly	
2150			2155			2160		
Thr Asn	Asp Ala	Asn Trp	Ser Ala	Met Asn	Lys Pro	Asn Val	Ala	
2165			2170			2175		
Lys Val	Val Asn	Ala Lys	Tyr Asp	Val Ile	Tyr Asn	Gly His	Thr	
2180			2185			2190		
Phe Ala	Thr Ser	Leu Pro	Ala Lys	Phe Val	Val Lys	Asp Val	Gln	
2195			2200			2205		
Pro Ala	Lys Pro	Thr Val	Thr Glu	Thr Ala	Ala Gly	Ala Ile	Thr	
2210			2215			2220		
Ile Ala	Pro Gly	Ala Asn	Gln Thr	Val Asn	Thr His	Ala Gly	Asn	
2225			2230			2235		
Val Thr	Thr Tyr	Ala Asp	Lys Leu	Val Ile	Lys Arg	Asn Gly	Asn	
2240			2245			2250		
Val Val	Thr Thr	Phe Thr	Arg Arg	Asn Asn	Thr Ser	Pro Trp	Val	
2255			2260			2265		
Lys Glu	Ala Ser	Ala Ala	Thr Val	Ala Gly	Ile Ala	Gly Thr	Asn	
2270			2275			2280		
Asn Gly	Ile Thr	Val Ala	Ala Ala	Gly Thr	Phe Asn	Pro Ala	Asp Thr	
2285			2290			2295		
Ile Gln	Val Val	Ala Thr	Gln Gly	Ser Gly	Glu Thr	Val Ser	Asp	
2300			2305			2310		
Glu Gln	Arg Ser	Asp Asp	Phe Thr	Val Val	Ala Pro	Gln Pro	Asn	
2315			2320			2325		
Gln Ala	Thr Thr	Lys Ile	Trp Gln	Asn Gly	His Ile	Asp Ile	Thr	
2330			2335			2340		
Pro Asn	Asn Pro	Ser Gly	His Leu	Ile Asn	Pro Thr	Gln Ala	Met	
2345			2350			2355		
Asp Ile	Ala Tyr	Thr Glu	Lys Val	Gly Asn	Gly Ala	Glu His	Ser	
2360			2365			2370		
Lys Thr	Ile Asn	Val Val	Arg Gly	Gln Asn	Asn Gln	Trp Thr	Ile	
2375			2380			2385		
Ala Asn	Lys Pro	Asp Tyr	Val Thr	Leu Asp	Ala Gln	Thr Gly	Lys	
2390			2395			2400		
Val Thr	Phe Asn	Ala Asn	Thr Ile	Lys Pro	Asn Ser	Ser Ile	Thr	
2405			2410			2415		
Ile Thr	Pro Lys	Ala Gly	Thr Gly	His Ser	Val Ser	Ser Asn	Pro	
2420			2425			2430		

-continued

---

Ser Thr 2435	Leu Thr Ala Pro 2440	Ala Ala His Thr Val 2445	Asn Thr Thr Glu 2445
Ile Val 2450	Lys Asp Tyr Gly Ser 2455	Asn Val Thr Ala Ala 2460	Glu Ile Asn 2460
Asn Ala 2465	Val Gln Val Ala Asn 2470	Lys Arg Thr Ala Thr 2475	Ile Lys Asn 2475
Gly Thr 2480	Ala Met Pro Thr Asn 2485	Leu Ala Gly Gly Ser 2490	Thr Thr Thr 2490
Ile Pro 2495	Val Thr Val Thr Tyr 2500	Asn Asp Gly Ser Thr 2505	Glu Glu Val 2505
Gln Glu 2510	Ser Ile Phe Thr Lys 2515	Ala Asp Lys Arg Glu 2520	Leu Ile Thr 2520
Ala Lys 2525	Asn His Leu Asp Asp 2530	Pro Val Ser Thr Glu 2535	Gly Lys Lys 2535
Pro Gly 2540	Thr Ile Thr Gln Tyr 2545	Asn Asn Ala Met His 2550	Asn Ala Gln 2550
Gln Gln 2555	Ile Asn Thr Ala Lys 2560	Thr Glu Ala Gln Gln 2565	Val Ile Asn 2565
Asn Glu 2570	Arg Ala Thr Pro Gln 2575	Gln Val Ser Asp Ala 2580	Leu Thr Lys 2580
Val Arg 2585	Ala Ala Gln Thr Lys 2590	Ile Asp Gln Ala Lys 2595	Ala Leu Leu 2595
Gln Asn 2600	Lys Glu Asp Asn Ser 2605	Gln Leu Val Thr Ser 2610	Lys Asn Asn 2610
Leu Gln 2615	Ser Ser Val Asn Gln 2620	Val Pro Ser Thr Ala 2625	Gly Met Thr 2625
Gln Gln 2630	Ser Ile Asp Asn Tyr 2635	Asn Ala Lys Lys Arg 2640	Glu Ala Glu 2640
Thr Glu 2645	Ile Thr Ala Ala Gln 2650	Arg Val Ile Asp Asn 2655	Gly Asp Ala 2655
Thr Ala 2660	Gln Gln Ile Ser Asp 2665	Glu Lys His Arg Val 2670	Asp Asn Ala 2670
Leu Thr 2675	Ala Leu Asn Gln Ala 2680	Lys His Asp Leu Thr 2685	Ala Asp Thr 2685
His Ala 2690	Leu Glu Gln Ala Val 2695	Gln Gln Leu Asn Arg 2700	Thr Gly Thr 2700
Thr Thr 2705	Gly Lys Lys Pro Ala 2710	Ser Ile Thr Ala Tyr 2715	Asn Asn Ser 2715
Ile Arg 2720	Ala Leu Gln Ser Asp 2725	Leu Thr Ser Ala Lys 2730	Asn Ser Ala 2730
Asn Ala 2735	Ile Ile Gln Lys Pro 2740	Ile Arg Thr Val Gln 2745	Glu Val Gln 2745
Ser Ala 2750	Leu Thr Asn Val Asn 2755	Arg Val Asn Glu Arg 2760	Leu Thr Gln 2760
Ala Ile 2765	Asn Gln Leu Val Pro 2770	Leu Ala Asp Asn Ser 2775	Ala Leu Lys 2775
Thr Ala 2780	Lys Thr Lys Leu Asp 2785	Glu Glu Ile Asn Lys 2790	Ser Val Thr 2790
Thr Asp 2795	Gly Met Thr Gln Ser 2800	Ser Ile Gln Ala Tyr 2805	Glu Asn Ala 2805
Lys Arg	Ala Gly Gln Thr Glu	Ser Thr Asn Ala Gln	Asn Val Ile

-continued

2810	2815	2820
Asn Asn Gly Asp Ala Thr 2825	Asp Gln Gln Ile Ala 2830	Ala Glu Lys Thr 2835
Lys Val Glu Glu Lys Tyr 2840	Asn Ser Leu Lys Gln 2845	Ala Ile Ala Gly 2850
Leu Thr Pro Asp Leu Ala 2855	Pro Leu Gln Thr Ala 2860	Lys Thr Gln Leu 2865
Gln Asn Asp Ile Asp Gln 2870	Pro Thr Ser Thr Thr 2875	Gly Met Thr Ser 2880
Ala Ser Ile Ala Ala Phe 2885	Asn Glu Lys Leu Ser 2890	Ala Ala Arg Thr 2895
Lys Ile Gln Glu Ile Asp 2900	Arg Val Leu Ala Ser 2905	His Pro Asp Val 2910
Ala Thr Ile Arg Gln Asn 2915	Val Thr Ala Ala Asn 2920	Ala Ala Lys Ser 2925
Ala Leu Asp Gln Ala Arg 2930	Asn Gly Leu Thr Val 2935	Asp Lys Ala Pro 2940
Leu Glu Asn Ala Lys Asn 2945	Gln Leu Gln His Ser 2950	Ile Asp Thr Gln 2955
Thr Ser Thr Thr Gly Met 2960	Thr Gln Asp Ser Ile 2965	Asn Ala Tyr Asn 2970
Ala Lys Leu Thr Ala Ala 2975	Arg Asn Lys Ile Gln 2980	Gln Ile Asn Gln 2985
Val Leu Ala Gly Ser Pro 2990	Thr Val Glu Gln Ile 2995	Asn Thr Asn Thr 3000
Ser Thr Ala Asn Gln Ala 3005	Lys Ser Asp Leu Asp 3010	His Ala Arg Gln 3015
Ala Leu Thr Pro Asp Lys 3020	Ala Pro Leu Gln Thr 3025	Ala Lys Thr Gln 3030
Leu Glu Gln Ser Ile Asn 3035	Gln Pro Thr Asp Thr 3040	Thr Thr Gly Met Thr 3045
Thr Ala Ser Leu Asn Ala 3050	Tyr Asn Gln Lys Leu 3055	Gln Ala Ala Arg 3060
Gln Lys Leu Thr Glu Ile 3065	Asn Gln Val Leu Asn 3070	Gly Asn Pro Thr 3075
Val Gln Asn Ile Asn Asp 3080	Lys Val Thr Glu Ala 3085	Asn Gln Ala Lys 3090
Asp Gln Leu Asn Thr Ala 3095	Arg Gln Gly Leu Thr 3100	Leu Asp Arg Gln 3105
Pro Ala Leu Thr Thr Leu 3110	His Gly Ala Ser Asn 3115	Leu Asn Gln Ala 3120
Gln Gln Asn Asn Phe Thr 3125	Gln Gln Ile Asn Ala 3130	Ala Gln Asn His 3135
Ala Ala Leu Glu Thr Ile 3140	Lys Ser Asn Ile Thr 3145	Ala Leu Asn Thr 3150
Ala Met Thr Lys Leu Lys 3155	Asp Ser Val Ala Asp 3160	Asn Asn Thr Ile 3165
Lys Ser Asp Gln Asn Tyr 3170	Thr Asp Ala Thr Pro 3175	Ala Asn Lys Gln 3180
Ala Tyr Asp Asn Ala Val 3185	Asn Ala Ala Lys Gly 3190	Val Ile Gly Glu 3195

-continued

Thr	Thr	Asn	Pro	Thr	Met	Asp	Val	Asn	Thr	Val	Asn	Gln	Lys	Ala
	3200					3205						3210		
Ala	Ser	Val	Lys	Ser	Thr	Lys	Asp	Ala	Leu	Asp	Gly	Gln	Gln	Asn
	3215					3220					3225			
Leu	Gln	Arg	Ala	Lys	Thr	Glu	Ala	Thr	Asn	Ala	Ile	Thr	His	Ala
	3230					3235					3240			
Ser	Asp	Leu	Asn	Gln	Ala	Gln	Lys	Asn	Ala	Leu	Thr	Gln	Gln	Val
	3245					3250					3255			
Asn	Ser	Ala	Gln	Asn	Val	Gln	Ala	Val	Asn	Asp	Ile	Lys	Gln	Thr
	3260					3265					3270			
Thr	Gln	Ser	Leu	Asn	Thr	Ala	Met	Thr	Gly	Leu	Lys	Arg	Gly	Val
	3275					3280					3285			
Ala	Asn	His	Asn	Gln	Val	Val	Gln	Ser	Asp	Asn	Tyr	Val	Asn	Ala
	3290					3295					3300			
Asp	Thr	Asn	Lys	Lys	Asn	Asp	Tyr	Asn	Asn	Ala	Tyr	Asn	His	Ala
	3305					3310					3315			
Asn	Asp	Ile	Ile	Asn	Gly	Asn	Ala	Gln	His	Pro	Val	Ile	Thr	Pro
	3320					3325					3330			
Ser	Asp	Val	Asn	Asn	Ala	Leu	Ser	Asn	Val	Thr	Ser	Lys	Glu	His
	3335					3340					3345			
Ala	Leu	Asn	Gly	Glu	Ala	Lys	Leu	Asn	Ala	Ala	Lys	Gln	Glu	Ala
	3350					3355					3360			
Asn	Thr	Ala	Leu	Gly	His	Leu	Asn	Asn	Leu	Asn	Asn	Ala	Gln	Arg
	3365					3370					3375			
Gln	Asn	Leu	Gln	Ser	Gln	Ile	Asn	Gly	Ala	His	Gln	Ile	Asp	Ala
	3380					3385					3390			
Val	Asn	Thr	Ile	Lys	Gln	Asn	Ala	Thr	Asn	Leu	Asn	Ser	Ala	Met
	3395					3400					3405			
Gly	Asn	Leu	Arg	Gln	Ala	Val	Ala	Asp	Lys	Asp	Gln	Val	Lys	Arg
	3410					3415					3420			
Thr	Glu	Asp	Tyr	Ala	Asp	Ala	Asp	Thr	Ala	Lys	Gln	Asn	Ala	Tyr
	3425					3430					3435			
Asn	Ser	Ala	Val	Ser	Ser	Ala	Glu	Thr	Ile	Ile	Asn	Gln	Thr	Thr
	3440					3445					3450			
Asn	Pro	Thr	Met	Ser	Val	Asp	Asp	Val	Asn	Arg	Ala	Thr	Ser	Ala
	3455					3460					3465			
Val	Thr	Ser	Asn	Lys	Asn	Ala	Leu	Asn	Gly	Tyr	Glu	Lys	Leu	Ala
	3470					3475					3480			
Gln	Ser	Lys	Thr	Asp	Ala	Ala	Arg	Ala	Ile	Asp	Ala	Leu	Pro	His
	3485					3490					3495			
Leu	Asn	Asn	Ala	Gln	Lys	Ala	Asp	Val	Lys	Ser	Lys	Ile	Asn	Ala
	3500					3505					3510			
Ala	Ser	Asn	Ile	Ala	Gly	Val	Asn	Thr	Val	Lys	Gln	Gln	Gly	Thr
	3515					3520					3525			
Asp	Leu	Asn	Thr	Ala	Met	Gly	Asn	Leu	Gln	Gly	Ala	Ile	Asn	Asp
	3530					3535					3540			
Glu	Gln	Thr	Thr	Leu	Asn	Ser	Gln	Asn	Tyr	Gln	Asp	Ala	Thr	Pro
	3545					3550					3555			
Ser	Lys	Lys	Thr	Ala	Tyr	Thr	Asn	Ala	Val	Gln	Ala	Ala	Lys	Asp
	3560					3565					3570			
Ile	Leu	Asn	Lys	Ser	Asn	Gly	Gln	Asn	Lys	Thr	Lys	Asp	Gln	Val
	3575					3580					3585			

-continued

---

Thr	Glu	Ala	Met	Asn	Gln	Val	Asn	Ser	Ala	Lys	Asn	Asn	Leu	Asp
3590						3595					3600			
Gly	Thr	Arg	Leu	Leu	Asp	Gln	Ala	Lys	Gln	Thr	Ala	Lys	Gln	Gln
3605						3610					3615			
Leu	Asn	Asn	Met	Thr	His	Leu	Thr	Thr	Ala	Gln	Lys	Thr	Asn	Leu
3620						3625					3630			
Thr	Asn	Gln	Ile	Asn	Ser	Gly	Thr	Thr	Val	Ala	Gly	Val	Gln	Thr
3635						3640					3645			
Val	Gln	Ser	Asn	Ala	Asn	Thr	Leu	Asp	Gln	Ala	Met	Asn	Thr	Leu
3650						3655					3660			
Arg	Gln	Ser	Ile	Ala	Asn	Lys	Asp	Ala	Thr	Lys	Ala	Ser	Glu	Asp
3665						3670					3675			
Tyr	Val	Asp	Ala	Asn	Asn	Asp	Lys	Gln	Thr	Ala	Tyr	Asn	Asn	Ala
3680						3685					3690			
Val	Ala	Ala	Ala	Glu	Thr	Ile	Ile	Asn	Ala	Asn	Ser	Asn	Pro	Glu
3695						3700					3705			
Met	Asn	Pro	Ser	Thr	Ile	Thr	Gln	Lys	Ala	Glu	Gln	Val	Asn	Ser
3710						3715					3720			
Ser	Lys	Thr	Ala	Leu	Asn	Gly	Asp	Glu	Asn	Leu	Ala	Ala	Ala	Lys
3725						3730					3735			
Gln	Asn	Ala	Lys	Thr	Tyr	Leu	Asn	Thr	Leu	Thr	Ser	Ile	Thr	Asp
3740						3745					3750			
Ala	Gln	Lys	Asn	Asn	Leu	Ile	Ser	Gln	Ile	Thr	Ser	Ala	Thr	Arg
3755						3760					3765			
Val	Ser	Gly	Val	Asp	Thr	Val	Lys	Gln	Asn	Ala	Gln	His	Leu	Asp
3770						3775					3780			
Gln	Ala	Met	Ala	Ser	Leu	Gln	Asn	Gly	Ile	Asn	Asn	Glu	Ser	Gln
3785						3790					3795			
Val	Lys	Ser	Ser	Glu	Lys	Tyr	Arg	Asp	Ala	Asp	Thr	Asn	Lys	Gln
3800						3805					3810			
Gln	Glu	Tyr	Asp	Asn	Ala	Ile	Thr	Ala	Ala	Lys	Ala	Ile	Leu	Asn
3815						3820					3825			
Lys	Ser	Thr	Gly	Pro	Asn	Thr	Ala	Gln	Asn	Ala	Val	Glu	Ala	Ala
3830						3835					3840			
Leu	Gln	Arg	Val	Asn	Asn	Ala	Lys	Asp	Ala	Leu	Asn	Gly	Asp	Ala
3845						3850					3855			
Lys	Leu	Ile	Ala	Ala	Gln	Asn	Ala	Ala	Lys	Gln	His	Leu	Gly	Thr
3860						3865					3870			
Leu	Thr	His	Ile	Thr	Thr	Ala	Gln	Arg	Asn	Asp	Leu	Thr	Asn	Gln
3875						3880					3885			
Ile	Ser	Gln	Ala	Thr	Asn	Leu	Ala	Gly	Val	Glu	Ser	Val	Lys	Gln
3890						3895					3900			
Asn	Ala	Asn	Ser	Leu	Asp	Gly	Ala	Met	Gly	Asn	Leu	Gln	Thr	Ala
3905						3910					3915			
Ile	Asn	Asp	Lys	Ser	Gly	Thr	Leu	Ala	Ser	Gln	Asn	Phe	Leu	Asp
3920						3925					3930			
Ala	Asp	Glu	Gln	Lys	Arg	Asn	Ala	Tyr	Asn	Gln	Ala	Val	Ser	Ala
3935						3940					3945			
Ala	Glu	Thr	Ile	Leu	Asn	Lys	Gln	Thr	Gly	Pro	Asn	Thr	Ala	Lys
3950						3955					3960			
Thr	Ala	Val	Glu	Gln	Ala	Leu	Asn	Asn	Val	Asn	Asn	Ala	Lys	His

-continued

3965	3970	3975
Ala Leu Asn Gly Thr Gln 3980	Asn Leu Asn Asn Ala 3985	Lys Gln Ala Ala 3990
Ile Thr Ala Ile Asn Gly 3995	Ala Ser Asp Leu Asn 4000	Gln Lys Gln Lys 4005
Asp Ala Leu Lys Ala Gln 4010	Ala Asn Gly Ala Gln 4015	Arg Val Ser Asn 4020
Ala Gln Asp Val Gln His 4025	Asn Ala Thr Glu Leu 4030	Asn Thr Ala Met 4035
Gly Thr Leu Lys His Ala 4040	Ile Ala Asp Lys Thr 4045	Asn Thr Leu Ala 4050
Ser Ser Lys Tyr Val Asn 4055	Ala Asp Ser Thr Lys 4060	Gln Asn Ala Tyr 4065
Thr Thr Lys Val Thr Asn 4070	Ala Glu His Ile Ile 4075	Ser Gly Thr Pro 4080
Thr Val Val Thr Thr Pro 4085	Ser Glu Val Thr Ala 4090	Ala Ala Asn Gln 4095
Val Asn Ser Ala Lys Gln 4100	Glu Leu Asn Gly Asp 4105	Glu Arg Leu Arg 4110
Glu Ala Lys Gln Asn Ala 4115	Asn Thr Ala Ile Asp 4120	Ala Leu Thr Gln 4125
Leu Asn Thr Pro Gln Lys 4130	Ala Lys Leu Lys Glu 4135	Gln Val Gly Gln 4140
Ala Asn Arg Leu Glu Asp 4145	Val Gln Thr Val Gln 4150	Thr Asn Gly Gln 4155
Ala Leu Asn Asn Ala Met 4160	Lys Gly Leu Arg Asp 4165	Ser Ile Ala Asn 4170
Glu Thr Thr Val Lys Thr 4175	Ser Gln Asn Tyr Thr 4180	Asp Ala Ser Pro 4185
Asn Asn Gln Ser Thr Tyr 4190	Asn Ser Ala Val Ser 4195	Asn Ala Lys Gly 4200
Ile Ile Asn Gln Thr Asn 4205	Asn Pro Thr Met Asp 4210	Thr Ser Ala Ile 4215
Thr Gln Ala Thr Thr Gln 4220	Val Asn Asn Ala Lys 4225	Asn Gly Leu Asn 4230
Gly Ala Glu Asn Leu Arg 4235	Asn Ala Gln Asn Thr 4240	Ala Lys Gln Asn 4245
Leu Asn Thr Leu Ser His 4250	Leu Thr Asn Asn Gln 4255	Lys Ser Ala Ile 4260
Ser Ser Gln Ile Asp Arg 4265	Ala Gly His Val Ser 4270	Glu Val Thr Ala 4275
Thr Lys Asn Ala Ala Thr 4280	Glu Leu Asn Thr Gln 4285	Met Gly Asn Leu 4290
Glu Gln Ala Ile His Asp 4295	Gln Asn Thr Val Lys 4300	Gln Ser Val Lys 4305
Phe Thr Asp Ala Asp Lys 4310	Ala Lys Arg Asp Ala 4315	Tyr Thr Asn Ala 4320
Val Ser Arg Ala Glu Ala 4325	Ile Leu Asn Lys Thr 4330	Gln Gly Ala Asn 4335
Thr Ser Lys Gln Asp Val 4340	Glu Ala Ala Ile Gln 4345	Asn Val Ser Ser 4350



-continued

Ala	Lys	Asn	Ala	Leu	Asn	Gly	Asp	Gln	Asn	Val	Thr	Asn	Ala	Lys
4355						4360						4365		
Asn	Ala	Ala	Lys	Asn	Ala	Leu	Asn	Asn	Leu	Thr	Ser	Ile	Asn	Asn
4370						4375						4380		
Ala	Gln	Lys	Arg	Asp	Leu	Thr	Thr	Lys	Ile	Asp	Gln	Ala	Thr	Thr
4385						4390					4395			
Val	Ala	Gly	Val	Glu	Ala	Val	Ser	Asn	Thr	Ser	Thr	Gln	Leu	Asn
4400						4405						4410		
Thr	Ala	Met	Ala	Asn	Leu	Gln	Asn	Gly	Ile	Asn	Asp	Lys	Thr	Asn
4415						4420						4425		
Thr	Leu	Ala	Ser	Glu	Asn	Tyr	His	Asp	Ala	Asp	Ser	Asp	Lys	Lys
4430						4435						4440		
Thr	Ala	Tyr	Thr	Gln	Ala	Val	Thr	Asn	Ala	Glu	Asn	Ile	Leu	Asn
4445						4450						4455		
Lys	Asn	Ser	Gly	Ser	Asn	Leu	Asp	Lys	Thr	Ala	Val	Glu	Asn	Ala
4460						4465						4470		
Leu	Ser	Gln	Val	Ala	Asn	Ala	Lys	Gly	Ala	Leu	Asn	Gly	Asn	His
4475						4480						4485		
Asn	Leu	Glu	Gln	Ala	Lys	Ser	Asn	Ala	Asn	Thr	Thr	Ile	Asn	Gly
4490						4495						4500		
Leu	Gln	His	Leu	Thr	Thr	Ala	Gln	Lys	Asp	Lys	Leu	Lys	Gln	Gln
4505						4510						4515		
Val	Gln	Gln	Ala	Gln	Asn	Val	Ala	Gly	Val	Asp	Thr	Val	Lys	Ser
4520						4525						4530		
Ser	Ala	Asn	Thr	Leu	Asn	Gly	Ala	Met	Gly	Thr	Leu	Arg	Asn	Ser
4535						4540						4545		
Ile	Gln	Asp	Asn	Thr	Ala	Thr	Lys	Asn	Gly	Gln	Asn	Tyr	Leu	Asp
4550						4555						4560		
Ala	Thr	Glu	Arg	Asn	Lys	Thr	Asn	Tyr	Asn	Asn	Ala	Val	Asp	Ser
4565						4570						4575		
Ala	Asn	Gly	Val	Ile	Asn	Ala	Thr	Ser	Asn	Pro	Asn	Met	Asp	Ala
4580						4585						4590		
Asn	Ala	Ile	Asn	Gln	Ile	Ala	Thr	Gln	Val	Thr	Ser	Thr	Lys	Asn
4595						4600						4605		
Ala	Leu	Asp	Gly	Thr	His	Asn	Leu	Thr	Gln	Ala	Lys	Gln	Thr	Ala
4610						4615						4620		
Thr	Asn	Ala	Ile	Asp	Gly	Ala	Thr	Asn	Leu	Asn	Lys	Ala	Gln	Lys
4625						4630						4635		
Asp	Ala	Leu	Lys	Ala	Gln	Val	Thr	Ser	Ala	Gln	Arg	Val	Ala	Asn
4640						4645						4650		
Val	Thr	Ser	Ile	Gln	Gln	Thr	Ala	Asn	Glu	Leu	Asn	Thr	Ala	Met
4655						4660						4665		
Gly	Gln	Leu	Gln	His	Gly	Ile	Asp	Asp	Glu	Asn	Ala	Thr	Lys	Gln
4670						4675						4680		
Thr	Gln	Lys	Tyr	Arg	Asp	Ala	Glu	Gln	Ser	Lys	Lys	Thr	Ala	Tyr
4685						4690						4695		
Asp	Gln	Ala	Val	Ala	Ala	Ala	Lys	Ala	Ile	Leu	Asn	Lys	Gln	Thr
4700						4705						4710		
Gly	Ser	Asn	Ser	Asp	Lys	Ala	Ala	Val	Asp	Arg	Ala	Leu	Gln	Gln
4715						4720						4725		
Val	Thr	Ser	Thr	Lys	Asp	Ala	Leu	Asn	Gly	Asp	Ala	Lys	Leu	Ala
4730						4735						4740		

-continued

---

Glu Ala	Lys Ala Ala Ala	Lys	Gln Asn Leu Gly	Thr	Leu Asn His
4745		4750		4755	
Ile Thr	Asn Ala Gln Arg	Thr	Asp Leu Glu Gly	Gln	Ile Asn Gln
4760		4765		4770	
Ala Thr	Thr Val Asp Gly	Val	Asn Thr Val Lys	Thr	Asn Ala Asn
4775		4780		4785	
Thr Leu	Asp Gly Ala Met	Asn	Ser Leu Gln Gly	Ser	Ile Asn Asp
4790		4795		4800	
Lys Asp	Ala Thr Leu Arg	Asn	Gln Asn Tyr Leu	Asp	Ala Asp Glu
4805		4810		4815	
Ser Lys	Arg Asn Ala Tyr	Thr	Gln Ala Val Thr	Ala	Ala Glu Gly
4820		4825		4830	
Ile Leu	Asn Lys Gln Thr	Gly	Gly Asn Thr Ser	Lys	Ala Asp Val
4835		4840		4845	
Asp Asn	Ala Leu Asn Ala	Val	Thr Arg Ala Lys	Ala	Ala Leu Asn
4850		4855		4860	
Gly Ala	Asp Asn Leu Arg	Asn	Ala Lys Thr Ser	Ala	Thr Asn Thr
4865		4870		4875	
Ile Asp	Gly Leu Pro Asn	Leu	Thr Gln Leu Gln	Lys	Asp Asn Leu
4880		4885		4890	
Lys His	Gln Val Glu Gln	Ala	Gln Asn Val Ala	Gly	Val Asn Gly
4895		4900		4905	
Val Lys	Asp Lys Gly Asn	Thr	Leu Asn Thr Ala	Met	Gly Ala Leu
4910		4915		4920	
Arg Thr	Ser Ile Gln Asn	Asp	Asn Thr Thr Lys	Thr	Ser Gln Asn
4925		4930		4935	
Tyr Leu	Asp Ala Ser Asp	Ser	Asn Lys Asn Asn	Tyr	Asn Thr Ala
4940		4945		4950	
Val Asn	Asn Ala Asn Gly	Val	Ile Asn Ala Thr	Asn	Asn Pro Asn
4955		4960		4965	
Met Asp	Ala Asn Ala Ile	Asn	Gly Met Ala Asn	Gln	Val Asn Thr
4970		4975		4980	
Thr Lys	Ala Ala Leu Asn	Gly	Ala Gln Asn Leu	Ala	Gln Ala Lys
4985		4990		4995	
Thr Asn	Ala Thr Asn Thr	Ile	Asn Asn Ala His	Asp	Leu Asn Gln
5000		5005		5010	
Lys Gln	Lys Asp Ala Leu	Lys	Thr Gln Val Asn	Asn	Ala Gln Arg
5015		5020		5025	
Val Ser	Asp Ala Asn Asn	Val	Gln His Thr Ala	Thr	Glu Leu Asn
5030		5035		5040	
Ser Ala	Met Thr Ala Leu	Lys	Ala Ala Ile Ala	Asp	Lys Glu Arg
5045		5050		5055	
Thr Lys	Ala Ser Gly Asn	Tyr	Val Asn Ala Asp	Gln	Glu Lys Arg
5060		5065		5070	
Gln Ala	Tyr Asp Ser Lys	Val	Thr Asn Ala Glu	Asn	Ile Ile Ser
5075		5080		5085	
Gly Thr	Pro Asn Ala Thr	Leu	Thr Val Asn Asp	Val	Asn Ser Ala
5090		5095		5100	
Ala Ser	Gln Val Asn Ala	Ala	Lys Thr Ala Leu	Asn	Gly Asp Asn
5105		5110		5115	
Asn Leu	Arg Val Ala Lys	Glu	His Ala Asn Asn	Thr	Ile Asp Gly

-continued

5120	5125	5130
Leu Ala Gln Leu Asn Asn 5135	Ala Gln Lys Ala Lys 5140	Leu Lys Glu Gln 5145
Val Gln Ser Ala Thr Thr 5150	Leu Asp Gly Val Gln 5155	Thr Val Lys Asn 5160
Ser Ser Gln Thr Leu Asn 5165	Thr Ala Met Lys Gly 5170	Leu Arg Asp Ser 5175
Ile Ala Asn Glu Ala Thr 5180	Ile Lys Ala Gly Gln 5185	Asn Tyr Thr Asp 5190
Ala Ser Pro Asn Asn Arg 5195	Asn Glu Tyr Asp Ser 5200	Ala Val Thr Ala 5205
Ala Lys Ala Ile Ile Asn 5210	Gln Thr Ser Asn Pro 5215	Thr Met Glu Pro 5220
Asn Thr Ile Thr Gln Val 5225	Thr Ser Gln Val Thr 5230	Thr Lys Glu Gln 5235
Ala Leu Asn Gly Ala Arg 5240	Asn Leu Ala Gln Ala 5245	Lys Thr Thr Ala 5250
Lys Asn Asn Leu Asn Asn 5255	Leu Thr Ser Ile Asn 5260	Asn Ala Gln Lys 5265
Asp Ala Leu Thr Arg Ser 5270	Ile Asp Gly Ala Thr 5275	Thr Val Ala Gly 5280
Val Asn Gln Glu Thr Ala 5285	Lys Ala Thr Glu Leu 5290	Asn Asn Ala Met 5295
His Ser Leu Gln Asn Gly 5300	Ile Asn Asp Glu Thr 5305	Gln Thr Lys Gln 5310
Thr Gln Lys Tyr Leu Asp 5315	Ala Glu Pro Ser Lys 5320	Lys Ser Ala Tyr 5325
Asp Gln Ala Val Asn Ala 5330	Ala Lys Ala Ile Leu 5335	Thr Lys Ala Ser 5340
Gly Gln Asn Val Asp Lys 5345	Ala Ala Val Glu Gln 5350	Ala Leu Gln Asn 5355
Val Asn Ser Thr Lys Thr 5360	Ala Leu Asn Gly Asp 5365	Ala Lys Leu Asn 5370
Glu Ala Lys Ala Ala Ala 5375	Lys Gln Thr Leu Gly 5380	Thr Leu Thr His 5385
Ile Asn Asn Ala Gln Arg 5390	Thr Ala Leu Asp Asn 5395	Glu Ile Thr Gln 5400
Ala Thr Asn Val Glu Gly 5405	Val Asn Thr Val Lys 5410	Ala Lys Ala Gln 5415
Gln Leu Asp Gly Ala Met 5420	Gly Gln Leu Glu Thr 5425	Ser Ile Arg Asp 5430
Lys Asp Thr Thr Leu Gln 5435	Ser Gln Asn Tyr Gln 5440	Asp Ala Asp Asp 5445
Ala Lys Arg Thr Ala Tyr 5450	Ser Gln Ala Val Asn 5455	Ala Ala Ala Thr 5460
Ile Leu Asn Lys Thr Ala 5465	Gly Gly Asn Thr Pro 5470	Lys Ala Asp Val 5475
Glu Arg Ala Met Gln Ala 5480	Val Thr Gln Ala Asn 5485	Thr Ala Leu Asn 5490
Gly Ile Gln Asn Leu Asp 5495	Arg Ala Lys Gln Ala 5500	Ala Asn Thr Ala 5505

-continued

---

Ile Thr	Asn Ala Ser Asp	Leu	Asn Thr Lys Gln	Lys	Glu Ala Leu
5510		5515		5520	
Lys Ala	Gln Val Thr Ser	Ala	Gly Arg Val Ser	Ala	Ala Asn Gly
5525		5530		5535	
Val Glu	His Thr Ala Thr	Glu	Leu Asn Thr Ala	Met	Thr Ala Leu
5540		5545		5550	
Lys Arg	Ala Ile Ala Asp	Lys	Ala Glu Thr Lys	Ala	Ser Gly Asn
5555		5560		5565	
Tyr Val	Asn Ala Asp Ala	Asn	Lys Arg Gln Ala	Tyr	Asp Glu Lys
5570		5575		5580	
Val Thr	Ala Ala Glu Asn	Ile	Val Ser Gly Thr	Pro	Thr Pro Thr
5585		5590		5595	
Leu Thr	Pro Ala Asp Val	Thr	Asn Ala Ala Thr	Gln	Val Thr Asn
5600		5605		5610	
Ala Lys	Thr Gln Leu Asn	Gly	Asn His Asn Leu	Glu	Val Ala Lys
5615		5620		5625	
Gln Asn	Ala Asn Thr Ala	Ile	Asp Gly Leu Thr	Ser	Leu Asn Gly
5630		5635		5640	
Pro Gln	Lys Ala Lys Leu	Lys	Glu Gln Val Gly	Gln	Ala Thr Thr
5645		5650		5655	
Leu Pro	Asn Val Gln Thr	Val	Arg Asp Asn Ala	Gln	Thr Leu Asn
5660		5665		5670	
Thr Ala	Met Lys Gly Leu	Arg	Asp Ser Ile Ala	Asn	Glu Ala Thr
5675		5680		5685	
Ile Lys	Ala Gly Gln Asn	Tyr	Thr Asp Ala Ser	Gln	Asn Lys Gln
5690		5695		5700	
Thr Asp	Tyr Asn Ser Ala	Val	Thr Ala Ala Lys	Ala	Ile Ile Gly
5705		5710		5715	
Gln Thr	Thr Ser Pro Ser	Met	Asn Ala Gln Glu	Ile	Asn Gln Ala
5720		5725		5730	
Lys Asp	Gln Val Thr Ala	Lys	Gln Gln Ala Leu	Asn	Gly Gln Glu
5735		5740		5745	
Asn Leu	Arg Thr Ala Gln	Thr	Asn Ala Lys Gln	His	Leu Asn Gly
5750		5755		5760	
Leu Ser	Asp Leu Thr Asp	Ala	Gln Lys Asp Ala	Val	Lys Arg Gln
5765		5770		5775	
Ile Glu	Gly Ala Thr His	Val	Asn Glu Val Thr	Gln	Ala Gln Asn
5780		5785		5790	
Asn Ala	Asp Ala Leu Asn	Thr	Ala Met Thr Asn	Leu	Lys Asn Gly
5795		5800		5805	
Ile Gln	Asp Gln Asn Thr	Ile	Lys Gln Gly Val	Asn	Phe Thr Asp
5810		5815		5820	
Ala Asp	Glu Ala Lys Arg	Asn	Ala Tyr Thr Asn	Ala	Val Thr Gln
5825		5830		5835	
Ala Glu	Gln Ile Leu Asn	Lys	Ala Gln Gly Pro	Asn	Thr Ser Lys
5840		5845		5850	
Asp Gly	Val Glu Thr Ala	Leu	Glu Asn Val Gln	Arg	Ala Lys Asn
5855		5860		5865	
Glu Leu	Asn Gly Asn Gln	Asn	Val Ala Asn Ala	Lys	Thr Thr Ala
5870		5875		5880	
Lys Asn	Ala Leu Asn Asn	Leu	Thr Ser Ile Asn	Asn	Ala Gln Lys
5885		5890		5895	

-continued

---

Glu Ala	Leu Lys Ser Gln Ile	Glu Gly Ala Thr Thr	Val Ala Gly
5900	5905	5910	
Val Asn	Gln Val Ser Thr Thr	Ala Ser Glu Leu Asn	Thr Ala Met
5915	5920	5925	
Ser Asn	Leu Gln Asn Gly Ile	Asn Asp Glu Ala Ala	Thr Lys Ala
5930	5935	5940	
Ala Gln	Lys Tyr Thr Asp Ala	Asp Arg Glu Lys Gln	Thr Ala Tyr
5945	5950	5955	
Asn Asp	Ala Val Thr Ala Ala	Lys Thr Leu Leu Asp	Lys Thr Ala
5960	5965	5970	
Gly Ser	Asn Asp Asn Lys Ala	Ala Val Glu Gln Ala	Leu Gln Arg
5975	5980	5985	
Val Asn	Thr Ala Lys Thr Ala	Leu Asn Gly Asp Glu	Arg Leu Asn
5990	5995	6000	
Glu Ala	Lys Asn Thr Ala Lys	Gln Gln Val Ala Thr	Met Ser His
6005	6010	6015	
Leu Thr	Asp Ala Gln Lys Ala	Asn Leu Thr Ser Gln	Ile Glu Ser
6020	6025	6030	
Gly Thr	Thr Val Ala Gly Val	Gln Gly Ile Gln Ala	Asn Ala Gly
6035	6040	6045	
Thr Leu	Asp Gln Ala Met Asn	Gln Leu Arg Gln Ser	Ile Ala Ser
6050	6055	6060	
Lys Asp	Ala Thr Lys Ser Ser	Glu Asp Tyr Gln Asp	Ala Asn Ala
6065	6070	6075	
Asp Leu	Gln Asn Ala Tyr Asn	Asp Ala Val Thr Asn	Ala Glu Gly
6080	6085	6090	
Ile Ile	Ser Ala Thr Asn Asn	Pro Glu Met Asn Pro	Asp Thr Ile
6095	6100	6105	
Asn Gln	Lys Ala Ser Gln Val	Asn Ser Ala Lys Ser	Ala Leu Asn
6110	6115	6120	
Gly Asp	Glu Lys Leu Ala Ala	Ala Lys Gln Thr Ala	Lys Ser Asp
6125	6130	6135	
Ile Gly	Arg Leu Thr Asp Leu	Asn Asn Ala Gln Arg	Thr Ala Ala
6140	6145	6150	
Asn Ala	Glu Val Asp Gln Ala	Pro Asn Leu Ala Ala	Val Thr Ala
6155	6160	6165	
Ala Lys	Asn Lys Ala Thr Ser	Leu Asn Thr Ala Met	Gly Asn Leu
6170	6175	6180	
Lys His	Ala Leu Ala Glu Lys	Asp Asn Thr Lys Arg	Ser Val Asn
6185	6190	6195	
Tyr Thr	Asp Ala Asp Gln Pro	Lys Gln Gln Ala Tyr	Asp Thr Ala
6200	6205	6210	
Val Thr	Gln Ala Glu Ala Ile	Thr Asn Ala Asn Gly	Ser Asn Ala
6215	6220	6225	
Asn Glu	Thr Gln Val Gln Ala	Ala Leu Asn Gln Leu	Asn Gln Ala
6230	6235	6240	
Lys Asn	Asp Leu Asn Gly Asp	Asn Lys Val Ala Gln	Ala Lys Glu
6245	6250	6255	
Ser Ala	Lys Arg Ala Leu Ala	Ser Tyr Ser Asn Leu	Asn Asn Ala
6260	6265	6270	
Gln Ser	Thr Ala Ala Ile Ser	Gln Ile Asp Asn Ala	Thr Thr Val

-continued

6275	6280	6285
Ala Gly Val Thr Ala Ala 6290	Gln Asn Thr Ala Asn 6295	Glu Leu Asn Thr 6300
Ala Met Gly Gln Leu Gln 6305	Asn Gly Ile Asn Asp 6310	Gln Asn Thr Val 6315
Lys Gln Gln Val Asn Phe 6320	Thr Asp Ala Asp 6325	Gln Gly Lys Lys Asp 6330
Ala Tyr Thr Asn Ala Val 6335	Thr Asn Ala Gln 6340	Gly Ile Leu Asp Lys 6345
Ala His Gly Gln Asn Met 6350	Thr Lys Ala Gln 6355	Val Glu Ala Ala Leu 6360
Asn Gln Val Thr Thr Ala 6365	Lys Asn Ala Leu 6370	Asn Gly Asp Ala Asn 6375
Val Arg Gln Ala Lys Ser 6380	Asp Ala Lys Ala 6385	Asn Leu Gly Thr Leu 6390
Thr His Leu Asn Asn Ala 6395	Gln Lys Gln Asp 6400	Leu Thr Ser Gln Ile 6405
Glu Gly Ala Thr Thr Val 6410	Asn Gly Val Asn 6415	Gly Val Lys Thr Lys 6420
Ala Gln Asp Leu Asp Gly 6425	Ala Met Gln Arg 6430	Leu Gln Ser Ala Ile 6435
Ala Asn Lys Asp Gln Thr 6440	Lys Ala Ser Glu 6445	Asn Tyr Ile Asp Ala 6450
Asp Pro Thr Lys Lys Thr 6455	Ala Phe Asp Asn 6460	Ala Ile Thr Gln Ala 6465
Glu Ser Tyr Leu Asn Lys 6470	Asp His Gly Ala 6475	Asn Lys Asp Lys Gln 6480
Ala Val Glu Gln Ala Ile 6485	Gln Ser Val Thr 6490	Ser Thr Glu Asn Ala 6495
Leu Asn Gly Asp Ala Asn 6500	Leu Gln Arg Ala 6505	Lys Thr Glu Ala Ile 6510
Gln Ala Ile Asp Asn Leu 6515	Thr His Leu Asn 6520	Thr Pro Gln Lys Thr 6525
Ala Leu Lys Gln Gln Val 6530	Asn Ala Ala Gln 6535	Arg Val Ser Gly Val 6540
Thr Asp Leu Lys Asn Ser 6545	Ala Thr Ser Leu 6550	Asn Asn Ala Met Asp 6555
Gln Leu Lys Gln Ala Ile 6560	Ala Asp His Asp 6565	Thr Ile Val Ala Ser 6570
Gly Asn Tyr Thr Asn Ala 6575	Ser Pro Asp Lys 6580	Gln Gly Ala Tyr Thr 6585
Asp Ala Tyr Asn Ala Ala 6590	Lys Asn Ile Val 6595	Asn Gly Ser Pro Asn 6600
Val Ile Thr Asn Ala Ala 6605	Asp Val Thr Ala 6610	Ala Thr Glu Arg Val 6615
Asn Asn Ala Glu Thr Gly 6620	Leu Asn Gly Asp 6625	Thr Asn Leu Ala Thr 6630
Ala Lys Gln Gln Ala Lys 6635	Asp Ala Leu Arg 6640	Gln Met Thr His Leu 6645
Ser Asp Ala Gln Lys Gln 6650	Ser Ile Thr Gly 6655	Gln Ile Asp Ser Ala 6660

-continued

Thr	Gln	Val	Thr	Gly	Val	Gln	Ser	Val	Lys	Asp	Asn	Ala	Thr	Asn
	6665					6670						6675		
Leu	Asp	Asn	Ala	Met	Asn	Gln	Leu	Arg	Asn	Ser	Ile	Ala	Asn	Lys
	6680					6685						6690		
Asp	Asp	Val	Lys	Ala	Ser	Gln	Pro	Tyr	Val	Asp	Ala	Asp	Arg	Asp
	6695					6700						6705		
Lys	Gln	Asn	Ala	Tyr	Asn	Thr	Ala	Val	Thr	Asn	Ala	Glu	Asn	Ile
	6710					6715						6720		
Ile	Asn	Ala	Thr	Ser	Gln	Pro	Thr	Leu	Asp	Pro	Ser	Ala	Val	Thr
	6725					6730						6735		
Gln	Ala	Ala	Asn	Gln	Val	Ser	Thr	Asn	Lys	Thr	Ala	Leu	Asn	Gly
	6740					6745						6750		
Ala	Gln	Asn	Leu	Ala	Asn	Lys	Lys	Gln	Glu	Thr	Thr	Ala	Asn	Ile
	6755					6760						6765		
Asn	Gln	Leu	Ser	His	Leu	Asn	Asn	Ala	Gln	Lys	Gln	Asp	Leu	Asn
	6770					6775						6780		
Thr	Gln	Val	Thr	Asn	Ala	Pro	Asn	Ile	Ser	Thr	Val	Asn	Gln	Val
	6785					6790						6795		
Lys	Thr	Lys	Ala	Glu	Gln	Leu	Asp	Gln	Ala	Met	Glu	Arg	Leu	Ile
	6800					6805						6810		
Asn	Gly	Ile	Gln	Asp	Lys	Asp	Gln	Val	Lys	Gln	Ser	Val	Asn	Phe
	6815					6820						6825		
Thr	Asp	Ala	Asp	Pro	Glu	Lys	Gln	Thr	Ala	Tyr	Asn	Asn	Ala	Val
	6830					6835						6840		
Thr	Ala	Ala	Glu	Asn	Ile	Ile	Asn	Gln	Ala	Asn	Gly	Thr	Asn	Ala
	6845					6850						6855		
Asn	Gln	Ser	Gln	Val	Glu	Ala	Ala	Leu	Ser	Thr	Val	Thr	Thr	Thr
	6860					6865						6870		
Lys	Gln	Ala	Leu	Asn	Gly	Asp	Arg	Lys	Val	Thr	Asp	Ala	Lys	Asn
	6875					6880						6885		
Asn	Ala	Asn	Gln	Thr	Leu	Ser	Thr	Leu	Asp	Asn	Leu	Asn	Asn	Ala
	6890					6895						6900		
Gln	Lys	Gly	Ala	Val	Thr	Gly	Asn	Ile	Asn	Gln	Ala	His	Thr	Val
	6905					6910						6915		
Ala	Glu	Val	Thr	Gln	Ala	Ile	Gln	Thr	Ala	Gln	Glu	Leu	Asn	Thr
	6920					6925						6930		
Ala	Met	Gly	Asn	Leu	Lys	Asn	Ser	Leu	Asn	Asp	Lys	Asp	Thr	Thr
	6935					6940						6945		
Leu	Gly	Ser	Gln	Asn	Phe	Ala	Asp	Ala	Asp	Pro	Glu	Lys	Lys	Asn
	6950					6955						6960		
Ala	Tyr	Asn	Glu	Ala	Val	His	Asn	Ala	Glu	Asn	Ile	Leu	Asn	Lys
	6965					6970						6975		
Ser	Thr	Gly	Thr	Asn	Val	Pro	Lys	Asp	Gln	Val	Glu	Ala	Ala	Met
	6980					6985						6990		
Asn	Gln	Val	Asn	Ala	Thr	Lys	Ala	Ala	Leu	Asn	Gly	Thr	Gln	Asn
	6995					7000						7005		
Leu	Glu	Lys	Ala	Lys	Gln	His	Ala	Asn	Thr	Ala	Ile	Asp	Gly	Leu
	7010					7015						7020		
Ser	His	Leu	Thr	Asn	Ala	Gln	Lys	Glu	Ala	Leu	Lys	Gln	Leu	Val
	7025					7030						7035		
Gln	Gln	Ser	Thr	Thr	Val	Ala	Glu	Ala	Gln	Gly	Asn	Glu	Gln	Lys
	7040					7045						7050		

-continued

---

Ala	Asn	Asn	Val	Asp	Ala	Ala	Met	Asp	Lys	Leu	Arg	Gln	Ser	Ile
7055						7060						7065		
Ala	Asp	Asn	Ala	Thr	Thr	Lys	Gln	Asn	Gln	Asn	Tyr	Thr	Asp	Ala
7070						7075						7080		
Ser	Gln	Asn	Lys	Lys	Asp	Ala	Tyr	Asn	Asn	Ala	Val	Thr	Thr	Ala
7085						7090						7095		
Gln	Gly	Ile	Ile	Asp	Gln	Thr	Thr	Ser	Pro	Thr	Leu	Asp	Pro	Thr
7100						7105						7110		
Val	Ile	Asn	Gln	Ala	Ala	Gly	Gln	Val	Ser	Thr	Thr	Lys	Asn	Ala
7115						7120						7125		
Leu	Asn	Gly	Asn	Glu	Asn	Leu	Glu	Ala	Ala	Lys	Gln	Gln	Ala	Ser
7130						7135						7140		
Gln	Ser	Leu	Gly	Ser	Leu	Asp	Asn	Leu	Asn	Asn	Ala	Gln	Lys	Gln
7145						7150						7155		
Thr	Val	Thr	Asp	Gln	Ile	Asn	Gly	Ala	His	Thr	Val	Asp	Glu	Ala
7160						7165						7170		
Asn	Gln	Ile	Lys	Gln	Asn	Ala	Gln	Asn	Leu	Asn	Thr	Ala	Met	Gly
7175						7180						7185		
Asn	Leu	Lys	Gln	Ala	Ile	Ala	Asp	Lys	Asp	Ala	Thr	Lys	Ala	Thr
7190						7195						7200		
Val	Asn	Phe	Thr	Asp	Ala	Asp	Gln	Ala	Lys	Gln	Gln	Ala	Tyr	Asn
7205						7210						7215		
Thr	Ala	Val	Thr	Asn	Ala	Glu	Asn	Ile	Ser	Lys	Ala	Asn	Gly	Asn
7220						7225						7230		
Ala	Thr	Gln	Ala	Glu	Val	Glu	Gln	Ala	Ile	Lys	Gln	Val	Asn	Ala
7235						7240						7245		
Ala	Lys	Gln	Ala	Leu	Asn	Gly	Asn	Ala	Asn	Val	Gln	His	Ala	Lys
7250						7255						7260		
Asp	Glu	Ala	Thr	Ala	Leu	Ile	Asn	Ser	Ser	Asn	Asp	Leu	Asn	Gln
7265						7270						7275		
Ala	Gln	Lys	Asp	Ala	Leu	Lys	Gln	Gln	Val	Gln	Asn	Ala	Thr	Thr
7280						7285						7290		
Val	Ala	Gly	Val	Asn	Asn	Val	Lys	Gln	Thr	Ala	Gln	Glu	Leu	Asn
7295						7300						7305		
Asn	Ala	Met	Thr	Gln	Leu	Lys	Gln	Gly	Ile	Ala	Asp	Lys	Glu	Gln
7310						7315						7320		
Thr	Lys	Ala	Asp	Gly	Asn	Phe	Val	Asn	Ala	Asp	Pro	Asp	Lys	Gln
7325						7330						7335		
Asn	Ala	Tyr	Asn	Gln	Ala	Val	Ala	Lys	Ala	Glu	Ala	Leu	Ile	Ser
7340						7345						7350		
Ala	Thr	Pro	Asp	Val	Val	Val	Thr	Pro	Ser	Glu	Ile	Thr	Ala	Ala
7355						7360						7365		
Leu	Asn	Lys	Val	Thr	Gln	Ala	Lys	Asn	Asp	Leu	Asn	Gly	Asn	Thr
7370						7375						7380		
Asn	Leu	Ala	Thr	Ala	Lys	Gln	Asn	Val	Gln	His	Ala	Ile	Asp	Gln
7385						7390						7395		
Leu	Pro	Asn	Leu	Asn	Gln	Ala	Gln	Arg	Asp	Glu	Tyr	Ser	Lys	Gln
7400						7405						7410		
Ile	Thr	Gln	Ala	Thr	Leu	Val	Pro	Asn	Val	Asn	Ala	Ile	Gln	Gln
7415						7420						7425		
Ala	Ala	Thr	Thr	Leu	Asn	Asp	Ala	Met	Thr	Gln	Leu	Lys	Gln	Gly



-continued

---

7430	7435	7440
Ile Ala Asn Lys Ala Gln 7445	Ile Lys Gly Ser Glu 7450	Asn Tyr His Asp 7455
Ala Asp Thr Asp Lys Gln 7460	Thr Ala Tyr Asp Asn 7465	Ala Val Thr Lys 7470
Ala Glu Glu Leu Leu Lys 7475	Gln Thr Thr Asn Pro 7480	Thr Met Asp Pro 7485
Asn Thr Ile Gln Gln Ala 7490	Leu Thr Lys Val Asn 7495	Asp Thr Asn Gln 7500
Ala Leu Asn Gly Asn Gln 7505	Lys Leu Ala Asp Ala 7510	Lys Gln Asp Ala 7515
Lys Thr Thr Leu Gly Thr 7520	Leu Asp His Leu Asn 7525	Asp Ala Gln Lys 7530
Gln Ala Leu Thr Thr Gln 7535	Val Glu Gln Ala Pro 7540	Asp Ile Ala Thr 7545
Val Asn Asn Val Lys Gln 7550	Asn Ala Gln Asn Leu 7555	Asn Asn Ala Met 7560
Thr Asn Leu Asn Asn Ala 7565	Leu Gln Asp Lys Thr 7570	Glu Thr Leu Asn 7575
Ser Ile Asn Phe Thr Asp 7580	Ala Asp Gln Ala Lys 7585	Lys Asp Ala Tyr 7590
Thr Asn Ala Val Ser His 7595	Ala Glu Gly Ile Leu 7600	Ser Lys Ala Asn 7605
Gly Ser Asn Ala Ser Gln 7610	Thr Glu Val Glu Gln 7615	Ala Met Gln Arg 7620
Val Asn Glu Ala Lys Gln 7625	Ala Leu Asn Gly Asn 7630	Asp Asn Val Gln 7635
Arg Ala Lys Asp Ala Ala 7640	Lys Gln Val Ile Thr 7645	Asn Ala Asn Asp 7650
Leu Asn Gln Ala Gln Lys 7655	Asp Ala Leu Lys Gln 7660	Gln Val Asp Ala 7665
Ala Gln Thr Val Ala Asn 7670	Val Asn Thr Ile Lys 7675	Gln Thr Ala Gln 7680
Asp Leu Asn Gln Ala Met 7685	Thr Gln Leu Lys Gln 7690	Gly Ile Ala Asp 7695
Lys Asp Gln Thr Lys Ala 7700	Asn Gly Asn Phe Val 7705	Asn Ala Asp Thr 7710
Asp Lys Gln Asn Ala Tyr 7715	Asn Asn Ala Val Ala 7720	His Ala Glu Gln 7725
Ile Ile Ser Gly Thr Pro 7730	Asn Ala Asn Val Asp 7735	Pro Gln Gln Val 7740
Ala Gln Ala Leu Gln Gln 7745	Val Asn Gln Ala Lys 7750	Gly Asp Leu Asn 7755
Gly Asn His Asn Leu Gln 7760	Val Ala Lys Asp Asn 7765	Ala Asn Thr Ala 7770
Ile Asp Gln Leu Pro Asn 7775	Leu Asn Gln Pro Gln 7780	Lys Thr Ala Leu 7785
Lys Asp Gln Val Ser His 7790	Ala Glu Leu Val Thr 7795	Gly Val Asn Ala 7800
Ile Lys Gln Asn Ala Asp 7805	Ala Leu Asn Asn Ala 7810	Met Gly Thr Leu 7815

-continued

---

Lys	Gln	Gln	Ile	Gln	Ala	Asn	Ser	Gln	Val	Pro	Gln	Ser	Val	Asp
	7820					7825					7830			
Phe	Thr	Gln	Ala	Asp	Gln	Asp	Lys	Gln	Gln	Ala	Tyr	Asn	Asn	Ala
	7835					7840					7845			
Ala	Asn	Gln	Ala	Gln	Gln	Ile	Ala	Asn	Gly	Ile	Pro	Thr	Pro	Val
	7850					7855					7860			
Leu	Thr	Pro	Asp	Thr	Val	Thr	Gln	Ala	Val	Thr	Thr	Met	Asn	Gln
	7865					7870					7875			
Ala	Lys	Asp	Ala	Leu	Asn	Gly	Asp	Glu	Lys	Leu	Ala	Gln	Ala	Lys
	7880					7885					7890			
Gln	Glu	Ala	Leu	Ala	Asn	Leu	Asp	Thr	Leu	Arg	Asp	Leu	Asn	Gln
	7895					7900					7905			
Pro	Gln	Arg	Asp	Ala	Leu	Arg	Asn	Gln	Ile	Asn	Gln	Ala	Gln	Ala
	7910					7915					7920			
Leu	Ala	Thr	Val	Glu	Gln	Thr	Lys	Gln	Asn	Ala	Gln	Asn	Val	Asn
	7925					7930					7935			
Thr	Ala	Met	Ser	Asn	Leu	Lys	Gln	Gly	Ile	Ala	Asn	Lys	Asp	Thr
	7940					7945					7950			
Val	Lys	Ala	Ser	Glu	Asn	Tyr	His	Asp	Ala	Asp	Ala	Asp	Lys	Gln
	7955					7960					7965			
Thr	Ala	Tyr	Thr	Asn	Ala	Val	Ser	Gln	Ala	Glu	Gly	Ile	Ile	Asn
	7970					7975					7980			
Gln	Thr	Thr	Asn	Pro	Thr	Leu	Asn	Pro	Asp	Glu	Ile	Thr	Arg	Ala
	7985					7990					7995			
Leu	Thr	Gln	Val	Thr	Asp	Ala	Lys	Asn	Gly	Leu	Asn	Gly	Glu	Ala
	8000					8005					8010			
Lys	Leu	Ala	Thr	Glu	Lys	Gln	Asn	Ala	Lys	Asp	Ala	Val	Ser	Gly
	8015					8020					8025			
Met	Thr	His	Leu	Asn	Asp	Ala	Gln	Lys	Gln	Ala	Leu	Lys	Gly	Gln
	8030					8035					8040			
Ile	Asp	Gln	Ser	Pro	Glu	Ile	Ala	Thr	Val	Asn	Gln	Val	Lys	Gln
	8045					8050					8055			
Thr	Ala	Thr	Ser	Leu	Asp	Gln	Ala	Met	Asp	Gln	Leu	Ser	Gln	Ala
	8060					8065					8070			
Ile	Asn	Asp	Lys	Ala	Gln	Thr	Leu	Ala	Asp	Gly	Asn	Tyr	Leu	Asn
	8075					8080					8085			
Ala	Asp	Pro	Asp	Lys	Gln	Asn	Ala	Tyr	Lys	Gln	Ala	Val	Ala	Lys
	8090					8095					8100			
Ala	Glu	Ala	Leu	Leu	Asn	Lys	Gln	Ser	Gly	Thr	Asn	Glu	Val	Gln
	8105					8110					8115			
Ala	Gln	Val	Glu	Ser	Ile	Thr	Asn	Glu	Val	Asn	Ala	Ala	Lys	Gln
	8120					8125					8130			
Ala	Leu	Asn	Gly	Asn	Asp	Asn	Leu	Ala	Asn	Ala	Lys	Gln	Gln	Ala
	8135					8140					8145			
Lys	Gln	Gln	Leu	Ala	Asn	Leu	Thr	His	Leu	Asn	Asp	Ala	Gln	Lys
	8150					8155					8160			
Gln	Ser	Phe	Glu	Ser	Gln	Ile	Thr	Gln	Ala	Pro	Leu	Val	Thr	Asp
	8165					8170					8175			
Val	Thr	Thr	Ile	Asn	Gln	Lys	Ala	Gln	Thr	Leu	Asp	His	Ala	Met
	8180					8185					8190			
Glu	Leu	Leu	Arg	Asn	Ser	Val	Ala	Asp	Asn	Gln	Thr	Thr	Leu	Ala
	8195					8200					8205			

-continued

---

Ser	Glu	Asp	Tyr	His	Asp	Ala	Thr	Ala	Gln	Arg	Gln	Asn	Asp	Tyr
8210						8215					8220			
Asn	Gln	Ala	Val	Thr	Ala	Ala	Asn	Asn	Ile	Ile	Asn	Gln	Thr	Thr
8225						8230					8235			
Ser	Pro	Thr	Met	Asn	Pro	Asp	Asp	Val	Asn	Gly	Ala	Thr	Thr	Gln
8240						8245					8250			
Val	Asn	Asn	Thr	Lys	Val	Ala	Leu	Asp	Gly	Asp	Glu	Asn	Leu	Ala
8255						8260					8265			
Ala	Ala	Lys	Gln	Gln	Ala	Asn	Asn	Arg	Leu	Asp	Gln	Leu	Asp	His
8270						8275					8280			
Leu	Asn	Asn	Ala	Gln	Lys	Gln	Gln	Leu	Gln	Ser	Gln	Ile	Thr	Gln
8285						8290					8295			
Ser	Ser	Asp	Ile	Ala	Ala	Val	Asn	Gly	His	Lys	Gln	Thr	Ala	Glu
8300						8305					8310			
Ser	Leu	Asn	Thr	Ala	Met	Gly	Asn	Leu	Ile	Asn	Ala	Ile	Ala	Asp
8315						8320					8325			
His	Gln	Ala	Val	Glu	Gln	Arg	Gly	Asn	Phe	Ile	Asn	Ala	Asp	Thr
8330						8335					8340			
Asp	Lys	Gln	Thr	Ala	Tyr	Asn	Thr	Ala	Val	Asn	Glu	Ala	Ala	Ala
8345						8350					8355			
Met	Ile	Asn	Lys	Gln	Thr	Gly	Gln	Asn	Ala	Asn	Gln	Thr	Glu	Val
8360						8365					8370			
Glu	Gln	Ala	Ile	Thr	Lys	Val	Gln	Thr	Thr	Leu	Gln	Ala	Leu	Asn
8375						8380					8385			
Gly	Asp	His	Asn	Leu	Gln	Val	Ala	Lys	Thr	Asn	Ala	Thr	Gln	Ala
8390						8395					8400			
Ile	Asp	Ala	Leu	Thr	Ser	Leu	Asn	Asp	Pro	Gln	Lys	Thr	Ala	Leu
8405						8410					8415			
Lys	Asp	Gln	Val	Thr	Ala	Ala	Thr	Leu	Val	Thr	Ala	Val	His	Gln
8420						8425					8430			
Ile	Glu	Gln	Asn	Ala	Asn	Thr	Leu	Asn	Gln	Ala	Met	His	Gly	Leu
8435						8440					8445			
Arg	Gln	Ser	Ile	Gln	Asp	Asn	Ala	Ala	Thr	Lys	Ala	Asn	Ser	Lys
8450						8455					8460			
Tyr	Ile	Asn	Glu	Asp	Gln	Pro	Glu	Gln	Gln	Asn	Tyr	Asp	Gln	Ala
8465						8470					8475			
Val	Gln	Ala	Ala	Asn	Asn	Ile	Ile	Asn	Glu	Gln	Thr	Ala	Thr	Leu
8480						8485					8490			
Asp	Asn	Asn	Ala	Ile	Asn	Gln	Ala	Ala	Thr	Thr	Val	Asn	Thr	Thr
8495						8500					8505			
Lys	Ala	Ala	Leu	His	Gly	Asp	Val	Lys	Leu	Gln	Asn	Asp	Lys	Asp
8510						8515					8520			
His	Ala	Lys	Gln	Thr	Val	Ser	Gln	Leu	Ala	His	Leu	Asn	Asn	Ala
8525						8530					8535			
Gln	Lys	His	Met	Glu	Asp	Thr	Leu	Ile	Asp	Ser	Glu	Thr	Thr	Arg
8540						8545					8550			
Thr	Ala	Val	Lys	Gln	Asp	Leu	Thr	Glu	Ala	Gln	Ala	Leu	Asp	Gln
8555						8560					8565			
Leu	Met	Asp	Ala	Leu	Gln	Gln	Ser	Ile	Ala	Asp	Lys	Asp	Ala	Thr
8570						8575					8580			
Arg	Ala	Ser	Ser	Ala	Tyr	Val	Asn	Ala	Glu	Pro	Asn	Lys	Lys	Gln

-continued

8585	8590	8595
Ser Tyr Asp Glu Ala Val 8600	Gln Asn Ala Glu Ser 8605	Ile Ile Ala Gly 8610
Leu Asn Asn Pro Thr Ile 8615	Asn Lys Gly Asn Val 8620	Ser Ser Ala Thr 8625
Gln Ala Val Ile Ser Ser 8630	Lys Asn Ala Leu Asp 8635	Gly Val Glu Arg 8640
Leu Ala Gln Asp Lys Gln 8645	Thr Ala Gly Asn Ser 8650	Leu Asn His Leu 8655
Asp Gln Leu Thr Pro Ala 8660	Gln Gln Gln Ala Leu 8665	Glu Asn Gln Ile 8670
Asn Asn Ala Thr Thr Arg 8675	Gly Glu Val Ala Gln 8680	Lys Leu Thr Glu 8685
Ala Gln Ala Leu Asn Gln 8690	Ala Met Glu Ala Leu 8695	Arg Asn Ser Ile 8700
Gln Asp Gln Gln Gln Thr 8705	Glu Ala Gly Ser Lys 8710	Phe Ile Asn Glu 8715
Asp Lys Pro Gln Lys Asp 8720	Ala Tyr Gln Ala Ala 8725	Val Gln Asn Ala 8730
Lys Asp Leu Ile Asn Gln 8735	Thr Asn Asn Pro Thr 8740	Leu Asp Lys Ala 8745
Gln Val Glu Gln Leu Thr 8750	Gln Ala Val Asn Gln 8755	Ala Lys Asp Asn 8760
Leu His Gly Asp Gln Lys 8765	Leu Ala Asp Asp Lys 8770	Gln His Ala Val 8775
Thr Asp Leu Asn Gln Leu 8780	Asn Gly Leu Asn Asn 8785	Pro Gln Arg Gln 8790
Ala Leu Glu Ser Gln Ile 8795	Asn Asn Ala Ala Thr 8800	Arg Gly Glu Val 8805
Ala Gln Lys Leu Ala Glu 8810	Ala Lys Ala Leu Asp 8815	Gln Ala Met Gln 8820
Ala Leu Arg Asn Ser Ile 8825	Gln Asp Gln Gln Gln 8830	Thr Glu Ser Gly 8835
Ser Lys Phe Ile Asn Glu 8840	Asp Lys Pro Gln Lys 8845	Asp Ala Tyr Gln 8850
Ala Ala Val Gln Asn Ala 8855	Lys Asp Leu Ile Asn 8860	Gln Thr Gly Asn 8865
Pro Thr Leu Asp Lys Ser 8870	Gln Val Glu Gln Leu 8875	Thr Gln Ala Val 8880
Thr Thr Ala Lys Asp Asn 8885	Leu His Gly Asp Gln 8890	Lys Leu Ala Arg 8895
Asp Gln Gln Gln Ala Val 8900	Thr Thr Val Asn Ala 8905	Leu Pro Asn Leu 8910
Asn His Ala Gln Gln Gln 8915	Ala Leu Thr Asp Ala 8920	Ile Asn Ala Ala 8925
Pro Thr Arg Thr Glu Val 8930	Ala Gln His Val Gln 8935	Thr Ala Thr Glu 8940
Leu Asp His Ala Met Glu 8945	Thr Leu Lys Asn Lys 8950	Val Asp Gln Val 8955
Asn Thr Asp Lys Ala Gln 8960	Pro Asn Tyr Thr Glu 8965	Ala Ser Thr Asp 8970

-continued

---

Lys	Lys	Glu	Ala	Val	Asp	Gln	Ala	Leu	Gln	Ala	Ala	Glu	Ser	Ile
	8975					8980						8985		
Thr	Asp	Pro	Thr	Asn	Gly	Ser	Asn	Ala	Asn	Lys	Asp	Ala	Val	Asp
	8990					8995						9000		
Gln	Val	Leu	Thr	Lys	Leu	Gln	Glu	Lys	Glu	Asn	Glu	Leu	Asn	Gly
	9005					9010						9015		
Asn	Glu	Arg	Val	Ala	Glu	Ala	Lys	Thr	Gln	Ala	Lys	Gln	Thr	Ile
	9020					9025						9030		
Asp	Gln	Leu	Thr	His	Leu	Asn	Ala	Asp	Gln	Ile	Ala	Thr	Ala	Lys
	9035					9040						9045		
Gln	Asn	Ile	Asp	Gln	Ala	Thr	Lys	Leu	Gln	Pro	Ile	Ala	Glu	Leu
	9050					9055						9060		
Val	Asp	Gln	Ala	Thr	Gln	Leu	Asn	Gln	Ser	Met	Asp	Gln	Leu	Gln
	9065					9070						9075		
Gln	Ala	Val	Asn	Glu	His	Ala	Asn	Val	Glu	Gln	Thr	Val	Asp	Tyr
	9080					9085						9090		
Thr	Gln	Ala	Asp	Ser	Asp	Lys	Gln	Asn	Ala	Tyr	Lys	Gln	Ala	Ile
	9095					9100						9105		
Ala	Asp	Ala	Glu	Asn	Val	Leu	Lys	Gln	Asn	Ala	Asn	Lys	Gln	Gln
	9110					9115						9120		
Val	Asp	Gln	Ala	Leu	Gln	Asn	Ile	Leu	Asn	Ala	Lys	Gln	Ala	Leu
	9125					9130						9135		
Asn	Gly	Asp	Glu	Arg	Val	Ala	Leu	Ala	Lys	Thr	Asn	Gly	Lys	His
	9140					9145						9150		
Asp	Ile	Asp	Gln	Leu	Asn	Ala	Leu	Asn	Asn	Ala	Gln	Gln	Asp	Gly
	9155					9160						9165		
Phe	Lys	Gly	Arg	Ile	Asp	Gln	Ser	Asn	Asp	Leu	Asn	Gln	Ile	Gln
	9170					9175						9180		
Gln	Ile	Val	Asp	Glu	Ala	Lys	Ala	Leu	Asn	Arg	Ala	Met	Asp	Gln
	9185					9190						9195		
Leu	Ser	Gln	Glu	Ile	Thr	Asp	Asn	Glu	Gly	Arg	Thr	Lys	Gly	Ser
	9200					9205						9210		
Thr	Asn	Tyr	Val	Asn	Ala	Asp	Thr	Gln	Val	Lys	Gln	Val	Tyr	Asp
	9215					9220						9225		
Glu	Thr	Val	Asp	Lys	Ala	Lys	Gln	Ala	Leu	Asp	Lys	Ser	Thr	Gly
	9230					9235						9240		
Gln	Asn	Leu	Thr	Ala	Lys	Gln	Val	Ile	Lys	Leu	Asn	Asp	Ala	Val
	9245					9250						9255		
Thr	Ala	Ala	Lys	Lys	Ala	Leu	Asn	Gly	Glu	Glu	Arg	Leu	Asn	Asn
	9260					9265						9270		
Arg	Lys	Ala	Glu	Ala	Leu	Gln	Arg	Leu	Asp	Gln	Leu	Thr	His	Leu
	9275					9280						9285		
Asn	Asn	Ala	Gln	Arg	Gln	Leu	Ala	Ile	Gln	Gln	Ile	Asn	Asn	Ala
	9290					9295						9300		
Glu	Thr	Leu	Asn	Lys	Ala	Ser	Arg	Ala	Ile	Asn	Arg	Ala	Thr	Lys
	9305					9310						9315		
Leu	Asp	Asn	Ala	Met	Gly	Ala	Val	Gln	Gln	Tyr	Ile	Asp	Glu	Gln
	9320					9325						9330		
His	Leu	Gly	Val	Ile	Ser	Ser	Thr	Asn	Tyr	Ile	Asn	Ala	Asp	Asp
	9335					9340						9345		
Asn	Leu	Lys	Ala	Asn	Tyr	Asp	Asn	Ala	Ile	Ala	Asn	Ala	Ala	His
	9350					9355						9360		

-continued

---

Glu Leu Asp Lys Val Gln Gly Asn Ala Ile Ala Lys Ala Glu Ala 9365 9370 9375
Glu Gln Leu Lys Gln Asn Ile Ile Asp Ala Gln Asn Ala Leu Asn 9380 9385 9390
Gly Asp Gln Asn Leu Ala Asn Ala Lys Asp Lys Ala Asn Ala Phe 9395 9400 9405
Val Asn Ser Leu Asn Gly Leu Asn Gln Gln Gln Gln Asp Leu Ala 9410 9415 9420
His Lys Ala Ile Asn Asn Ala Asp Thr Val Ser Asp Val Thr Asp 9425 9430 9435
Ile Val Asn Asn Gln Ile Asp Leu Asn Asp Ala Met Glu Thr Leu 9440 9445 9450
Lys His Leu Val Asp Asn Glu Ile Pro Asn Ala Glu Gln Thr Val 9455 9460 9465
Asn Tyr Gln Asn Ala Asp Asp Asn Ala Lys Thr Asn Phe Asp Asp 9470 9475 9480
Ala Lys Arg Leu Ala Asn Thr Leu Leu Asn Ser Asp Asn Thr Asn 9485 9490 9495
Val Asn Asp Ile Asn Gly Ala Ile Gln Ala Val Asn Asp Ala Ile 9500 9505 9510
His Asn Leu Asn Gly Asp Gln Arg Leu Gln Asp Ala Lys Asp Lys 9515 9520 9525
Ala Ile Gln Ser Ile Asn Gln Ala Leu Ala Asn Lys Leu Lys Glu 9530 9535 9540
Ile Glu Ala Ser Asn Ala Thr Asp Gln Asp Lys Leu Ile Ala Lys 9545 9550 9555
Asn Lys Ala Glu Glu Leu Ala Asn Ser Ile Ile Asn Asn Ile Asn 9560 9565 9570
Lys Ala Thr Ser Asn Gln Ala Val Ser Gln Val Gln Thr Ala Gly 9575 9580 9585
Asn His Ala Ile Glu Gln Val His Ala Asn Glu Ile Pro Lys Ala 9590 9595 9600
Lys Ile Asp Ala Asn Lys Asp Val Asp Lys Gln Val Gln Ala Leu 9605 9610 9615
Ile Asp Glu Ile Asp Arg Asn Pro Asn Leu Thr Asp Lys Glu Lys 9620 9625 9630
Gln Ala Leu Lys Asp Arg Ile Asn Gln Ile Leu Gln Gln Gly His 9635 9640 9645
Asn Gly Ile Asn Asn Ala Met Thr Lys Glu Glu Ile Glu Gln Ala 9650 9655 9660
Lys Ala Gln Leu Ala Gln Ala Leu Gln Asp Ile Lys Asp Leu Val 9665 9670 9675
Lys Ala Lys Glu Asp Ala Lys Gln Asp Val Asp Lys Gln Val Gln 9680 9685 9690
Ala Leu Ile Asp Glu Ile Asp Gln Asn Pro Asn Leu Thr Asp Lys 9695 9700 9705
Glu Lys Gln Ala Leu Lys Tyr Arg Ile Asn Gln Ile Leu Gln Gln 9710 9715 9720
Gly His Asn Asp Ile Asn Asn Ala Leu Thr Lys Glu Glu Ile Glu 9725 9730 9735
Gln Ala Lys Ala Gln Leu Ala Gln Ala Leu Gln Asp Ile Lys Asp

-continued

9740	9745	9750
Leu Val Lys Ala Lys Glu Asp 9755	Ala Lys Asn Ala Ile 9760	Lys Ala Leu 9765
Ala Asn Ala Lys Arg Asp 9770	Gln Ile Asn Ser Asn Pro 9775	Asp Leu Thr 9780
Pro Glu Gln Lys Ala Lys 9785	Ala Leu Lys Glu Ile 9790	Asp Glu Ala Glu 9795
Lys Arg Ala Leu Gln Asn 9800	Val Glu Asn Ala Gln Thr 9805	Ile Asp Gln 9810
Leu Asn Arg Gly Leu Asn 9815	Leu Gly Leu Asp Asp 9820	Ile Arg Asn Thr 9825
His Val Trp Glu Val Asp 9830	Glu Gln Pro Ala Val Asn 9835	Glu Ile Phe 9840
Glu Ala Thr Pro Glu Gln 9845	Ile Leu Val Asn Gly Glu 9850	Leu Ile Val 9855
His Arg Asp Asp Ile Ile 9860	Thr Glu Gln Asp Ile 9865	Leu Ala His Ile 9870
Asn Leu Ile Asp Gln Leu 9875	Ser Ala Glu Val Ile Asp 9880	Thr Pro Ser 9885
Thr Ala Thr Ile Ser Asp 9890	Ser Leu Thr Ala Lys Val 9895	Glu Val Thr 9900
Leu Leu Asp Gly Ser Lys 9905	Val Ile Val Asn Val Pro 9910	Val Lys Val 9915
Val Glu Lys Glu Leu Ser 9920	Val Val Lys Gln Gln Ala 9925	Ile Glu Ser 9930
Ile Glu Asn Ala Ala Gln 9935	Gln Lys Ile Asn Glu Ile 9940	Asn Asn Ser 9945
Val Thr Leu Thr Leu Glu 9950	Gln Lys Glu Ala Ala Ile 9955	Ala Glu Val 9960
Asn Lys Leu Lys Gln Gln 9965	Ala Ile Asp His Val Asn 9970	Asn Ala Pro 9975
Asp Val His Ser Val Glu 9980	Glu Ile Gln Gln Gln Glu 9985	Gln Ala His 9990
Ile Glu Gln Phe Asn Pro 9995	Glu Gln Phe Thr Ile Glu 10000	Gln Ala Lys 10005
Ser Asn Ala Ile Lys Ser 10010	Ile Glu Asp Ala Ile Gln 10015	His Met Ile 10020
Asp Glu Ile Lys Ala Arg 10025	Thr Asp Leu Thr Asp Lys 10030	Glu Lys Gln 10035
Glu Ala Ile Ala Lys Leu 10040	Asn Gln Leu Lys Glu Gln 10045	Ala Ile Gln 10050
Ala Ile Gln Arg Ala Gln 10055	Ser Ile Asp Glu Ile Ser 10060	Glu Gln Leu 10065
Glu Gln Phe Lys Ala Gln 10070	Met Lys Ala Ala Asn Pro 10075	Thr Ala Lys 10080
Glu Leu Ala Lys Arg Lys 10085	Gln Glu Ala Ile Ser Arg 10090	Ile Lys Asp 10095
Phe Ser Asn Glu Lys Ile 10100	Asn Ser Ile Arg Asn Ser 10105	Glu Ile Gly 10110
Thr Ala Asp Glu Lys Gln 10115	Ala Ala Met Asn Gln Ile 10120	Asn Glu Ile 10125

-continued

---

Val Leu	Glu Thr Ile Arg	Asp	Ile Asn Asn Ala His	Thr Leu Gln
10130		10135		10140
Gln Val	Glu Ala Ala Leu	Asn	Asn Gly Ile Ala Arg	Ile Ser Ala
10145		10150		10155
Val Gln	Ile Val Thr Ser	Asp	Arg Ala Lys Gln Ser	Ser Ser Thr
10160		10165		10170
Gly Asn	Glu Ser Asn Ser	His	Leu Thr Ile Gly Tyr	Gly Thr Ala
10175		10180		10185
Asn His	Pro Phe Asn Ser	Ser	Thr Ile Gly His Lys	Lys Lys Leu
10190		10195		10200
Asp Glu	Asp Asp Asp Ile	Asp	Pro Leu His Met Arg	His Phe Ser
10205		10210		10215
Asn Asn	Phe Gly Asn Val	Ile	Lys Asn Ala Ile Gly	Val Val Gly
10220		10225		10230
Ile Ser	Gly Leu Leu Ala	Ser	Phe Trp Phe Phe Ile	Ala Lys Arg
10235		10240		10245
Arg Arg	Lys Glu Asp Glu	Glu	Glu Glu Leu Glu Ile	Arg Asp Asn
10250		10255		10260
Asn Lys	Asp Ser Ile Lys	Glu	Thr Leu Asp Asp Thr	Lys His Leu
10265		10270		10275
Pro Leu	Leu Phe Ala Lys	Arg	Arg Arg Lys Glu Asp	Glu Glu Asp
10280		10285		10290
Val Thr	Val Glu Glu Lys	Asp	Ser Leu Asn Asn Gly	Glu Ser Leu
10295		10300		10305
Asp Lys	Val Lys His Thr	Pro	Phe Phe Leu Pro Lys	Arg Arg Arg
10310		10315		10320
Lys Glu	Asp Glu Glu Asp	Val	Glu Val Thr Asn Glu	Asn Thr Asp
10325		10330		10335
Glu Lys	Val Leu Lys Asp	Asn	Glu His Ser Pro Leu	Leu Phe Ala
10340		10345		10350
Lys Arg	Arg Lys Asp Lys	Glu	Glu Asp Val Glu Thr	Thr Thr Ser
10355		10360		10365
Ile Glu	Ser Lys Asp Glu	Asp	Val Pro Leu Leu Leu	Ala Lys Lys
10370		10375		10380
Lys Asn	Gln Lys Asp Asn	Gln	Ser Lys Asp Lys Lys	Ser Ala Ser
10385		10390		10395
Lys Asn	Thr Ser Lys Lys	Val	Ala Ala Lys Lys Lys	Lys Lys Lys
10400		10405		10410
Ala Lys	Lys Asn Lys Lys			
10415				

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 340

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 25

Met Lys Lys Lys Leu Leu Val Leu Thr Met Ser Thr Leu Phe Ala Thr
1 5 10 15

Gln Ile Met Asn Ser Asn His Ala Lys Ala Ser Val Thr Glu Ser Val
20 25 30

Asp Lys Lys Phe Val Val Pro Glu Ser Gly Ile Asn Lys Ile Ile Pro
35 40 45



-continued

---

Ala Tyr Asp Glu Phe Lys Asn Ser Pro Lys Val Asn Val Ser Asn Leu  
50 55 60

Thr Asp Asn Lys Asn Phe Val Ala Ser Glu Asp Lys Leu Asn Lys Ile  
65 70 75 80

Ala Asp Ser Ser Ala Ala Ser Lys Ile Val Asp Lys Asn Phe Val Val  
85 90 95

Pro Glu Ser Lys Leu Gly Asn Ile Val Pro Glu Tyr Lys Glu Ile Asn  
100 105 110

Asn Arg Val Asn Val Ala Thr Asn Asn Pro Ala Ser Gln Gln Val Asp  
115 120 125

Lys His Phe Val Ala Lys Gly Pro Glu Val Asn Arg Phe Ile Thr Gln  
130 135 140

Asn Lys Val Asn His His Phe Ile Thr Thr Gln Thr His Tyr Lys Lys  
145 150 155 160

Val Ile Thr Ser Tyr Lys Ser Thr His Val His Lys His Val Asn His  
165 170 175

Ala Lys Asp Ser Ile Asn Lys His Phe Ile Val Lys Pro Ser Glu Ser  
180 185 190

Pro Arg Tyr Thr His Pro Ser Gln Ser Leu Ile Ile Lys His His Phe  
195 200 205

Ala Val Pro Gly Tyr His Ala His Lys Phe Val Thr Pro Gly His Ala  
210 215 220

Ser Ile Lys Ile Asn His Phe Cys Val Val Pro Gln Ile Asn Ser Phe  
225 230 235 240

Lys Val Ile Pro Pro Tyr Gly His Asn Ser His Arg Met His Val Pro  
245 250 255

Ser Phe Gln Asn Asn Thr Thr Ala Thr His Gln Asn Ala Lys Val Asn  
260 265 270

Lys Ala Tyr Asp Tyr Lys Tyr Phe Tyr Ser Tyr Lys Val Val Lys Gly  
275 280 285

Val Lys Lys Tyr Phe Ser Phe Ser Gln Ser Asn Gly Tyr Lys Ile Gly  
290 295 300

Lys Pro Ser Leu Asn Ile Lys Asn Val Asn Tyr Gln Tyr Ala Val Pro  
305 310 315 320

Ser Tyr Ser Pro Thr His Tyr Val Pro Glu Phe Lys Gly Ser Leu Pro  
325 330 335

Ala Pro Arg Val  
340

<210> SEQ ID NO 26  
 <211> LENGTH: 130  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 26

Met Asn Phe Asn Asp Ile Glu Thr Met Val Lys Ser Lys Phe Lys Asp  
1 5 10 15

Ile Lys Lys His Ala Glu Glu Ile Ala His Glu Ile Glu Val Arg Ser  
20 25 30

Gly Tyr Leu Arg Lys Ala Glu Gln Tyr Lys Arg Leu Glu Phe Asn Leu  
35 40 45

Ser Phe Ala Leu Asp Asp Ile Glu Ser Thr Ala Lys Asp Val Gln Thr  
50 55 60

-continued

---

Ala Lys Ser Ser Ala Asn Lys Asp Ser Val Thr Val Lys Gly Lys Ala  
65 70 75 80

Pro Asn Thr Leu Tyr Ile Glu Lys Arg Asn Leu Met Lys Gln Lys Leu  
85 90 95

Glu Met Leu Gly Glu Asp Ile Asp Lys Asn Lys Glu Ser Leu Gln Lys  
100 105 110

Ala Lys Glu Ile Ala Gly Glu Lys Ala Ser Glu Tyr Phe Asn Lys Ala  
115 120 125

Met Asn  
130

<210> SEQ ID NO 27  
<211> LENGTH: 636  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 27

Met Lys Lys Gln Ile Ile Ser Leu Gly Ala Leu Ala Val Ala Ser Ser  
1 5 10 15

Leu Phe Thr Trp Asp Asn Lys Ala Asp Ala Ile Val Thr Lys Asp Tyr  
20 25 30

Ser Gly Lys Ser Gln Val Asn Ala Gly Ser Lys Asn Gly Thr Leu Ile  
35 40 45

Asp Ser Arg Tyr Leu Asn Ser Ala Leu Tyr Tyr Leu Glu Asp Tyr Ile  
50 55 60

Ile Tyr Ala Ile Gly Leu Thr Asn Lys Tyr Glu Tyr Gly Asp Asn Ile  
65 70 75 80

Tyr Lys Glu Ala Lys Asp Arg Leu Leu Glu Lys Val Leu Arg Glu Asp  
85 90 95

Gln Tyr Leu Leu Glu Arg Lys Lys Ser Gln Tyr Glu Asp Tyr Lys Gln  
100 105 110

Trp Tyr Ala Asn Tyr Lys Lys Glu Asn Pro Arg Thr Asp Leu Lys Met  
115 120 125

Ala Asn Phe His Lys Tyr Asn Leu Glu Glu Leu Ser Met Lys Glu Tyr  
130 135 140

Asn Glu Leu Gln Asp Ala Leu Lys Arg Ala Leu Asp Asp Phe His Arg  
145 150 155 160

Glu Val Lys Asp Ile Lys Asp Lys Asn Ser Asp Leu Lys Thr Phe Asn  
165 170 175

Ala Ala Glu Glu Asp Lys Ala Thr Lys Glu Val Tyr Asp Leu Val Ser  
180 185 190

Glu Ile Asp Thr Leu Val Val Ser Tyr Tyr Gly Asp Lys Asp Tyr Gly  
195 200 205

Glu His Ala Lys Glu Leu Arg Ala Lys Leu Asp Leu Ile Leu Gly Asp  
210 215 220

Thr Asp Asn Pro His Lys Ile Thr Asn Glu Arg Ile Lys Lys Glu Met  
225 230 235 240

Ile Asp Asp Leu Asn Ser Ile Ile Asp Asp Phe Phe Met Glu Thr Lys  
245 250 255

Gln Asn Arg Pro Lys Ser Ile Thr Lys Tyr Asn Pro Thr Thr His Asn  
260 265 270

Tyr Lys Thr Asn Ser Asp Asn Lys Pro Asn Phe Asp Lys Leu Val Glu  
275 280 285

-continued

---

Glu Thr Lys Lys Ala Val Lys Glu Ala Asp Asp Ser Trp Lys Lys Lys  
 290 295 300  
 Thr Val Lys Lys Tyr Gly Glu Thr Glu Thr Lys Ser Pro Val Val Lys  
 305 310 315 320  
 Glu Glu Lys Lys Val Glu Glu Pro Gln Ala Pro Lys Val Asp Asn Gln  
 325 330 335  
 Gln Glu Val Lys Thr Thr Ala Gly Lys Ala Glu Glu Thr Thr Gln Pro  
 340 345 350  
 Val Ala Gln Pro Leu Val Lys Ile Pro Gln Gly Thr Ile Thr Gly Glu  
 355 360 365  
 Ile Val Lys Gly Pro Glu Tyr Pro Thr Met Glu Asn Lys Thr Val Gln  
 370 375 380  
 Gly Glu Ile Val Gln Gly Pro Asp Phe Leu Thr Met Glu Gln Ser Gly  
 385 390 395 400  
 Pro Ser Leu Ser Asn Asn Tyr Thr Asn Pro Pro Leu Thr Asn Pro Ile  
 405 410 415  
 Leu Glu Gly Leu Glu Gly Ser Ser Ser Lys Leu Glu Ile Lys Pro Gln  
 420 425 430  
 Gly Thr Glu Ser Thr Leu Lys Gly Thr Gln Gly Glu Ser Ser Asp Ile  
 435 440 445  
 Glu Val Lys Pro Gln Ala Thr Glu Thr Thr Glu Ala Ser Gln Tyr Gly  
 450 455 460  
 Pro Arg Pro Gln Phe Asn Lys Thr Pro Lys Tyr Val Lys Tyr Arg Asp  
 465 470 475 480  
 Ala Gly Thr Gly Ile Arg Glu Tyr Asn Asp Gly Thr Phe Gly Tyr Glu  
 485 490 495  
 Ala Arg Pro Arg Phe Asn Lys Pro Ser Glu Thr Asn Ala Tyr Asn Val  
 500 505 510  
 Thr Thr His Ala Asn Gly Gln Val Ser Tyr Gly Ala Arg Pro Thr Tyr  
 515 520 525  
 Lys Lys Pro Ser Glu Thr Asn Ala Tyr Asn Val Thr Thr His Ala Asn  
 530 535 540  
 Gly Gln Val Ser Tyr Gly Ala Arg Pro Thr Gln Asn Lys Pro Ser Lys  
 545 550 555 560  
 Thr Asn Ala Tyr Asn Val Thr Thr His Gly Asn Gly Gln Val Ser Tyr  
 565 570 575  
 Gly Ala Arg Pro Thr Gln Asn Lys Pro Ser Lys Thr Asn Ala Tyr Asn  
 580 585 590  
 Val Thr Thr His Ala Asn Gly Gln Val Ser Tyr Gly Ala Arg Pro Thr  
 595 600 605  
 Tyr Lys Lys Pro Ser Lys Thr Asn Ala Tyr Asn Val Thr Thr His Ala  
 610 615 620  
 Asp Gly Thr Ala Thr Tyr Gly Pro Arg Val Thr Lys  
 625 630 635

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 745

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 28

Ala Glu Gln His Thr Pro Met Lys Ala His Ala Val Thr Thr Ile Asp  
 1 5 10 15

-continued

---

Lys Ala Thr Thr Asp Lys Gln Gln Val Pro Pro Thr Lys Glu Ala Ala  
 20 25 30  
 His His Ser Gly Lys Glu Ala Ala Thr Asn Val Ser Ala Ser Ala Gln  
 35 40 45  
 Gly Thr Ala Asp Asp Thr Asn Ser Lys Val Thr Ser Asn Ala Pro Ser  
 50 55 60  
 Asn Lys Pro Ser Thr Val Val Ser Thr Lys Val Asn Glu Thr Arg Asp  
 65 70 75 80  
 Val Asp Thr Gln Gln Ala Ser Thr Gln Lys Pro Thr His Thr Ala Thr  
 85 90 95  
 Phe Lys Leu Ser Asn Ala Lys Thr Ala Ser Leu Ser Pro Arg Met Phe  
 100 105 110  
 Ala Ala Asn Ala Pro Gln Thr Thr Thr His Lys Ile Leu His Thr Asn  
 115 120 125  
 Asp Ile His Gly Arg Leu Ala Glu Glu Lys Gly Arg Val Ile Gly Met  
 130 135 140  
 Ala Lys Leu Lys Thr Val Lys Glu Gln Glu Lys Pro Asp Leu Met Leu  
 145 150 155 160  
 Asp Ala Gly Asp Ala Phe Gln Gly Leu Pro Leu Ser Asn Gln Ser Lys  
 165 170 175  
 Gly Glu Glu Met Ala Lys Ala Met Asn Ala Val Gly Tyr Asp Ala Met  
 180 185 190  
 Ala Val Gly Asn His Glu Phe Asp Phe Gly Tyr Asp Gln Leu Lys Lys  
 195 200 205  
 Leu Glu Gly Met Leu Asp Phe Pro Met Leu Ser Thr Asn Val Tyr Lys  
 210 215 220  
 Asp Gly Lys Arg Ala Phe Lys Pro Ser Thr Ile Val Thr Lys Asn Gly  
 225 230 235 240  
 Ile Arg Tyr Gly Ile Ile Gly Val Thr Thr Pro Glu Thr Lys Thr Lys  
 245 250 255  
 Thr Arg Pro Glu Gly Ile Lys Gly Val Glu Phe Arg Asp Pro Leu Gln  
 260 265 270  
 Ser Val Thr Ala Glu Met Met Arg Ile Tyr Lys Asp Val Asp Thr Phe  
 275 280 285  
 Val Val Ile Ser His Leu Gly Ile Asp Pro Ser Thr Gln Glu Thr Trp  
 290 295 300  
 Arg Gly Asp Tyr Leu Val Lys Gln Leu Ser Gln Asn Pro Gln Leu Lys  
 305 310 315 320  
 Lys Arg Ile Thr Val Ile Asp Gly His Ser His Thr Val Leu Gln Asn  
 325 330 335  
 Gly Gln Ile Tyr Asn Asn Asp Ala Leu Ala Gln Thr Gly Thr Ala Leu  
 340 345 350  
 Ala Asn Ile Gly Lys Ile Thr Phe Asn Tyr Arg Asn Gly Glu Val Ser  
 355 360 365  
 Asn Ile Lys Pro Ser Leu Ile Asn Val Lys Asp Val Glu Asn Val Thr  
 370 375 380  
 Pro Asn Lys Ala Leu Ala Glu Gln Ile Asn Gln Ala Asp Gln Thr Phe  
 385 390 395 400  
 Arg Ala Gln Thr Ala Glu Val Ile Ile Pro Asn Asn Thr Ile Asp Phe  
 405 410 415  
 Lys Gly Glu Arg Asp Asp Val Arg Thr Arg Glu Thr Asn Leu Gly Asn  
 420 425 430

-continued

---

Ala Ile Ala Asp Ala Met Glu Ala Tyr Gly Val Lys Asn Phe Ser Lys  
 435 440 445

Lys Thr Asp Phe Ala Val Thr Asn Gly Gly Gly Ile Arg Ala Ser Ile  
 450 455 460

Ala Lys Gly Lys Val Thr Arg Tyr Asp Leu Ile Ser Val Leu Pro Phe  
 465 470 475 480

Gly Asn Thr Ile Ala Gln Ile Asp Val Lys Gly Ser Asp Val Trp Thr  
 485 490 495

Ala Phe Glu His Ser Leu Gly Ala Pro Thr Thr Gln Lys Asp Gly Lys  
 500 505 510

Thr Val Leu Thr Ala Asn Gly Gly Leu Leu His Ile Ser Asp Ser Ile  
 515 520 525

Arg Val Tyr Tyr Asp Ile Asn Lys Pro Ser Gly Lys Arg Ile Asn Ala  
 530 535 540

Ile Gln Ile Leu Asn Lys Glu Thr Gly Lys Phe Glu Asn Ile Asp Leu  
 545 550 555 560

Lys Arg Val Tyr His Val Thr Met Asn Asp Phe Thr Ala Ser Gly Gly  
 565 570 575

Asp Gly Tyr Ser Met Phe Gly Gly Pro Arg Glu Glu Gly Ile Ser Leu  
 580 585 590

Asp Gln Val Leu Ala Ser Tyr Leu Lys Thr Ala Asn Leu Ala Lys Tyr  
 595 600 605

Asp Thr Thr Glu Pro Gln Arg Met Leu Leu Gly Lys Pro Ala Val Ser  
 610 615 620

Glu Gln Pro Ala Lys Gly Gln Gln Gly Ser Lys Gly Ser Lys Ser Gly  
 625 630 635 640

Lys Asp Thr Gln Pro Ile Gly Asp Asp Lys Val Met Asp Pro Ala Lys  
 645 650 655

Lys Pro Ala Pro Gly Lys Val Val Leu Leu Leu Ala His Arg Gly Thr  
 660 665 670

Val Ser Ser Gly Thr Glu Gly Ser Gly Arg Thr Ile Glu Gly Ala Thr  
 675 680 685

Val Ser Ser Lys Ser Gly Lys Gln Leu Ala Arg Met Ser Val Pro Lys  
 690 695 700

Gly Ser Ala His Glu Lys Gln Leu Pro Lys Thr Gly Thr Asn Gln Ser  
 705 710 715 720

Ser Ser Pro Glu Ala Met Phe Val Leu Leu Ala Gly Ile Gly Leu Ile  
 725 730 735

Ala Thr Val Arg Arg Arg Lys Ala Ser  
 740 745

&lt;210&gt; SEQ ID NO 29

&lt;211&gt; LENGTH: 628

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 29

Met Ser Asp Arg Phe Ile Lys Phe Asn Asp Glu Gln Leu Asp Ala Lys  
 1 5 10 15

Gln Val Met Met Leu Gln Asp Leu Ala Arg Leu Leu Leu Lys Asn Glu  
 20 25 30

Gln Thr Gln Val Lys Ile Gln Lys Phe Pro Tyr Tyr Asn Pro Val Gln  
 35 40 45

-continued

---

Asn Val Leu Ile Thr Ser Trp Phe Trp Ser His Arg Pro Ser His Ile  
 50 55 60  
 Glu Met Ala Gly Leu Lys Thr Asp Val Met Leu Ala Ala Tyr Gly Tyr  
 65 70 75 80  
 His Met Met Asp Val Gln Ile Val Asn Glu Val Val Gln Asp Lys Thr  
 85 90 95  
 Phe Lys His Pro Lys Phe Tyr Gln Gln Leu Phe Lys Leu Leu Glu Asp  
 100 105 110  
 Met Arg Val Leu Asn Ser Ile Lys Val Glu Arg Pro Ser Thr Ala Lys  
 115 120 125  
 Leu Ile Asp Leu Arg Leu Asp Thr Arg Ile Ser Tyr Thr Glu Ser Gln  
 130 135 140  
 Ile Lys Val Tyr Arg Thr Lys Thr Gln Tyr Thr Asp Leu Leu Phe Leu  
 145 150 155 160  
 Tyr Leu Glu His Ala Phe Leu Ser Gln Asp Phe Phe Asp Ile Pro Ser  
 165 170 175  
 Ile His Ser Asp Leu Asp Asp Ile Leu Val Asn Met Phe Leu Tyr Leu  
 180 185 190  
 Pro Asn Phe Phe Gln Asn Gln Asn Ser Glu Asp Asn Met Tyr Leu Ala  
 195 200 205  
 Gln Arg Ile Met Tyr Gln Val Asp Asp Ile Leu Lys Glu Asp Met Leu  
 210 215 220  
 Asn Glu Tyr Tyr Tyr Leu Pro Lys Thr Leu Tyr Asn Thr Leu Ala Ser  
 225 230 235 240  
 Pro Glu Phe Asp Asp Leu Lys Arg Thr Asp Ala Ser Gln Val Asp Gly  
 245 250 255  
 Gln Asp Asp Thr Ser Glu Asp Asp Asp Asn Glu Ser Glu Lys Ala Asp  
 260 265 270  
 Ser Lys Ser Ala Asp Ser Glu Ser Lys Gly Gly Ala Tyr Leu Glu Met  
 275 280 285  
 Glu Leu His Glu Gly Gln Asn Ser Glu Thr Leu Gly Asn Asp Glu Ala  
 290 295 300  
 Arg Glu Gly Asp Ala Thr Asp Asp Met Thr Asp Met Met Thr Lys Lys  
 305 310 315 320  
 Gly Lys Gly Ser Asn Asp Thr Leu Asn Arg Glu Glu Gly Asp Ala Val  
 325 330 335  
 Gly Gln Ser Gln Ala Phe Gln Leu Asp Gly Val Asn Lys Asn Val Glu  
 340 345 350  
 Ile Lys Trp Gln Ile Pro Glu Ile Glu Pro Gln Tyr Val Leu Glu Tyr  
 355 360 365  
 Gln Glu Ser Lys Gln Asp Val Gln Tyr Glu Ile Lys Asp Leu Ile Gln  
 370 375 380  
 Ile Ile Lys Lys Thr Ile Glu Arg Glu Gln Arg Asp Ala Arg Phe Asn  
 385 390 395 400  
 Leu Thr Lys Gly Arg Leu Gln Lys Asp Leu Ile Asn Trp Phe Ile Asp  
 405 410 415  
 Asp Gln Tyr Lys Leu Phe Tyr Lys Lys Gln Asp Leu Ser Lys Ser Phe  
 420 425 430  
 Asp Ala Thr Phe Thr Leu Leu Ile Asp Ala Ser Ala Ser Met His Asp  
 435 440 445  
 Lys Met Ala Glu Thr Lys Lys Gly Val Val Leu Phe His Glu Thr Leu

-continued

---

450	455	460
Lys Ala Leu Asn Ile 465	Lys His Glu Ile 470	Leu Ser Phe Ser Glu Asp Ala 475
Phe Asp Ser Asp 485	Glu His Ala Gln Pro 490	Asn Ile Ile Asn Glu Ile Ile 495
Asn Tyr Asp Tyr Ser Thr 500	Phe Glu Lys Asp Gly 505	Pro Arg Ile Met Ala 510
Leu Glu Pro Gln Asp Asp 515	Asn Arg Asp Gly Val 520	Ala Ile Arg Val Ala 525
Ser Glu Arg Leu Met Arg 530	Arg Asn Gln His 535	Gln Arg Phe Leu Ile Val 540
Phe Ser Asp Gly Glu Pro 545	Ser Ala Phe Asn Tyr 550	Ser Gln Asp Gly Ile 555
Ile Asp Thr Tyr Glu Ala Val 565	Glu Met Ser Arg Lys 570	Phe Gly Ile Glu 575
Val Phe Asn Val Phe Leu Ser 580	Gln Asp Pro Ile Thr 585	Glu Asp Val Glu 590
Gln Thr Ile His Asn Ile Tyr 595	Gly Gln Tyr Ala 600	Ile Phe Val Glu Gly 605
Val Ala His Leu Pro Gly 610	His Leu Ser Pro 615	Leu Leu Lys Lys Leu Leu 620
Leu Lys Ser Leu 625		

<210> SEQ ID NO 30  
<211> LENGTH: 154  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 30

Ala Glu Ile Asn Lys Gln Thr Thr Ser Gln Gly Val Thr Thr Glu Lys 1 5 10 15
Asn Asn Gly Ile Ala Val Leu Glu Gln Asp Val Ile Thr Pro Thr Val 20 25 30
Lys Pro Gln Ala Lys Gln Asp Ile Ile Gln Ala Val Thr Thr Arg Lys 35 40 45
Gln Gln Ile Lys Lys Ser Asn Ala Ser Leu Gln Asp Glu Lys Asp Val 50 55 60
Ala Asn Asp Lys Ile Gly Lys Ile Glu Thr Lys Ala Ile Lys Asp Ile 65 70 75 80
Asp Ala Ala Thr Thr Asn Ala Gln Val Glu Ala Ile Lys Thr Lys Ala 85 90 95
Ile Asn Asp Ile Asn Gln Thr Thr Pro Ala Thr Thr Ala Lys Ala Ala 100 105 110
Ala Leu Glu Glu Phe Asp Glu Val Val Gln Ala Gln Ile Asp Gln Ala 115 120 125
Pro Leu Asn Pro Asp Thr Thr Asn Glu Glu Val Ala Glu Ala Ile Glu 130 135 140
Arg Ile Asn Ala Ala Lys Val Ser Gly Val 145 150

<210> SEQ ID NO 31  
<211> LENGTH: 584  
<212> TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 31

```

Met Lys Phe Lys Ser Leu Ile Thr Thr Thr Leu Ala Leu Gly Val Leu
1           5           10           15
Ala Ser Thr Gly Ala Asn Phe Asn Asn Asn Glu Ala Ser Ala Ala Ala
20           25           30
Lys Pro Leu Asp Lys Ser Ser Ser Ser Leu His His Gly Tyr Ser Lys
35           40           45
Val His Val Pro Tyr Ala Ile Thr Val Asn Gly Thr Ser Gln Asn Ile
50           55           60
Leu Ser Ser Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp
65           70           75           80
Leu Glu Asp Arg Val Lys Ser Val Leu Lys Ser Asp Arg Gly Ile Ser
85           90           95
Asp Ile Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Tyr Phe
100          105          110
Lys Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ala Gly Ile Tyr Thr
115          120          125
Ala Asp Leu Ile Asn Thr Ser Glu Ile Lys Ala Ile Asn Ile Asn Val
130          135          140
Asp Thr Lys Lys Gln Val Glu Asp Lys Lys Lys Asp Lys Ala Asn Tyr
145          150          155          160
Gln Val Pro Tyr Thr Ile Thr Val Asn Gly Thr Ser Gln Asn Ile Leu
165          170          175
Ser Asn Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp Leu
180          185          190
Glu Asp Lys Val Lys Ser Val Leu Glu Ser Asn Arg Gly Ile Thr Asp
195          200          205
Val Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Asn Phe Lys
210          215          220
Asn Gly Thr Lys Lys Val Ile Asp Leu Lys Ser Gly Ile Tyr Thr Ala
225          230          235          240
Asn Leu Ile Asn Ser Ser Asp Ile Lys Ser Ile Asn Ile Asn Val Asp
245          250          255
Thr Lys Lys His Ile Glu Asn Lys Ala Lys Arg Asn Tyr Gln Val Pro
260          265          270
Tyr Ser Ile Asn Leu Asn Gly Thr Ser Thr Asn Ile Leu Ser Asn Leu
275          280          285
Ser Phe Ser Asn Lys Pro Trp Thr Asn Tyr Lys Asn Leu Thr Ser Gln
290          295          300
Ile Lys Ser Val Leu Lys His Asp Arg Gly Ile Ser Glu Gln Asp Leu
305          310          315          320
Lys Tyr Ala Lys Lys Ala Tyr Tyr Thr Val Tyr Phe Lys Asn Gly Gly
325          330          335
Lys Arg Ile Leu Gln Leu Asn Ser Lys Asn Tyr Thr Ala Asn Leu Val
340          345          350
His Ala Lys Asp Val Lys Arg Ile Glu Ile Thr Val Lys Thr Gly Thr
355          360          365
Lys Ala Lys Ala Asp Arg Tyr Val Pro Tyr Thr Ile Ala Val Asn Gly
370          375          380
Thr Ser Thr Pro Ile Leu Ser Asp Leu Lys Phe Thr Gly Asp Pro Arg

```



-continued

---

```

385             390             395             400
Val Gly Tyr Lys Asp Ile Ser Lys Lys Val Lys Ser Val Leu Lys His
                405
Asp Arg Gly Ile Gly Glu Arg Glu Leu Lys Tyr Ala Lys Lys Ala Thr
                420             425             430
Tyr Thr Val His Phe Lys Asn Gly Thr Lys Lys Val Ile Asn Ile Asn
                435             440             445
Ser Asn Ile Ser Gln Leu Asn Leu Leu Tyr Val Gln Asp Ile Lys Lys
                450             455             460
Ile Asp Ile Asp Val Lys Thr Gly Thr Lys Ala Lys Ala Asp Ser Tyr
465             470             475
Val Pro Tyr Thr Ile Ala Val Asn Gly Thr Ser Thr Pro Ile Leu Ser
                485             490             495
Lys Leu Lys Ile Ser Asn Lys Gln Leu Ile Ser Tyr Lys Tyr Leu Asn
                500             505             510
Asp Lys Val Lys Ser Val Leu Lys Ser Glu Arg Gly Ile Ser Asp Leu
515             520             525
Asp Leu Lys Phe Ala Lys Gln Ala Lys Tyr Thr Val Tyr Phe Lys Asn
530             535             540
Gly Lys Lys Gln Val Val Asn Leu Lys Ser Asp Ile Phe Thr Pro Asn
545             550             555             560
Leu Phe Ser Ala Lys Asp Ile Lys Lys Ile Asp Ile Asp Val Lys Gln
                565             570             575
Tyr Thr Lys Ser Lys Lys Asn Lys
                580

<210> SEQ ID NO 32
<211> LENGTH: 508
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 32
Met Lys Asn Lys Leu Leu Val Leu Ser Leu Gly Ala Leu Cys Val Ser
1             5             10             15
Gln Ile Trp Glu Ser Asn Arg Ala Ser Ala Val Val Ser Gly Glu Lys
                20             25             30
Asn Pro Tyr Val Ser Glu Ser Leu Lys Leu Thr Asn Asn Lys Asn Lys
                35             40             45
Ser Arg Thr Val Glu Glu Tyr Lys Lys Ser Leu Asp Asp Leu Ile Trp
50             55             60
Ser Phe Pro Asn Leu Asp Asn Glu Arg Phe Asp Asn Pro Glu Tyr Lys
65             70             75             80
Glu Ala Met Lys Lys Tyr Gln Gln Arg Phe Met Ala Glu Asp Glu Ala
                85             90             95
Leu Lys Lys Phe Phe Ser Glu Glu Lys Lys Ile Lys Asn Gly Asn Thr
100            105            110
Asp Asn Leu Asp Tyr Leu Gly Leu Ser His Glu Arg Tyr Glu Ser Val
115            120            125
Phe Asn Thr Leu Lys Lys Gln Ser Glu Glu Phe Leu Lys Glu Ile Glu
130            135            140
Asp Ile Lys Lys Asp Asn Pro Glu Leu Lys Asp Phe Asn Glu Glu Glu
145            150            155            160
Gln Leu Lys Cys Asp Leu Glu Leu Asn Lys Leu Glu Asn Gln Ile Leu

```

-continued

165					170					175					
Met	Leu	Gly	Lys	Thr	Phe	Tyr	Gln	Asn	Tyr	Arg	Asp	Asp	Val	Glu	Ser
			180					185					190		
Leu	Tyr	Ser	Lys	Leu	Asp	Leu	Ile	Met	Gly	Tyr	Lys	Asp	Glu	Glu	Arg
		195					200					205			
Ala	Asn	Lys	Lys	Ala	Val	Asn	Lys	Arg	Met	Leu	Glu	Asn	Lys	Lys	Glu
		210					215					220			
Asp	Leu	Glu	Thr	Ile	Ile	Asp	Glu	Phe	Phe	Ser	Asp	Ile	Asp	Lys	Thr
		225					230					235			240
Arg	Pro	Asn	Asn	Ile	Pro	Val	Leu	Glu	Asp	Glu	Lys	Gln	Glu	Lys	
				245					250					255	
Asn	His	Lys	Asn	Met	Ala	Gln	Leu	Lys	Ser	Asp	Thr	Glu	Ala	Ala	Lys
			260					265					270		
Ser	Asp	Glu	Ser	Lys	Arg	Ser	Lys	Arg	Ser	Lys	Arg	Ser	Leu	Asn	Thr
		275					280					285			
Gln	Asn	His	Lys	Pro	Ala	Ser	Gln	Glu	Val	Ser	Glu	Gln	Gln	Lys	Ala
		290					295					300			
Glu	Tyr	Asp	Lys	Arg	Ala	Glu	Glu	Arg	Lys	Ala	Arg	Phe	Leu	Asp	Asn
		305					310					315			320
Gln	Lys	Ile	Lys	Lys	Thr	Pro	Val	Val	Ser	Leu	Glu	Tyr	Asp	Phe	Glu
			325						330					335	
His	Lys	Gln	Arg	Ile	Asp	Asn	Glu	Asn	Asp	Lys	Lys	Leu	Val	Val	Ser
			340					345					350		
Ala	Pro	Thr	Lys	Lys	Pro	Thr	Ser	Pro	Thr	Thr	Tyr	Thr	Glu	Thr	Thr
			355				360						365		
Thr	Gln	Val	Pro	Met	Pro	Thr	Val	Glu	Arg	Gln	Thr	Gln	Gln	Gln	Ile
		370					375					380			
Ile	Tyr	Asn	Ala	Pro	Lys	Gln	Leu	Ala	Gly	Leu	Asn	Gly	Glu	Ser	His
		385					390					395			400
Asp	Phe	Thr	Thr	Thr	His	Gln	Ser	Pro	Thr	Thr	Ser	Asn	His	Thr	His
				405					410					415	
Asn	Asn	Val	Val	Glu	Phe	Glu	Glu	Thr	Ser	Ala	Leu	Pro	Gly	Arg	Lys
			420					425					430		
Ser	Gly	Ser	Leu	Val	Gly	Ile	Ser	Gln	Ile	Asp	Ser	Ser	His	Leu	Thr
		435					440					445			
Glu	Arg	Glu	Lys	Arg	Val	Ile	Lys	Arg	Glu	His	Val	Arg	Glu	Ala	Gln
		450					455					460			
Lys	Leu	Val	Asp	Asn	Tyr	Lys	Asp	Thr	His	Ser	Tyr	Lys	Asp	Arg	Ile
		465					470					475			480
Asn	Ala	Gln	Gln	Lys	Val	Asn	Thr	Leu	Ser	Glu	Gly	His	Gln	Lys	Arg
				485					490					495	
Phe	Asn	Lys	Gln	Ile	Asn	Lys	Val	Tyr	Asn	Gly	Lys				
			500					505							

&lt;210&gt; SEQ ID NO 33

&lt;211&gt; LENGTH: 520

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 33

Met	Leu	Thr	Leu	Gln	Ile	His	Thr	Gly	Gly	Ile	Asn	Leu	Lys	Lys	Lys
1			5					10					15		

Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile Ala Ser Val Thr

-continued

20				25				30								
Leu	Gly	Thr	35	Leu	Leu	Ile	Ser	Gly	Gly	Val	Thr	Pro	Ala	Ala	Asn	Ala
Ala	Gln	His	50	Asp	Glu	Ala	Gln	Asn	Ala	Phe	Tyr	Gln	Val	Leu	Asn	
Met	Pro	Asn	65	Leu	Asn	Ala	Asp	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu
Lys	Asp	Asp	85	Pro	Ser	Gln	Ser	Ala	Asn	Val	Leu	Gly	Glu	Ala	Gln	Lys
Leu	Asn	Asp	100	Ser	Gln	Ala	Pro	Lys	Ala	Asp	Ala	Gln	Gln	Asn	Asn	Phe
Asn	Lys	Asp	115	Gln	Gln	Ser	Ala	Phe	Tyr	Glu	Ile	Leu	Asn	Met	Pro	Asn
Leu	Asn	Glu	130	Ala	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp
Pro	Ser	Gln	145	Ser	Thr	Asn	Val	Leu	Gly	Glu	Ala	Lys	Lys	Leu	Asn	Glu
Ser	Gln	Ala	165	Pro	Lys	Ala	Asp	Asn	Asn	Phe	Asn	Lys	Glu	Gln	Gln	Asn
Ala	Phe	Tyr	180	Glu	Ile	Leu	Asn	Met	Pro	Asn	Leu	Asn	Glu	Glu	Gln	Arg
Asn	Gly	Phe	195	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser	Gln	Ser	Ala	Asn
Leu	Leu	Ser	210	Glu	Ala	Lys	Lys	Leu	Asn	Glu	Ser	Gln	Ala	Pro	Lys	Ala
Asp	Asn	Lys	225	Phe	Asn	Lys	Glu	Gln	Gln	Asn	Ala	Phe	Tyr	Glu	Ile	Leu
His	Leu	Pro	245	Asn	Leu	Asn	Glu	Glu	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser
Leu	Lys	Asp	260	Asp	Pro	Ser	Gln	Ser	Ala	Asn	Leu	Leu	Ala	Glu	Ala	Lys
Lys	Leu	Asn	275	Asp	Ala	Gln	Ala	Pro	Lys	Ala	Asp	Asn	Lys	Phe	Asn	Lys
Glu	Gln	Gln	290	Asn	Ala	Phe	Tyr	Glu	Ile	Leu	His	Leu	Pro	Asn	Leu	Thr
Glu	Glu	Gln	305	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser
Val	Ser	Lys	325	Glu	Ile	Leu	Ala	Glu	Ala	Lys	Lys	Leu	Asn	Asp	Ala	Gln
Ala	Pro	Lys	340	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Gly	Asn	
Lys	Pro	Gly	355	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Lys
Lys	Pro	Gly	370	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Asn
Lys	Pro	Gly	385	Lys	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Lys
Lys	Pro	Gly	405	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Gly	Asn
Lys	Pro	Gly	420	Lys	Glu	Asp	Gly	Asn	Gly	Val	His	Val	Val	Lys	Pro	Gly

-continued

---

```

Asp Thr Val Asn Asp Ile Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys
    435                                440                                445

Ile Ala Ala Asp Asn Lys Leu Ala Asp Lys Asn Met Ile Lys Pro Gly
    450                                455                                460

Gln Glu Leu Val Val Asp Lys Lys Gln Pro Ala Asn His Ala Asp Ala
465                                470                                475                                480

Asn Lys Ala Gln Ala Leu Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile
                                485                                490                                495

Gly Thr Thr Val Phe Gly Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu
    500                                505                                510

Leu Ala Gly Arg Arg Arg Glu Leu
    515                                520

```

```

<210> SEQ ID NO 34
<211> LENGTH: 291
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

```

```

<400> SEQUENCE: 34

```

```

Ala Gln His Asp Glu Ala Lys Lys Asn Ala Phe Tyr Gln Val Leu Asn
 1                                5                                10                                15

Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu
                                20                                25                                30

Lys Ala Ala Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys
 35                                40                                45

Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln Gln Asn Asn Phe
 50                                55                                60

Asn Lys Asp Lys Lys Ser Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn
65                                70                                75                                80

Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Ala Ala
 85                                90                                95

Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys Leu Asn Glu
100                                105                                110

Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys Glu Lys Lys Asn
115                                120                                125

Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn Glu Glu Gln Arg
130                                135                                140

Asn Gly Phe Ile Gln Ser Leu Lys Ala Ala Pro Ser Gln Ser Ala Asn
145                                150                                155                                160

Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala
165                                170                                175

Asp Asn Lys Phe Asn Lys Glu Lys Lys Asn Ala Phe Tyr Glu Ile Leu
180                                185                                190

His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser
195                                200                                205

Leu Lys Ala Ala Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys
210                                215                                220

Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys
225                                230                                235                                240

Glu Lys Lys Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Thr
245                                250                                255

Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Ala Ala Pro Ser
260                                265                                270

```

-continued

---

Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln  
 275 280 285

Ala Pro Lys  
 290

<210> SEQ ID NO 35  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 35

gctgcacata tggcgcaaca cgatgaagct caac 34

<210> SEQ ID NO 36  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 36

agtggatcct tatgctttgt tagcatctgc 30

<210> SEQ ID NO 37  
 <211> LENGTH: 19  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 37

Met Gly Ser Ser His His His His His His Ser Ser Gly Leu Val Pro  
 1 5 10 15

Arg Gly Ser

<210> SEQ ID NO 38  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 38

aacatatggt caacaaagat caacaaagc 29

<210> SEQ ID NO 39  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 39

aaggatccag attcgtttaa ttttttagc 29

<210> SEQ ID NO 40  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 40

cttcattcaa agtcttaaag cgcgcccaag ccaaagcact aac 43

<210> SEQ ID NO 41  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 41

---

-continued

---

gtttagtgctt tggcttgggg cggctttaag actttgaatg aag 43

<210> SEQ ID NO 42  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 42

catatgttca acaaagataa aaaaagcgcc ttctatgaaa tc 42

<210> SEQ ID NO 43  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 43

gatttcatag aaggcgcttt ttttatcttt gttgaacata tg 42

<210> SEQ ID NO 44  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 44

catatgttca acaaagatgg aggaagcgcc ttctatgaaa tc 42

<210> SEQ ID NO 45  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 45

gatttcatag aaggcgcttc ctccatcttt gttgaacata tg 42

<210> SEQ ID NO 46  
<211> LENGTH: 52  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 46

ggggacaagt ttgtacaaaa aagcaggctg atgactaagt tgaaaaaaga ag 52

<210> SEQ ID NO 47  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 47

aaggatcccc tccaaaatgt aattgccc 28

<210> SEQ ID NO 48  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 48

aaggatccgt ttgtaactct atccaaagac 30

<210> SEQ ID NO 49  
<211> LENGTH: 49  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

---

-continued

---

<400> SEQUENCE: 49

ggggaccact ttgtacaaga aagctgggtg acacctattg cagattcg 49

<210> SEQ ID NO 50

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 50

ggggacaagt ttgtacaaaa aagcaggctc agatagcgat tcagattcag 50

<210> SEQ ID NO 51

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 51

aaggatccct gtattttctc cttaattttc c 31

<210> SEQ ID NO 52

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 52

aaggatccca tggctgcaaa gcaaataatg 30

<210> SEQ ID NO 53

<211> LENGTH: 51

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 53

ggggaccact ttgtacaaga aagctgggtg ccctggtgta acaaatttat g 51

<210> SEQ ID NO 54

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 54

gaaggatccg tttattctag ttaatata gttaatg 37

<210> SEQ ID NO 55

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 55

gaactgcagc tgtatgtctt tggatagagt tac 33

<210> SEQ ID NO 56

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 56

gaaggatccg gtggcttttt tacttggatt ttc 33

<210> SEQ ID NO 57

---

-continued

---

<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 57  
gaactgcagc gacaaactca ttatttgctt tgc 33

<210> SEQ ID NO 58  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 58  
gaactcgagt ctagcttatt tacatgg 27

<210> SEQ ID NO 59  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 59  
gaactcgaga tagaaggcag aatagtaaca aaggattata gtggg 45

<210> SEQ ID NO 60  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 60  
gtaggatcct gggatagagt tacaaac 27

<210> SEQ ID NO 61  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 61  
gaactcgagg cattatgtgt atcacaaatt tggg 34

<210> SEQ ID NO 62  
<211> LENGTH: 43  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 62  
gaactcgaga tagaaggcag agtggtttct ggggagaaga atc 43

<210> SEQ ID NO 63  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 63  
gaactcgagg cagccatgca ttaattattt gcc 33

<210> SEQ ID NO 64  
<211> LENGTH: 677  
<212> TYPE: PRT  
<213> ORGANISM: *Staphylococcus* sp.  
<400> SEQUENCE: 64  
Met Lys Ser Asn Leu Arg Tyr Gly Ile Arg Lys His Lys Leu Gly Ala



-continued

1	5	10	15
Ala Ser Val Phe Leu Gly Thr Met Ile Val Val Gly Met Gly Gln Glu	20	25	30
Lys Glu Ala Ala Ala Ser Glu Gln Asn Asn Thr Thr Val Glu Glu Ser	35	40	45
Gly Ser Ser Ala Thr Glu Ser Lys Ala Ser Glu Thr Gln Thr Thr Thr	50	55	60
Asn Asn Val Asn Thr Ile Asp Glu Thr Gln Ser Tyr Ser Ala Thr Ser	65	70	75
Thr Glu Gln Pro Ser Gln Ser Thr Gln Val Thr Thr Glu Glu Ala Pro	85	90	95
Lys Thr Val Gln Ala Pro Lys Val Glu Thr Ser Arg Val Asp Leu Pro	100	105	110
Ser Glu Lys Val Ala Asp Lys Glu Thr Thr Gly Thr Gln Val Asp Ile	115	120	125
Ala Gln Pro Ser Asn Val Ser Glu Ile Lys Pro Arg Met Lys Arg Ser	130	135	140
Thr Asp Val Thr Ala Val Ala Glu Lys Glu Val Val Glu Glu Thr Lys	145	150	155
Ala Thr Gly Thr Asp Val Thr Asn Lys Val Glu Val Glu Glu Gly Ser	165	170	175
Glu Ile Val Gly His Lys Gln Asp Thr Asn Val Val Asn Pro His Asn	180	185	190
Ala Glu Arg Val Thr Leu Lys Tyr Lys Trp Lys Phe Gly Glu Gly Ile	195	200	205
Lys Ala Gly Asp Tyr Phe Asp Phe Thr Leu Ser Asp Asn Val Glu Thr	210	215	220
His Gly Ile Ser Thr Leu Arg Lys Val Pro Glu Ile Lys Ser Thr Asp	225	230	235
Gly Gln Val Met Ala Thr Gly Glu Ile Ile Gly Glu Arg Lys Val Arg	245	250	255
Tyr Thr Phe Lys Glu Tyr Val Gln Glu Lys Lys Asp Leu Thr Ala Glu	260	265	270
Leu Ser Leu Asn Leu Phe Ile Asp Pro Thr Thr Val Thr Gln Lys Gly	275	280	285
Asn Gln Asn Val Glu Val Lys Leu Gly Glu Thr Thr Val Ser Lys Ile	290	295	300
Phe Asn Ile Gln Tyr Leu Gly Gly Val Arg Asp Asn Trp Gly Val Thr	305	310	315
Ala Asn Gly Arg Ile Asp Thr Leu Asn Lys Val Asp Gly Lys Phe Ser	325	330	335
His Phe Ala Tyr Met Lys Pro Asn Asn Gln Ser Leu Ser Ser Val Thr	340	345	350
Val Thr Gly Gln Val Thr Lys Gly Asn Lys Pro Gly Val Asn Asn Pro	355	360	365
Thr Val Lys Val Tyr Lys His Ile Gly Ser Asp Asp Leu Ala Glu Ser	370	375	380
Val Tyr Ala Lys Leu Asp Asp Val Ser Lys Phe Glu Asp Val Thr Asp	385	390	395
Asn Met Ser Leu Asp Phe Asp Thr Asn Gly Gly Tyr Ser Leu Asn Phe	405	410	415

-continued

---

Asn	Asn	Leu	Asp	Gln	Ser	Lys	Asn	Tyr	Val	Ile	Lys	Tyr	Glu	Gly	Tyr
			420					425					430		
Tyr	Asp	Ser	Asn	Ala	Ser	Asn	Leu	Glu	Phe	Gln	Thr	His	Leu	Phe	Gly
		435					440					445			
Tyr	Tyr	Asn	Tyr	Tyr	Tyr	Thr	Ser	Asn	Leu	Thr	Trp	Lys	Asn	Gly	Val
		450				455					460				
Ala	Phe	Tyr	Ser	Asn	Asn	Ala	Gln	Gly	Asp	Gly	Lys	Asp	Lys	Leu	Lys
465					470					475					480
Glu	Pro	Ile	Ile	Glu	His	Ser	Thr	Pro	Ile	Glu	Leu	Glu	Phe	Lys	Ser
				485					490						495
Glu	Pro	Pro	Val	Glu	Lys	His	Glu	Leu	Thr	Gly	Thr	Ile	Glu	Glu	Ser
			500					505					510		
Asn	Asp	Ser	Lys	Pro	Ile	Asp	Phe	Glu	Tyr	His	Thr	Ala	Val	Glu	Gly
		515					520					525			
Ala	Glu	Gly	His	Ala	Glu	Gly	Thr	Ile	Glu	Thr	Glu	Glu	Asp	Ser	Ile
		530				535					540				
His	Val	Asp	Phe	Glu	Glu	Ser	Thr	His	Glu	Asn	Ser	Lys	His	His	Ala
545					550					555					560
Asp	Val	Val	Glu	Tyr	Glu	Glu	Asp	Thr	Asn	Pro	Gly	Gly	Gly	Gln	Val
				565					570						575
Thr	Thr	Glu	Ser	Asn	Leu	Val	Glu	Phe	Asp	Glu	Asp	Ser	Thr	Lys	Gly
			580					585						590	
Ile	Val	Thr	Gly	Ala	Val	Ser	Asp	His	Thr	Thr	Ile	Glu	Asp	Thr	Lys
		595					600					605			
Glu	Tyr	Thr	Thr	Glu	Ser	Asn	Leu	Ile	Glu	Leu	Val	Asp	Glu	Leu	Pro
		610					615				620				
Glu	Glu	His	Gly	Gln	Ala	Gln	Gly	Pro	Ile	Glu	Glu	Ile	Thr	Glu	Asn
625					630					635					640
Asn	His	His	Ile	Ser	His	Ser	Gly	Leu	Gly	Thr	Glu	Asn	Gly	His	Gly
				645					650					655	
Asn	Tyr	Gly	Val	Ile	Glu	Glu	Ile	Glu	Glu	Asn	Ser	His	Val	Asp	Ile
			660					665					670		
Lys	Ser	Glu	Leu	Gly											
		675													

---

What is claimed is:

1. An isolated polypeptide comprising a variant Protein A (SpA) having (a) at least one amino acid substitution that disrupts Fc binding and (b) at least a second amino acid substitution that disrupts VH3 binding and (c) an amino acid sequence that is at least 70% identical to the amino acid sequence of SEQ ID NO:2.

2. The isolated polypeptide of claim 1, wherein the variant SpA comprises a variant domain D segment.

3. The isolated polypeptide of claim 2, wherein the SpA variant has one or more amino acid substitution at amino acid position 9 or 10 of SEQ ID NO:2 and having an amino acid sequence that is at least 80% identical to the amino acid sequence of SEQ ID NO:2.

4. The isolated polypeptide of claim 3, wherein the amino acid substitution is a lysine residue for a glutamine residue.

5. The isolated polypeptide of claim 3, further comprising an amino acid substitution at amino acid positions 36 and 37 of SEQ ID NO:2.

6. The isolated polypeptide of claim 5, wherein the amino acid sequence of the domain D comprises an alanine residue substitution at amino acid position 36 and 37 of SEQ ID NO:2.

7. The isolated polypeptide of claim 2, further comprising one or more variants of an SpA E domain, A domain, B domain, or C domain.

8. The isolated polypeptide of claim 2, comprising two or more D domain segments.

9. The isolated polypeptide of claim 1, further comprising a non-Protein A segment.

10. The isolated polypeptide of claim 9, wherein the non-Protein A segment is a second antigen segment.

11. The isolated polypeptide of claim 10, wherein the second antigen segment is a staphylococcal antigen segment.

12. The isolated polypeptide of claim 11, wherein the staphylococcal antigen segment is an Emp, EsxA, EsxB, EsaC, Eap, Ebh, EsaB, Coa, vWbp, vWh, Hla, SdrC, SdrD, SdrE, IsdA, IsdB, IsdC, ClfA, ClfB, and/or SasF segment.

13. A peptide composition comprising, in a pharmaceutically acceptable composition, a non-toxicogenic variant Protein A (SpA) peptide having one or more mutations that attenuate the binding of the Protein A domain D to IgG, Fc $\gamma$ , VH3 F(ab)<sub>2</sub>, von Willebrand factor (vWF), and tumor necrosis factor  $\alpha$  receptor 1 (TNFR1), wherein the composition is capable of stimulating an immune response in a subject in need thereof.

14. The composition of claim 13, wherein the SpA variant comprises one or more amino acid substitutions at amino acid positions 9 and 10 of SEQ ID NO:2.

15. The composition of claim 14, wherein the SpA variant comprises a substitution of lysine at amino acid positions 9 and 10 of SEQ ID NO:2.

16. The composition of claim 14, further comprising amino acid substitutions at amino acid positions 36 and 37 of SEQ ID NO:2.

17. The composition of claim 15, wherein the SpA variant comprises an amino acid sequence at least 70% identical to SEQ ID NO:2.

18. The composition of claim 13, wherein the SpA variant comprises the amino acid sequence of SEQ ID NO:8.

19. The composition of claim 13, further comprising at least a second staphylococcal antigen.

20. The composition of claim 19, wherein the second antigen selected from an EsaB, Emp, EsxA, EsxB, EsaC, Eap, Ebh, Coa, vWbp, vWh, Hla, SdrC, SdrD, SdrE, IsdA, IsdB, IsdC, ClfA, ClfB, and/or SasF peptide.

21. The composition of claim 13, wherein the composition contains less than 1% by weight of staphylococcal bacterial components other than the peptide comprising the Protein A domain D peptide.

22. The composition of claim 13, wherein the composition further comprises an adjuvant.

23. The composition of claim 22, wherein the SpA variant is coupled to an adjuvant.

24. An immunogenic composition comprising an isolated peptide comprising a Protein A (SpA) variant having an amino acid substitution at amino acid positions 9, 10, 36, and 37 of SEQ ID NO:2.

25. The composition of any one of claims 1-24, further comprising at least one other staphylococcal antigen or immunogenic fragment thereof selected from the group consisting of: Emp, EsxA, EsxB, EsaC, Eap, Ebh, EsaB, Coa, vWbp, vWh, Hla, SdrC, SdrD, SdrE, IsdA, IsdB, IsdC, ClfA, ClfB, and SasF.

26. The composition of any one of claims 1-25, further comprising a PIA polysaccharide or oligosaccharide.

27. The composition of any one of claims 1-26 further comprising a type V and/or type VIII capsular polysaccharide or oligosaccharide from *S. aureus*.

28. The immunogenic composition of any one of claims 1-27 wherein a staphylococcal capsular polysaccharide is conjugated to a protein carrier.

29. The immunogenic composition of claim 28 wherein the protein carrier is selected from the group consisting of tetanus toxoid, diphtheria toxoid, CRM197, *Haemophilus influenzae* protein D, pneumolysin and alpha toxoid.

30. A vaccine comprising the composition of any one of claims 1-29 and a pharmaceutically acceptable excipient.

31. A method of making a vaccine comprising the steps of mixing antigens to make the composition of any one of claims 1-29 and adding a pharmaceutically acceptable excipient.

32. A method of preventing or treating staphylococcal infection comprising the step of administering the vaccine of claim 30 to a patient in need thereof.

33. A use of the composition of any one of claims 1-29 in the manufacture of a vaccine for treatment or prevention of staphylococcal infection.

34. A method of preparing an immunoglobulin for use in prevention or treatment of staphylococcal infection comprising the steps of immunizing a recipient with the vaccine of claim 30 and isolating immunoglobulin from the recipient.

35. An immunoglobulin prepared by the method of claim 34.

36. A pharmaceutical composition comprising the immunoglobulin of claim 35 and a pharmaceutically acceptable excipient.

37. A method for treatment or prevention of staphylococcal infection comprising a step of administering to a patient an effective amount of the pharmaceutical preparation of claim 36.

38. A use of the pharmaceutical preparation of claim 36 in the manufacture of a medicament for the treatment or prevention of staphylococcal infection.

39. A method for eliciting an immune response against a *staphylococcus* bacterium in a subject comprising providing to the subject an effective amount of a composition comprising a Protein A (SpA) variant having an amino acid substitution at amino acid positions 9 and 10 of SEQ ID NO:2.

40. The method of claim 39, wherein the subject is provided with an effective amount of an SpA variant by administering to the subject a composition comprising:

- i) an isolated SpA variant having an amino acid substitution at amino acid positions 9 and 10 of SEQ ID NO:2, or
- ii) at least one isolated recombinant nucleic acid molecule encoding a SpA variant having an amino acid substitution at amino acid positions 9 and 10 of SEQ ID NO:2.

41. The method of claim 40, wherein the composition comprises an isolated SpA variant.

42. The method of claim 39, where the subject is also administered an adjuvant.

43. The method of claim 41, wherein the composition comprises an adjuvant.

44. The method of claim 41, wherein the SpA variant is coupled to an adjuvant.

45. The method of claim 39, wherein the SpA variant is at least 70% identical to SEQ ID NO:2 and having an amino acid substitution at amino acid position 9 and 10 of SEQ ID NO:2.

46. The method of claim 41, wherein the composition is formulated in a pharmaceutically acceptable composition.

47. The method of claim 39, wherein the *staphylococcus* bacterium is a *S. aureus* bacterium.

48. The method of claim 39, wherein the *staphylococcus* bacterium is resistant to one or more treatments.

49. The method of claim 48, wherein the bacterium is methicillin resistant.

50. The method of claim 39, further comprising administering the composition more than one time to the subject.

51. The method of claim 39, wherein the composition is administered orally, parenterally, subcutaneously, intramuscularly, or intravenously.

52. The method of claim 39, further comprising administering to the subject a composition comprising a second staphylococcal antigen.

53. The method of claim 52, wherein the second staphylococcal antigen is one or more of Emp, EsxA, EsxB, EsaC,

Eap, Ebh, EsaB, Coa, vWbp, vWh, Hla, SdrC, SdrD, SdrE, IsdA, IsdB, IsdC, ClfA, ClfB, and SasF.

54. The method of claim 40, wherein the composition comprises a recombinant nucleic acid molecule encoding the SpA variant.

55. The method of claim 54, wherein the composition includes a recombinant, non-*staphylococcus* bacterium expressing the SpA variant.

56. The method of claim 55, wherein the recombinant non-*staphylococcus* bacterium is a *Salmonella*.

57. The method of claim 39, wherein the subject is a mammal.

58. The method of claim 39, wherein the subject is human.

59. The method of claim 39, wherein the immune response is a protective immune response.

60. A method for treating a staphylococcal infection in a subject comprising providing to a subject having, suspected of having or at risk of developing a staphylococcal infection an effective amount of an isolated peptide comprising a Protein A variant having an amino acid substitution at amino acid positions 9 and 10 of SEQ ID NO:2.

61. The method of claim 60, wherein the subject is diagnosed with a persistent staphylococcal infection.

62. The method of claim 60, wherein the SpA variant elicits production of an antibody that binds Protein A in the subject.

63. The method of claim 62, wherein the SpA variant is at least 80% identical to SEQ ID NO:2 and has an amino acid substitution at amino acid positions 9 and 10 of SEQ ID NO:2.

64. The method of claim 62, wherein the SpA variant has an amino acid sequence of SEQ ID NO:8.

65. The method of claim 62, wherein the SpA variant is administered with an adjuvant.

66. The method of claim 65, wherein the SpA variant is coupled to an adjuvant.

67. The method of claim 60, wherein the SpA variant is formulated in a pharmaceutically acceptable composition.

68. The method of claim 60, further comprising administering a second staphylococcal antigen.

69. The method of claim 68, wherein the second staphylococcal antigen is administered concurrently with the SpA variant.

70. The method of claim 69, wherein the second staphylococcal antigen and the SpA variant are administered in the same composition.

71. The method of claim 69, wherein the second staphylococcal antigen is fused with the SpA variant.

72. The method of claim 68, wherein the second staphylococcal antigen is one or more of Emp, EsxA, EsxB, EsxC, Eap, Ebh, EsaB, Coa, vWbp, vWh, Hla, SdrC, SdrD, SdrE, IsdA, IsdB, IsdC, ClfA, ClfB, and SasF peptide.

73. The method of claim 60, wherein the staphylococcal infection is a *Staphylococcus aureus* infection.

74. The method of claim 60, wherein the peptide is administered orally, parenterally, transdermally, transmucosally, subcutaneously, intramuscularly, or by inhalation.

75. The method of claim 60, further comprising administering the peptide more than one time to the subject.

76. The method of claim 60, wherein the subject is a mammal.

77. The method of claim 60, wherein the subject is human.

\* \* \* \* \*