



US007452705B2

(12) **United States Patent**  
**Kazmierczak et al.**

(10) **Patent No.:** **US 7,452,705 B2**  
(45) **Date of Patent:** **Nov. 18, 2008**

(54) **N4 VIRION SINGLE-STRANDED DNA  
DEPENDENT RNA POLYMERASE**

(75) Inventors: **Krystyna M. Kazmierczak**, Park Ridge,  
IL (US); **Elena K. Davydova**, Chicago,  
IL (US); **Lucia B. Rothman-Denes**,  
Chicago, IL (US)

(73) Assignee: **The University of Chicago**, Chicago, IL  
(US)

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/153,219**

(22) Filed: **May 22, 2002**

(65) **Prior Publication Data**

US 2003/0096349 A1 May 22, 2003

**Related U.S. Application Data**

(60) Provisional application No. 60/292,845, filed on May  
22, 2001.

(51) **Int. Cl.**

**C12N 9/12** (2006.01)

**C12N 1/20** (2006.01)

**C12N 15/63** (2006.01)

**C07H 21/04** (2006.01)

(52) **U.S. Cl.** ..... **435/194**; 435/183; 435/252.3;  
435/320.1; 435/6; 536/23.2

(58) **Field of Classification Search** ..... 435/194,  
435/320.1, 252.3, 183, 8; 536/23.1, 23.2;  
530/350

See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

5,700,667 A 12/1997 Marble et al. .... 435/91.3  
5,869,320 A 2/1999 Studier et al. .... 435/252.33  
5,981,247 A \* 11/1999 Hagedorn et al. .... 435/194  
6,218,145 B1 4/2001 Bogosian et al. .... 435/69.1

**FOREIGN PATENT DOCUMENTS**

WO WO 00/11160 3/2000  
WO WO 01/75067 10/2001

**OTHER PUBLICATIONS**

Ngo et al., Computational Complexity, Protein Structure Prediction,  
and the Levinthal Paradox, in The Protein Folding Problem and  
Tertiary Structure Prediction, 1994, Merz et al. (ed.), Birkhauser,  
Boston, MA, pp. 433 and 492-495.\*

GenEMBL Database Accession No. Z82091, Nov. 1996.\*

Abrevaya and Rothman-Denes, "N4 RNA polymerase II sites of  
transcription initiation," *J. Mol. Biol.*, 211:359-372, 1990.

Butler and Chamberlin, "Bacteriophage SP6-specific RNA  
polymerase. I. Isolation and characterization of the enzyme," *J. Biol.  
Chem.*, 257: 5772-5778, 1982.

Cermakian et al., "Sequences homologous to the yeast mitochondrial  
and bacteriophage T3 and T7 RNA polymerases are widespread  
throughout the eukaryotic lineage," *Nuc. Acids Res.*, 24:648-654,  
1996.

Cheetham and Steitz, "Insights into transcription: structure and func-  
tion of single-subunit DNA-dependent RNA polymerases," *Curr. Op.  
In Struc. Biol.*, 10:117-123, 2000.

Cramer et al., "Structural basis of transcription: RNA polymerase II  
at 2.8 Ångstrom resolution," *Science*, 292(5523):1863-1876, 2001.

Dai and Rothman-Denes, "Sequence and DNA structural determi-  
nants of N4 virion RNA polymerase-promoter recognition," *Genes  
Devepmt.*, 12:2782-2790, 1998.

Delarue et al., "An attempt to unify the structure of polymerases,"  
*Protein Engineering*, 3:461-467, 1990.

Falco and Rothman-Denes, "Bacteriophage N4-induced transcribing  
activities in *E. coli*: I. Detection and characterization in cell extracts,"  
*Virology*, 95:454-465, 1979.

Falco and Rothman-Denes, "Bacteriophage N4-induced transcribing  
activities in *E. coli*: II. Association of the N4 transcriptional apparatus  
with the cytoplasmic membrane," *Virology*, 95:466-475, 1979.

Falco et al., "Virion-associated RNA polymerase required for  
bacteriophage N4 development," *Proc. Natl. Acad. Sci. USA*, 74:520-  
523, 1977.

Falco et al., "DNA-dependent RNA polymerase from bacteriophage  
N4 varions: purification and characterization," *J. Biol. Chem.*, 255:  
4339-4347, 1980.

Falco et al., "Novel template requirements of N4 virion RNA  
polymerase," *Proc. Natl. Acad. Sci. USA*, 75: 3220-3224, 1978.

Glucksmann et al., "Specific sequences and a hairpin structure in the  
template strand are required for N4 virion RNA polymerase-pro-  
moter recognition," *Cell*, 70:491-500, 1992.

Glucksmann-Kuis et al., "*E. coli* SSB activation of N4 virion RNA  
polymerase: specific stabilization of an essential DNA hairpin  
required for promoter recognition," *Cell*, 84:147-154, 1996.

Gross et al., "The functional and regulatory roles of sigma factors in  
transcription," In: *Mechanisms of Transcription*, Cold Spring Harbor  
*Symp. Quant. Biol.*, 63: 141-155, 1998.

Haynes and Rothman-Denes, "N4 virion RNA polymerase sites of  
transcription initiation," *Cell*, 41:597-605, 1985.

Hedtke et al., "Mitochondrial and chloroplast phage-type RNA  
polymerases in Arabidopsis," *Science*, 277:809-811, 1997.

(Continued)

*Primary Examiner*—Richard G Hutson

(74) *Attorney, Agent, or Firm*—Fulbright & Jaworski L.L.P.

(57) **ABSTRACT**

A histidine-tagged, deletion mutant of bacteriophage  
N4-coded, virion RNA polymerase (mini-vRNAP) which is  
active has been developed. The his-tagged mini-vRNAP has  
been cloned under the control of the pBAD promoter, is stable  
and is purified in a single step yielding large amounts (10  
mg/liter of *E. coli* expressing cells). This RNA polymerase  
uses single-stranded DNA containing 17 bases (the promoter)  
upstream of the transcribed regions as a template. In the  
presence of *E. coli* SSB protein, it transcribes this template  
efficiently, providing a unique system to synthesize RNAs of  
the desired sequence using single-stranded DNA templates.  
The enzyme incorporates derivatized nucleoside triphos-  
phates with high efficiency. A mutant of mini-vRNAP has  
been generated that incorporates deoxynucleoside triphos-  
phates. In addition, the inventors have developed an in vivo  
system to express RNAs and proteins under mini vRNA poly-  
merase promoter control.

OTHER PUBLICATIONS

Hochschild and Dove, "Protein-protein contacts that activate and repress prokaryotic transcription," *Cell*, 92:597-600, 1998.

Malone et al., "Cloning and generation of a genetic map of bacteriophage N4 DNA," *Virology*, 162: 328-336, 1988.

Markiewicz et al., "*E. coli* single-stranded DNA binding (SSB) protein is a supercoiled-template dependent transcriptional activator of N4 virion RNA polymerase," *Genes and Dev.*, 6:2010-2019, 1992.

Miller et al., "RNA polymerase  $\beta'$  subunit: a target of DNA binding-independent activation," *Science*, 275:1655-1657, 1997.

Zivin and Zehring, "Transcriptional map of bacteriophage N4: location and polarity of N4 RNAs," *J. Mol. Biol.*, 152:335-356, 1981.

Dai et al., "Supercoil-induced extrusion of a regulatory DNA hairpin," *Proc. Natl. Acad. Sci. USA*, 94:2174-2179, 1997.

Kazmierczak et al., "The phage N4 virion RNA polymerase catalytic domain is related to single-subunit RNA polymerases," *EMBO Journal*, 21:5815-5823, 2002.

Database Accession No. ABG11811.

\* cited by examiner

# Bacteriophage N4 vRNAP promoters




FIG. 1

# N4 vRNAP and generation of mini-vRNAP




FIG. 2

# Identification of the minimal active domain of N4 vRNAP by proteolytic cleavage.





FIG. 3



Plasmid name: pMini-vRNAP  
Plasmid size: 7426 bp  
Constructed by: K. M. Kazmierczak  
Construction date: 2/2000  
Comments: Insert cloned into Invitrogen  
pBAD B expression plasmid

FIG. 4

Purification of cloned vRNAP and mini-vRNAP




FIG. 5

Activation of N4 vRNAP transcription by Eco SSB  
at different ssDNA concentrations




FIG. 6

**Effect of Eco SSB on ssDNA template recycling**

	A	B	C	D
Extra ssDNA	-	+	+	+
Eco SSB	-	-	+	-
time at 37°C	10' 20' 30' 40'	10' 20' 30' 40'	10' 20' 30' 40'	10' 20' 30' 40'

**FIG. 7**

# Effect of *Eco* SSB on the state of template DNA and product RNA in vRNAP transcription




FIG. 8

# Transcription initiation by vRNAP and mini-vRNAP




FIG. 9

Effect of Eco SSB on transcription of vRNAP and mini-RNAP





FIG. 10

# Determination of mini-vRNAP promoter contacts



**B**




FIG. 11

### Binding affinities of stem-length promoter mutants




FIG. 12



## UV crosslinking of mutant mini-vRNAPases to promoter oligonucleotides




FIG. 14

# Run-off transcription by mutant mini-vRNAPases




FIG. 15

# Mutant mini-vRNAPases transcription initiation

Motif B RgiaKnplitiYG  
          ▼          ▼  
          A          F




FIG. 16

# Detection of *in vivo* activities of N4 vRNAP and mini-vRNAP




FIG. 17

## N4 VIRION SINGLE-STRANDED DNA DEPENDENT RNA POLYMERASE

This application claims the priority of U.S. Provisional Patent Application Serial No. 60/292,845, filed May 22, 2001, the entire disclosure of which is specifically incorporated herein by reference.

The government may own rights in the present invention pursuant to grant number R01 A1 12575 from the National Institute of Health.

### BACKGROUND OF THE INVENTION

#### I. Field of the Invention

The present invention relates generally to an RNA polymerase. More particularly, it provides a bacteriophage N4 virion RNA polymerase for synthesis of RNAs of desired sequences using single-stranded DNA templates.

#### II. Description of Related Art

The expression of a protein-encoding gene in a host cell involves transcription of messenger RNA (mRNA) from DNA by an RNA polymerase enzyme. Subsequently the mRNA is processed, involving recognition of a region of the 3' UTR and addition of a tail of polyadenylate nucleotides to the 3' end of the mRNA by polyadenylation enzymes. After transcription, the mRNA encounters ribosomes which associate with a region of the 5' UTR of the mRNA and translocate in a 3'-ward direction along the mRNA. During translocation, amino acids are added to one another in sequence to form the polypeptide product of the protein-encoding gene. For prokaryotic transcription-translation, the Shine-Dalgarno sequence of the bacterial mRNA located about six to nine nucleotides before the initiation site for translation may be used for ribosome loading. This sequence is complementary to a sequence on the 3' end of the 16S rRNA and stimulates ribosome binding to the mRNA. The base pairing between the Shine-Dalgarno sequence and the mRNA sequences serves to align the initiating AUG for decoding.

Transcription of DNA into mRNA is regulated by the promoter region of the DNA. The promoter region contains a sequence of bases that signals RNA polymerase to associate with the DNA, and to initiate the transcription of mRNA using one of the DNA strands as a template to make a corresponding complementary strand of RNA. RNA polymerases from different species typically recognize promoter regions comprised of different sequences. In order to express a protein-encoding gene in a host cell, either the promoter driving transcription of the protein-encoding gene must be recognized by a host RNA polymerase, or an RNA polymerase which recognizes the promoter driving transcription of the protein-encoding gene must be provided to the host cell (U.S. Pat. No. 6,218,145).

Most DNA-dependent RNA polymerases read double-stranded DNA, limiting RNA synthesis to systems in which a double-stranded DNA template is available. The synthesis of RNA using single-stranded DNA is not as common. Synthesizing RNA using a single-stranded DNA template immobilized on a solid support is described in U.S. Pat. No. 5,700,667.

Therefore, this invention provides an RNA polymerase that reads single-stranded DNA. Also provided is an RNA polymerase for which the promoter sequence is present upstream of the transcription initiation site and therefore is not transcribed by the polymerase.

### SUMMARY OF THE INVENTION

The invention provides a novel N4 virion RNA polymerase (vRNAP) and a mini-vRNA polymerase and method of use thereof. The novel polymerases are described by an isolated nucleic acid comprising a region encoding a polypeptide having the amino sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15. The nucleic acid may comprise the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14. The vRNAP and mini-vRNA polymerase transcribe nucleic acid operatively linked to an N4 promoter such as a P2 promoter of SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29. The promoter of SEQ ID NO:16 or SEQ ID NO:28 is preferred.

An aspect of the current invention comprises a recombinant host cell comprising a DNA segment encoding a N4 virion RNA polymerase. The DNA segment is either single- or double-stranded and the polypeptide encoded by the DNA segment is preferably SEQ ID NO:4 or SEQ ID NO:6. The recombinant host cell may be an *E. coli* cell. Another aspect of the current invention comprises a recombinant vector comprising a DNA segment encoding a N4 virion RNA polymerase polypeptide under the control of a promoter.

Yet another aspect of the current invention comprises an isolated polynucleotide comprising a sequence identical or complementary to at least 14 contiguous nucleotides of SEQ ID NO:1. The polynucleotide may comprise at least 20, 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000, 2000, 3000, 3300 or more contiguous nucleotides of SEQ ID NO:1. The polynucleotide may comprise all contiguous nucleotides of SEQ ID NO:3 or all contiguous nucleotides of SEQ ID NO:1. Similarly, the polynucleotide may comprise at least 20, 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000, 2000, 3000, 3300 or more nucleotides complementary to at least 20, 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000, 2000, 3000, 3300 or more contiguous nucleotides of SEQ ID NO:1.

Another aspect of the current invention comprises a purified N4 virion RNA polymerase comprising at least 20 contiguous amino acids of SEQ ID NO:2. It is preferred that the polymerase contain at least 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000 or more contiguous amino acids of SEQ ID NO:2.

Yet another aspect of the current invention comprises an isolated nucleic acid comprising a region encoding a polypeptide comprising at least 6 contiguous amino acids of SEQ ID NO:2, wherein the polypeptide has RNA polymerase activity under appropriate reaction conditions. It is preferred that this polypeptide comprises at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 75, 100, 150, 200, 250, 300, 400, 600, 800, 1000 or more contiguous amino acids of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, or SEQ ID NO:15. The encoded polypeptide may have at least one hexahistidine tag or other tag. The polypeptide may be a mutant of the peptide found in SEQ ID NO:2 or SEQ ID NO:4, such as an enzyme possessing an amino acid substitution at position Y678.

An embodiment of the current invention comprises a method of making RNA. This method comprises: (a) obtaining a N4 virion RNA polymerase (i.e. the polypeptide); (b) obtaining DNA wherein the DNA preferably contains a N4 virion RNA polymerase promoter sequence; (c) admixing the RNA polymerase and the DNA; and (d) culturing the RNA polymerase and the DNA under conditions effective to allow RNA synthesis. Optionally, the method may comprise syn-

thesizing polynucleotides containing modified ribonucleotides or deoxyribonucleotides. The DNA is preferably single-stranded DNA or denatured double-stranded DNA. Step (c) may occur in a host cell such as an *E. coli* host cell.

The amino acid sequence of the RNA polymerase is preferably the sequence essentially as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:15, or a mutant form of the polymerase of SEQ ID NO:4 or SEQ ID NO:6. The mutation may be, for example, at position number Y678. The RNA transcript may contain derivatized nucleotides.

An aspect of the current invention comprises using an N4 vRNAP promoter to direct transcription. The promoter is preferentially an N4 promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29. The P2 promoter of SEQ ID NO:16 or SEQ ID NO:28 is preferred. The promoter sequence may be upstream of the transcription initiation site. The promoter may comprise a set of inverted repeats forming a hairpin with a 2-7 base pair long stem and 3-5 base loop having purines in the central and/or next to the central position of the loop.

The preferred conditions of the transcription method claimed herein includes a pH in step (c) of between 6 and 9, with a pH of between 7.5 and 8.5 more preferred. Mg<sup>+2</sup> or Mn<sup>+2</sup>, preferably Mg<sup>+2</sup> may be admixed. Preferred temperatures for the reaction are 25° C. to 50° C. with the range of 30° C. to 45° C. being more preferred and the range of 32° C. to 42° C. being most preferred. The admixing may occur in vivo or in vitro.

An aspect of the current invention also includes translation of the RNA after transcription. A reporter gene such as an  $\alpha$ -peptide of  $\beta$ -galactosidase may be used. It is preferred the transcription comprises admixing an *E. coli* single-stranded binding protein (EcoSSB), a SSB protein homologous to EcoSSB or another naturally occurring or chimeric SSB protein homologous to EcoSSB with the polymerase and DNA.

The DNA admixed with the RNA polymerase of the current invention may be single-stranded linear DNA or single-stranded circular DNA such as bacteriophage M13 DNA. The DNA may be denatured DNA, such as single-stranded, double-stranded linear or double-stranded circular denatured DNA. The DNA may also be double-stranded DNA under certain conditions. The RNA may be pure RNA or may contain modified nucleotides. Mixed RNA-DNA oligonucleotides may also be synthesized with the Y678F mutant mini-vRNAP (SEQ ID NO:8) of the current invention.

Yet another aspect of the current invention is the transcription method in which no EcoSSB is admixed with the RNA polymerase and DNA; the product of this method is a DNA/RNA hybrid.

The synthesized RNA may comprise a detectable label such as a fluorescent tag, biotin, digoxigenin, 2'-fluoro nucleoside triphosphate, or a radiolabel such as a <sup>35</sup>S- or <sup>32</sup>P-label. The synthesized RNA may be adapted for use as a probe for blotting experiments or in-situ hybridization. Nucleoside triphosphates (NTPs) or derivatized NTPs may be incorporated into the RNA, and may optionally have a detectable label. Deoxynucleoside triphosphates may be incorporated into the RNA.

The RNA may be adapted for use for NMR structural determination. Short RNAs such as those between 10 and 1000 bases or between 10 and 300 bases may be used. The RNA may be adapted for use in spliceosome assembly, splicing reactions or antisense experiments. Also, the RNA may be adapted for use in probing for a complementary nucleotide sequence or for use as a probe in RNase protection studies.

Yet another aspect of the current invention comprises delivering RNA into a cell after transcription of the RNA. The delivery may be by microinjection. Another aspect of the invention comprises amplifying the RNA after transcription.

Another embodiment of the current invention comprises a method of making RNA comprising: (a) obtaining a N4 virion RNA polymerase; (b) obtaining a single-stranded DNA oligonucleotide wherein the oligonucleotide contains a N4 virion RNA polymerase promoter sequence; (c) admixing the RNA polymerase and the oligonucleotide; and (d) culturing the RNA polymerase and the oligonucleotide under conditions effective to allow RNA synthesis. The polymerase preferentially has the amino sequence set forth in SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8. In this embodiment, it is preferred that the DNA has between 20 and 200 bases.

Yet another embodiment of the invention comprises a method of making RNA comprising: (a) obtaining a N4 virion RNA polymerase; (b) obtaining a single-stranded DNA wherein the DNA contains a N4 virion RNA polymerase promoter sequence; (c) obtaining a ribonucleoside triphosphate (XTP) or a derivatized ribonucleoside triphosphate; (d) admixing the RNA polymerase, the DNA and the XTP; and (e) culturing the RNA polymerase and the oligonucleotide under conditions effective to allow RNA synthesis wherein the RNA is a derivatized RNA. The RNA polymerase preferentially has the amino sequence set forth in SEQ ID NO:4 or SEQ ID NO:6 or a mutant of the polymerase of SEQ ID NO:4 or SEQ ID NO:6, such as a mutant with a mutation at position number Y678 or the polymerase of SEQ ID NO:8.

Another embodiment of the invention comprises a method for in vivo or in vitro protein synthesis comprising: (a) obtaining an RNA polymerase having the amino sequence set forth in SEQ ID NO:4, SEQ ID NO:6 or a mutant thereof; (b) obtaining DNA wherein the DNA contains a N4 virion RNA polymerase promoter sequence; (c) admixing the RNA polymerase and the DNA; (d) culturing the RNA polymerase and the DNA under conditions effective to allow RNA synthesis; and (e) culturing the RNA in vivo or in vitro under conditions effective to allow protein synthesis. Step (e) may comprise using a two plasmid system or a one plasmid system in which a reporter gene and the RNA polymerase gene are located on the same plasmid.

Yet another embodiment of the invention comprises a method of making a N4 mini-vRNAP comprising: (a) expressing vRNAP, wherein the vRNAP has the amino sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:15 or a mutant thereof; and (b) purifying the vRNAP. The expression of vRNAP may occur in a bacteria, yeast, CHO, Cos, HeLa, NIH3T3, Jurkat, 293, Saos, or a PC12 host cell. A promoter such as pBAD may be used for making the vRNAP in bacterial cells. Any other promoter appropriate to the host cell line used can be employed when expressing vRNAP in other host cells. The polymerase may have a specific recombinant sequence that can be used in purification of the polymerase. The vRNAP may have at least one hexahistidine, FLAG, hemagglutinin or c-myc tag, or may not have a tag.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1—Bacteriophage N4 vRNAP promoters on single-stranded templates. These promoters are characterized by conserved sequences and a 5 bp stem, 3 base loop hairpin structure.

FIG. 2A and FIG. 2B—N4 vRNAP and generation of mini-vRNAP. FIG. 2A shows a schematic of the N4 vRNAP protein with three motifs: the T/DxxGR motif found in DNA-dependent polymerases, the P-loop, an ATP/GTP-binding motif present in some nucleotide-binding proteins, and motif B (Rx<sub>3</sub>Kx<sub>6-7</sub>YG), one of three motifs common to the Pol I and Pol α DNA polymerases and the T7-like RNA polymerases (SEQ ID NOS: 39 and 40). FIG. 2B shows the mini-vRNAP.

FIG. 3A and FIG. 3B—Identification of the minimal transcriptionally active domain of N4 vRNAP by proteolytic cleavage. FIG. 3A, SDS-PAGE analysis of the products of vRNAP digestion with trypsin. FIG. 3B N-terminal sequencing of the three initial proteolytic fragments indicated that the stable active polypeptide (mini-vRNAP) corresponds to the middle 1/3 of vRNAP, the region containing the three motifs described in FIG. 2A.

FIG. 4—ORFs for full length polymerase, mini-vRNAP and mutants thereof were cloned under pBAD control with an N-terminal hexahistidine tag.

FIG. 5—Purification of cloned vRNAP and mini-vRNAP. The left hand side shows the relative amounts of full size and mini-vRNAP proteins purified on TALON columns from the same volume of induced cells. Further concentration on a monoQ column reveals that, in contrast to full size vRNAP, mini-vRNAP is stable after induction (right).

FIG. 6—Activation of N4 vRNAP transcription by EcoSSB at three different ssDNA concentrations. The extent of EcoSSB activation is template-concentration dependent, with highest activation at low DNA template concentration.

FIG. 7A, FIG. 7B, FIG. 7C, and FIG. 7D—Effect of EcoSSB on ssDNA template recycling. In the absence of EcoSSB, no increase in transcription was observed beyond 10 min of incubation (FIG. 7A). Addition of template at 20 min to the reaction carried out in the absence of EcoSSB led to a dramatic increase in RNA synthesis (FIG. 7B). RNA synthesis increased linearly throughout the period of incubation (FIG. 7C). Addition of EcoSSB at 20 min led to a slow rate of transcriptional recovery (FIG. 7D).

FIG. 8—Effect of EcoSSB on the state of template DNA and product RNA in vRNAP transcription. Native gel electrophoresis was carried out in the absence and in the presence of EcoSSB. Transcription was performed at an intermediate (5 nM) DNA concentration, at which only a 2-fold effect of EcoSSB is observed. Either <sup>32</sup>P-labeled template (right panel) or labeled NTPs (left panel) were used to analyze the state of the template (right panel) or RNA product (left panel) in the absence or presence of EcoSSB.

FIG. 9A, FIG. 9B, and FIG. 9C—Transcription initiation by vRNAP and mini-vRNAP. The initiation properties of the full length and mini-vRNA polymerases were compared at similar molar concentrations (FIG. 9A) using the catalytic autolabeling assay and two reaction conditions: using a template containing +1C, the benzaldehyde derivative of GTP and α<sup>32</sup>P-ATP, or a template containing +1T, the benzaldehyde derivative of ATP and α<sup>32</sup>P-GTP. Comparison of the results in FIGS. 9B and 9C demonstrates that mini-vRNAP exhibits initiation properties similar to full size vRNAP.

FIG. 10—Effect of EcoSSB on transcription of vRNAP and mini-vRNAP. The elongation and termination properties of vRNAP and mini-vRNAP are compared.

FIG. 11A and FIG. 11B—Determination of mini-vRNAP promoter contacts. A 20-base oligonucleotide containing wild type promoter P2 sequence binds with a 1 nM Kd (FIG.

11A) (nucleotides 13-32 of SEQ ID NO:16). Most oligonucleotides substituted with 5-Iodo-dU at specific positions showed close to wild type affinity except for the oligonucleotides substituted at positions -11 (at the center of the loop) and -8, indicating that these positions are essential for promoter recognition (FIG. 11B). UV crosslinking indicates that mini-vRNAP primarily contacts the -11 position.

FIG. 12—Binding affinities of stem-length promoter mutants. Wild type promoter P2 with a 5 bp stem has a Kd of 1 nM (top). (nucleotides 18-37 of SEQ ID NO:17). The stem was shortened by removal of 3' bases (left). (SEQ ID NOS: 35-38). The stem can be shortened by two base pairs without change in the binding affinity. The effect of lengthening the stem by addition of 3' bases is shown (right). The stem can be lengthened by two base pairs without change in the binding affinity.

FIG. 13A and FIG. 13B—Identification of the transcription start site by catalytic autolabeling. A series of templates were constructed with a single C placed at different distances from the center of the hairpin (position -11) by addition or deletion of the tract of As present at promoter P2 (FIG. 13A) (nucleotides 18-37 of SEQ ID NO:17). The affinity of mini-vRNAP for these promoters was measured by filter binding, and transcription initiation was measured by catalytic autolabeling of mini-vRNAP. All templates showed similar binding affinities. However, only the template with a C positioned 12 bases downstream from the center of the hairpin was able to support transcription initiation (FIG. 13B).

FIG. 14—UV crosslinking of mutant mini-vRNAPases to promoter oligonucleotides. Two mutants (K670A and Y678F) were tested for their ability to bind to wild type promoters. Both mutant RNA polymerases bound to promoter DNA with wild type affinities and crosslinked to 5-Iodo-dU substituted P2 DNA templates at positions -11 and +3 as well as the wild type enzyme, indicating that these polymerase mutations do not affect promoter binding.

FIG. 15—Run-off transcription by mutant mini-vRNAPases. The wild type and Y678F (SEQ ID NO:8) enzymes displayed similar activities at both template excess and template-limiting conditions, while the K670A enzyme exhibited decreased activity under both conditions. Under limiting template conditions, all three enzymes were activated by EcoSSB (right panel). However, the Y678F enzyme showed reduced discrimination between incorporation of ribo- and deoxyribonucleoside triphosphates.

FIG. 16—Mutant mini-vRNAPases in transcription initiation. The initiation properties of the three enzymes were compared using catalytic autolabeling. The K670A enzyme displays significantly reduced activity with the GTP derivative. The Y678F enzyme, in contrast to wild type polymerase, incorporates dATP as efficiently as rATP in a single round of phosphodiester bond formation.

FIG. 17A, FIG. 17B, and FIG. 17C—Detection of in vivo activities of N4 vRNAP and mini-vRNAP. Transcription of β-galactosidase α-peptide by full size and mini-vRNAP was assayed on inducing-Xgal media (FIG. 17A). Plasmid (pA-CYC) templates were constructed with a reporter gene (α-peptide of β-galactosidase) under the control of vRNAP promoter P2 cloned in either of two orientations (FIG. 17B). Induction of mini-vRNAP led to production and accumulation of detectable levels of the protein, whereas full-length vRNAP was degraded (FIG. 17C).

## DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

The present invention overcomes deficiencies in the art by providing a stable RNA polymerase that uses single-stranded DNA and provides a unique system to synthesize RNAs of a desired sequence. The RNA polymerase and mini-vRNA polymerase can be used to synthesize RNA for use as probes in RNase protection studies of DNAs or RNAs, in situ hybridization studies, and in Southern and Northern blot analysis, for the synthesis of defined RNA:DNA hybrids, for NMR structure determination of RNAs, for in vitro studies of spliceosome assembly, splicing reactions and antisense experiments, for in vitro translation or microinjection, and for nucleic acid amplification. The present invention allows for the synthesis of derivatized RNA and can use ssDNA in the form of single-stranded oligonucleotides, denatured DNA or DNA cloned into M13 templates.

## I. RNA Polymerases

## a. Structure and Promoter Recognition of DNA-Dependent RNA Polymerases

Inspection of the sequences of phage, archaeobacterial, eubacterial, eukaryotic and viral DNA-dependent RNA polymerases has revealed the existence of two enzyme families. The eubacterial, eukaryotic, archaeobacterial, chloroplast and the vaccinia virus RNA polymerases are complex multisubunit enzymes (5-14 subunits) composed of two large subunits, one to several subunits of intermediate molecular weight (30-50-kDa) and none to several subunits of small molecular weight (<30-kDa) (Archambault, et al., 1993; Record, et al., 1995). Eubacterial RNA polymerases are the simplest with an  $\alpha_2\beta\beta'$  core structure. Sequence comparison of the genes coding for the different subunits of these enzymes has revealed: 1-sequence homology in eight segments (A to H) between  $\beta'$  and the largest subunit of other RNA polymerases, 2-sequence homology in nine segments (A to I) between  $\beta$  and the next largest subunit of other RNA polymerases, 3-sequence homology in 3 segments (1.1, 1.2 and 2) between  $\alpha$  and a subunit in RNA polymerases I, II and III (Puhler, et al., 1989; Sweetser, et al., 1987). Not surprisingly, the crystal structures of yeast RNAP II and *E. coli* RNAP core revealed remarkable similarities (Zhang, et al., 1999; Cramer, et al., 2001).

In contrast, members of the phage T7-like (T3, SP6) family of RNA polymerases consist of a single (~100 kDa) polypeptide which catalyzes all functions required for accurate transcription (Cheetham, et al., 2000). The heterodimeric bacteriophage N4 RNAP II, nuclear-coded mitochondrial, and Arabidopsis chloroplast RNA polymerases show sequence similarity to the phage RNA polymerases (Cermakian, et al., 1996; Hedtke, et al., 1997; Zehring, et al., 1983). Three sequence motifs -A and C, which contain the two aspartic acids required for catalysis, and motif B- are conserved in polymerases that use DNA as a template (Delarue, et al., 1990). The crystal structure of T7 RNAP resembles a "cupped right hand" with "palm," "fingers" and "thumb" subdomains (Sousa, et al., 1993). The two catalytic aspartates are present in the "palm" of the structure. This structure is shared by the polymerase domains of *E. coli* DNA polymerase I and HIV reverse transcriptase (Sousa, 1996). Genetic, biochemical and structural information indicates that T7 RNA polymerase contains additional structures dedicated to nascent RNA binding, promoter recognition, dsDNA unwinding and RNA:DNA hybrid unwinding (Cheetham, et al., 2000; Sousa, 1996)

Both Class I and Class II RNA polymerases recognize specific sequences, called promoters, on B form double-

stranded DNA. Eubacterial promoters (except those recognized by  $\sigma^{54}$ ) are characterized by two regions of sequence homology: the -10 and the -35 hexamers (Gross, et al., 1998). Specificity of promoter recognition is conferred to the core enzyme by the  $\sigma$  subunit, which makes specific interactions with the -10 and -35 sequences through two distinct DNA binding domains (Gross, et al., 1998). This modular promoter structure is also present at the promoters for eukaryotic RNA polymerases I, II and III. Transcription factors TFIIB and TFIIC direct recognition of RNAP III to two separate sequences (boxes A and C, separated by defined spacing) at the 5S gene promoter, while transcription factors TFIIB and TFIIC direct recognition of this enzyme to blocks A and B, separated by variable distance (31-74 bp) at the tRNA promoters (Paule, et al., 2000). Sequences important for RNAP I transcription initiation at the human rRNA promoters are also restricted to two regions: the "core" region located at -40 to +1 and the "upstream" region present at -160 to -107 (Paule, et al., 2000). Assembly of the initiation complex at RNAP II promoters requires several general transcription factors (TFIIA, TFIIB, TFIID, TFIIE, TFIIIF and TFIIH). Recognition involves three core elements: the TATA box located at position -30 and recognized by TBP, the initiator element located near -1, and the downstream promoter element near +30 (Roeder, 1996).

Promoters for the T7-like and mitochondrial RNAPases are simpler. The T7-type RNAP promoters span a continuous highly conserved 23 bp region extending from position -17 to +6 relative to the start site of transcription (+1) (Rong, et al., 1998). The yeast mitochondrial RNAP promoters are even smaller, extending from -8 to +1 (Shadel, et al., 1993). One exception are the promoters for N4 RNAP II, which are restricted to two blocks of conserved sequence: a/tTTTA at +1 and AAGACCTG present 18-26 bp upstream of +1 (Abravaya, et al., 1990).

The activity of the multisubunit class of RNA polymerases is enhanced by activators at weak promoters. Transcription activators generally bind at specific sites on double-stranded DNA upstream of the -35 region (with the exception of the T4 sliding clamp activator), or at large distances in the cases of enhancers (Sanders, et al., 1997). Activators modulate transcription by increasing the binding (formation of closed complex) or isomerization (formation of open complex) steps of transcription through interactions with the  $\alpha$  or  $\sigma$  subunits of RNAP (Hochschild, et al., 1998). An exception is N4SSB, the activator of *E. coli* RNAP $\sigma^{70}$  at the bacteriophage N4 late promoters, which activates transcription through direct interactions with the  $\beta'$  subunit of RNAP in the absence of DNA binding (Miller, et al., 1997).

Proteins that bind to ssDNAs with high affinity but without sequence specificity have been purified and characterized from several prokaryotes, eukaryotes, and their viruses (Chase, et al., 1986). These proteins (SSBs), which are required for replication, recombination and repair, bind stoichiometrically and, in many cases, cooperatively to ssDNA to cover the transient single-stranded regions of DNA that normally arise in vivo as a result of replication, repair and recombination. Binding to DNA results in the removal of hairpin structures found on ssDNA, providing an extended conformation for proteins involved in DNA metabolism. Several lines of evidence suggest that single-stranded DNA binding proteins play a more dynamic role in cellular processes. Genetic and biochemical evidence indicates that these proteins are involved in a multitude of protein-protein interactions including transcription activation (Rothman-Denes, et al., 1999).

### b. The Bacteriophage N4 Virion RNA Polymerase

Bacteriophage N4 virion RNA polymerase (N4 vRNAP) is present in N4 virions and is injected into the *E. coli* cell at the beginning of infection, where it is responsible for transcription of the N4 early genes (Falco, et al., 1977; Falco, et al., 1979; Malone, et al., 1988). The N4 vRNAP gene maps to the late region of the N4 genome (Zivin, et al., 1981). N4 vRNAP purified from virions is composed of a single polypeptide with an apparent molecular mass of approximately 320,000 kDa (Falco, et al., 1980). In contrast to other DNA-dependent RNAPases, N4 vRNAP recognizes promoters on single-stranded templates (Falco, et al., 1978). These promoters are characterized by conserved sequences and a 5 bp stem, 3 base loop hairpin structure (FIG. 1) (Haynes, et al., 1985; Glucksmann, et al., 1992). In vivo, *E. coli* gyrase and single-stranded binding protein are required for transcription by N4 vRNAP (Falco, et al., 1980; Markiewicz, et al., 1992).

Sequencing of the N4 vRNAP gene revealed an ORF coding for a protein 3,500 amino acids in length (SEQ ID NO:1-2). Inspection of the sequence revealed no extensive homology to either the multisubunit or the T7-like families of RNA polymerases. However, three motifs are present (FIG. 2A): the T/DxxGR motif found in DNA-dependent polymerases, and Motif B (Rx<sub>3</sub>Kx<sub>6-7</sub>YG), one of three motifs common to the Pol I and Pol  $\alpha$  DNA polymerases and the T7-like RNA polymerases.

### c. Transcription Using N4 vRNAP

RNA synthesis requires RNA polymerase, a DNA template, an activated precursor (the ribonucleoside triphosphates ATP, GTP, UTP and CTP (XTP)), and divalent metal ions such as Mg<sup>2+</sup> or Mn<sup>2+</sup>. The metal ion Mg<sup>2+</sup> is strongly preferred. Synthesis of RNA begins at the promoter site on the DNA. This site contains a sequence which the RNA polymerase recognizes and binds. The RNA synthesis proceeds until a termination site is reached. N4 vRNAP termination signals comprise a hairpin loop that forms in the newly synthesized RNA which is followed by a string of uracils (poly U). The sequence of the terminator signals for vRNAP present in the N4 genome include SEQ ID NOS:21-26. These N4 vRNAP termination signals possess all of the characteristics of eubacterial sequence-dependent terminators.

The ribonucleoside triphosphate may be derivatized with, for example, biotin. Derivatized XTPs can be used for the preparation of derivatized RNA. Exemplary methods for making derivatized XTPs are disclosed in detail in Rashtchian et al. (1992), herein incorporated by reference.

Single-stranded DNA of varying lengths can be used as a template for RNA synthesis using the N4 vRNAP or mini-vRNAP. Oligonucleotides and polynucleotides of intermediate length may be used. One particular single-stranded DNA that may be used is M13 DNA. M13 genomic DNA exists temporarily inside infected *E. coli* cells as a double-stranded DNA plasmid and is packaged as a small, single-stranded circular DNA into phage particles. M13 phage particles are secreted by an infected cell and single-stranded DNA can be purified from these particles for use as a transcription template. Initially M13 phage vectors required a working knowledge of phage biology and were primarily used for creating single-strand DNA molecules for DNA sequencing. M13-derived cloning vectors called "phagemids" take advantage of M13 replication to produce single-strand molecules, but can be propagated as conventional ColE1-based replicating double-strand plasmids.

EcoSSB is essential for N4 vRNAP transcription in vivo (Falco et al., 1978; Glucksmann, et al., 1992, herein incorporated by reference). EcoSSB is a specific activator of N4 vRNAP on single-stranded and supercoiled double-stranded

DNA templates. EcoSSB, unlike other SSBs, does not melt the N4 vRNAP promoter hairpin structure (Glucksmann-Kuis, et al., 1996). EcoSSB has a high specificity for N4 vRNAP and mini-vRNAP resulting from EcoSSB's ability to stabilize the template-strand hairpin, whereas the nontemplate strand hairpin is destabilized. Other single-stranded DNA binding proteins destabilize the template-strand hairpin (Glucksmann-Kuis et al., 1996; Dai et al., 1998). When EcoSSB is not used in N4 vRNAP transcription in vitro, a DNA:RNA hybrid is formed, preventing template reutilization.

## II. Genes and DNA Segments

Important aspects of the present invention concern isolated DNA segments and recombinant vectors encoding N4 vRNAP or more particularly mini-vRNAP or a mutant of mini-vRNAP and the creation and use of recombinant host cells through the application of DNA technology, that express a wild type, polymorphic or mutant vRNAP. Other aspects of the present invention concern isolated nucleic acid segments and recombinant vectors encoding vRNAP. Sequences of SEQ ID NO:1, 3, 5, 7, 14 and biologically functional equivalents thereof are used in the current invention. Single-stranded DNA oligonucleotides and polynucleotides can be used as DNA templates.

The present invention concerns isolated nucleic acid segments that are capable of expressing a protein, polypeptide or peptide that has RNA polymerase activity. As used herein, the term "nucleic acid segment" refers to a nucleic acid molecule that has been isolated free of total genomic DNA of a particular species. Therefore, a nucleic acid segment encoding vRNAP refers to a nucleic acid segment that contains wild-type, polymorphic or mutant vRNAP coding sequences yet is isolated away from, or purified free from, total bacterial or N4 phage genomic DNA. Included within the term "nucleic acid segment," are nucleic acid segments and smaller fragments of such segments, and also recombinant vectors, including, for example, plasmids, cosmids, phage, viruses, and the like.

Similarly, a nucleic acid segment comprising an isolated or purified vRNAP gene refers to a nucleic acid segment including vRNAP protein, polypeptide or peptide coding sequences and, in certain aspects, regulatory sequences, isolated substantially away from other naturally occurring genes or protein encoding sequences. In this respect, the term "gene" is used for simplicity to refer to a functional protein, polypeptide or peptide encoding unit. As will be understood by those of skill in the art, this functional term includes both genomic sequences, cDNA sequences and engineered segments that express, or may be adapted to express, proteins, polypeptides, domains, peptides, vRNAPs and mutants of vRNAP encoding sequences.

"Isolated substantially away from other coding sequences" means that the gene of interest, in this case the vRNAP, or more particularly mini-vRNAP genes, forms the significant part of the coding region of the nucleic acid segment, and that the nucleic acid segment does not contain large portions of naturally-occurring coding DNA, such as large chromosomal fragments or other functional genes or cDNA coding regions. Of course, this refers to the DNA segment as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

The term "a sequence essentially as set forth in SEQ ID NO:2 means, for example, that the sequence substantially corresponds to a portion of SEQ ID NO:2 and has relatively few amino acids that are not identical to, or a biologically functional equivalent of, the amino acids of SEQ ID NO:2.

This applies with respect to all peptide and protein sequences herein, such as those of SEQ ID NO:4, 6, 8 and 15.

The term "biologically functional equivalent" is well understood in the art and is further defined in detail herein. Accordingly, sequences that have about 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or about 99%, and any range derivable therein, such as, for example, about 70% to about 80%, and more preferably about 81% and about 90%; or even more preferably, between about 91% and about 99%; of amino acids that are identical or functionally equivalent to the amino acids of SEQ ID NO:2 will be sequences that are "essentially as set forth in SEQ ID NO:2, provided the biological activity of the protein is maintained. In particular embodiments, the biological activity of a vRNAP protein, polypeptide or peptide, or a biologically functional equivalent, comprises transcription. A preferred transcriptional activity that may be possessed by a vRNAP protein, polypeptide or peptide, or a biologically functional equivalent, is RNA synthesis using single-stranded N4 vRNAP promoter-containing DNA as a template.

In certain other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors that include within their sequence a nucleic acid sequence essentially as set forth in SEQ ID NO:1. The term "essentially as set forth in SEQ ID NO:1" is used in the same sense as described above and means that the nucleic acid sequence substantially corresponds to a portion of SEQ ID NO:1 and has relatively few codons that are not identical, or functionally equivalent, to the codons of SEQ ID NO:1. Again, nucleic acid segments that encode proteins, polypeptide or peptides exhibiting RNAP activity will be most preferred.

The term "functionally equivalent codon" is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine and serine, and also refers to codons that encode biologically equivalent amino acids. For optimization of expression of vRNAP in human cells, the codons are shown in Table 1 in preference of use from left to right. Thus, the most preferred codon for alanine is thus "GCC," and the least is "GCG" (see Table 1 below). Codon usage for various organisms and organelles can be found at the website <http://www.kazusa.or.jp/codon/>, incorporated herein by reference, allowing one of skill in the art to optimize codon usage for expression in various organisms using the disclosures herein. Thus, it is contemplated that codon usage may be optimized for other animals, as well as other organisms such as a prokaryote (e.g., an eubacteria), an archaea, an eukaryote (e.g., a protist, a plant, a fungus, an animal), a virus and the like, as well as organelles that contain nucleic acids, such as mitochondria or chloroplasts, based on the preferred codon usage as would be known to those of ordinary skill in the art.

TABLE 1

<u>Preferred Human DNA Codons</u>		
Amino Acids		Codons
Alanine	Ala A	GCC GCT GCA GCG
Cysteine	Cys C	TGC TGT
Aspartic acid	Asp D	GAG GAT
Glutamic acid	Glu E	GAG GAA
Phenylalanine	Phe F	TTC TTT

TABLE 1-continued

<u>Preferred Human DNA Codons</u>		
Amino Acids		Codons
Glycine	Gly G	GGC GGG GGA GGT
Histidine	His H	CAC CAT
Isoleucine	Ile I	ATC ATT ATA
Lysine	Lys K	AAG AAA
Leucine	Leu L	CTG CTC TTG CTT CTA TTA
Methionine	Met M	ATG
Asparagine	Asn N	AAC AAT
Proline	Pro P	CCC CCT CCA CCG
Glutamine	Gln Q	CAG CAA
Arginine	Arg R	CGC AGG CGG AGA CGA CGT
Serine	Ser S	AGC TCC TCT AGT TCA TCG
Threonine	Thr T	ACC ACA ACT ACG
Valine	Val V	GTG GTC GTT GTA
Tryptophan	Trp W	TGG
Tyrosine	Tyr Y	TAC TAT

It will also be understood that amino acid and nucleic acid sequences may include additional residues, such as additional N- or C-terminal amino acids or 5' or 3' sequences, and yet still be essentially as set forth in one of the sequences disclosed herein, so long as the sequence meets the criteria set forth above, including the maintenance of biological protein, polypeptide or peptide activity. The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region or may include various internal sequences, i.e., introns, which are known to occur within genes.

Excepting intronic or flanking regions, and allowing for the degeneracy of the genetic code, sequences that have about 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or about 99%, and any range derivable therein, such as, for example, about 50% to about 80%, and more preferably about 81% and about 90%; or even more preferably, between about 91% and about 99%; of nucleotides that are identical to the nucleotides of SEQ ID NO:1 will be sequences that are "essentially as set forth in SEQ ID NO:1".

a. Nucleic Acid Hybridization

The nucleic acid sequences disclosed herein also have a variety of uses. Contiguous sequences from vRNAP nucleic acid sequences can be used, for example, as templates to synthesize vRNAP.

Naturally, the present invention also encompasses DNA segments that are complementary, or essentially complementary, to the sequence set forth in SEQ ID NO:1, 3, 5, 7 and 14. Nucleic acid sequences that are "complementary" are those that are capable of base-pairing according to the standard Watson-Crick complementary rules. As used herein, the term "complementary sequences" means nucleic acid sequences that are complementary, as may be assessed by the same nucleotide comparison set forth above, or as defined as being capable of hybridizing to the nucleic acid segment of SEQ ID NO:1 under stringent conditions such as those described herein.

As used herein, a "DNA/RNA hybrid" is understood to mean that a single strand of RNA is hybridized to a single strand of DNA.

The term "appropriate reaction conditions" as described herein mean that temperature, pH, buffer, and other parameters are adjusted to optimize the reaction rate and yield.

As used herein, "hybridization," "hybridizes" or "capable of hybridizing" is understood to mean the forming of a double or triple stranded molecule or a molecule with partial double or triple stranded nature. The term "hybridization," "hybridize(s)" or "capable of hybridizing" encompasses the terms "stringent condition(s)" or "high stringency" and the terms "low stringency" or "low stringency condition(s)."

As used herein "stringent condition(s)" or "high stringency" are those conditions that allow hybridization between or within one or more nucleic acid strand(s) containing complementary sequence(s), but precludes hybridization of random sequences. Stringent conditions tolerate little, if any, mismatch between a nucleic acid and a target strand. Such conditions are well known to those of ordinary skill in the art, and are preferred for applications requiring high selectivity. Non-limiting applications include isolating a nucleic acid, such as a gene or a nucleic acid segment thereof, or detecting at least one specific mRNA transcript or a nucleic acid segment thereof, and the like.

Stringent conditions may comprise low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.15 M NaCl at temperatures of about 50° C. to about 70° C. It is understood that the temperature and ionic strength of a desired stringency are determined in part by the length of the particular nucleic acid(s), the length and nucleobase content of the target sequence(s), the charge composition of the nucleic acid(s), and to the presence or concentration of formamide, tetramethylammonium chloride or other solvent(s) in a hybridization mixture.

It is also understood that these ranges, compositions and conditions for hybridization are mentioned by way of non-limiting examples only, and that the desired stringency for a particular hybridization reaction is often determined empirically by comparison to one or more positive or negative controls. Depending on the application envisioned it is preferred to employ varying conditions of hybridization to achieve varying degrees of selectivity of a nucleic acid towards a target sequence. In a non-limiting example, identification or isolation of a related target nucleic acid that does not hybridize to a nucleic acid under stringent conditions may be achieved by hybridization at low temperature and/or high ionic strength. For example, a medium stringency condition could be provided by about 0.1 to 0.25 M NaCl at temperatures of about 37° C. to about 55° C. Under these conditions, hybridization may occur even though the sequences of probe and target strand are not perfectly complementary, but are mismatched at one or more positions. In another example, a low stringency condition could be provided by about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20° C. to about 55° C. Of course, it is within the skill of one in the art to further modify the low or high stringency conditions to suit a particular application. For example, in other embodiments, hybridization may be achieved under conditions of 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl<sub>2</sub>, 1.0 mM dithiothreitol, at temperatures between approximately 20° C. to about 37° C. Other hybridization conditions utilized could include approximately 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, at temperatures ranging from approximately 40° C. to about 72° C.

Accordingly, the nucleotide sequences of the disclosure may be used for their ability to selectively form duplex molecules with complementary stretches of genes or RNAs or to provide primers for amplification of DNA or RNA from tissues. Depending on the application envisioned, it is preferred

to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence.

The nucleic acid segments of the present invention, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, enhancers, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol.

For example, nucleic acid fragments may be prepared that include a contiguous stretch of nucleotides identical to or complementary to SEQ ID NO:1, 3, 5, 7 or 14. Nucleic acid fragments for use as a DNA transcription template may also be prepared. These fragments may be short or of intermediate lengths, such as, for example, about 8, about 10 to about 14, or about 15 to about 20 nucleotides, and that are chromosome-sized pieces, up to about 35,000, about 30,000, about 25,000, about 20,000, about 15,000, about 10,000, or about 5,000 base pairs in length, as well as DNA segments with total lengths of about 1,000, about 500, about 200, about 100 and about 50 base pairs in length (including all intermediate lengths of these lengths listed above, i.e., any range derivable therein and any integer derivable therein such a range) are also contemplated to be useful.

For example, it will be readily understood that "intermediate lengths," in these contexts, means any length between the quoted ranges, such as 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 130, 140, 150, 160, 170, 180, 190, including all integers through the 200-500; 500-1,000; 1,000-2,000; 2,000-3,000; 3,000-5,000; 5,000-10,000 ranges, up to and including sequences of about 12,001, 12,002, 13,001, 13,002, 15,000, 20,000 and the like.

Various nucleic acid segments may be designed based on a particular nucleic acid sequence, and may be of any length. By assigning numeric values to a sequence, for example, the first residue is 1, the second residue is 2, etc., an algorithm defining all nucleic acid segments can be created:

$$n \text{ to } n+y$$

where n is an integer from 1 to the last number of the sequence and y is the length of the nucleic acid segment minus one, where n+y does not exceed the last number of the sequence. Thus, for a 10-mer, the nucleic acid segments correspond to bases 1 to 10, 2 to 11, 3 to 12 . . . and/or so on. For a 15-mer, the nucleic acid segments correspond to bases 1 to 15, 2 to 16, 3 to 17 . . . and/or so on. For a 20-mer, the nucleic segments correspond to bases 1 to 20, 2 to 21, 3 to 22 . . . and/or so on. In certain embodiments, the nucleic acid segment may be a probe or primer. As used herein, a "probe" generally refers to a nucleic acid used in a detection method or composition. As used herein, a "primer" generally refers to a nucleic acid used in an extension or amplification method or composition.

The use of a hybridization probe of between 17 and 100 nucleotides in length, or in some aspect of the invention even up to 1-2 Kb or more in length, allows the formation of a duplex molecule that is both stable and selective. Molecules having complementary sequences over stretches greater than 20 bases in length are generally preferred, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of particular hybrid molecules obtained.

One will generally prefer to design nucleic acid molecules having complementary sequences over stretches of 20 to 30 nucleotides, or even longer where desired. Such fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means or by introducing selected sequences into recombinant vectors for recombinant production.

In general, it is envisioned that the hybridization probes described herein will be useful both as reagents in solution hybridization, as in PCR<sup>TM</sup>, for detection of expression of corresponding genes, as well as in embodiments employing a solid phase. In embodiments involving a solid phase, the test DNA (or RNA) is adsorbed or otherwise affixed to a selected matrix or surface. This fixed, single-stranded nucleic acid is then subjected to hybridization with selected probes under desired conditions. The selected conditions will depend on the particular circumstances based on the particular criteria required (depending, for example, on the G+C content, type of target nucleic acid, source of nucleic acid, size of hybridization probe, etc.). Following washing of the hybridized surface to remove non-specifically bound probe molecules, hybridization is detected, or even quantified, by means of the label.

#### b. Nucleic Acid Amplification

Nucleic acid used as a template for amplification is isolated from cells contained in the biological sample, according to standard methodologies (Sambrook et al., 1989). The nucleic acid may be genomic DNA or fractionated or whole cell RNA. Where RNA is used, it may be desired to convert the RNA to a complementary DNA. In one embodiment, the RNA is whole cell RNA and is used directly as the template for amplification.

Pairs of primers that selectively hybridize to nucleic acids are contacted with the isolated nucleic acid under conditions that permit selective hybridization. The term "primer," as defined herein, is meant to encompass any nucleic acid that is capable of priming the synthesis of a nascent nucleic acid in a template-dependent process. Typically, primers are oligonucleotides from ten to twenty or thirty base pairs in length, but longer sequences can be employed. Primers may be provided in double-stranded or single-stranded form, although the single-stranded form is preferred.

Once hybridized, the nucleic acid:primer complex is contacted with one or more enzymes that facilitate template-dependent nucleic acid synthesis. Multiple rounds of amplification, also referred to as "cycles," are conducted until a sufficient amount of amplification product is produced.

Next, the amplification product is detected. In certain applications, the detection may be performed by visual means. Alternatively, the detection may involve indirect identification of the product via chemiluminescence, radioactive scintigraphy of incorporated radiolabel or fluorescent label, or even via a system using electrical or thermal impulse signals (Affymax technology).

A number of template dependent processes are available to amplify the marker sequences present in a given template sample. One of the best known amplification methods is the polymerase chain reaction (referred to as PCR<sup>TM</sup>) which is described in detail in U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,800,159, each incorporated herein by reference in its entirety.

Briefly, in PCR<sup>TM</sup>, two primer sequences are prepared that are complementary to regions on opposite complementary strands of the marker sequence. An excess of deoxynucleoside triphosphates are added to a reaction mixture along with a DNA polymerase, e.g., Taq polymerase. If the marker sequence is present in a sample, the primers will bind to the

marker and the polymerase will cause the primers to be extended along the marker sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the marker to form reaction products, excess primers will bind to the marker and to the reaction products, and the process is repeated.

A reverse transcriptase PCR<sup>TM</sup> amplification procedure may be performed in order to quantify the amount of mRNA amplified. Methods of reverse transcribing RNA into cDNA are well known and described in Sambrook et al., 1989. Alternative methods for reverse transcription utilize thermostable, RNA-dependent DNA polymerases. These methods are described in WO 90/07641, filed Dec. 21, 1990, incorporated herein by reference. Polymerase chain reaction methodologies are well known in the art.

Another method for amplification is the ligase chain reaction ("LCR"), disclosed in EPA No. 320 308, incorporated herein by reference in its entirety. In LCR, two complementary probe pairs are prepared, and in the presence of the target sequence, each pair will bind to opposite complementary strands of the target such that they abut. In the presence of a ligase, the two probe pairs will link to form a single unit. By temperature cycling, as in PCR<sup>TM</sup>, bound ligated units dissociate from the target and then serve as "target sequences" for ligation of excess probe pairs. U.S. Pat. No. 4,883,750 describes a method similar to LCR for binding probe pairs to a target sequence.

Qbeta Replicase, described in PCT Application No. PCT/US87/00880, incorporated herein by reference, may also be used as still another amplification method in the present invention. In this method, a replicative sequence of RNA that has a region complementary to that of a target is added to a sample in the presence of an RNA polymerase. The polymerase will copy the replicative sequence, which can then be detected.

An isothermal amplification method, in which restriction endonucleases and ligases are used to achieve the amplification of target molecules that contain nucleotide 5'-[alpha-thio]-triphosphates in one strand of a restriction site may also be useful in the amplification of nucleic acids in the present invention.

Strand Displacement Amplification (SDA) is another method of carrying out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement and synthesis, i.e., nick translation. A similar method, called Repair Chain Reaction (RCR), involves annealing several probes throughout a region targeted for amplification, followed by a repair reaction in which only two of the four bases are present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA. Target specific sequences can also be detected using a cyclic probe reaction (CPR). In CPR, a probe having 3' and 5' sequences of non-specific DNA and a middle sequence of specific RNA is hybridized to DNA that is present in a sample. Upon hybridization, the reaction is treated with RNase H, and the products of the probe identified as distinctive products that are released after digestion. The original template is annealed to another cycling probe and the reaction is repeated.

Still another amplification method described in GB Application No. 2 202 328, and in PCT Application No. PCT/US89/01025, each of which is incorporated herein by reference in its entirety, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template- and enzyme-dependent synthesis. The primers may be modified by labeling with

a capture moiety (e.g., biotin) and/or a detector moiety (e.g., enzyme). In the latter application, an excess of labeled probes are added to a sample. In the presence of the target sequence, the probe binds and is cleaved catalytically. After cleavage, the target sequence is released intact to be bound by excess probe. Cleavage of the labeled probe signals the presence of the target sequence.

Other nucleic acid amplification procedures include transcription-based amplification systems (TAS), including nucleic acid sequence based amplification (NASBA) and 3SR (Gingeras et al., PCT Application WO 88/10315, incorporated herein by reference). In NASBA, the nucleic acids can be prepared for amplification by standard phenol/chloroform extraction, heat denaturation of a clinical sample, treatment with lysis buffer and minispin columns for isolation of DNA and RNA or guanidinium chloride extraction of RNA. These amplification techniques involve annealing a primer which has target specific sequences. Following polymerization, DNA/RNA hybrids are digested with RNase H while double-stranded DNA molecules are heat denatured again. In either case, the single-stranded DNA is made fully double-stranded by addition of second target specific primer, followed by polymerization. The double-stranded DNA molecules are then multiply transcribed by an RNA polymerase such as T7 or SP6. In an isothermal cyclic reaction, the RNAs are reverse transcribed into single-stranded DNA, which is then converted to double-stranded DNA, and then transcribed once again with an RNA polymerase such as T7 or SP6. The resulting products, whether truncated or complete, indicate target specific sequences.

Davey et al., EPA No. 329 822 (incorporated herein by reference in its entirety) disclose a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (ds-DNA), which may be used in accordance with the present invention. The ssRNA is a template for a first primer oligonucleotide, which is elongated by reverse transcriptase (RNA-dependent DNA polymerase). The RNA is then removed from the resulting DNA:RNA duplex by the action of ribonuclease H (RNase H, an RNase specific for RNA in duplex with either DNA or RNA). The resultant ssDNA is a template for a second primer, which also includes the sequences of an RNA polymerase promoter (exemplified by T7 RNA polymerase) 5' to its homology to the template. This primer is then extended by DNA polymerase (exemplified by the large "Klenow" fragment of *E. coli* DNA polymerase I), resulting in a double-stranded DNA ("dsDNA") molecule, having a sequence identical to that of the original RNA between the primers and having additionally, at one end, a promoter sequence. This promoter sequence can be used by the appropriate RNA polymerase to make many RNA copies of the DNA. These copies can then re-enter the cycle leading to very swift amplification. With proper choice of enzymes, this amplification can be done isothermally without addition of enzymes at each cycle. Because of the cyclical nature of this process, the starting sequence can be chosen to be in the form of either DNA or RNA.

Miller et al., PCT Application WO 89/06700 (incorporated herein by reference in its entirety) disclose a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. This scheme is not cyclic, i.e., new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" and "one-sided PCR" (Frohman, 1990, incorporated herein by reference).

Methods based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide," thereby amplifying the di-oligonucleotide, may also be used in the amplification step of the present invention.

#### c. Nucleic Acid Detection

In certain embodiments, it will be advantageous to employ nucleic acid sequences of the present invention such as all or part of SEQ ID NO:1, 3, 5, 7, 14 or a mutant thereof in combination with an appropriate means, such as a label, for hybridization assays, RNase protection and Northern hybridization. A wide variety of appropriate indicator means are known in the art, including fluorescent, radioactive, enzymatic or other ligands, such as avidin/biotin, which are capable of being detected. In preferred embodiments, one may desire to employ a fluorescent label or an enzyme tag such as urease, alkaline phosphatase or peroxidase, instead of radioactive or other environmentally undesirable reagents. In the case of enzyme tags, colorimetric indicator substrates are known that can be employed to provide a detection means visible to the human eye or spectrophotometrically, to identify specific hybridization with complementary nucleic acid-containing samples.

In embodiments wherein nucleic acids are amplified, it may be desirable to separate the amplification product from the template and the excess primer for the purpose of determining whether specific amplification has occurred. In one embodiment, amplification products are separated by agarose, agarose-acrylamide or polyacrylamide gel electrophoresis using standard methods (Sambrook et al., 1989).

Alternatively, chromatographic techniques may be employed to effect separation. There are many kinds of chromatography which may be used in the present invention: adsorption, partition, ion-exchange and molecular sieve, and many specialized techniques for using them including column, paper, thin-layer and gas chromatography.

Amplification products must be visualized in order to confirm amplification of the marker sequences. One typical visualization method involves staining of a gel with ethidium bromide and visualization under UV light. Alternatively, if the amplification products are integrally labeled with radio- or fluorometrically-labeled nucleotides, the amplification products can then be exposed to x-ray film or visualized under the appropriate stimulating spectra, following separation.

In one embodiment, visualization is achieved indirectly. Following separation of amplification products, a labeled, nucleic acid probe is brought into contact with the amplified marker sequence. The probe preferably is conjugated to a chromophore but may be radiolabeled. In another embodiment, the probe is conjugated to a binding partner, such as an antibody or biotin, and the other member of the binding pair carries a detectable moiety.

In one embodiment, detection is by Southern blotting and hybridization with a labeled probe. The techniques involved in Southern blotting are well known to those of skill in the art and can be found in many standard books on molecular protocols (see Sambrook et al., 1989). Briefly, amplification products are separated by gel electrophoresis. The gel is then contacted with a membrane, such as nitrocellulose, permitting transfer of the nucleic acid and non-covalent binding. Subsequently, the membrane is incubated with a chromophore-conjugated probe that is capable of hybridizing with a target amplification product. Detection is by exposure of the membrane to x-ray film or ion-emitting detection devices.

One example of the foregoing is described in U.S. Pat. No. 5,279,721, incorporated by reference herein, which discloses

an apparatus and method for the automated electrophoresis and transfer of nucleic acids. The apparatus permits electrophoresis and blotting without external manipulation of the gel and is ideally suited to carrying out methods according to the present invention.

Other methods for genetic screening to accurately detect mutations in genomic DNA, cDNA or RNA samples may be employed, depending on the specific situation.

Historically, a number of different methods have been used to detect point mutations, including denaturing gradient gel electrophoresis ("DGGE"), restriction enzyme polymorphism analysis, chemical and enzymatic cleavage methods, and others. The more common procedures currently in use include direct sequencing of target regions amplified by PCR<sup>TM</sup> (see above) and single-strand conformation polymorphism analysis ("SSCP").

Another method of screening for point mutations is based on RNase cleavage of base pair mismatches in RNA/DNA and RNA/RNA heteroduplexes. As used herein, the term "mismatch" is defined as a region of one or more unpaired or mispaired nucleotides in a double-stranded RNA/RNA, RNA/DNA or DNA/DNA molecule. This definition thus includes mismatches due to insertion/deletion mutations, as well as single and multiple base point mutations.

U.S. Pat. No. 4,946,773 describes an RNase A mismatch cleavage assay that involves annealing single-stranded DNA or RNA test samples to an RNA probe, and subsequent treatment of the nucleic acid duplexes with RNase A. After the RNase cleavage reaction, the RNase is inactivated by proteolytic digestion and organic extraction, and the cleavage products are denatured by heating and analyzed by electrophoresis on denaturing polyacrylamide gels. For the detection of mismatches, the single-stranded products of the RNase A treatment, electrophoretically separated according to size, are compared to similarly treated control duplexes. Samples containing smaller fragments (cleavage products) not seen in the control duplex are scored as positive.

Currently available RNase mismatch cleavage assays, including those performed according to U.S. Pat. No. 4,946,773, require the use of radiolabeled RNA probes. Myers and Maniatis in U.S. Pat. No. 4,946,773 describe the detection of base pair mismatches using RNase A. Other investigators have described the use of an *E. coli* enzyme, RNase I, in mismatch assays. Because it has broader cleavage specificity than RNase A, RNase I would be a desirable enzyme to employ in the detection of base pair mismatches if components can be found to decrease the extent of non-specific cleavage and increase the frequency of cleavage of mismatches. The use of RNase I for mismatch detection is described in literature from Promega Biotech. Promega markets a kit containing RNase I that is shown in their literature to cleave three out of four known mismatches, provided the enzyme level is sufficiently high.

The RNase Protection assay was first used to detect and map the ends of specific mRNA targets in solution. The assay relies on being able to easily generate high specific activity radiolabeled RNA probes complementary to the mRNA of interest by in vitro transcription. Originally, the templates for in vitro transcription were recombinant plasmids containing bacteriophage promoters. The probes are mixed with total cellular RNA samples to permit hybridization to their complementary targets, then the mixture is treated with RNase to degrade excess unhybridized probe. Also, as originally intended, the RNase used is specific for single-stranded RNA, so that hybridized double-stranded probe is protected from degradation. After inactivation and removal of the RNase, the protected probe (which is proportional in amount

to the amount of target mRNA that was present) is recovered and analyzed on a polyacrylamide gel.

The RNase Protection assay was adapted for detection of single base mutations. In this type of RNase A mismatch cleavage assay, radiolabeled RNA probes transcribed in vitro from wild-type sequences are hybridized to complementary target regions derived from test samples. The test target generally comprises DNA (either genomic DNA or DNA amplified by cloning in plasmids or by PCR<sup>TM</sup>), although RNA targets (endogenous mRNA) have occasionally been used. If single nucleotide (or greater) sequence differences occur between the hybridized probe and target, the resulting disruption in Watson-Crick hydrogen bonding at that position ("mismatch") can be recognized and cleaved in some cases by single-strand specific ribonuclease. To date, RNase A has been used almost exclusively for cleavage of single-base mismatches, although RNase I has recently been shown as useful also for mismatch cleavage. There are recent descriptions of using the MutS protein and other DNA-repair enzymes for detection of single-base mismatches.

Nuclease S1 analysis of reaction products can be used to measure RNA. An exemplary procedure for S1 analysis involves hybridization reaction with the RNA of interest (0.005-0.1 mg) and an excess of S1 probe which comprises a labeled oligonucleotide complementary to 20-80 or more sequential nucleotides of a specific RNA in S1 hybridization buffer (80% formamide, 0.4 M NaCl, 1 mM EDTA, 40 mM Pipes, pH 6.4). After denaturation for 4 min at 94° C., overnight hybridization at 30° C. and precipitation with ethanol, the S1 probe/RNA mixture is resuspended in S1 buffer (0.26 M NaCl, 0.05 M sodium acetate, pH 4.6, and 4.5 mM zinc sulfate). The sample is divided into two volumes and 100 units of S1 nuclease (Sigma Chemical Company) is added to one tube. The samples are incubated for 60 minutes at 37° C.; then EDTA (10 mM final concentration) and 15 g polyI-polyC RNA are added and the sample is extracted with phenol/chloroform and precipitated in ethanol. The samples are then subjected to polyacrylamide gel electrophoresis.

One method to produce a radiolabeled RNA probe with high specific activity includes admixing a radiolabeled NTP during transcription. Suitable isotopes for radiolabeling include <sup>35</sup>S- and <sup>32</sup>P-labeled UTP, GTP, CTP or ATP. For optimal results, a gel-purified radiolabeled RNA probe which is preferentially 300-500 bases in length, with a specific activity of 1-3 X10<sup>8</sup> cpm/μg should be generated using the RNA polymerase of the current invention. In order to produce this in vitro transcript, it is often advisable to use a high specific activity (e.g., [<sup>α-32</sup>P]CTP at 3,000Ci/mmol) NTP. To prevent background hybridization, it is important to remove plasmid template DNA by digestion which can be done with, for example, RQ1 RNase-Free DNase followed by phenol:chloroform:isoamyl alcohol extraction and ethanol precipitation.

Another method for producing radiolabeled probes includes using a riboprobe system which can produce high specific activity, radiolabeled RNA probes or microgram quantities of in vitro transcript. Riboprobes are useful with radiolabeled RNA probes in many applications including RNase protection, Northern hybridization, S1 analysis and in situ hybridization assays. The principle components of an in vitro transcription are the riboprobe, an RNA polymerase, a DNA template which includes a phage RNA polymerase promoter and ribonucleotide triphosphates.

#### d. Cloning vRNAP Genes

The present invention contemplates cloning vRNAP, or more particularly mini-vRNAP genes. A technique often employed by those skilled in the art of protein production today is to obtain a so-called "recombinant" version of the

protein, to express it in a recombinant cell and to obtain the protein, polypeptide or peptide from such cells. These techniques are based upon the "cloning" of a nucleic acid molecule encoding the protein from a DNA library, i.e., on obtaining a specific DNA molecule distinct from other portions of DNA. This can be achieved by, for example, cloning a cDNA molecule, or cloning a genomic-like DNA molecule.

The first step in such cloning procedures is the screening of an appropriate DNA library, such as, for example, from a phage, bacteria, yeast, fungus, mouse, rat, monkey or human. The screening protocol may utilize nucleotide segments or probes that are designed to hybridize to cDNA or genomic sequences of vRNAPs from protists. Additionally, antibodies designed to bind to the expressed vRNAP proteins, polypeptides, or peptides may be used as probes to screen an appropriate viral, eubacterial, archaeobacterial or eukaryotic DNA expression library. Alternatively, activity assays may be employed. The operation of such screening protocols are well known to those of skill in the art and are described in detail in the scientific literature, for example, in Sambrook et al. (1989), incorporated herein by reference. Moreover, as the present invention encompasses the cloning of genomic segments as well as cDNA molecules, it is contemplated that suitable genomic cloning methods, as known to those in the art, may also be used.

Encompassed by the invention are DNA segments encoding relatively small peptides, such as, for example, peptides of from about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 35, about 40, about 45, to about 50 amino acids in length, and more preferably, of from about 15 to about 30 amino acids in length; as set forth in SEQ ID NO:2, 4, 6, 8 or 15 and also larger polypeptides up to and including proteins corresponding to the full-length sequences set forth in SEQ ID NO:2 and SEQ ID NO:15, and any range derivable therein and any integer derivable in such a range. In addition to the "standard" DNA and RNA nucleotide bases, modified bases are also contemplated for use in particular applications of the present invention. A table of exemplary, but not limiting, modified bases is provided herein below.

TABLE 2

Modified Bases			
Abbr.	Modified base description	Abbr.	Modified base description
ac4c	4-acetylcytidine	Mam5s2u	5-methoxyaminomethyl-2-thiouridine
chm5u	5-(carboxyhydroxymethyl)uridine	Man q	Beta,D-mannosylqueosine
Cm	2'-O-methylcytidine	Mcm5s2u	5-methoxycarbonylmethyl-2-thiouridine
Cmm5s2u	5-carboxymethylaminomethyl-2-thiouridine	Mcm5u	5-methoxycarbonylmethyluridine
Cmm5u	5-carboxymethylaminomethyluridine	Mo5u	5-methoxyuridine
D	Dihydrouridine	Ms2i6a	2-methylthio-N6-isopentenyladenosine
Fm	2'-O-methylpseudouridine	Ms2t6a	N-((9-beta-D-ribofuranosyl-2-methylthiopurine-6-yl)carbamoyl)threonine

TABLE 2-continued

Modified Bases			
Abbr.	Modified base description	Abbr.	Modified base description
gal q	Beta,D-galactosylqueosine	Mt6a	N-((9-beta-D-ribofuranosylpurine-6-yl)N-methylcarbamoyl)threonine
Gm	2'-O-methylguanosine	Mv	Uridine-5-oxycetic acid methyl ester
I	Inosine	o5u	Uridine-5-oxycetic acid (v)
I6a	N6-isopentenyladenosine	Osyw	Wybutoxosine
m1a	1-methyladenosine	P	Pseudouridine
m1f	1-methylpseudouridine	Q	Queosine
m1g	1-methylguanosine	s2c	2-thiocytidine
m1l	1-methylinosine	s2t	5-methyl-2-thiouridine
m22g	2,2-dimethylguanosine	s2u	2-thiouridine
m2a	2-methyladenosine	s4u	4-thiouridine
m2g	2-methylguanosine	T	5-methyluridine
m3c	3-methylcytidine	t6a	N-((9-beta-D-ribofuranosylpurine-6-yl)carbamoyl)threonine
m5c	5-methylcytidine	Tm	2'-O-methyl-5-methyluridine
m6a	N6-methyladenosine	Um	2'-O-methyluridine
m7g	7-methylguanosine	Yw	Wybutosine
Mam5u	5-methylaminomethyluridine	X	3-(3-amino-3-carboxypropyl)uridine, (acp3)u

### III. Recombinant Vectors, Promoters, Host Cells and Expression

Recombinant vectors form an important further aspect of the present invention. The term "expression vector or construct" means any type of genetic construct containing a nucleic acid coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed. The transcript may be translated into a proteinaceous molecule, but it need not be, such as in the case of mini-vRNAP transcribing an RNA using a single-stranded DNA template. Thus, in certain embodiments, expression includes both transcription of a single-stranded DNA and translation of an RNA into the protein product. In other embodiments, expression only includes transcription of the nucleic acid. A recombinant vector can also be used for delivery of the RNA of the current invention.

Particularly useful vectors are contemplated to be those vectors in which the coding portion of the DNA segment, whether encoding a full length protein or smaller polypeptide or peptide, is positioned under the transcriptional control of a promoter. A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. The phrases "operatively positioned," "under control" or "under transcriptional control" means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the gene.

One particularly useful vector is pBAD. The pBAD expression vectors allow for greater control of bacterial expression of recombinant proteins and allow tight regulation for turning expression on or off. pBAD vectors allow for dose dependent induction for modulation of expression levels. The pBAD expression system helps overcome two of the most common problems of heterologous protein expression in bacteria: tox-

icity of the recombinant protein to the host and insolubility of the recombinant protein when it is expressed at high, uncontrolled levels. In both cases, a tightly-regulated expression system is critical for maximizing recombinant protein yields. The pBAD expression system is based on the araBAD operon which controls the arabinose metabolic pathway in *E. coli* and allows for precise modulation of heterologous expression to levels that are optimal for recovering high yields of the protein of interest (Guzman et al., 1995).

#### a. Promoters

Any promoters normally found in a host cell in the native state can be used in the present invention to drive expression of N4 vRNA or mini-vRNA polymerase. Also, promoters not normally found in the host cell in the native state that are recognized by a native, normally native host cell RNA polymerase, or non-native RNA polymerase expressed in the cell can be used in the present invention to drive expression of the RNA polymerase. Other promoters may be selected from a nucleic acid sequence database accessible to those of skill in the art, e.g., GenBank, or the promoter can be isolated by a screening method. A promoter recognized by the host cell can be operably linked to the gene or genes encoding the N4 RNA polymerase. The operable linkage can be constructed using any known techniques for DNA manipulation, as referred to herein.

Promoters are described as either constitutive or inducible. Constitutive promoters actively drive expression of genes under their control. Inducible promoters, in contrast, are activated in response to specific environmental stimuli. Both constitutive and inducible promoters can be used in the present invention for expressing non-host genes in a host cell.

Inducible promoters include, but are not limited to, trp, tac, lac, ara, reca,  $\lambda$ Pr, and  $\lambda$ P1. These promoters and others that can be used in the present invention for expression of the N4 vRNA or mini-vRNA polymerase, in embodiments in which the host cell is *E. coli*, are described by Makrides, *Microbiological Reviews*, (1996), 60, 512-538, herein incorporated by reference. Further, in embodiments of the present invention wherein the host cell is a microbe other than *E. coli*, such as *Saccharomyces*, *Bacillus*, and *Pseudomonas*, any inducible promoter known to those skilled in the art to be active in the host cell can be used to drive expression of the heterologous RNA polymerase. (U.S. Pat. No. 6,218,145).

The promoter may be in the form of the promoter that is naturally associated with N4 vRNA or mini-vRNA polymerase, as may be obtained by isolating the 5' non-coding sequences located upstream of the coding segment or exon, for example, using recombinant cloning and/or PCR<sup>TM</sup> technology, in connection with the compositions disclosed herein (PCR<sup>TM</sup> technology is disclosed in U.S. Pat. No. 4,683,202 and U.S. Pat. No. 4,682,195, each incorporated herein by reference).

In other embodiments, it is contemplated that certain advantages will be gained by positioning the coding DNA segment under the control of a recombinant, or heterologous, promoter. As used herein, a recombinant or heterologous promoter is intended to refer to a promoter that is not normally associated with N4 vRNA or mini-vRNA polymerase in its natural environment. Such promoters may include promoters normally associated with other genes, and/or promoters isolated from any other bacterial, viral, eukaryotic, protist, or mammalian cell, and/or promoters made by the hand of man that are not "naturally occurring," i.e., containing different elements from different promoters, or mutations that increase, decrease, or alter expression.

Naturally, it will be important to employ a promoter that effectively directs the expression of the DNA segment in the

cell type, organism, or even animal, chosen for expression. The use of promoter and cell type combinations for protein expression is generally known to those of skill in the art of molecular biology, for example, see Sambrook et al. (1989), incorporated herein by reference. The promoters employed may be constitutive, or inducible, and can be used under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins, polypeptides or peptides.

At least one module in a promoter generally functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation.

Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the thymidine kinase promoter, the spacing between promoter elements can be increased to 50 base pairs apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either co-operatively or independently to activate transcription.

The particular promoter that is employed to control the expression of a nucleic acid is not believed to be critical, so long as it is capable of expressing the nucleic acid in the targeted cell. Thus, where a human cell is targeted, it is preferable to position the nucleic acid coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a human or viral promoter.

In various other embodiments, the human cytomegalovirus (CMV) immediate early gene promoter, the SV40 early promoter and the Rous sarcoma virus long terminal repeat can be used to obtain high-level expression of the instant nucleic acids. The use of other viral or mammalian cellular or bacterial phage promoters which are well-known in the art to achieve expression are contemplated as well, provided that the levels of expression are sufficient for a given purpose. Tables 3 and 4 below list several elements/promoters which may be employed, in the context of the present invention, to regulate the expression of a vRNAP gene. This list is not intended to be exhaustive of all the possible elements involved in the promotion of expression but, merely, to be exemplary thereof.

In certain embodiments of the invention, promoter sequences may be used that are recognized specifically by a DNA-dependent RNA polymerase, such as, but not limited to, those described by Chamberlin and Ryan (1982) and by Jorgensen et al., (1991). These promoters can be used to express a wild-type or mutant form of a miniV RNA polymerase of the invention. Several RNA polymerase promoter sequences are especially useful, including, but not limited to, promoters derived from SP6 (e.g., Zhou and Doetsch, 1993), T7 (e.g., Martin, and Coleman, 1987) and T3 (e.g., McGraw et al., 1985). An RNA polymerase promoter sequence derived from *Thermus thermophilus* can also be used (see, e.g., Wendt et al., 1990; Faraldo et al., 1992; Hartmann et al., 1987; Hartmann et al., 1991). The length of the promoter sequence will vary depending upon the promoter chosen. For example,

the T7 RNA polymerase promoter can be only about 25 bases in length and act as a functional promoter, while other promoter sequences require 50 or more bases to provide a functional promoter.

In other embodiments of the invention, a promoter is used that is recognized by an RNA polymerase from a T7-like bacteriophage. The genetic organization of all T7-like phages that have been examined has been found to be essentially the same as that of T7. Examples of T7-like phages according to the invention include, but are not limited to *Escherichia coli* phages T3, .phi.I, .phi.II, W31, H, Y, A1, 122, cro, C21, C22, and C23; *Pseudomonas putida* phage gh-1; *Salmonella typhimurium* phage SP6; *Serratia marcescens* phages IV; Citrobacter phage VIII; and Klebsiella phage No. 11 (Hausmann, 1976; Korsten et al., 1975; Dunn, et al. 1971; Towle, et al., 1975; Butler and Chamberlin, 1982).

When a T7 RNA polymerase promoter, or another T7-like RNA polymerase promoter is used to express a wild-type or mutant form of a gene for a miniV RNA polymerase of the invention, the gene can be expressed in a host cell which expresses the T7 RNA polymerase, or the corresponding T7-like RNA polymerase for the promoter used, wherein the RNA polymerase for the promoter is expressed either constitutively, or more preferably, from an inducible promoter. By way of example, a T7 RNA polymerase expression system, such as, but not limited to, the expression systems disclosed in, for example, U.S. Pat. Nos. 5,693,489 and 5,869,320, the disclosures of which are incorporated herein by reference in their entirety.

b. Enhancers

Enhancers were originally detected as genetic elements that increased transcription from a promoter located at a distant position on the same molecule of DNA. This ability to act over a large distance had little precedent in classic studies of prokaryotic transcriptional regulation. Subsequent work showed that regions of DNA with enhancer activity are organized much like promoters. That is, they are composed of many individual elements, each of which binds to one or more transcriptional proteins.

The basic distinction between enhancers and promoters is operational. An enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have one or more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a very similar modular organization.

Additionally any promoter/enhancer combination (as per the Eukaryotic Promoter Data Base EPDB, <http://www.epd.isb-sib.ch/>) could also be used to drive expression. Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic expression construct.

TABLE 3

Promoter and Enhancer Elements	
Promoter/Enhancer	References
Immunoglobulin Heavy Chain	Banerji et al., 1983; Gilles et al., 1983; Grosschedl and Baltimore, 1985; Atchinson and Perry, 1986, 1987; Imler et al., 1987; Weinberger et al., 1984; Kiledjian et al., 1988; Porton et al.; 1990
Immunoglobulin Light Chain	Queen and Baltimore, 1983; Picard and Schaffner, 1984

TABLE 3-continued

Promoter and Enhancer Elements	
Promoter/Enhancer	References
5 T-Cell Receptor	Luria et al., 1987; Winoto and Baltimore, 1989; Redondo et al.; 1990
10 HLA DQ a and DQ β	Sullivan and Peterlin, 1987
β-Interferon	Goodbourn et al., 1986; Fujita et al., 1987; Goodbourn and Maniatis, 1988
Interleukin-2	Greene et al., 1989
15 Interleukin-2 Receptor	Greene et al., 1989; Lin et al., 1990
MHC Class II 5	Koch et al., 1989
MHC Class II HLA-Dra	Sherman et al., 1989
β-Actin	Kawamoto et al., 1988; Ng et al.; 1989
Muscle Creatine Kinase	Jaynes et al., 1988; Horlick and Benfield, 1989; Johnson et al., 1989
Prealbumin (Transthyretin)	Costa et al., 1988
20 Elastase I	Ornitz et al., 1987
Metallothionein	Karin et al., 1987; Culotta and Hamer, 1989
Collagenase	Pinkert et al., 1987; Angel et al., 1987
Albumin Gene	Pinkert et al., 1987; Tronche et al., 1989, 1990
α-Fetoprotein	Godbout et al., 1988; Campere and Tilghman, 1989
25 t-Globin	Bodine and Ley, 1987; Perez-Stable and Constantini, 1990
β-Globin	Trudel and Constantini, 1987
e-fos	
c-HA-ras	Deschamps et al., 1985
30 Insulin	Edlund et al., 1985
Neural Cell Adhesion Molecule (NCAM)	Hirsh et al., 1990
α <sub>1</sub> -Antitrypsin	Latimer et al., 1990
H2B (TH2B) Histone	Hwang et al., 1990
Mouse or Type I Collagen	Ripe et al., 1989
35 Glucose-Regulated Proteins (GRP94 and GRP78)	Chang et al., 1989
Rat Growth Hormone	Larsen et al., 1986
Human Serum Amyloid A (SAA)	Edbrooke et al., 1989
Troponin I (TN I)	Yutzey et al., 1989
40 Platelet-Derived Growth Factor	Pech et al., 1989
Duchenne Muscular Dystrophy SV40	Klamut et al., 1990
	Banerji et al., 1981; Moreau et al., 1981; Sleigh and Lockett, 1985; Firak and Subramanian, 1986; Herr and Clarke, 1986; Imbra and Karin, 1986; Kadesch and Berg, 1986; Wang and Calame, 1986; Ondek et al., 1987; Kuhl et al., 1987; Schaffner et al., 1988
45 Polyoma	Swartzendruber and Lehman, 1975; Vasseur et al., 1980; Katinka et al., 1980, 1981; Tyndell et al., 1981; Dandolo et al., 1983; de Villiers et al., 1984; Hen et al., 1986; Satake et al., 1988; Campbell and Villarreal, 1988
50 Retroviruses	Kriegler and Botchan, 1982, 1983; Levinson et al., 1982; Kriegler et al., 1983, 1984a, b, 1988; Bosze et al., 1986; Miksicek et al., 1986; Celander and Haseltine, 1987; Thiesen et al., 1988; Celander et al., 1988; Choi et al., 1988; Reisman and Rotter, 1989
55 Papilloma Virus	Campo et al., 1983; Lusky et al., 1983; Spandidos and Wilkie, 1983; Spalholz et al., 1985; Lusky and Botchan, 1986; Cripe et al., 1987; Gloss et al., 1987; Hirochika et al., 1987; Stephens and Hentschel, 1987
Hepatitis B Virus	Bulla and Siddiqui, 1986; Jameel and Siddiqui, 1986; Shaul and Ben-Levy, 1987; Spandau and Lee, 1988; Vannice and Levinson, 1988
65	

TABLE 3-continued

<u>Promoter and Enhancer Elements</u>	
Promoter/Enhancer	References
Human Immunodeficiency Virus	Muesing et al., 1987; Hauber and Cullan, 1988; Jakobovits et al., 1988; Feng and Holland, 1988; Takebe et al., 1988; Rosen et al., 1988; Berkhout et al., 1989; Laspia et al., 1989; Sharp and Marciniak, 1989; Braddock et al., 1989
Cytomegalovirus	Weber et al., 1984; Boshart et al., 1985; Foecking and Hofstetter, 1986
Gibbon Ape Leukemia Virus	Holbrook et al., 1987; Quinn et al., 1989

TABLE 4

<u>Inducible Elements</u>		
Element	Inducer	References
MT II	Phorbol Ester (TFA) Heavy metals	Palmiter et al., 1982; Haslinger and Karin, 1985; Searle et al., 1985; Stuart et al., 1985; Imagawa et al., 1987; Karin et al., 1987; Angel et al., 1987b; McNeall et al., 1989
MMTV (mouse mammary tumor virus)	Glucocorticoids	Huang et al., 1981; Lee et al., 1981; Majors and Varmus, 1983; Chandler et al., 1983; Lee et al., 1984; Ponta et al., 1985; Sakai et al., 1988
$\beta$ -Interferon	Poly(rI)x and Poly(rc)	Tavemier et al., 1983
Adenovirus 5 E2	Ela	Imperiale and Nevins, 1984
Collagenase	Phorbol Ester (TPA)	Angel et al., 1987a
Stromelysin	Phorbol Ester (TPA)	Angel et al., 1987b
SV40	Phorbol Ester (TPA)	Angel et al., 1987b
Murine MX Gene	Interferon, Newcastle Disease Virus	
GRP78 Gene	A23187	Resendez et al., 1988
$\alpha$ -2-Macroglobulin	IL-6	Kunz et al., 1989
Vimentin	Serum	Rittling et al., 1989
MHC Class I Gene H-2kb	Interferon	Blonar et al., 1989
HSP70	Ela, SV40 Large T Antigen	Taylor et al., 1989; Taylor and Kingston, 1990a, b
Proliferin	Phorbol Ester-TPA	Mordacq and Linzer, 1989
Tumor Necrosis Factor	FMA	Hensel et al., 1989
Thyroid Stimulating Hormone a Gene	Thyroid Hormone	Chatterjee et al., 1989

Turning to the expression of the proteinaceous molecules after transcription using the vRNAP, mini-vRNAP, or mutants thereof of the present invention, once a suitable clone or clones have been obtained, whether they be cDNA based or genomic, one may proceed to prepare an expression system. The engineering of DNA segment(s) for expression in a prokaryotic or eukaryotic system may be performed by techniques generally known to those of skill in recombinant expression. It is believed that virtually any expression system may be employed in the expression of the proteinaceous molecules of the present invention.

Both cDNA and genomic sequences are suitable for eukaryotic expression, as the host cell will generally process

the genomic transcripts to yield functional mRNA for translation into proteinaceous molecules. Generally speaking, it may be more convenient to employ as the recombinant gene a cDNA version of the gene. It is believed that the use of a cDNA version will provide advantages in that the size of the gene will generally be much smaller and more readily employed to transfect the targeted cell than will a genomic gene, which will typically be up to an order of magnitude or more larger than the cDNA gene. However, it is contemplated that a genomic version of a particular gene may be employed where desired.

In expression, one will typically include a polyadenylation signal to effect proper polyadenylation of the transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed. Preferred embodiments include the SV40 polyadenylation signal and the bovine growth hormone polyadenylation signal, convenient and known to function well in various target cells. Also contemplated as an element of the expression cassette is a terminator. These elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

#### c. Antisense and Ribozymes

In some embodiments of the invention the vRNA polymerase can be used to synthesize antisense RNA or ribozymes.

The term "antisense nucleic acid" is intended to refer to the oligonucleotides complementary to the base sequences of DNA and RNA. Antisense oligonucleotides, when introduced into a target cell, specifically bind to their target nucleic acid and interfere with transcription, RNA processing, transport, translation, and/or stability. Targeting double-stranded (ds) DNA with oligonucleotides leads to triple-helix formation; targeting RNA will lead to double-helix formation. An antisense nucleic acid may be complementary to SEQ ID NO:1, 3, 5, 7 or 14, complementary to a mini-vRNAP encoding sequence or to mini-vRNAP non-coding sequences. Antisense RNA constructs, or DNA encoding such antisense RNAs, may be employed to inhibit gene transcription or translation or both within a host cell, either in vitro or in vivo, such as within a host animal, including a human subject.

Antisense constructs may be designed to bind to the promoter and other control regions, exons, introns or even exon-intron boundaries (splice junctions) of a gene. It is contemplated that the most effective antisense constructs may include regions complementary to intron/exon splice junctions. Thus, antisense constructs with complementary regions within 50-200 bases of an intron-exon splice junction may be used. It has been observed that some exon sequences can be included in the construct without seriously affecting the target selectivity thereof. The amount of exonic material included will vary depending on the particular exon and intron sequences used. One can readily test whether too much exon DNA is included simply by testing the constructs in vitro to determine whether normal cellular function is affected or whether the expression of related genes having complementary sequences is affected.

As stated above, "complementary" or "antisense" means polynucleotide sequences that are substantially complementary over their entire length and have very few base mismatches. For example, sequences of fifteen bases in length may be termed complementary when they have complementary nucleotides at thirteen or fourteen positions. Naturally, sequences which are completely complementary will be sequences which are entirely complementary throughout their entire length and have no base mismatches. Other sequences with lower degrees of homology also are contem-

plated. For example, an antisense construct which has limited regions of high homology, but also contains a non-homologous region (e.g., ribozyme) could be designed. These molecules, though having less than 50% homology, would bind to target sequences under appropriate conditions.

It may be advantageous to combine portions of genomic DNA with cDNA or synthetic sequences to generate specific constructs. For example, where an intron is desired in the ultimate construct, a genomic clone will need to be used. The cDNA or a synthesized polynucleotide may provide more convenient restriction sites for the remaining portion of the construct and, therefore, would be used for the rest of the sequence.

While all or part of the gene sequence may be employed in the context of antisense construction, statistically, any sequence 17 bases long should occur only once in the human genome and, therefore, suffice to specify a unique target sequence. Although shorter oligomers are easier to make and increase in vivo accessibility, numerous other factors are involved in determining the specificity of hybridization. Both binding affinity and sequence specificity of an oligonucleotide to its complementary target increases with increasing length. It is contemplated that oligonucleotides of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more base pairs will be used. One can readily determine whether a given antisense nucleic acid is effective at targeting of the corresponding host cell gene simply by testing the constructs in vivo to determine whether the endogenous gene's function is affected or whether the expression of related genes having complementary sequences is affected.

In certain embodiments, one may wish to employ antisense constructs which include other elements, for example, those which include C-5 propyne pyrimidines. Oligonucleotides which contain C-5 propyne analogues of uridine and cytidine have been shown to bind RNA with high affinity and to be potent antisense inhibitors of gene expression (Wagner et al., 1993).

As an alternative to targeted antisense delivery, targeted ribozymes may be used. The term "ribozyme" refers to an RNA-based enzyme capable of targeting and cleaving particular base sequences in oncogene DNA and RNA. Ribozymes either can be targeted directly to cells, in the form of RNA oligonucleotides incorporating ribozyme sequences, or introduced into the cell as an expression construct encoding the desired ribozymal RNA. Ribozymes may be used and applied in much the same way as described for antisense nucleic acids. Sequences for ribozymes may be included in the DNA template to eliminate undesired 5' end sequences in RNAs generated through T7 RNA polymerase transcription.

Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, 1987; Gerlack et al., 1987; Forster and Symons, 1987). For example, a large number of ribozymes accelerate phosphoester transfer reactions with a high degree of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech et al., 1981; Michel and Westhof, 1990; Reinhold-Hurek and Shub, 1992). This specificity has been attributed to the requirement that the substrate bind via specific base-pairing interactions to the internal guide sequence ("IGS") of the ribozyme prior to chemical reaction.

Ribozyme catalysis has primarily been observed as part of sequence specific cleavage/ligation reactions involving nucleic acids (Joyce, 1989; Cech et al., 1981). For example, U.S. Pat. No. 5,354,855 reports that certain ribozymes can act as endonucleases with a sequence specificity greater than that

of known ribonucleases and approaching that of the DNA restriction enzymes. Thus, sequence-specific ribozyme-mediated inhibition of gene expression may be particularly suited to therapeutic applications (Scanlon et al., 1991; Sarver et al., 1990; Sioud et al., 1992). Recently, it was reported that ribozymes elicited genetic changes in some cell lines to which they were applied; the altered genes included the oncogenes H-ras, c-fos and genes of HIV. Most of this work involved the modification of a target mRNA, based on a specific mutant codon that is cleaved by a specific ribozyme. In light of the information included herein and the knowledge of one of ordinary skill in the art, the preparation and use of additional ribozymes that are specifically targeted to a given gene will now be straightforward.

Several different ribozyme motifs have been described with RNA cleavage activity (reviewed in Symons, 1992). Examples of ribozymes include sequences from the Group I self-splicing introns including tobacco ringspot virus (Prody, et al., 1986), avocado sunblotch viroid (Palukaitis, et al., 1979; Symons, 1981), and Lucerne transient streak virus (Forster and Symons, 1987). Sequences from these and related viruses are referred to as hammerhead ribozymes based on a predicted folded secondary structure.

Other suitable ribozymes include sequences from RNase P with RNA cleavage activity (Yuan, et al., 1992; Yuan and Altman, 1994), hairpin ribozyme structures (Berzal-Herranz, et al., 1992; Chowrira et al., 1993) and hepatitis  $\delta$  virus based ribozymes (Perrotta and Been, 1992). The general design and optimization of ribozyme directed RNA cleavage activity has been discussed in detail (Haseloff and Gerlach, 1988; Symons, 1992; Chowrira, et al., 1994; and Thompson, et al., 1995).

The other variable on ribozyme design is the selection of a cleavage site on a given target RNA. Ribozymes are targeted to a given sequence by virtue of annealing to a site by complementary base pair interactions. Two stretches of homology are required for this targeting. These stretches of homologous sequences flank the catalytic ribozyme structure defined above. Each stretch of homologous sequence can vary in length from 7 to 15 nucleotides. The only requirement for defining the homologous sequences is that, on the target RNA, they are separated by a specific sequence which is the cleavage site. For hammerhead ribozymes, the cleavage site is a dinucleotide sequence on the target RNA, uracil (U) followed by either an adenine, cytosine or uracil (A, C or U; Perriman, et al., 1992; Thompson, et al., 1995). The frequency of this dinucleotide occurring in any given RNA is statistically 3 out of 16. Therefore, for a given target messenger RNA of 1000 bases, 187 dinucleotide cleavage sites are statistically possible.

Designing and testing ribozymes for efficient cleavage of a target RNA is a process well known to those skilled in the art. Examples of scientific methods for designing and testing ribozymes are described by Chowrira et al. (1994) and Lieber and Strauss (1995), each incorporated by reference. The identification of operative and preferred sequences for use in ribozymes is simply a matter of preparing and testing a given sequence, and is a routinely practiced "screening" method known to those of skill in the art.

A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon and adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be "in-frame" with the reading frame of the

desired coding sequence to ensure translation of the entire insert. The exogenous translational control signals and initiation codons can be either natural or synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements.

#### d. Host Cells

Host cells may be derived from prokaryotes or eukaryotes, including yeast cells, insect cells, and mammalian cells, depending upon whether the desired result is replication of the vector or expression of part or all of the vector-encoded nucleic acid sequences. Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection (ATCC), which is an organization that serves as an archive for living cultures and genetic materials ([www.atcc.org](http://www.atcc.org)). An appropriate host can be determined by one of skill in the art based on the vector backbone and the desired result. A plasmid or cosmid, for example, can be introduced into a prokaryotic host cell for replication of many vector copies. Bacterial cells used as host cells for vector replication and/or expression include DH5 $\alpha$ , BL 21, JM109, and KC8, as well as a number of commercially available bacterial hosts such as SURE<sup>®</sup> Competent Cells and SOLOPACK Gold Cells (STRAT-AGENE<sup>®</sup>, La Jolla, Calif.). Alternatively, bacterial cells such as *E. coli* LE392 could be used as host cells. Appropriate yeast cells include *Saccharomyces cerevisiae*, *Saccharomyces pombe*, and *Pichia pastoris*.

Examples of eukaryotic host cells for replication and/or expression of a vector include HeLa, NIH3T3, Jurrat, 293, Cos, CHO, Saos, BHK, C127 and PC12. Many host cells from various cell types and organisms are available and would be known to one of skill in the art. Similarly, a viral vector may be used in conjunction with either a eukaryotic or prokaryotic host cell, particularly one that is permissive for replication or expression of the vector.

Some vectors may employ control sequences that allow it to be replicated and/or expressed in both prokaryotic and eukaryotic cells. One of skill in the art would further understand the conditions under which to incubate all of the above described host cells to maintain them and to permit replication of a vector. Also understood and known are techniques and conditions that would allow large-scale production of vectors, as well as production of the nucleic acids encoded by vectors and/or their cognate polypeptides, proteins, or peptides.

It is proposed that vRNAP, or more particularly mini-vRNAP may be co-expressed with other selected proteinaceous molecules such as EcoSSB and other proteins of interest, wherein the proteinaceous molecules may be co-expressed in the same cell or vRNAP gene may be provided to a cell that already has another selected proteinaceous molecule. Co-expression may be achieved by co-transfecting the cell with two distinct recombinant vectors, each bearing a copy of either of the respective DNAs. Alternatively, a single recombinant vector may be constructed to include the coding regions for both of the proteinaceous molecules, which could then be expressed in cells transfected with the single vector. In either event, the term "co-expression" herein refers to the expression of both the vRNAP gene and the other selected proteinaceous molecules in the same recombinant cell.

As used herein, the terms "engineered" and "recombinant" cells or host cells are intended to refer to a cell into which an exogenous DNA segment or gene, such as a cDNA or gene encoding vRNAP, mini-vRNAP or a mutant thereof, has been introduced. Therefore, engineered cells are distinguishable from naturally occurring cells which do not contain a recombinantly introduced exogenous DNA segment or gene. Engi-

neered cells are thus cells having a gene or genes introduced through the hand of man. Recombinant cells include those having an introduced cDNA or genomic gene, and also include genes positioned adjacent to a promoter not naturally associated with the particular introduced gene.

To express a recombinant vRNAP, whether mutant or wild-type, in accordance with the present invention one would prepare an expression vector that comprises a wild-type, or mutant vRNAP proteinaceous molecule-encoding nucleic acid under the control of one or more promoters. To bring a coding sequence "under the control of" a promoter, one positions the 5' end of the transcription initiation site of the transcriptional reading frame generally between about 1 and about 50 nucleotides "downstream" of (i.e., 3' of) the chosen promoter. The "upstream" promoter directs transcription of the DNA and promotes expression of the encoded recombinant protein, polypeptide or peptide. This is the meaning of "recombinant expression" in this context.

Many standard techniques are available to construct expression vectors containing the appropriate nucleic acids and transcriptional/translational control sequences in order to achieve protein, polypeptide or peptide expression in a variety of host expression systems. Cell types available for expression include, but are not limited to, bacteria, such as *E. coli* and *B. subtilis*, transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors.

Certain examples of prokaryotic hosts are *E. coli* strain RR1, *E. coli* LE392, *E. coli* B, *E. coli* X 1776 (ATCC No. 31537) as well as *E. coli* W3110 (F-, lambda-, prototrophic, ATCC No. 273325); bacilli such as *Bacillus subtilis*; and other enterobacteriaceae such as *Salmonella typhimurium*, *Serratia marcescens*, and various *Pseudomonas* species.

In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication origin, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is often transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species. pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, promoters which can be used by the microbial organism for expression of its own proteins.

In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, the phage lambda GEM<sup>™</sup>-11 may be utilized in making a recombinant phage vector which can be used to transform host cells, such as *E. coli* LE392.

Further useful vectors include pIN vectors (Inouye et al., 1985); and pGEX vectors, for use in generating glutathione S-transferase (GST) soluble proteins for later purification and separation or cleavage.

The following details concerning recombinant protein production in bacterial cells, such as *E. coli*, are provided by way of exemplary information on recombinant protein production in general, the adaptation of which to a particular recombinant expression system will be known to those of skill in the art.

Bacterial cells, for example, *E. coli*, containing the expression vector are grown in any of a number of suitable media, for example, LB. The expression of the recombinant proteinaceous molecule may be induced, e.g., by adding IPTG or any appropriate inducer to the media or by switching incubation

to a higher temperature, depending on the regulated promoter used. After culturing the bacteria for a further period, generally of between 2 and 24 hours, the cells are collected by centrifugation and washed to remove residual media.

The bacterial cells are then lysed, for example, by disruption in a cell homogenizer, by sonication or cell press and centrifuged to separate the dense inclusion bodies and cell membranes from the soluble cell components. This centrifugation can be performed under conditions whereby the dense inclusion bodies are selectively enriched by incorporation of sugars, such as sucrose, into the buffer and centrifugation at a selective speed.

If the recombinant proteinaceous molecule is expressed in the inclusion bodies, as is the case in many instances, these can be washed in any of several solutions to remove some of the contaminating host proteins, then solubilized in solutions containing high concentrations of urea (e.g., 8M) or chaotropic agents such as guanidine hydrochloride in the presence of reducing agents, such as  $\beta$ -mercaptoethanol or DTT (dithiothreitol).

Under some circumstances, it may be advantageous to incubate the proteinaceous molecule for several hours under conditions suitable for the proteinaceous molecule to undergo a refolding process into a conformation which more closely resembles that of the native proteinaceous molecule. Such conditions generally include low proteinaceous molecule concentrations, less than 500 mg/ml, low levels of reducing agent, concentrations of urea less than 2 M and often the presence of reagents such as a mixture of reduced and oxidized glutathione which facilitate the interchange of disulfide bonds within the proteinaceous molecule.

The refolding process can be monitored, for example, by SDS-PAGE, or with antibodies specific for the native molecule (which can be obtained from animals vaccinated with the native molecule or smaller quantities of recombinant proteinaceous molecule). Following refolding, the proteinaceous molecule can then be purified further and separated from the refolding mixture by chromatography on any of several supports including ion exchange resins, gel permeation resins or on a variety of affinity columns.

For expression in *Saccharomyces*, the plasmid YRp7, for example, is commonly used. This plasmid already contains the *trp1* gene which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example ATCC No. 44076 or PEP4-1. The presence of the *trp1* lesion as a characteristic of the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

Suitable promoter sequences in yeast vectors include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate protein, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. In constructing suitable expression plasmids, the termination sequences associated with these genes are also ligated into the expression vector 3' of the sequence desired to be expressed to provide polyadenylation of the mRNA and termination.

In addition to micro-organisms, cultures of cells derived from multicellular organisms may also be used as hosts. In principle, any such cell culture is workable, whether from vertebrate or invertebrate culture. In addition to mammalian cells, these include insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus); and plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic

virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing one or more RNAP coding sequences.

Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteinaceous molecules. Appropriate cells lines or host systems can be chosen to ensure the correct modification and processing of the foreign proteinaceous molecule expressed.

A number of viral-based expression systems may be utilized, for example, commonly used promoters are derived from polyoma, Adenovirus 2, and most frequently Simian Virus 40 (SV40). The early and late promoters of SV40 virus are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication. Smaller or larger SV40 fragments may also be used, provided there is included the approximately 250 bp sequence extending from the HindIII site toward the BglI site located in the viral origin of replication.

In cases where an adenovirus is used as an expression vector, the coding sequences may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1, E3, or E4) will result in a recombinant virus that is viable and capable of expressing an RNA in infected hosts.

Specific initiation signals may also be used for more efficient translation using the vRNAP of the current invention. These signals include the ATG initiation codon and adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may additionally need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be in-frame (or in-phase) with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements and transcription terminators.

In eukaryotic expression, one will also typically desire to incorporate into the transcriptional unit an appropriate polyadenylation site (e.g., 5'-AATAAA-3') if one was not contained within the original cloned segment. Typically, the poly A addition site is placed about 30 to 2000 nucleotides "downstream" of the termination site of the proteinaceous molecule at a position prior to transcription termination.

For long-term, high-yield production of a recombinant vRNAP protein, polypeptide or peptide, stable expression is preferred. For example, cell lines that stably express constructs encoding a vRNAP protein, polypeptide or peptide may be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with vectors controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines.

A number of selection systems may be used, including, but not limited to, the herpes simplex virus thymidine kinase (tk), hypoxanthine-guanine phosphoribosyltransferase (hgprt) and adenine phosphoribosyltransferase (aprt) genes, in tk<sup>-</sup>, hgprt<sup>-</sup> or aprt<sup>-</sup> cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dihydrofolate reductase (dhfr), that confers resistance to methotrexate; gpt, that confers resistance to mycophenolic acid; neomycin (neo), that confers resistance to the aminoglycoside G-418; and hygromycin (hygro), that confers resistance to hygromycin.

Large scale suspension culture of bacterial cells in stirred tanks is a common method for production of recombinant proteinaceous molecules. Two suspension culture reactor designs are in wide use—the stirred reactor and the airlift reactor. The stirred design has successfully been used on an 8000 liter capacity for the production of interferon. Cells are grown in a stainless steel tank with a height-to-diameter ratio of 1:1 to 3:1. The culture is usually mixed with one or more agitators, based on bladed disks or marine propeller patterns. Agitator systems offering less shear forces than blades have been described. Agitation may be driven either directly or indirectly by magnetically coupled drives. Indirect drives reduce the risk of microbial contamination through seals on stirrer shafts.

The airlift reactor for microbial fermentation relies on a gas stream to both mix and oxygenate the culture. The gas stream enters a riser section of the reactor and drives circulation. Gas disengages at the culture surface, causing denser liquid free of gas bubbles to travel downward in the downcomer section of the reactor. The main advantage of this design is the simplicity and lack of need for mechanical mixing. Typically, the height-to-diameter ratio is 10:1. The airlift reactor scales up relatively easily, has good mass transfer of gases and generates relatively low shear forces.

It is contemplated that the vRNAP proteins, polypeptides or peptides of the invention may be “overexpressed,” i.e., expressed in increased levels relative to its natural expression in cells. Such overexpression may be assessed by a variety of methods, including radio-labeling and/or proteinaceous molecule purification. However, simple and direct methods are preferred, for example, those involving SDS/PAGE and proteinaceous composition staining or western blotting, followed by quantitative analyses, such as densitometric scanning of the resultant gel or blot. A specific increase in the level of the recombinant protein, polypeptide or peptide in comparison to the level in natural cells is indicative of overexpression, as is a relative abundance of the specific proteinaceous molecule in relation to the other proteins produced by the host cell and, e.g., visible on a gel.

#### IV. Methods of Gene Transfer

In order to mediate the effect of transgene expression in a cell, it will be necessary to transfer the expression constructs (e.g., a therapeutic construct) of the present invention into a cell. Such transfer may employ viral or non-viral methods of gene transfer. This section provides a discussion of methods and compositions of gene or nucleic acid transfer, including transfer of antisense sequences.

The vRNAP genes are incorporated into a viral vector to mediate gene transfer to a cell. Additional expression constructs encoding EcoSSB and other therapeutic agents as described herein may also be transferred via viral transduction using infectious viral particles, for example, by transformation with an adenovirus vector of the present invention. Alternatively, a retrovirus, bovine papilloma virus, an adeno-associated virus (AAV), a lentiviral vector, a vaccinia virus, a

polyoma virus, or an infective virus that has been engineered to express a specific binding ligand may be used. Similarly, nonviral methods which include, but are not limited to, direct delivery of DNA such as by injection, electroporation, calcium phosphate precipitation, liposome mediated transfection, and microprojectile bombardment may be employed. Thus, in one example, viral infection of cells is used in order to deliver therapeutically significant genes to a cell. Typically, the virus simply will be exposed to the appropriate host cell under physiologic conditions, permitting uptake of the virus.

Microinjection can be used for delivery into a cell. Microinjection involves the insertion of a substance such as RNA into a cell through a microelectrode. Typical applications include the injection of drugs, histochemical markers (such as horseradish peroxidase or lucifer yellow) and RNA or DNA in molecular biological studies. To extrude the substances through the very fine electrode tips, either hydrostatic pressure (pressure injection) or electric currents (iontophoresis) is employed.

#### V. Proteinaceous Compositions

In certain embodiments, the present invention concerns novel compositions or methods comprising at least one proteinaceous molecule. The proteinaceous molecule may have a sequence essentially as set forth in SEQ ID NO:2, 4, 6, 8 or 15. The proteinaceous molecule may be a vRNAP or more preferably a mini-vRNAP, or a delivery agent. The proteinaceous molecule may also be a mutated mini-vRNAP.

As used herein, a “proteinaceous molecule,” “proteinaceous composition,” “proteinaceous compound,” “proteinaceous chain” or “proteinaceous material” generally refers to, but is not limited to, a protein of greater than about 200 amino acids or the full length endogenous sequence translated from a gene; a polypeptide of greater than about 100 amino acids; and/or a peptide of from about 3 to about 100 amino acids. All the “proteinaceous” terms described above may be used interchangeably herein.

In certain embodiments the size of the at least one proteinaceous molecule may comprise, but is not limited to, about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, about 50, about 51, about 52, about 53, about 54, about 55, about 56, about 57, about 58, about 59, about 60, about 61, about 62, about 63, about 64, about 65, about 66, about 67, about 68, about 69, about 70, about 71, about 72, about 73, about 74, about 75, about 76, about 77, about 78, about 79, about 80, about 81, about 82, about 83, about 84, about 85, about 86, about 87, about 88, about 89, about 90, about 91, about 92, about 93, about 94, about 95, about 96, about 97, about 98, about 99, about 100, about 110, about 120, about 130, about 140, about 150, about 160, about 170, about 180, about 190, about 200, about 210, about 220, about 230, about 240, about 250, about 275, about 300, about 325, about 350, about 375, about 400, about 425, about 450, about 475, about 500, about 525, about 550, about 575, about 600, about 625, about 650, about 675, about 700, about 725, about 750, about 775, about 800, about 825, about 850, about 875, about 900, about 925, about 950, about 975, about 1000, about 1100, about 1200, about 1300, about 1400, about 1500, about 1750, about 2000, about 2250, about 2500 or greater amino molecule residues, and any range derivable therein.

As used herein, an "amino molecule" refers to any amino acid, amino acid derivative or amino acid mimic as would be known to one of ordinary skill in the art. In certain embodiments, the residues of the proteinaceous molecule are sequential, without any non-amino molecule interrupting the sequence of amino molecule residues. In other embodiments, the sequence may comprise one or more non-amino molecule moieties. In particular embodiments, the sequence of residues of the proteinaceous molecule may be interrupted by one or more non-amino molecule moieties.

Accordingly, the term "proteinaceous composition" encompasses amino molecule sequences comprising at least one of the 20 common amino acids in naturally synthesized proteins, or at least one modified or unusual amino acid, including but not limited to those shown on Table 5 below.

TABLE 5

Modified and Unusual Amino Acids			
Abbr.	Amino Acid	Abbr.	Amino Acid
Aad	2-Aminoadipic acid	EtAsn	N-Ethylasparagine
Baad	3-Aminoadipic acid	Hyl	Hydroxylysine
Bala	$\beta$ -alanine, $\beta$ -Amino-propionic acid	AHyl	allo-Hydroxylysine
Abu	2-Aminobutyric acid	3Hyp	3-Hydroxyproline
4Abu	4-Aminobutyric acid, piperidinic acid	4Hyp	4-Hydroxyproline
Acp	6-Aminocaproic acid	Ide	Isodesmosine
Ahe	2-Aminoheptanoic acid	Alle	allo-Isoleucine
Aib	2-Aminoisobutyric acid	MeGly	N-Methylglycine, sarcosine
Baib	3-Aminoisobutyric acid	MeIle	N-Methylisoleucine
Apm	2-Aminopimelic acid	MeLys	6-N-Methyllysine
Dbu	2,4-Diaminobutyric acid	MeVal	N-Methylvaline
Des	Desmosine	Nva	Norvaline
Dpm	2,2'-Diaminopimelic acid	Nle	Norleucine
Dpr	2,3-Diaminopropionic acid	Orn	Ornithine
EtGly	N-Ethylglycine		

In certain embodiments the proteinaceous composition comprises at least one protein, polypeptide or peptide, such as vRNAP or mini-vRNAP. In further embodiments the proteinaceous composition comprises a biocompatible protein, polypeptide or peptide. As used herein, the term "biocompatible" refers to a substance which produces no significant untoward effects when applied to, or administered to, a given organism according to the methods and amounts described herein. Such untoward or undesirable effects are those such as significant toxicity or adverse immunological reactions. In preferred embodiments, biocompatible protein, polypeptide or peptide containing compositions will generally be mammalian proteins or peptides or synthetic proteins or peptides each essentially free from toxins, pathogens and harmful immunogens.

Proteinaceous compositions may be made by any technique known to those of skill in the art, including the expression of proteins, polypeptides or peptides through standard molecular biological techniques, the isolation of proteinaceous compounds from natural sources, or the chemical synthesis of proteinaceous materials. The nucleotide and protein, polypeptide and peptide sequences for various genes have been previously disclosed, and may be found at computerized databases known to those of ordinary skill in the art. One such database is the National Center for Biotechnology Information's Genbank and GenPept databases (<http://www.ncbi.nlm.nih.gov/>). The coding regions for these known genes may be amplified and/or expressed using the techniques disclosed herein or as would be known to those of ordinary skill in the art.

Alternatively, various commercial preparations of proteins, polypeptides and peptides are known to those of skill in the art.

In certain embodiments, a proteinaceous compound may be purified. Generally, "purified" will refer to a specific or desired protein, polypeptide, or peptide composition that has been subjected to fractionation to remove various other proteins, polypeptides, or peptides, and which composition substantially retains its activity, as may be assessed, for example, by the protein assays, as would be known to one of ordinary skill in the art for the specific or desired protein, polypeptide or peptide.

In certain embodiments, the proteinaceous composition may comprise at least one antibody. A mini-vRNAP antibody may comprise all or part of an antibody that specifically recognizes mini-vRNAP. As used herein, the term "antibody" is intended to refer broadly to any immunologic binding agent such as IgG, IgM, IgA, IgD and IgE. Generally, IgG and/or IgM are preferred because they are the most common antibodies in the physiological situation and because they are most easily made in a laboratory setting.

The term "antibody" is used to refer to any antibody-like molecule that has an antigen binding region, and includes antibody fragments such as Fab', Fab, F(ab')<sub>2</sub>, single domain antibodies (DABs), Fv, scFv (single chain Fv), and the like. The techniques for preparing and using various antibody-based constructs and fragments are well known in the art. Means for preparing and characterizing antibodies are also well known in the art (See, e.g., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988; incorporated herein by reference).

It is contemplated that virtually any protein, polypeptide or peptide containing component may be used in the compositions and methods disclosed herein. However, it is preferred that the proteinaceous material is biocompatible. In certain embodiments, it is envisioned that the formation of a more viscous composition will be advantageous in that the high viscosity will allow the composition to be more precisely or easily applied to the tissue and to be maintained in contact with the tissue throughout the procedure. In such cases, the use of a peptide composition, or more preferably, a polypeptide or protein composition, is contemplated. Ranges of viscosity include, but are not limited to, about 40 to about 100 poise. In certain aspects, a viscosity of about 80 to about 100 poise is preferred.

Proteins and peptides suitable for use in this invention may be autologous proteins or peptides, although the invention is clearly not limited to the use of such autologous proteins. As used herein, the term "autologous protein, polypeptide or peptide" refers to a protein, polypeptide or peptide which is derived or obtained from an organism. Organisms that may be used include, but are not limited to, a bovine, a reptilian, an amphibian, a piscine, a rodent, an avian, a canine, a feline, a fungal, a plant, or a prokaryotic organism, with a selected animal or human subject being preferred. The "autologous protein, polypeptide or peptide" may then be used as a component of a composition intended for application to the selected animal or human subject. In certain aspects, the autologous proteins or peptides are prepared, for example from whole plasma of the selected donor. The plasma is placed in tubes and placed in a freezer at about -80° C. for at least about 12 hours and then centrifuged at about 12,000 times g for about 15 minutes to obtain the precipitate. The precipitate, such as fibrinogen may be stored for up to about one year (Oz, 1990).

## VI. Protein Purification

To prepare a composition comprising a vRNAP or mini-vRNAP, it is desirable to purify the components or variants thereof. Purification of the mini-vRNAP (SEQ ID NO:4) can be done in two steps using affinity columns. The mini-vRNAP of SEQ ID NO:6 has been modified to comprise a His tag such that purification can be done in a single step when using metal affinity columns such as those which employ nickel, cobalt or zinc. The full length vRNAP of SEQ ID NO:15 is also His tagged for purification.

According to one embodiment of the present invention, purification of a peptide comprising vRNAP can be utilized ultimately to operatively link this domain with a selective agent. Protein purification techniques are well known to those of skill in the art. These techniques involve, at one level, the crude fractionation of the cellular milieu to polypeptide and non-polypeptide fractions. Having separated the polypeptide from other proteins, the polypeptide of interest may be further purified using chromatographic and electrophoretic techniques to achieve partial or complete purification (or purification to homogeneity). Analytical methods particularly suited to the preparation of a pure peptide are ion-exchange chromatography, exclusion chromatography; polyacrylamide gel electrophoresis; isoelectric focusing. A particularly efficient method of purifying peptides is affinity chromatography.

A tag may be used for protein or peptide purification and detection such as hexahistidine (6-His, HHHHHH), FLAG (DYKDDDDK), hemagglutinin (HA, YPYDVPDYA) and c-myc (EQKLISEEDL). Other tags also have been generated, most of which are very small, comprising only a few amino acids, and are therefore likely to have little to no effect on the conformation of the mature protein or peptide. These small tags do not require any special conformation to be recognized by antibodies. Systems for protein purification using these tags include NTA resin (6-His) or the FLAG fusion system marketed by IBI (FLAG) where the fusion protein is affinity-purified on an antibody column.

Certain aspects of the present invention concern the purification, and in particular embodiments, the substantial purification, of an encoded protein or peptide, such as a vRNAP. The term "purified protein or peptide" as used herein, is intended to refer to a composition, isolatable from other components, wherein the protein or peptide is purified to any degree relative to its naturally-obtainable state. A purified protein or peptide therefore also refers to a protein or peptide, free from the environment in which it may naturally occur.

Generally, "purified" will refer to a protein or peptide composition, such as the vRNAP, that has been subjected to fractionation to remove various other components, and which composition substantially retains its expressed biological activity. Where the term "substantially purified" is used, this designation will refer to a composition in which the protein or peptide forms the major component of the composition, such as constituting about 50%, about 60%, about 70%, about 80%, about 90%, about 95% or more of the proteins in the composition.

Various methods for quantifying the degree of purification of the protein or peptide will be known to those of skill in the art in light of the present disclosure. These include, for example, determining the specific activity of an active fraction, or assessing the amount of polypeptides within a fraction by SDS/PAGE analysis. A preferred method for assessing the purity of a fraction is to calculate the specific activity of the fraction, to compare it to the specific activity of the initial extract, and to thus calculate the degree of purity, herein assessed by a "-fold purification" number. The actual units

used to represent the amount of activity will, of course, be dependent upon the particular assay technique chosen to follow the purification and whether or not the expressed protein or peptide exhibits a detectable activity.

Various techniques suitable for use in protein purification will be well known to those of skill in the art. These include, for example, precipitation with ammonium sulphate, PEG, antibodies and the like or by heat denaturation, followed by centrifugation; chromatography steps such as ion exchange, gel filtration, reverse phase, hydroxylapatite and affinity chromatography; isoelectric focusing; gel electrophoresis; and combinations of such and other techniques. As is generally known in the art, it is believed that the order of conducting the various purification steps may be changed, or that certain steps may be omitted, and still result in a suitable method for the preparation of a substantially purified protein or peptide.

There is no general requirement that the protein or peptide always be provided in their most purified state. Indeed, it is contemplated that less substantially purified products will have utility in certain embodiments. Partial purification may be accomplished by using fewer purification steps in combination, or by utilizing different forms of the same general purification scheme. For example, it is appreciated that a cation-exchange column chromatography performed utilizing an HPLC apparatus will generally result in a greater "-fold" purification than the same technique utilizing a low pressure chromatography system. Methods exhibiting a lower degree of relative purification may have advantages in total recovery of protein product, or in maintaining the activity of an expressed protein.

It is known that the migration of a polypeptide can vary, sometimes significantly, with different conditions of SDS/PAGE (Capaldi et al., 1977). It will therefore be appreciated that under differing electrophoresis conditions, the apparent molecular weights of purified or partially purified expression products may vary.

Ion exchange chromatography is a preferred method of separation. Using column resins such as the metal affinity chromatography resin TALON are also preferred. TALON resin has an enhanced resolving power for polyhistidine-tagged proteins. This results in greater purity with less effort. TALON employs cobalt, an electropositive metal with a remarkably high affinity for polyhistidine-tagged proteins and a low affinity for other proteins. Often, no discernible binding of host proteins occurs and a separate wash step is not required. The binding properties of cobalt allow protein elution under mild pH conditions that protect protein integrity.

Further concentration of the proteins can be done on an anion exchange column, such as the MonoQ column, a high resolution, anion exchange column. This column works at pressures less than 5 MPa, has a high capacity and gives very high chromatographic resolution.

High Performance Liquid Chromatography (HPLC) is characterized by a very rapid separation with extraordinary resolution of peaks. This is achieved by the use of very fine particles and high pressure to maintain an adequate flow rate. Separation can be accomplished in a matter of minutes, or at most an hour. Moreover, only a very small volume of the sample is needed because the particles are so small and close-packed that the void volume is a very small fraction of the bed volume. Also, the concentration of the sample need not be very great because the bands are so narrow that there is very little dilution of the sample.

Gel chromatography, or molecular sieve chromatography, is a special type of partition chromatography that is based on molecular size. The theory behind gel chromatography is that the column, which is prepared with tiny particles of an inert

substance that contain small pores, separates larger molecules from smaller molecules as they pass through or around the pores, depending on their size. As long as the material of which the particles are made does not adsorb the molecules, the sole factor determining rate of flow is the size. Hence, molecules are eluted from the column in decreasing size, so long as the shape is relatively constant. Gel chromatography is unsurpassed for separating molecules of different size because separation is independent of all other factors such as pH, ionic strength, temperature, etc. There also is virtually no adsorption, less zone spreading and the elution volume is related in a simple matter to molecular weight.

Affinity chromatography, a particularly efficient method of purifying peptides, is a chromatographic procedure that relies on the specific affinity between a substance to be isolated and a molecule that it can specifically bind to. This is a receptor-ligand type interaction. The column material is synthesized by covalently coupling one of the binding partners to an insoluble matrix. The column material is then able to specifically adsorb the substance from the solution. Elution occurs by changing the conditions to those in which binding will not occur (e.g., alter pH, ionic strength, and temperature). Tags, as described herein above, can be used in affinity chromatography.

The matrix should be a substance that itself does not adsorb molecules to any significant extent and that has a broad range of chemical, physical and thermal stability. The ligand should be coupled in such a way as to not affect its binding properties. The ligand also should provide relatively tight binding, and it should be possible to elute the substance without destroying the sample or the ligand. One of the most common forms of affinity chromatography is immunoaffinity chromatography. The generation of antibodies that would be suitable for use in accordance with the present invention is discussed below.

An affinity column may have an N4 promoter which the vRNAP or mini-vRNAP proteins recognize attached to a matrix. This column would be suitable for use for the purification of polymerases with no additional tags such as histidine tags.

#### VII. Separation, Quantitation, and Identification Methods

Following synthesis of the RNA, it may be desirable to separate the amplification products of several different lengths from each other and from the template and the excess primer.

##### a. Gel Electrophoresis

In one embodiment, amplification products are separated by agarose, agarose-acrylamide or polyacrylamide gel electrophoresis using standard methods (Sambrook et al., 1989).

##### b. Chromatographic Techniques

Alternatively, chromatographic techniques may be employed to effect separation. There are many kinds of chromatography which may be used in the present invention: adsorption, partition, ion-exchange and molecular sieve, and many specialized techniques for using them including column, paper, thin-layer and gas chromatography (Freifelder, 1982). In yet another alternative, labeled cDNA products, such as biotin-labeled or antigen-labeled, can be captured with beads bearing avidin or antibody, respectively.

##### c. Microfluidic Techniques

Microfluidic techniques include separation on a platform such as microcapillaries, designed by ACLARA BioSciences Inc., or the LabChip™ "liquid integrated circuits" made by Caliper Technologies Inc. These microfluidic platforms require only nanoliter volumes of sample, in contrast to the microliter volumes required by other separation technolo-

gies. Miniaturizing some of the processes involved in genetic analysis has been achieved using microfluidic devices. For example, published PCT Application No. WO 94/05414, to Northrup and White, incorporated herein by reference, reports an integrated micro-PCR™ apparatus for collection and amplification of nucleic acids from a specimen. U.S. Pat. Nos. 5,304,487 to Wilding et al., and 5,296,375 to Kricka et al., discuss devices for collection and analysis of cell containing samples and are incorporated herein by reference. U.S. Pat. No. 5,856,174 describes an apparatus which combines the various processing and analytical operations involved in nucleic acid analysis and is incorporated herein by reference.

##### d. Capillary Electrophoresis

In some embodiments, it may be desirable to provide an additional, or alternative means for analyzing the amplified genes. In these embodiments, micro capillary arrays are contemplated to be used for the analysis.

Microcapillary array electrophoresis generally involves the use of a thin capillary or channel which may or may not be filled with a particular separation medium. Electrophoresis of a sample through the capillary provides a size based separation profile for the sample. The use of microcapillary electrophoresis in size separation of nucleic acids has been reported in, e.g., Woolley and Mathies, 1994. Microcapillary array electrophoresis generally provides a rapid method for size-based sequencing, PCR™ product analysis and restriction fragment sizing. The high surface to volume ratio of these capillaries allows for the application of higher electric fields across the capillary without substantial thermal variation across the capillary, consequently allowing for more rapid separations. Furthermore, when combined with confocal imaging methods, these methods provide sensitivity in the range of attomoles, which is comparable to the sensitivity of radioactive sequencing methods. Microfabrication of microfluidic devices including microcapillary electrophoretic devices has been discussed in detail in, e.g., Jacobsen et al., 1994; Effenhauser et al., 1994; Harrison et al., 1993; Effenhauser et al., 1993; Manz et al., 1992; and U.S. Pat. No. 5,904,824. Typically, these methods comprise photolithographic etching of micron scale channels on a silica, silicon or other crystalline substrate or chip, and can be readily adapted for use in the present invention. In some embodiments, the capillary arrays may be fabricated from the same polymeric materials described for the fabrication of the body of the device, using the injection molding techniques described herein.

Tsuda et al., 1990, describes rectangular capillaries, an alternative to the cylindrical capillary glass tubes. Some advantages of these systems are their efficient heat dissipation due to the large height-to-width ratio and, hence, their high surface-to-volume ratio and their high detection sensitivity for optical on-column detection modes. These flat separation channels have the ability to perform two-dimensional separations, with one force being applied across the separation channel, and with the sample zones detected by the use of a multi-channel array detector.

In many capillary electrophoresis methods, the capillaries, e.g., fused silica capillaries or channels etched, machined or molded into planar substrates, are filled with an appropriate separation/sieving matrix. Typically, a variety of sieving matrices are known in the art may be used in the microcapillary arrays. Examples of such matrices include, e.g., hydroxyethyl cellulose, polyacrylamide, agarose and the like. Generally, the specific gel matrix, running buffers and running conditions are selected to maximize the separation characteristics of the particular application, e.g., the size of the nucleic acid fragments, the required resolution, and the presence of

native or undenatured nucleic acid molecules. For example, running buffers may include denaturants, chaotropic agents such as urea or the like, to denature nucleic acids in the sample.

#### e. Mass Spectrometry

Mass spectrometry provides a means of "weighing" individual molecules by ionizing the molecules in vacuo and making them "fly" by volatilization. Under the influence of combinations of electric and magnetic fields, the ions follow trajectories depending on their individual mass (m) and charge (z). For low molecular weight molecules, mass spectrometry has been part of the routine physical-organic repertoire for analysis and characterization of organic molecules by the determination of the mass of the parent molecular ion. In addition, by arranging collisions of this parent molecular ion with other particles (e.g., argon atoms), the molecular ion is fragmented forming secondary ions by the so-called collision induced dissociation (CID). The fragmentation pattern/pathway very often allows the derivation of detailed structural information. Other applications of mass spectrometric methods known in the art can be found summarized in *Methods in Enzymology*, Vol. 193: "Mass Spectrometry" (J. A. McCloskey, editor), 1990, Academic Press, New York.

Due to the apparent analytical advantages of mass spectrometry in providing high detection sensitivity, accuracy of mass measurements, detailed structural information by CID in conjunction with an MS/MS configuration and speed, as well as on-line data transfer to a computer, there has been considerable interest in the use of mass spectrometry for the structural analysis of nucleic acids. Reviews summarizing this field include K. H. Schram (1990); and P. F. Crain (1990). The biggest hurdle to applying mass spectrometry to nucleic acids is the difficulty of volatilizing these very polar biopolymers. Therefore, "sequencing" had been limited to low molecular weight synthetic oligonucleotides by determining the mass of the parent molecular ion and through this, confirming the already known sequence, or alternatively, confirming the known sequence through the generation of secondary ions (fragment ions) via CID in an MS/MS configuration utilizing, in particular, for the ionization and volatilization, the method of fast atomic bombardment (FAB mass spectrometry) or plasma desorption (PD mass spectrometry). As an example, the application of FAB to the analysis of protected dimeric blocks for chemical synthesis of oligodeoxynucleotides has been described (Koster et al. 1987).

Two ionization/desorption techniques are electrospray/ion spray (ES) and matrix-assisted laser desorption/ionization (MALDI). ES mass spectrometry was introduced by Fenn et al. 1984; WO 90/14148 and its applications are summarized in review articles (R. D. Smith et al. 1990; B. Ardrey, 1992). As a mass analyzer, a quadrupole is most frequently used. The determination of molecular weights in femtomole amounts of sample is very accurate due to the presence of multiple ion peaks, which all could be used for the mass calculation.

MALDI mass spectrometry, in contrast, can be particularly attractive when a time-of-flight (TOF) configuration is used as a mass analyzer. The MALDI-TOF mass spectrometry has been introduced by Hillenkamp et al. (1990). Since, in most cases, no multiple molecular ion peaks are produced with this technique, the mass spectra, in principle, look simpler compared to ES mass spectrometry. DNA molecules up to a molecular weight of 410,000 Daltons could be desorbed and volatilized (Williams et al., 1989). More recently, the use of infra red lasers (IR) in this technique (as opposed to UV-lasers) has been shown to provide mass spectra of larger nucleic acids such as synthetic DNA, restriction enzyme frag-

ments of plasmid DNA, and RNA transcripts up to a size of 2180 nucleotides (Berkenkamp et al., 1998). Berkenkamp et al., 1998, also describe how DNA and RNA samples can be analyzed by limited sample purification using MALDI-TOF IR.

In Japanese Patent No. 59-131909, an instrument is described which detects nucleic acid fragments separated either by electrophoresis, liquid chromatography or high speed gel filtration. Mass spectrometric detection is achieved by incorporating into the nucleic acids atoms which normally do not occur in DNA such as S, Br, I or Ag, Au, Pt, Os, Hg.

#### f. Energy Transfer

Labeling hybridization oligonucleotide probes with fluorescent labels is a well known technique in the art and is a sensitive, nonradioactive method for facilitating detection of probe hybridization. More recently developed detection methods employ the process of fluorescence energy transfer (FET) rather than direct detection of fluorescence intensity for detection of probe hybridization. FET occurs between a donor fluorophore and an acceptor dye (which may or may not be a fluorophore) when the absorption spectrum of one (the acceptor) overlaps the emission spectrum of the other (the donor) and the two dyes are in close proximity. Dyes with these properties are referred to as donor/acceptor dye pairs or energy transfer dye pairs. The excited-state energy of the donor fluorophore is transferred by a resonance dipole-induced dipole interaction to the neighboring acceptor. This results in quenching of donor fluorescence. In some cases, if the acceptor is also a fluorophore, the intensity of its fluorescence may be enhanced. The efficiency of energy transfer is highly dependent on the distance between the donor and acceptor, and equations predicting these relationships have been developed (Forster, 1948). The distance between donor and acceptor dyes at which energy transfer efficiency is 50% is referred to as the Forster distance ( $R_0$ ). Other mechanisms of fluorescence quenching are also known including, for example, charge transfer and collisional quenching.

Energy transfer and other mechanisms which rely on the interaction of two dyes in close proximity to produce quenching are an attractive means for detecting or identifying nucleotide sequences, as such assays may be conducted in homogeneous formats. Homogeneous assay formats are simpler than conventional probe hybridization assays which rely on detection of the fluorescence of a single fluorophore label, as heterogeneous assays generally require additional steps to separate hybridized label from free label. Several formats for FET hybridization assays are reviewed in *Nonisotopic DNA Probe Techniques* (1992).

Homogeneous methods employing energy transfer or other mechanisms of fluorescence quenching for detection of nucleic acid amplification have also been described. Higuchi et al. (1992) disclose methods for detecting DNA amplification in real-time by monitoring increased fluorescence of ethidium bromide as it binds to double-stranded DNA. The sensitivity of this method is limited because binding of the ethidium bromide is not target specific and background amplification products are also detected. Lee, et al. (1993) disclose a real-time detection method in which a doubly-labeled detector probe is cleaved in a target amplification-specific manner during PCR<sup>TM</sup>. The detector probe is hybridized downstream of the amplification primer so that the 5'-3' exonuclease activity of Taq polymerase digests the detector probe, separating two fluorescent dyes which form an energy transfer pair. Fluorescence intensity increases as the probe is cleaved. WO 96/21144 discloses continuous fluorometric assays in which enzyme-mediated cleavage of nucleic acids results in increased fluorescence. Fluorescence energy trans-

fer is suggested for use in the methods, but only in the context of a method employing a single fluorescent label which is quenched by hybridization to the target.

Signal primers or detector probes which hybridize to the target sequence downstream of the hybridization site of the amplification primers have been described for use in detection of nucleic acid amplification (U.S. Pat. No. 5,547,861). The signal primer is extended by the polymerase in a manner similar to extension of the amplification primers. Extension of the amplification primer displaces the extension product of the signal primer in a target amplification-dependent manner, producing a double-stranded secondary amplification product which may be detected as an indication of target amplification. The secondary amplification products generated from signal primers may be detected by means of a variety of labels and reporter groups, restriction sites in the signal primer which are cleaved to produce fragments of a characteristic size, capture groups, and structural features such as triple helices and recognition sites for double-stranded DNA binding proteins.

Many donor/acceptor dye pairs known in the art and may be used in the present invention. These include, for example, fluorescein isothiocyanate (FITC)/tetramethylrhodamine isothiocyanate (TRITC), FITC/Texas Red (Molecular Probes), FITC/N-hydroxysuccinimidyl 1-pyrenebutyrate (PYB), FITC/eosin isothiocyanate (EITC), N-hydroxysuccinimidyl 1-pyrenesulfonate (PYS)/FITC, FITC/Rhodamine X, FITC/tetramethylrhodamine (TAMRA), and others. The selection of a particular donor/acceptor fluorophore pair is not critical. For energy transfer quenching mechanisms, it is only necessary that the emission wavelengths of the donor fluorophore overlap the excitation wavelengths of the acceptor, i.e., there must be sufficient spectral overlap between the two dyes to allow efficient energy transfer, charge transfer or fluorescence quenching. P-(dimethyl aminophenylazo) benzoic acid (DABCYL) is a non-fluorescent acceptor dye which effectively quenches fluorescence from an adjacent fluorophore, e.g., fluorescein or 5-(2'-aminoethyl) aminonaphthalene (EDANS). Any dye pair which produces fluorescence quenching in the detector nucleic acids of the invention are suitable for use in the methods of the invention, regardless of the mechanism by which quenching occurs. Terminal and internal labeling methods are both known in the art and may be routinely used to link the donor and acceptor dyes at their respective sites in the detector nucleic acid.

#### g. In Vitro Studies

The synthesized RNA of the current invention may be used for in vitro studies of spliceosome assembly, splicing reactions, or antisense experiments.

The spliceosome is a large, multisubunit complex consisting of small, nuclear ribonucleoprotein particles (snRNPs). There are a total of 5 snRNAs: U1, U2, U4, U5, and U6 which are small and uridine rich. Each snRNP has 1 or 2 of these RNAs. In addition to catalyzing the splicing reaction, the spliceosome retains intermediate products, positions splice sites for precise joining of the exons, and prevents exons from diffusing away after cleavage and before ligation. Spliceosome catalysis involves concerted cleavage/ligation reactions in which the 2'-OH of branch site A attacks the 5' splice site to form a 2'-5' phosphodiester bond with the first nucleotide of the intron. The resulting 3'-OH at the end of the 5' exon attacks the 3' splice site to release the lariat form of the intron and join the two exons together with a normal 3'-5' phosphodiester bond. At least 50 different proteins are involved in spliceosome assembly and function. In the group I and group II introns, splicing is improved (in velocity and accuracy) by protein factors (Coetze et al., 1994; Mohr et al., 1994).

#### VIII. Kits

Any of the compositions described herein may be comprised in a kit. In a non-limiting example, a vRNAP or more preferably a mini-vRNAP, a derivatized mini-vRNAP, a mutant vRNAP and/or additional agent, may be comprised in a kit. The kits will thus comprise, in suitable container means, a vRNAP, mini-vRNAP, a derivatized mini-vRNAP, a mutant vRNAP and/or an additional agent of the present invention. The inventors envisage other components that may be included in a kit. These include but are not limited to immunodetection agents such as peroxidase and alkaline phosphatase linked monoclonal and polyclonal antibodies, immunoprecipitation reagents such as protein A- or protein G-linked beads, immune cell purification reagents such as a TALON or monoQ column, cloning reagents for the purpose of manipulating an expression vector, and protein expression reagents including prokaryotic and eukaryotic cells lines for the purpose of protein expression.

The kits may comprise a suitably aliquoted vRNAP, mini-vRNAP, a derivatized mini-vRNAP, a mutant vRNAP and/or additional agent compositions of the present invention, whether labeled or unlabeled, as may be used to prepare a standard curve for a detection assay. The components of the kits may be packaged either in aqueous media or in lyophilized form. The container means of the kits will generally include at least one vial, test tube, flask, bottle, syringe or other container means, into which a component may be placed, and preferably, suitably aliquoted. Where there is more than one component in the kit, the kit also will generally contain a second, third or other additional container into which the additional components may be separately placed. However, various combinations of components may be comprised in a vial. The kits of the present invention also will typically include a means for containing the vRNAP, lipid, additional agent, and any other reagent containers in close confinement for commercial sale. Such containers may include injection or blow-molded plastic containers into which the desired vials are retained.

However, the components of the kit may be provided as dried powder(s). When reagents and/or components are provided as a dry powder, the powder can be reconstituted by the addition of a suitable solvent. It is envisioned that the solvent may also be provided in another container means.

The kits of the present invention will also typically include a means for containing the vials in close confinement for commercial sale, such as, e.g., injection and/or blow-molded plastic containers into which the desired vials are retained.

As used herein in the specification, "a" or "an" may mean one or more. As used herein in the claim(s), when used in conjunction with the word "comprising," the words "a" or "an" may mean one or more than one. As used herein "another" may mean at least a second or more.

#### IX. EXAMPLES

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

## Example 1

Identification of a Transcriptionally Active Domain  
of N4 Virion RNA Polymerase

To determine the minimal domain possessing RNA polymerase activity, controlled proteolysis was performed followed by catalytic (transcriptional) autolabeling (Hartmann, et al., 1988). Upon incubation of RNA polymerase with a benzaldehyde derivative of the initiating nucleotide, the benzaldehyde group forms a Schiff-base with the  $\epsilon$ -amino group of lysines located within 12 Å of the nucleotide-binding site. The crosslinking step was performed in the presence of DNA template because it stimulates binding of the initiating nucleotide. The unstable Schiff-base is converted to a stable secondary amine by reduction under mild conditions with sodium borohydride, with concomitant reduction of any non-reacted benzaldehyde derivative. Addition of the next template-directed  $\alpha$ - $^{32}\text{P}$  labeled NTP leads to phosphodiester bond formation and catalytic autolabeling of the transcriptionally active polypeptide. Controlled trypsin proteolysis of vRNAP was performed, followed by catalytic autolabeling and analysis on SDS-PAGE (FIG. 3A). Initially, three proteolytic fragments are generated, of which the smaller two are catalytically active. Upon further incubation with trypsin, a single stable, transcriptionally active product approximately 1,100 amino acids in length remains. N-terminal sequencing of the three initial proteolytic fragments (FIG. 3B) indicated that the stable active polypeptide (mini-vRNAP) corresponds to the middle  $\frac{1}{3}$  of vRNAP, the region containing the three motifs described above (FIG. 2A, SEQ ID NOS:3-4).

## Example 2

## Cloning and Purification of N4 mini-vRNAP

The full-size vRNAP and the mini-vRNAP (SEQ ID NOS:6 and 15) ORFs were cloned under pBAD control with an N-terminal hexahistidine tag (FIG. 4). The mini-vRNAP domain was cloned into the pBAD B expression plasmid, which was purchased from Invitrogen. Five restriction enzyme sites within pBAD B have been altered; the SnaI site was converted to a HpaI site, and the PflMI and EcoRV sites were destroyed, all by site-directed mutagenesis. The BstBI and HindIII sites were destroyed by enzyme digestion followed by Klenow treatment and re-ligation. FIG. 5 (left) shows the relative amounts of full-length and mini-vRNAP proteins purified on TALON columns from the same volume of *E. coli* BL21 induced cells. Cloned mini-vRNAP is expressed at 100-fold higher levels than cloned full size vRNAP. Further concentration on a MonoQ column reveals that, in contrast to full size vRNAP, mini-vRNAP is stable after induction (FIG. 5, right). At least 10 mg of mini-vRNAP at a 20 mg/ml concentration are obtained from 1 L of induced cells in just two purification steps: TALON and MonoQ mini-columns. A non-histagged version of mini-vRNAP has also been cloned (SEQ ID NO:4). In this case, the enzyme is purified from a crude extract of induced cells in two steps: a promoter DNA-affinity column and MonoQ.

Mini-vRNAP possesses a high binding affinity ( $K_d=1$  nM) for N4 promoter-containing DNA oligonucleotides. This property was used for purification of non-his tagged mini-vRNAP (SEQ ID NO:4) on a DNA-affinity column. The column was prepared by adsorbing a 5' biotinylated N4 promoter-containing DNA oligonucleotide onto the matrix of a 1 ml HiTrap Streptavidin column (Pharmacia/Amersham Cat.#17-5112-01) according to the manufacturer's instruc-

tions. A debris-free sonicate of bacterial cells expressing mini-vRNAP was passed through the column. To bind mini vRNAP to the DNA-affinity column, the pH in the extract and binding/washing buffer should be between 5 to 9, and the NaCl concentration should be between 50 mM and 2M. Nucleases in the extract are inhibited by addition of 2 mM EDTA. After washing the column, mini-vRNAP was eluted with warm (25° C.) water; the elution temperature was raised from 4° C. to 25° C. to increase mini-vRNAP recovery. For complete elution, the temperature can be raised up to 43° C. without significant change in the quality of the preparation. Elution under these conditions occurs due to the removal of metal ions and consequent melting of the promoter hairpin and dissociation of mini-vRNAP. Different DNA oligonucleotides containing variants of the P2 promoter (SEQ ID NOS: 16-19), were used in DNA-affinity columns and tested in mini-vRNAP affinity purification. The best yield was achieved using the DNA oligonucleotide of SEQ ID NO:16. However, the DNA oligonucleotides of SEQ ID NOS:19-20 require a lower temperature than the DNA oligonucleotide of SEQ ID NO:16 for complete elution of the protein, in agreement with the lower thermal stability of the respective promoter hairpins.

Up to 1 mg of mini-vRNAP of 90% purity is obtained from a crude extract of 100 ml *E. coli* culture expressing mini-vRNAP in a single purification step using a 1 ml DNA-affinity column. The binding capacity of the DNA-affinity column was not detectably decreased by multiple use.

## Example 3

Effect of EcoSSB on Transcription of  
Single-Stranded Templates

Inventors have previously shown that EcoSSB is required for N4 vRNAP transcription in vivo (Glucksmann, et al, 1992). EcoSSB is unique in that, unlike other SSBs whose effect on vRNAP transcription was tested, it does not melt the promoter hairpin structure (Glucksmann-Kuis, et al., 1996). Recently, inventors have reinvestigated the effect of EcoSSB on vRNAP transcription of single-stranded templates. FIG. 6 shows transcription in the absence and presence of Eco SSB at three different ssDNA template concentrations. The extent of EcoSSB activation is template-concentration dependent, with highest activation at low DNA template concentration. These results suggest that EcoSSB overcomes template limitation on ssDNA templates.

To further explore this hypothesis, the effect of addition of template or EcoSSB to transcription reactions after 20 min incubation in the absence of EcoSSB was tested. The transcription reaction mixtures (5-50  $\mu\text{l}$ ) contained 20 mM Tris-HCl (pH 7.9 at 25° C.), 10 mM  $\text{MgCl}_2$ , 50 mM NaCl, 1 mM dithiothreitol, 0.01-1  $\mu\text{M}$  mini-vRNAP, 1-100 nM ssDNA template (30-100 nt long, synthesized by Integrated DNA Technologies), 1 mM each of 3 non-labeled NTPs, 0.1 mM  $\alpha$ - $^{32}\text{P}$  NTP (1-2 Ci/mmol, NEN), and 1-10  $\mu\text{M}$  *E. coli* SSB. Incubation was for 1 to 80 min at 37° C. at the indicated temperature. In the presence of EcoSSB, RNA synthesis increased linearly throughout the period of incubation (FIG. 7C). In the absence of EcoSSB, no increase in transcription was observed beyond 10 min of incubation (FIG. 7A). Addition of template at 20 min to the reaction carried out in the absence of EcoSSB led to a dramatic increase in RNA synthesis (FIG. 7B). Addition of EcoSSB at 20 min led to a slow rate of transcriptional recovery (FIG. 7D). These results sug-

gest that EcoSSB converts the template from a transcriptionally inactive RNA:DNA hybrid to transcriptionally active single-stranded DNA.

To test this hypothesis, the physical states of the DNA template and the RNA product were analyzed by native gel electrophoresis in the absence and in the presence of EcoSSB. In order to have effective transcription in the absence of EcoSSB, transcription was performed at an intermediate (5 nM) DNA concentration, at which only a 2-fold effect of EcoSSB is observed.

The results of this experiment are shown in FIG. 8. Either  $^{32}\text{P}$ -labeled template (right panel) or labeled NTPs (left panel) were used to analyze the state of the template (right panel) or RNA product (left panel) in the absence or presence of EcoSSB. After transcription, the mixtures were split further into 3 samples: a control sample with no additions, a sample to which RNase H was added to specifically degrade RNA in RNA:DNA hybrids, and a third sample to which Nuclease S1 was added to degrade single-stranded nucleic acids. In the absence of EcoSSB, both the DNA template and the RNA product are in RNA:DNA hybrids, since the RNA product is RNase H sensitive while the DNA-containing bands show altered mobility after RNase H treatment. In the presence of EcoSSB, a significant portion of the RNA product is RNase H resistant and therefore free, although an RNase sensitive band is present that corresponds to an intermediate RNA:DNA:SSB complex. Under these conditions, the DNA is in an SSB:DNA complex. These results indicate that EcoSSB stimulates transcription through template recycling.

To define regions of EcoSSB essential for vRNAP transcription activation on single-stranded templates, the inventors have tested the effect of human mitochondrial SSB (HmtSSB), which shows extensive sequence and structural homology to EcoSSB. The N-terminus of EcoSSB contains DNA binding and tetramerization determinants while the C-terminus is involved in interaction with other replication proteins. Hmt SSB has no effect on vRNAP transcription although it does not melt the promoter hairpin. Interestingly, preliminary results using mutant EcoSSBs and EcoSSB-Hmt SSB chimeras suggest that the C-terminal region of EcoSSB is essential for vRNAP transcriptional activation.

#### Example 4

##### Characterization of mini-vRNAP Transcription Properties

The initiation properties of the full length RNA polymerase and mini-vRNAP were compared at similar molar concentrations (FIG. 9A) using the catalytic autolabeling assay and two reaction conditions: 1- using a template containing +1C, the benzaldehyde derivative of GTP and  $\alpha^{32}\text{P}$ -ATP, or 2- a template containing +1T, the benzaldehyde derivative of ATP and  $\alpha^{32}\text{P}$ -GTP. Comparison of the results in FIGS. 9B and 9C demonstrates that mini-vRNAP exhibits initiation properties similar to full-length vRNAP. In addition, both enzymes discriminate against dATP incorporation to the same extent. Mini-vRNAP does not synthesize abortive products when the first four nucleotides of the transcript are comprised of 50% or more G or C nucleotides.

The elongation and termination properties of both enzymes are compared in FIG. 10. Similar run-off and terminated transcripts are synthesized. Moreover, EcoSSB activates transcription by both enzymes to the same levels. This result indicates that, if there are any sites of specific contact between vRNAP and EcoSSB, they reside in the mini-vRNAP domain.

The sequence of the terminator signals for vRNAP present in the N4 genome include SEQ ID NOS:21-26. The signals of SEQ ID NO:21 and 22 have been tested in vitro on single-stranded templates.

The rate of mini-vRNAP transcription has been compared to the rate of T7 RNA polymerase under the same conditions using the same DNA template. The template used was linearized pET11 containing the original T7 promoter and the N4 vRNAP P2 promoter that was introduced through cloning. The DNA template was denatured before performing transcription using N4 mini-vRNAP. The concentrations of T7 RNAP (Promega, Cat.#P2075) and mini-vRNAP were compared using SDS-PAGE. Transcription reactions contained 50 nM of polymerase, 100 nM of DNA template, 5 $\times$  transcription buffer provided with the T7 RNAP, and 1 mM of each ATP, GTP and CTP and 0.1 mM of [ $^{32}\text{P}$ ]-UTP (1 Ci/mmol). Each reaction mixture was split in two, and *E. coli* SSB was added to one half. The mixtures were incubated at 37 $^{\circ}$  C. and aliquots were taken at different time points. Transcription products were electrophoresed on a 6% sequencing gel and the amount of radioactively-labeled RNA was quantitated by phosphorimaging. The results showed that: (a) transcription of T7 RNAP was not affected by the presence of *E. coli* SSB and (b) N4 mini-vRNAP synthesized 1.5 to 5 fold more RNA in the presence of EcoSSB than T7 RNAP at different time points of incubation.

The optimal temperature for mini-vRNAP transcription is 37 $^{\circ}$  C. It exhibits 70% activity at 30 $^{\circ}$  C., 65% at 45 $^{\circ}$  C., and only 20% at 50 $^{\circ}$  C.

The average error frequency was estimated by determining the misincorporation frequency of each of four [ $^{32}\text{P}$ ]- $\alpha$  NTPs into RNA products using template ssDNAs missing the corresponding template nucleotide in the transcribed region. The following values were obtained:  $\frac{1}{5}\times 10^4$  for misincorporation of G and U using "no C" (SEQ ID NO:10) and "no A" (SEQ ID NO:11) ssDNA templates, respectively;  $\frac{1}{4}\times 10^4$  for misincorporation of C using the "no G" (SEQ ID NO:12) template, and  $\frac{1}{2}\times 10^4$  for misincorporation of A using the "no T" (SEQ ID NO:13) template. For comparison, the average error frequency for T7 RNAP is  $\frac{1}{2}\times 10^4$  (Huang, et al., 2000). Using the method for detection of mispair formation described by Huang, et al. (2000), no misincorporation by mini-vRNAP was detected.

The ability of mini-vRNAP to incorporate derivatized nucleotides was measured. Transcription by mini-vRNAP in the presence of 0.1-1 mM Digoxigenin-11-UTP (cat# 1209256, Roche), Biotin-16-UTP (cat# 1388908, Roche) or underivatized UTP, yielded comparable amounts of product RNA using "control" ssDNA (SEQ ID NO:9) as a transcription template. The product RNAs synthesized in the presence of derivatized UTP have higher molecular mass than those synthesized in the presence of underivatized UTP, and the difference corresponds to the mass difference of the UTPs used. Several derivatives (i.e. 2'Fluoro-ribonucleoside triphosphates, dideoxynucleoside triphosphates) are being tested. The fluorescent analog Fluorescein-12-UTP (Roche catalog #1427857) has been tested using a template which encodes a 51 nucleotide transcript containing a run of 4 Us, and a nucleotide mix containing ATP, CTP, GTP and Fluorescein-12-UTP only. Transcription was only 3% of that achieved with UTP, biotin-6-UTP or digoxigenin-11-UTP under the same reaction conditions. However, incorporation of the fluorescent analog at higher yields is expected to occur in the presence of underivatized UTP or on templates with other sequence compositions.

## Example 5

## Sequence Determinants of mini-vRNAP Promoter Binding

The three N4 early promoters present in the N4 genome contain a pair of Cs separated by 4 nucleotides from the base of the 5 bp promoter stem. In the preferred promoter P2, these 4 bases are As and the Cs are followed by a T. Preferably, mini-vRNAP uses a 17 nucleotide promoter sequence located immediately upstream of the transcription initiation site. Promoters for N4 vRNA polymerase are described by Haynes et al, (1985) and Dai et al., (1998), herein incorporated by reference. vRNAP-promoter recognition and activity require specific sequences and a hairpin structure on the template strand. The vRNAP promoters of SEQ ID NOS:27-29 assume a hairpin structure comprised of a 5-7 bp stem (the inverted repeats are underlined in Table 6) and 3 b purine-containing loop (shown in bold in Table 6). The -11 position corresponds to the center of the loop; +1 indicates the transcription start site.

TABLE 6

Promoter Sequences			
	-11	+1	
P1	3' - <u>CAACCGAAGCGTTG</u> AATACCT-5'		SEQ ID NO: 27
	-11	+1	
P2	3' - <u>TTCTTCGAGGCGAAGAAA</u> ACCT-5'		SEQ ID NO: 28
	-11	+1	
P3	3' - <u>CGACCGAAGCGTCGAAA</u> ACCA-5'		SEQ ID NO: 29

Other possible vRNAP promoters of the current invention include a set of any inverted repeats forming a hairpin with a 2-7 bp long stem and 3-5 b loop having purines in the central and/or next to the central position of the loop.

To study the sequence determinants of promoter binding, 20 base-long promoter oligonucleotides, containing the wild type vRNAP promoter P2 sequence and substituted at every position with a single 5-Iodo-dU, were used. Whenever substitutions were made in the stem, the corresponding pairing base was changed to A. These oligonucleotides were <sup>32</sup>P end-labeled and used to determine the enzyme's affinity for promoter DNAs by a filter binding assay and the ability to crosslink to mini-vRNAP upon UV irradiation at 320 nm. A 20-base oligonucleotide with wild type promoter P2 sequence binds with a 1 nM Kd. Most oligonucleotides showed close to wild type affinity except for the oligonucleotides substituted at positions -11 (at the center of the loop) and -8, indicating that these positions are essential for promoter recognition (FIG. 11). Surprisingly, UV crosslinking was most effective at position -11, in spite of the low binding affinity, indicating a specific contact at this position to mini-vRNAP. Crosslinking was also observed to positions +1, +2 and +3, indicating non-specific contacts with this region of the template, since 5-Iodo-dU substituted oligonucleotides at these positions showed wild type binding affinity.

The effect of changes in the stem length of the hairpin on the ability of mini-vRNAP to bind P2 promoter DNA was analyzed. As shown above, wild type promoter P2 with a 5 bp stem has a Kd of 1 nM (FIG. 12, top). The stem was shortened by removal of 3' bases as shown in FIG. 12 (left). The stem can be shortened by two base pairs without change in the binding affinity. If two or one loop-closing base pairs remain, the binding affinity of templates is still substantial (2-10 nM).

This result, although surprising, is not unexpected since it has been shown that the oligonucleotide 3'd(CGAGGCG)5' forms an unusually stable minihairpin (Yoshizawa, et al., 1997). No binding is observed if one more nucleotide is removed and the loop cannot form. These results indicate that formation of a loop is essential for vRNAP-promoter recognition.

The effect of lengthening the stem by addition of 3' bases is shown in FIG. 12 (right). The stem can be lengthened by two base pairs without change in the binding affinity. On the other hand, base pairing at -2 reduces binding affinity by two orders of magnitude, with a further one order of magnitude reduction caused by base pairing at -1 and +1. These results indicate that single-strandedness of the template at positions -2, -1 and +1 is required for efficient template binding.

All three N4 early promoters present in the N4 genome contain a pair of Cs separated by 4 nucleotides from the base of the 5 bp promoter stem. In promoter P2, these 4 bases are As and the Cs are followed by a T. To identify the determinants of the site of transcription initiation, a series of templates were constructed with a single C placed at different distances from position -11 of the hairpin by addition or deletion of the tract of As present at promoter P2 (FIG. 13). The affinity of mini-vRNAP for these promoters was measured by filter binding and transcription initiation was measured by catalytic autolabeling of mini-vRNAP. All templates showed similar binding affinities. However, only the template with a C positioned 12 bases downstream from the center of the hairpin was able to support transcription initiation. This result indicates that mini-vRNAP utilizes this position as the transcription start site (+1).

## Example 6

## Identification of Sequence Motifs Essential for mini-vRNAP Activity

As shown in FIG. 2A, vRNAP contains the sequence Rx<sub>3</sub>Kx<sub>6</sub>YG, designated Motif B in the Pol I and Pol α DNA polymerases and the T7-like RNA polymerases. To determine the relevance of this motif to vRNAP activity, two mutants K670A and Y678F (SEQ ID NO:8) (position numbers in mini-vRNAP) were constructed by site-specific mutagenesis of mini-vRNAP. These two positions were chosen because, in T7-like RNA polymerases, the lysine is involved in nucleotide binding and the tyrosine in discrimination against deoxynucleoside triphosphates (Maksimova, et al., 1991 Bonner, et al., 1992; Osumi-Davis, et al., 1992). The His-tagged Y678F mini-vRNAP gene (SEQ ID NO:7) differs from that of the mini-vRNAP domain sequence (SEQ ID NO:3) at two positions: nucleotide 2033 (A) was changed to a T, and nucleotide 2034 (T) was changed to a C.

These RNA polymerase mutants were cloned under pBAD control, purified and tested for their ability to bind to wild type promoters. Both mutant polymerases bound to promoter DNA with wild type affinities and crosslinked to 5-Iodo-dU substituted P2 DNA templates at positions -11 and +3 with wild type affinities (FIG. 14), indicating that these mutations do not affect promoter binding.

The mutant enzymes were tested for their ability to support run-off transcription. The wild type enzyme and Y678F enzyme (SEQ ID NO:8) displayed similar activities at both template excess and template-limiting conditions, while the K670A enzyme exhibited decreased activity under both conditions (FIG. 15). Under limiting template conditions, all three enzymes were activated by Eco SSB (right panel). How-

ever, the Y678F enzyme showed reduced discrimination between ribo- and deoxyribonucleoside triphosphates.

The initiation properties of the three enzymes were compared using catalytic autolabeling (FIG. 16). The K670A enzyme displays significantly reduced activity with the GTP derivative. The Y678F enzyme, in contrast to wild type polymerase, incorporates dATP as efficiently as rATP in a single round of phosphodiester bond formation.

Therefore, the behavior of the K670A and Y678F mutant enzymes indicates that Motif B is involved in catalysis, with the lysine probably required for NTP binding and the tyrosine responsible for dNTP discrimination. These results suggest that, despite its lack of extensive sequence similarity, vRNAP is a Class II T7-like RNA polymerase. Results of recent experiments revealed the location of the two carboxylates (aspartates) involved in catalysis.

#### Example 7

##### Development of an In Vivo System Using Mini-vRNAP and N4 vRNAP Promoters for in Vivo Expression of RNAs and Proteins

Plasmid templates were constructed with a reporter gene ( $\alpha$ -peptide of  $\beta$ -galactosidase) cloned under the control of vRNAP promoter P2 present in either of two orientations (FIG. 17B). The reporter construct was generated by cloning a cassette into plasmid pACYC177, which was obtained from New England Biolabs. The cassette contains an approximately 30 bp long fragment originating from pT7Ac (purchased from United States Biochemical), a N4 promoter, and sequence encoding the alpha fragment of lacZ (*lacZ'*). The N4 promoter and *lacZ'* were generated by oligonucleotide annealing and PCR™ amplification, respectively. This cassette replaces the pACYC177 sequence located between the cleavage sites for restriction enzymes ApaI and BamHI. These reporter plasmids and recombinant full-length or mini-vRNAP expressing plasmids were introduced into *E. coli* DH5 $\alpha$  ( $\Delta$ M15), a strain that encodes the  $\beta$ -galactosidase  $\omega$ -peptide. Expression of the reporter gene  $\alpha$ -peptide in this strain results in the synthesis of active  $\beta$ -galactosidase and consequent production of blue colonies on X-gal plates. Transcription of  $\alpha$ -peptide by full-length and mini-vRNAP was assayed on inducing-Xgal media and shown in FIG. 17A. Induction of full-length polymerase results in small colonies with no  $\beta$ -galactosidase activity. This is not surprising since full-length vRNAP is degraded in these cells (FIG. 17C). In contrast, induction of mini-vRNAP led to detectable levels of the protein (FIG. 17C) and to  $\beta$ -galactosidase activity only from the plasmid containing promoter P2 in the proper orientation (FIG. 17A). These results indicate that this system will be suitable for in vivo expression of RNAs and proteins under mini-N4 vRNAP promoter control.

All of the methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are

deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

#### REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- 5 EPA 320,308
- 10 EPA 329,822
- GB 2202,328
- PCT Application No. US87/00880
- PCT Application No. US89/01025
- 15 U.S. Pat. No. 6,218,145
- U.S. Pat. No. 4,682,195
- U.S. Pat. No. 4,683,195
- U.S. Pat. No. 4,683,202
- U.S. Pat. No. 4,684,611
- 20 U.S. Pat. No. 4,797,368
- U.S. Pat. No. 4,800,159
- U.S. Pat. No. 4,883,750
- U.S. Pat. No. 4,946,773
- U.S. Pat. No. 4,952,500
- 25 U.S. Pat. No. 5,082,592
- U.S. Pat. No. 5,139,941
- U.S. Pat. No. 5,279,721
- U.S. Pat. No. 5,302,523
- U.S. Pat. No. 5,322,783
- 30 U.S. Pat. No. 5,354,855
- U.S. Pat. No. 5,384,253
- U.S. Pat. No. 5,464,765
- U.S. Pat. No. 5,538,877
- U.S. Pat. No. 5,538,880
- 35 U.S. Pat. No. 5,550,318
- U.S. Pat. No. 5,563,055
- U.S. Pat. No. 5,580,859
- U.S. Pat. No. 5,589,466
- U.S. Pat. No. 5,591,616
- 40 U.S. Pat. No. 5,610,042
- U.S. Pat. No. 5,656,610
- U.S. Pat. No. 5,670,488
- U.S. Pat. No. 5,672,344
- U.S. Pat. No. 5,693,489
- 45 U.S. Pat. No. 5,702,932
- U.S. Pat. No. 5,736,524
- U.S. Pat. No. 5,780,448
- U.S. Pat. No. 5,789,215
- U.S. Pat. No. 5,869,320
- 50 U.S. Pat. No. 5,945,100
- U.S. Pat. No. 5,981,274
- U.S. Pat. No. 5,994,136
- U.S. Pat. No. 5,994,624
- U.S. Pat. No. 5,610,287
- 55 U.S. Pat. No. 6,013,516
- U.S. Pat. No. 6,046,173
- WO 88/10315
- WO 89/06700
- 60 WO 90/07641
- WO 94/09699
- WO 95/06128
- Abravaya, K. and Rothman-Denes, L. B. (1990) N4 RNA polymerase II sites of transcription initiation. *J. Mol. Biol.* 211: 359-372.
- 65 Angel et al., *Cell*, 49:729, 1987b.
- Angel et al., *Mol. Cell. Biol.*, 7:2256, 1987a.

- Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, 1988
- Archambault, J. and Friesen, J. D. (1993) Genetics of eukaryotic RNA polymerases I, II and III. *Microbiol. Rev.* 57: 703-724.
- Atchison and Perry, *Cell*, 48:121, 1987.
- Baichwal and Sugden, In: *Gene Transfer*, Kucherlapati R, ed., New York, Plenum Press, pp. 117-148, 1986.
- Banerji et al., *Cell*, 27:299, 1981.
- Banerji et al, *Cell*, 35:729, 1983.
- Benvenisty and Neshif, *Proc. Nat. Acad. Sci. USA*, 83:9551-9555, 1986.
- Berkhout et al., *Cell*, 59:273, 1989.
- Berzal-Herranz et al., *Genes and Devel.*, 6:129-134, 1992.
- Blanar et al., *EMBO J.*, 8:1139, 1989.
- Bodine and Ley, *EMBO J.*, 6:2997, 1987.
- Bonner, G., Patra, D., Lafer, E. M., and Sousa R. (1992) Mutations in T7 RNA polymerase that support the proposal for a common polymerase active site structure. *EMBO J.* 11: 3767-3775.
- Boshart et al., *Cell*, 41:521, 1985.
- Bosze et al., *EMBO J.*, 5:1615, 1986.
- Braddock et al., *Cell*, 58:269, 1989.
- Bulla and Siddiqui, *J. Virol.*, 62:1437, 1986.
- Butler and Chamberlin, *J. Biol. Chem.*, 257: 5772-5778, 1982.
- Campbell and Villarreal, *Mol. Cell. Biol.*, 8:1993, 1988.
- Campere and Tilghman, *Genes and Dev.*, 3:537, 1989.
- Campo et al., *Nature*, 303:77, 1983.
- Capaldi et al., *Biochem. Biophys. Res. Comm.*, 76:425, 1977.
- Carter and Flotte, *Ann. N.Y. Acad. Sci.*, 770:79-90, 1995.
- Cech et al., "In vitro splicing of the ribosomal RNA precursor of Tetrahymena: involvement of a guanosine nucleotide in the excision of the intervening sequence," *Cell*, 27:487-496, 1981.
- Celander and Haseltine, *J. Virology*, 61:269, 1987.
- Celander et al., *J. Virology*, 62:1314, 1988.
- Cermakian, N., Ikeda, T. M., Cedergren, R., and Gray, M. W. (1996) Sequences homologous to the yeast mitochondrial and bacteriophage T3 and T7 RNA polymerases are widespread throughout the eukaryotic lineage. *Nuc. Acids Res.* 24: 648-654.
- Chamberlin and Ryan, In: *The Enzymes*. San Diego, Calif., Academic Press, 15: 87-108, 1982
- Chandler et al., *Cell*, 33:489, 1983.
- Chang et al., "Foreign gene delivery and expression in hepatocytes using a hepatitis B virus vector," *Hepatology*, 14:134A, 1991.
- Chang et al., *Mol. Cell. Biol.*, 9:2153, 1989.
- Chase, J. W. and Williams, K. R. (1986) Single-stranded DNA binding proteins required for DNA replication. *Ann. Rev. Biochem.* 55: 130-136.
- Chatterjee et al., *Proc. Natl. Acad. Sci. USA.*, 86:9114, 1989.
- Cheetham, G. M. T. and Steitz, T. A. (2000) Insights into transcription: structure and function of single-subunit DNA-dependent RNA polymerases. *Curr. Op. In Struc. Biol.* 10: 117-123.
- Cheetham, G. M., Jeruzalmi, D. and Steitz, T. A. 1999. Structural basis for initiation of transcription from an RNA polymerase-promoter complex. *Nature* 399: 80-83.
- Chen and Okayama, *Mol. Cell Biol.*, 7:2745-2752, 1987.
- Choi et al., *Cell*, 53:519, 1988.
- Chowrira et al., "In vitro and in vivo comparison of hammerhead, hairpin, and hepatitis delta virus self-processing ribozyme cassettes," *J. Biol. Chem.*, 269:25856-25864, 1994.
- Chowrira et al., *Biochemistry*, 32:1088-1095, 1993.

- Clark, Voulgaropoulou, Fraley, and Johnson, "Cell lines for the production of recombinant adeno-associated virus," *Human Gene Therapy*, 6:1329-1341, 1995.
- Coetze et al., 1994 *Genes & Develop.* 8, 1575.
- 5 Coffin, "Retroviridae and their replication," In: *Virology*, Fields et al. (eds.), New York: Raven Press, pp. 1437-1500, 1990.
- Costa et al., *Mol. Cell. Biol.*, 8:81, 1988.
- Couch et al., "Immunization with types 4 and 7 adenovirus by selective infection of the intestinal tract," *Am. Rev. Resp. Dis.*, 88:394-403, 1963.
- Cramer, P., Bushnell, D. A., and Kornberg, R. D. (2001) Structural basis of transcription: RNA polymerase II at 2.8 Å resolution *Scienceexpress*, www.scienceexpress.org. 19
- 15 April
- Cramer, P., Bushnell, D. A., Fu, J., Gnat, A. L., Maier-Davis, B., Thompson, N. E., Burgess, R. R., Edwards, A. M., David, P. R., and Kornberg, R. D. 2000. *Science* 288: 640-649.
- 20 Cripe et al., *EMBO J.*, 6:3745, 1987.
- Culotta and Hamer, *Mol. Cell. Biol.*, 9:1376, 1989.
- Dai, X and Rothman-Denes, L. B. 1998. Sequence and DNA structural determinants of N4 virion RNA polymerases-promoter recognition. *Genes Develpment*. 12:2782-2790.
- 25 Dandolo et al., *J. Virology*, 47:55, 1983.
- De Villiers et al, *Nature*, 312:242, 1984.
- Delarue, M., Poch, O., Tordo, N., Moras, D., and Argos, P. (1990) An attempt to unify the structure of polymerases. *Protein Engineering*. 3: 461-467.
- 30 Deschamps et al., *Science*, 230:1174, 1985.
- Di Chiara et al., *Trends Pharmacol Sci.*, 13:185, 1992.
- Dubensky et al., *Proc. Nat. Acad. Sci. USA*, 81:7529-7533, 1984.
- Dunn et al., *Nature New Biology*, 230: 94-96, 1971
- 35 Edbrooke et al., *Mol. Cell. Biol.*, 9:1908, 1989.
- Edlund et al., *Science*, 230:912, 1985.
- Elliott, Hynansky, Inturrisi, "Dextromethorphan attenuates and reverses analgesic tolerance to morphine," *Pain*, 59:361-368, 1994.
- 40 Falco, S. C. and Rothman-Denes, L. B. 1979. Bacteriophage N4-induced transcribing activities in *E. coli*: I. Detection and characterization in cell extracts. *Virology* 95: 454-465.
- Falco, S. C. , Vander Laan, K., and Rothman-Denes, L. B. 1977. Virion-associated RNA polymerase required for bacteriophage N4 development. *Proc. Natl. Acad.Sci. (USA)* 74: 520-523.
- Falco, S.C. , Zehring, W.A., and Rothman-Denes, L. B. 1980. DNA-dependent RNA polymerase from bacteriophage N4 virions: purification and characterization. *J. Biol. Chem.* 255: 4339-4347.
- 50 Falco, S. C. , Zivin, R., and Rothman-Denes, L. B. 1978. Novel template requirements of N4 virion RNA polymerase. *Proc. Natl. Acad. Sci. (USA)* 75: 3220-3224.
- Faraldo et al., *J. Bact.*, 174: 7458-7462, 1992
- 55 Fechheimer et al., *Proc. Natl. Acad. Sci. USA*, 84:8463-8467, 1987.
- Feng and Holland, *Nature*, 334:6178, 1988.
- Ferkol et al., *FASEB J.*, 7:1081-1091, 1993.
- Firak and Subramanian, *Mol. Cell. Biol.*, 6:3667, 1986.
- 60 Foecking M K, Hofstetter H. *Gene*. 45(1):101-105, 1986.
- Foley, "Opioid analgesics in clinical pain management. In: *Handbook of Experimental Pharmacology*, Herz, (Ed.), Vol. Vol. 104/II: Opioids II., Springer-Verlag, Berlin, pp. 693-743, 1993.
- 65 Forster and Symons, "Self-cleavage of plus and minus RNAs of a virusoid and a structural model for the active sites," *Cell*, 49:211-220, 1987.

- Fraley et al., Proc. Natl. Acad. Sci. USA, 76:3348-3352, 1979.
- Frohman, In: *PCR Protocols: A Guide To Methods And Applications*, Academic Press, N.Y., 1990.
- Fujita et al., Cell, 49:357, 1987.
- Gerlach et al., "Construction of a plant disease resistance gene from the satellite RNA of tobacco ringspot virus," *Nature (London)*, 328:802-805, 1987.
- Ghosh and Bachhawat, In: *Liver diseases, targeted diagnosis and therapy using specific receptors and ligands*, (Wu G, Wu C ed.), New York: Marcel Dekker, pp. 87-104, 1991.
- Gilles et al., Cell, 33:717, 1983.
- Gloss et al., EMBO J., 6:3735, 1987.
- Glucksmann, M. A., Markiewicz, P., Malone, C., and Rothman-Denes, L. B. 1992.  
Specific sequences and a hairpin structure in the template strand are required for N4 virion RNA polymerase-promoter recognition. Cell 70, 491-500.
- Glucksmann-Kuis, A. M., Dai, X., Markiewicz, P. M. and Rothman-Denes, L. B. 1996. *E. coli* SSB activation of N4 virion RNA polymerase: specific stabilization of an essential DNA hairpin required for promoter recognition. Cell 84, 147-154.
- Godbout et al., Mol. Cell. Biol., 8:1169, 1988.
- Gomez-Flores and Weber, "Differential effects of buprenorphine and morphine on immune and neuroendocrine functions following acute administration in the rat mesencephalon periaqueductal gray," *Immunopharm,m*, 48:145-156, 2000.
- Goodbourn and Maniatis, Proc. Natl. Acad. Sci. USA, 85:1447, 1988.
- Goodbourn et al., Cell, 45:601, 1986.
- Gopal, Mol. Cell Biol., 5:1188-1190, 1985.
- Graham and Prevec, "Adenovirus-based expression vectors and recombinant vaccines," *Biotechnology*, 20:363-390, 1992.
- Graham and Prevec, "Manipulation of adenovirus vectors," In: *Gene Transfer and Expression Protocols*, Murray, E. J., ed., Humana, N.J., vol. 7, 109-128, 1991.
- Graham and Van Der Eb, *Virology*, 52:456-467, 1973.
- Greene et al., *Immunology Today*, 10:272, 1989.
- Gross, C. A., Chan, C., Dombroski, A., Gruber, T., Sharp, M., Tupy, J., and Young, B. (1998) The functional and regulatory roles of sigma factors in transcription. In: *Mechanisms of Transcription*. Cold Spring Harbor Symp. Quant. Biol. 63: 141-156.
- Grosschedl and Baltimore, Cell, 41:885, 1985.
- Guzman, L. M. et al. (1995) *J. Bact.* 177: 4121-4130.
- Hartmann et al., *Nucl. Acids Res.*, 19: 5957-5964, 1991
- Hartmann et al., *Biochem*, 69: 1097-1104, 1987
- Hartmann, G. R., Biebricker, C., Glaser, S. J., Grosse, F., Katzameyer, M., Lindner, A. J., Mosig, H., Nasheuer, H. P., Rothman-Denes, L. B., Schaffner, A. R., Schneider, G., Stetter, K -D., and Thomm, M . 1988. Initiation of transcription—a general tool for affinity labelling of RNA polymerases by autocatalysis. *Biol. Chem. Hoppe-Seyler* 369: 775-788.
- Haseloff and Gerlach, "Simple RNA enzymes with new and highly specific endoribonuclease activities," *Nature*, 334: 585-591, 1988.
- Haslinger and Karin, Proc. Natl. Acad. Sci. USA., 82:8572, 1985.
- Hauber and Cullen, *J. Virology*, 62:673, 1988.
- Hausmann *Current Topics in Microbiology and Immunology*, 75: 77-109, 1976

- Haynes, L. L. and Rothman-Denes, L. B. 1985. N4 virion RNA polymerase sites of transcription initiation. Cell 41: 597-605.
- Hedtke, B., Borner, T., and Weihe, A. (1997) Mitochondrial and chloroplast phage-type RNA polymerases in Arabidopsis. *Science*. 277: 809-811.
- Hen et al., *Nature*, 321:249, 1986.
- Hensel et al., *Lymphokine Res.*, 8:347, 1989.
- Herr and Clarke, Cell, 45:461, 1986.
- Higuchi, et al., *Biotechnology* 10:413-417 1992.
- Hirochikaeta., *J. Virol.*, 61:2599, 1987.
- Hirsch et al., *Mol. Cell. Biol.*, 10:1959, 1990.
- Hochschild, A. and Dove, S. L. (1998) Protein-protein contacts that activate and repress prokaryotic transcription. Cell 92:597-600.
- Holbrook et al., *Virology*, 157:211, 1987.
- Horlick and Benfield, *Mol. Cell. Biol.*, 9:2396, 1989.
- Huang et al., Cell, 27:245, 1981.
- Huang, J., Briebe, L. G., and Sousa, R. (2000) Misincorporation by Wild-Type and Mutant T7 RNA Polymerases: Identification of Interactions That Reduce Misincorporation Rates by Stabilizing the Catalytically Incompetent Open Conformation *Biochemistry*, 38: 11571-11580.
- Hughes, J.; Smith, T. W; Kosterlitz, H.; Fothergill, L.; Morgan, B. and Morris, H. Identification of two related pentapeptides from the brain with potent opiate agonist activity. *Nature* 1975, 258: 577-579.
- Hwang et al., *Mol. Cell. Biol.*, 10:585, 1990.
- Imagawa et al., Cell, 51:251, 1987.
- Imbra and Karin, *Nature*, 323:555, 1986.
- Imler et al., *Mol. Cell. Biol.*, 7:2558, 1987.
- Imperiale and Nevins, *Mol. Cell. Biol.*, 4:875, 1984.
- Jakovovits et al., *Mol. Cell. Biol.*, 8:2555, 1988.
- Jameel and Siddiqui, *Mol. Cell. Biol.*, 6:710, 1986.
- Jaynes et al., *Mol. Cell. Biol.*, 8:62, 1988.
- Johnson et al., *Mol. Cell. Biol.*, 9:3393, 1989.
- Jones and Shenk, "Isolation of deletion and substitution mutants of adenovirus type 5," *Cell*, 13:181-188, 1978.
- Jorgensen et al., *J. Biol. Chem.*, 266: 645-655, 1991.
- Joyce, "RNA evolution and the origins of life," *Nature*, 338: 217-244, 1989.
- Kadesch and Berg, *Mol. Cell. Biol.*, 6:2593, 1986.
- Kaneda et al., *Science*, 243:375-378, 1989.
- Karin et al., *Mol. Cell. Biol.*, 7:606, 1987.
- Kase et al., A potent inhibitor of protein kinase C from microbial origin," *J. Antibiot.*, (8):1059-1065, 1986
- Katinka et al., Cell, 20:393, 1980.
- Katinka et al., *Nature*, 290:720, 1981.
- Kato et al, *J. Biol. Chem.*, 266:3361-3364, 1991.
- Kawamoto et al., *Mol. Cell. Biol.*, 8:267, 1988.
- Kazmierczak, K. M. 2001. PhD thesis. The University of Chicago.
- Kiledjian et al., *Mol. Cell. Biol.*, 8:145, 1988.
- Kim and Cech, "Three dimensional model of the active site of the self-splicing rRNA precursor of Tetrahymena," *Proc. Natl. Acad. Sci. USA*, 84:8788-8792, 1987.
- Klamut et al., *Mol. Cell. Biol.*, 10:193, 1990.
- Klein et al., *Nature*, 327:70-73, 1987.
- Koch et al., *Mol. Cell. Biol.*, 9:303, 1989.
- Korsten et al., *J. Gen. Virol.*, 43: 57-73, 1975
- Kriegler and Botchan, In: *Eukaryotic Viral Vectors*, Y. Gluzman, ed., Cold Spring Harbor: Cold Spring Harbor Laboratory, NY, 1982.
- Kriegler and Botchan, *Mol. Cell. Biol.*, 3:325, 1983.
- Kriegler et al., Cell, 38:483, 1984a.
- Kriegler et al., Cell, 53:45, 1988.

- Kriegler et al., In: *Cancer Cells 2/Oncogenes and Viral Genes*, Van de Woude et al., eds, Cold Spring Harbor, Cold Spring Harbor Laboratory, 1984b.
- Kriegler et al., In: *Gene Expression*, D. Hamer and M. Rosenberg, eds., New York: Alan R. Liss, 1983.
- Kuhl et al., *Cell*, 50:1057, 1987.
- Kunz et al., *Nucl. Acids Res.*, 17:1121, 1989.
- Larsen et al., *Proc. Natl. Acad. Sci. USA.*, 83:8283, 1986.
- Laspia et al., *Cell*, 59:283, 1989.
- Latimer et al., *Mol. Cell. Biol.*, 10:760, 1990.
- Le Gal La Salle et al., "An adenovirus vector for gene transfer into neurons and glia in the brain," *Science*, 259:988-990, 1993.
- Lee et al., *Nature*, 294:228, 1981.
- Lee, Tomasetto, Sager, *Proc. Natl. Acad. Sci. USA*, 88:2825, 1991.
- Levinson et al., *Nature*, 295:79, 1982.
- Lieber and Strauss, "Selection of efficient cleavage sites in target RNAs by using a ribozyme expression library." *Mol. Cell. Biol.*, 15: 540-551, 1995.
- Lin et al., *Mol. Cell. Biol.*, 10:850, 1990.
- Luria et al., *EMBO J.*, 6:3307, 1987.
- Lusky and Botchan, *Proc. Natl. Acad. Sci. USA.*, 83:3609, 1986.
- Lusky et al., *Mol. Cell. Biol.*, 3:1108, 1983.
- Lutz et al., *J. Receptor Res.*, 12:267, 1992.
- Majors and Varmus, *Proc. Natl. Acad. Sci. USA.*, 80:5866, 1983.
- Maksimova, T. G., Mustayev, A. A., Zaychikov, E. F., Lyaikhov, D. L., Tunitskaya, V. L., Akbarov, A. K., Luchin, S. V., Rechinsky, V. O., Chernov, B. K., and Kochetkov, S. N. (1991) Lys631 residue in the active site of the bacteriophage T7 RNA polymerase. Affinity labeling and site-directed mutagenesis. *Eur J Biochem.* 195: 841-847.
- Malone, C., Spellman, S., Hyman, D., and Rothman-Denes, L. B. 1988. Cloning and generation of a genetic map of bacteriophage N4. *Virology* 162: 328-336.
- Mann et al., "Mammalian protein serine/threonine phosphatase 2C: cDNA cloning and comparative analysis of amino acid sequences," *Biochim. Biophys. Acta*, 1130:100-104, 1992.
- Markiewicz, P., Malone, C., Chase, J. W. and Rothman-Denes, L. B. 1992, *E. coli* single-stranded DNA binding (SSB) protein is a supercoiled-template dependent transcriptional activator of N4 virion RNA polymerase. *Genes and Dev.*, 6: 2010-2019.
- Martin and Coleman, *Biochemistry*, 26: 2690-2696, 1987
- McGraw et al., *Nucl. Acid. Res.*, 13: 6753-6766, 1985
- McLaughlin, Collis, Hermonat, and Muzyczka, "Adeno-Associated Virus General Transduction Vectors: Analysis of Proviral Structures," *J. Virol.*, 62:1963-1973, 1988.
- McNeall et al., *Gene*, 76:81, 1989.
- Michel and Westhof, "Modeling of the three-dimensional architecture of group I catalytic introns based on comparative sequence analysis," *J. Mol. Biol.*, 216:585-610, 1990.
- Miksicek et al., *Cell*, 46:203, 1986.
- Miller, A. A., Wood, D., Ebright, R. E., and Rothman-Denes, L. B. (1997) RNA polymerase  $\beta'$  subunit: a target of DNA binding-independent activation. *Science*. 275: 1655-1657.
- Miller, *Curr. Top. Microbiol. Immunol.*, 158:1, 1992.
- Miyomoto and Takemore, "Inhibition of naloxone-precipitated withdrawal jumping by i.c.v. and i.t. administration of saline in morphine-dependent mice," *Life Sci.*, 52(13): 1129-1134, 1993b.
- Miyomoto and Takemore, "Relative involvement of supraspinal and spinal mu opioid receptors in morphine dependence in mice," *Life Sci.*, 52(12):1039-1044, 1993a.

- Mohr et al., 1994 *Nature* 370, 147.
- Mordacq and Linzer, *Genes and Dev.*, 3:760, 1989.
- Moreau et al., *Nucl. Acids Res.*, 9:6047, 1981.
- Muesing et al., *Cell*, 48:691, 1987.
- 5 Narita, Nartia, Mizoguchi, Tseng, "Inhibition of Protein Kinase C, but not of Protein Kinase A, blocks the development of acute antinociceptive tolerance to an intrathecally administered  $\mu$ -opioid receptor agonist in the mouse," *European Pharmacology*, 280:R1-R3, 1995.
- 10 Ng et al., *Nuc. Acids Res.*, 17:601, 1989.
- Nicolas and Rubinstein, "Retroviral vectors," In: *Vectors: A survey of molecular cloning vectors and their uses*, Rodriguez and Denhardt (eds.), Stoneham: Butterworth, 494-513, 1988.
- 15 Nicolau and Sene, *Biochim. Biophys. Acta*, 721:185-190, 1982.
- Nicolau et al., *Methods Enzymol.*, 149:157-176, 1987.
- Olson et al., *Peptides*, 10:1253, 1988.
- Ondek et al., *EMBO J.*, 6:1017, 1987.
- 20 Ornitz et al., *Mol. Cell. Biol.*, 7:3466, 1987.
- Osumi-Davis, P. A., de Aguilera, M. C., Woody, R. W., and Woody, A. Y. (1992) Asp537, Asp812 are essential and Lys631, His811 are catalytically significant in bacteriophage T7 RNA polymerase activity. *J Mol Biol.*226:37-45.
- 25 Palmiter et al., *Nature*, 300:611, 1982.
- Palukaitis et al., "Characterization of a viroid associated with avocado sunblotch disease," *Virology*, 99:145-151, 1979.
- Paule, M. and White, R. J. (2000) Transcription by RNA polymerases I and III. *Nuc. Acids Res.* 28:1283-1298.
- 30 Pech et al., *Mol. Cell. Biol.*, 9:396, 1989.
- Perales et al., *Proc. Natl. Acad. Sci.* 91:4086-4090, 1994.
- Perez-Stable and Constantini, *Mol. Cell. Biol.*, 10:1116, 1990.
- 35 Perriman et al., "Extended target-site specificity for a hammerhead ribozyme," *Gene*, 113: 157-163, 1992.
- Pert and Snyder, "Opiate receptor; demonstration in nervous tissue," *Science*, 179:1011-1014, 1973.
- Picard and Schaffner, *Nature*, 307:83, 1984.
- 40 Pick, Roques, Gacel, Pasternak, "Supraspinal  $\mu_2$  receptors mediate spinal/supraspinal morphine synergy," *Eur. J Pharmacol.*, 220:275-277, 1992a.
- Pinkert et al., *Genes and Dev.*, 1:268, 1987.
- Ponta et al., *Proc. Natl. Acad. Sci. USA.*, 82: 1020, 1985.
- 45 Porton et al., *Mol. Cell. Biol.*, 10:1076, 1990.
- Potter et al., *Proc. Nat. Acad. Sci. USA*, 81:7161-7165, 1984.
- Prody et al., "Autolytic processing of dimeric plant virus satellite RNA." *Science*, 231:1577-1580, 1986.
- Puhler, G., Leffers, H., Gropp, F., Palm, P., Klenk, H. -P., Lottspeich, F., Garrett, R. A., and Zillig, W. (1989) Archaeobacterial DNA-dependent RNA polymerases testify to the evolution of the eukaryotic nuclear genome. *Proc. Natl. Acad. Sci. USA.* 86: 4569-4573
- Queen and Baltimore, *Cell*, 35:741, 1983.
- 55 Quinn et al., *Mol. Cell. Biol.*, 9:4713, 1989.
- Radler et al., *Science*, 275:810-814, 1997.
- Rashtchian and Mackey, "Labeling and Detection of Nucleic Acids," in "Nonradioactive Labeling and Detection of Biomolecules," C. Kessler, Ed., Springer-Verlag, New York, 1992, pp. 70-84
- 60 Record, M. T., Reznikoff, W. S., and Schlaw, P. J. (1995) *E. coli* RNA polymerase (E-sigma70), promoters and the kinetics of the steps of transcription initiation, In *E. coli and Salmonella typhimurium: Cell and Molecular Biology* 1:792-821. F. Neidhardt ed. ASM
- Redondo et al., *Science*, 247:1225, 1990.

- Reinhold-Hurek and Shub, "Self-splicing introns in tRNA genes of widely divergent bacteria," *Nature*, 357:173-176, 1992.
- Reisman and Rotter, *Mol. Cell. Biol.*, 9:3571, 1989.
- Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990.
- Resendez Jr. et al., *Mol. Cell. Biol.*, 8:4579, 1988.
- Ripe et al., *Mol. Cell. Biol.*, 9:2224, 1989.
- Rippe et al., *Mol. Cell Biol.*, 10:689-695, 1990.
- Ritling et al., *Nucl. Acids Res.*, 17:1619, 1989.
- Roeder, R. (1996) The role of general transcription factors in transcription by RNA polymerase II. *Trends Biochem. Sci.* 21: 327-335.
- Rong, M., He, B., McAllister, W. T. and Durbin, R. K. (1998) Promoter specificity determinants of T7 RNA polymerase *Proc. Natl. Acad. Sci. USA* 95: 515-519.
- Rosen et al., *Cell*, 41:813, 1988.
- Rossi, Pasternak, Bodnar, "Synergistic brainstem interactions for morphine analgesia," *Brain Res.*, 624:171-180, 1993.
- Rothman-Denes, L. B., Dai, X., Davydova, E., Carter, R., and Kazmierczak, K. 1999. Transcriptional Regulation by DNA Structural Transitions and Single-Stranded DNA Binding Proteins. 63rd Cold Spring Harbor Symp. Quant. Biol. 63: 63-73.
- Roux et al., "A versatile and potentially general approach to the targeting of specific cell types by retroviruses: Application to the infection of human cells by means of major histocompatibility complex class I and class II antigens by mouse ecotropic murine leukemia virus-derived viruses," *Proc. Natl. Acad. Sci. USA*, 86:9079-9083, 1989.
- Running, J. A. et al., *BioTechniques* 8:276-277, 1990.
- Sakai et al., *Genes and Dev.*, 2:1144, 1988.
- Saldanha et al., 1993 *FASEB J.* 7, 15.
- Sambrook, Fritsch, Maniatis, In: *Molecular Cloning: A Laboratory Manual* 2 rev.ed., Cold Spring Harbor: Cold Spring Harbor Laboratory Press, 1989.
- Sanders, G. M., Kassavetis, G. A., and Geiduschek, E. P. (1997) Dual targets of a transcriptional activator that tracks on DNA. *EMBO Journal*. 16: 3124-3132.
- Sarver et al., "Ribozymes as potential anti-HIV-1 therapeutic agents," *Science*, 247:1222-1225, 1990.
- Satake et al, *J. Virology*, 62:970, 1988.
- Scanlon et al., "Ribozyme-mediated cleavages of c-fos mRNA reduce gene expression of DNA synthesis enzymes and metallothionein," *Proc Natl Acad Sci USA*, 88:10591-10595, 1991.
- Schaffner et al., *J. Mol. Biol.*, 201:81, 1988.
- Searle et al., *Mol. Cell. Biol.*, 5:1480, 1985.
- Shadel, G. S. and Clayton, D. A. (1993) Mitochondrial transcription initiation, variation and conservation. *J. Biol. Chem.* 268: 16083-16086.
- Sharma, Klee, Nirenberg, "Opiate-dependent modulation of adenylate cyclase," *Proc. Natl. Acad. Sci. U.S.A.*, 74:3365-3369, 1977.
- Sharp and Marciniak, *Cell*, 59:229, 1989.
- Sharp and Yaksh, "Pain killers of the immune system," *Nature Medicine*, 3:831-832, 1997.
- Shaul and Ben-Levy, *EMBO J.*, 6:1913, 1987.
- Shelling and Smith, "Targeted integration of transfected and infected adeno-associated virus vectors containing the neomycin resistance gene," *Gene Therapy*, 1:165-169, 1994.
- Sherman et al., *Mol. Cell. Biol.*, 9:50, 1989.
- Simon, *Medicinal Res. Rev.*, 11:357, 1991.
- Sioud et al., "Preformed ribozyme destroys tumour necrosis factor mRNA in human cells," *J Mol. Biol.*, 223:831-835, 1992.

- Sleigh and Lockett, *J. EMBO*, 4:3831, 1985.
- Sousa, R. (1996) Structural and mechanistic relationships between nucleic acid polymerases. *Trends in Biochem. Sci.* 21: 186-190.
- Sousa, R., Chung, Y., Rose, J. P., and Wang, B. -C. (1993) Crystal structure of bacteriophage T7 RNA polymerase. *Nature*. 364: 593-599.
- Spalholz et al., *Cell*, 42:183, 1985.
- Spandau and Lee, *J. Virology*, 62:427, 1988.
- Spandidos and Wilkie, *EMBO J.*, 2:1193, 1983.
- Stephens and Hentschel, *Biochem. J.*, 248:1, 1987.
- Stuart et al., *Nature*, 317:828, 1985.
- Sullivan and Peterlin, *Mol. Cell. Biol.*, 7:3315, 1987.
- Swartzendruber and Lehman, *J. Cell. Physiology*, 85:179, 1975.
- Sweetser, D., Nonet, M., and Young, R. A. (1987) Prokaryotic and eukaryotic RNA polymerases have homologous core subunits. *Proc. Natl. Acad. Sci. USA*. 84: 1192-1196.
- Symons, "Avocado sunblotch viroid: primary sequence and proposed secondary structure." *Nucl. Acids Res.*, 9:6527-6537, 1981.
- Symons, "Small catalytic RNAs." *Annu. Rev. Biochem.*, 61:641-671, 1992.
- Takebe et al., *Mol. Cell. Biol.*, 8:466, 1988.
- Tavernier et al., *Nature*, 301:634, 1983.
- Taylor and Kingston, *Mol. Cell. Biol.*, 10: 165, 1990a.
- Taylor and Kingston, *Mol. Cell. Biol.*, 10: 176, 1990b.
- Taylor et al., *J. Biol. Chem.*, 264:15160, 1989.
- Thiesen et al., *J. Virology*, 62:614, 1988.
- Thompson et al., "Ribozymes in gene therapy." *Nature Medicine*, 1:277-278, 1995.
- Top et al., "Immunization with live types 7 and 4 adenovirus vaccines. II. Antibody response and protective effect against acute respiratory disease due to adenovirus type 7," *J. Infect. Dis.*, 124:155-160, 1971.
- Towle et al., *J. Biol. Chem.*, 250:1723-1733, 1975
- Tronche et al., *Mol. Biol. Med.*, 7:173, 1990.
- Tronche et al., *Mol. Cell. Biol.*, 9:4759, 1989.
- Trudel and Constantini, *Genes and Dev.*, 6:954, 1987.
- Tur-Kaspaetal., *Mol. Cell Biol.*, 6:716-718, 1986.
- Tyndall et al., *Nuc. Acids. Res.*, 9:6231, 1981.
- Ueda, Miyamae, Hayashi, Watanabe, Fukushima, Sasaki, Iwamura, Misu, "Protein kinase C involvement in homologous desensitization of  $\alpha$ -opioid receptor coupled to G<sub>i</sub> phospholipase C activation in xenopus oocytes," *Journal of Neuroscience*, 15:7485-7499, 1995.
- Vannice and Levinson, *J. Virology*, 62:1305, 1988.
- Vasseur et al., *Proc. Natl. Acad. Sci. USA.*, 77:1068, 1980.
- Wagner et al., *Proc. Natl. Acad. Sci.* 87(9):3410-3414, 1990.
- Wagner et a., *Science*, 260:1510-1513, 1993.
- Walsh, Nienhuis, Samulski, Brown, Miller, Young, and Liu, "Phenotypic correction of Fanconi anemia in human hematopoietic cells with a recombinant adeno-associated virus vector," *J. Clin. Invest.* 94:1440-1448, 1994.
- Wang and Calame, *Cell*, 47:241, 1986.
- Wang, Johnson, Persico, Hawkins, Griffin, Uhl, "Human  $\alpha$  opiate receptor: cDNA and genomic clones, pharmacologic characterization and chromosomal assignment," *FEBS Letters*, 338: 217-222, 1994.
- Watt et al., *Proc. Natl Acad. Sci.*, 83(2): 3166-3170, 1986.
- Weber et al., *Cell*, 36:983, 1984.
- Weinberger et al., *Mol. Cell. Biol.*, 8:988, 1984.
- Wendt et al., *Eur. J. Biochem.*, 191: 467-472, 1990
- Whistler, J. and vonZastrow, M. (1998) *Proc. Natl. Acad. Sci.* 95, 9914-9.
- Winoto and Baltimore, *Cell*, 59:649, 1989.
- Wong et al., *Gene*, 10:87-94, 1980.

- Wu and Wu, *Adv. Drug Delivery Rev.*, 12:159-167, 1993.  
 Wu and Wu, *Biochem.*, 27:887-892, 1988.  
 Wu and Wu, *J. Biol. Chem.*, 262:4429-4432, 1987.  
 Yang et al., *Proc. Natl. Acad. Sci USA*, 87:9568-9572, 1990.  
 Yang, Chen, Trempe, "Characterization of cell lines that  
 inducibly express the adeno-associated virus Rep pro- 5  
 teins," *J. Virol.*, 68:4847-4856, 1994.  
 Yoshizawa, S., Kawai, G., Watanabe, K., Miura, K., and  
 Hirao, I. (1997) GNA trinucleotide loop sequences produc- 10  
 ing extraordinarily stable DNA mini hairpins. *Biochemis-  
 try* 36, 4761-4767.  
 Yuan and Altman, "Selection of guide sequences that direct  
 efficient cleavage of mRNA by human ribonuclease P,"  
*Science*, 263:1269-1273, 1994.  
 Yuan et al., "Targeted cleavage of mRNA by human RNase P," 15  
*Proc. Natl. Acad. Sci. USA*, 89:8006-8010, 1992.

- Yutzey et al., *Mol. Cell. Biol.*, 9:1397, 1989.  
 Zehring, W. A. and Rothman-Denes, L. B. (1983) Purification  
 and characterization of coliphage N4 RNA polymerase II  
 activity from infected cell extracts. *J. Biol. Chem.* 258:  
 8074-8080.  
 Zhang, G., Campbell, E. A., Minakhin, L., Richter, C., Sev-  
 erinov, K. and Darst, S. A. (1999) Crystal structure of  
 Thermus aquaticus core RNA polymerase at 3.3 Å resolu-  
 tion. *Cell* 98: 811-824.  
 Zhou and Doetsch, *Proc. Nat. Acad. Sci. USA*, 90: 6601-  
 6605, 1993  
 Zivin, R., Zehring, W. A., and Rothman-Denes, L. B. 1981.  
 Transcriptional map of bacteriophage N4: location and  
 polarity of N4 RNAs. *J. Mol. Biol.* 152: 335-356.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 29

<210> SEQ ID NO 1

<211> LENGTH: 10506

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 1

```

atgtcagtat ttgatagact ggctgggttc gcagacagcg taaccaatgc aaagcaagtt      60
gacgtctcta ctgcaaccgc ccagaagaaa gctgaacaag gtgtcactac tctctttggt      120
tctcctgatg ctgcttatca aatgcaagct gcccgactg gtaatgttg ggctaattgca      180
tttgaaccag ggacagtgc atcagatttc atgaatctga cccaatgca aatcatgaat      240
aagtatgggg ttgagcaagg cttacaactt atcaatgctc gtgctgatgc agggaaaccag      300
gtattcaatg attcagttac tacaagaact cctggggaag aactggggga tattgctact      360
gggtgtggcc ttggttttgt taataccctt gggggcattg gtgctcttgg ggcaggctta      420
ctcaacgatg atgcaggctc tgtgtttgct caacaattga gtaagtttaa tgatgctgtt      480
catgctaccc aaagccaggc attacaagat aaacgtaagc tctttgctgc tcgtaactta      540
atgaatgaag tagagagtga acgtcagtat caaacagata agaaagaagg cactaatgac      600
atagtagctt ccttatctaa atttggacgt gattttgtag gttcaattga gaatgctgct      660
caaaactgact ctattatttc tgatgggtta gcagaagggg taggttctct attaggtgct      720
ggctctgtat taagggtgct atctttactg ggtaaagcag ttgttccagc aaatactctt      780
cgtagtctg cttggctggt tgctattgat gcaggactg gtactcagtc actggctcgt      840
attgcctcta ctgtaggtag agctgcaccg ggtatggttg gtgttggtgc aatggaagct      900
gggtgtgcat accaacaaac tgctgatgaa attatgaaga tgagtcttaa agacttagag      960
aagtctcctg tttatcagca acatattaaa gatggtatgt cccctgaaca ggctcgtcgt      1020
cagactgcat ctgaaactgg tcttactgct gctgctattc aattacctat tgctgctgca      1080
accggtcctc tggtatcccc ttttgagatg gctcctttcc gtgctggctc tttagggtgct      1140
gtaggtatga accttgcccc tgaaacagtg gaagaaggtg ttcagggtgc tacaggccaa      1200
ctggctcaga atattgcaca gcaacaaaac attgataaga accaagacct gcttaaggt      1260
gtcggtacac aggctggttt aggtgctctt tatggctttg gttctgctgg tgtgttacag      1320

```

-continued

---

gctccggctg	gtgctgctcg	tttagcaggt	gctgcaactg	ctcctgtatt	gcgtaccaca	1380
atggctggty	ttaaagctgc	tggtagtgtg	gcaggtaagg	ttgtttctcc	tattaagaat	1440
actttagtag	ctcgtgggta	acgggttatg	aagcagaatg	aagaagcadc	tcctggtgct	1500
gatgactatg	ttgcacaggc	agcacaagaa	gctatggctc	aagcaccaga	agcagaagtt	1560
actattcgtg	atgctgttga	agcaactgat	gctactccag	aacagaaagt	tgacgacac	1620
cagtatgttt	ctgacttaac	gaatgctact	cgttttaatc	ctgaaaatta	tcaggaagca	1680
ccagagcata	ttcgtaatgc	tgtagctggg	tctactgacc	aagtacaggt	tattcagaag	1740
tttagcagact	tagttaaac	attagatgaa	tctaactctc	aagcactgat	ggaagctgca	1800
tcttatatgt	atgatgctgt	ttcagagttt	gagcagttca	ttaaccgtga	ccctgctgca	1860
ctggatagca	ttcctaagaa	ttctccggct	attgagttac	tcaaccgtta	tacgaatctg	1920
acagctaata	ttcagaacac	accaaagta	attggtgcac	tgaatgttat	taatcgaatg	1980
attaatgaat	ctgctcagaa	tggttctttg	aatgtgactg	aagaatccag	tcacacaggaa	2040
atgcagaacg	tagcattagc	tgctgaagta	gccctgaaa	agctcaatcc	agagtctgta	2100
aatgttggtc	ttaaacatgc	tgctgatggg	cgtattaaac	tgaataatcg	ccagattgct	2160
gccctccaga	atgctgctgc	aatcctgaag	ggggcacggg	aatatgatgc	agaagctgcc	2220
cgtcttggat	tacgtctca	agacattgtg	agtaaacaga	ttaaacggga	tgagagcaga	2280
actcaggaag	gacaatactc	tgctgtgcaa	catgcgaata	ggattcggtc	tgctgataac	2340
tctggaatt	tcgagttggc	ctccgcttac	ctgaacgact	ttatgcagtt	cgcccagcac	2400
atgcagaata	aggttggagc	gttgaatgag	catcttgta	cggggaatgc	ggataagaat	2460
aagtctgtcc	actaccaagc	tcttactgct	gacagagaat	gggttcgtag	ccgtaccgga	2520
ttgggggtca	atccctatga	cactaagtcg	gttaaatttg	cccagcaagt	tgctctgaa	2580
gcgaaaacgg	tagcggatata	tgctaatagcc	ctcgtctcgg	cttaccggga	actgaaggtc	2640
agtcataata	aagttactcc	attggattca	cgtcttaacg	ctcctgctgc	tgaggtggtc	2700
aaggcattcc	gtcaaggcaa	tcgagacggt	gcttcttctc	aaccgaaagc	tgactccgtg	2760
aatcagggtta	aagaaactcc	tgttacaaaa	caggaaccag	ttacatctac	tgtacagact	2820
aagactcctg	ttagtgaatc	tgttaaaaca	gaacctacta	ctaaagagtc	tagcccacag	2880
gctataaaaag	aacctgtgaa	ccagctgtaa	aaacaggatg	ttaaccttac	taatgaggac	2940
aacatcaagc	aacctactga	atctgttaaa	gaaactgaaa	cttctacaaa	agaaagtaca	3000
gttacagaag	aattaaaga	aggatttgat	gctgtttacc	cttcattggt	aggtaactgct	3060
gattctaaag	cagaggggat	taagaactat	ttcaaattgt	cctttacctt	accagaagaa	3120
cagaaatccc	gtactgttgg	ttcagaagca	cctctaaaag	atgtagcca	agctctgtct	3180
tctcgtgctc	gttatgaact	ctttactgag	aaagaaactg	ctaaccctgc	ttttaatggg	3240
gaagttatta	agcgatacaa	agaactcatg	gaacatgggg	aaggattgct	tgatattctt	3300
cgtcccgtc	tggtcaagtt	ccttaacact	aaggatgttg	gtaaacgttt	tgctcaaggt	3360
acagaagcca	accgttgggt	agggtgtaag	ttacttaaca	ttgttgagca	ggatggggat	3420
acctttaagt	acaacgaaca	attgctacag	actgctgtat	tagcaggtct	tcaatggaga	3480
cttactgcta	ccagcaatac	tgctatcaaa	gatgcaaaag	atgttctgctc	tattactggt	3540
attgaccaag	ctctgctgcc	agaaggttta	gtagagcaat	ttgatactgg	tatgacactc	3600
actgaagcag	ttagttccct	ggctcagaaa	attgagctct	actggggatt	atctcgtaat	3660

-continued

---

ccaaatgctc cattgggcta taccaaaggc atccctacag caatggctgc tgaattctg	3720
gctgcatttg tagagtctac tgatgttgta gagaacatcg tggatgatgc agaaattgac	3780
ccagataaca agaagactat tggctctgtac accattactg aactggattc cttcgacca	3840
attaatagct tccctactgc tattgaagaa gctgttttag tgaatcctac agagaagatg	3900
ttctttggty atgacattcc tcctgtagct aatactcagc ttcgtaacct tgctgttcgt	3960
aatactccag aacagaaggc tgcattgaaa gcagagcagg ctacagagtt ctatgtacac	4020
accccaatgg ttcaattcta tgagacgtta ggtaaagacc gtattctcga actgatgggt	4080
gctggtaactc tgaataaaga gttacttaat gataaccatg ctaaatctct ggaaggttaag	4140
aaccgttcag tagaggactc ttacaaccaa ctgttctccg tcattgagca ggtaagagca	4200
cagagcgaag acatctctac tgtacctatt cactatgcat acaatgatgac ccgtgttggt	4260
cgtagcaga tgttaggtaa atacaatcct caatcagcca aactggttcg tgaggccatc	4320
ttacctacta aagctacttt ggatttatcg aaccagaaca atgaagactt ctctgcattc	4380
cagttaggtc tggctcagcg attggacatt aaagtccata ctatgactcg tgaggttatg	4440
tctgacgagt tgactaaatt actggaaggt aatctgaac cagccattga tatgatggtt	4500
gagtttaata cactgggttc cttaccagaa aacgcagttg atgttctgaa tacagatta	4560
ggagatagga agtcattcgt agcattgatg gctcttatgg agtattcccg ttacttagta	4620
gcagaggata aatctgcatt tgtaactcca ctgtatgtag aagcagatgg tgttactaat	4680
ggtccaatca atgccatgat gctaatagaca ggccgtctgt ttactcctga ctggattcgt	4740
aatattgcca aagggggctt gttcattggt tctccaaata agaccatgaa tgagcatcgc	4800
tctactgctg acaataatga tttatatcaa gcatccacta atgctttgat ggaatcgttg	4860
ggtaagttac gtagtaacta tgccctctaat atgctattc agtctcagat agacagtctt	4920
ctttctctga tggatttgtt tttaccggat attaactctg gtgagaatgg tgctttagaa	4980
cttaaacgtg gtattgctaa gaaccactg actattacca tctatggttc tggtgctcgt	5040
ggtattgcag gtaagctggt tagttctggt actgatgcca tctatgagcg tatgtctgat	5100
gtactgaaag ctctgtctaa agaccctaat atctctgctg ctatggcaat gtttggttaag	5160
caagctgctt cagaagcaca tgctgaagaa cttcttgccc gtttcctgaa agatatggaa	5220
acactgactt ctactgttcc tgttaaactg aaaggtgtac tggaaactaca atccacaggt	5280
acaggagcca aaggaataat caatcctaag acctatacca ttaagggcga gcaactgaag	5340
gcacttcagg aaaatagct gcacttcttt gtagaaccac tacgtaatgg tattactcag	5400
actgtaggtg aaagtctggt gtactctact gaacaattac agaaagctac tcagattcaa	5460
tctgtagtgc tggaaagatat gttcaaacag cgagtacaag agaagctggc agagaaggct	5520
aaagacccaa catggaagaa aggtgatttc cttactcaga aagaactgaa tgatattcag	5580
gcttctctga ataacttagc ccctatgatt gagactggt ctcagacttt ctacattgct	5640
ggttcagaaa atgcagaagt agcaaatcag gtattagcta ctaacctga tgaccgtatg	5700
cggtgaccaa tgagtatcta tgctccagca caggccggty tagcaggtat tccatttatg	5760
actattggta ctgggtgatgg catgatgatg caaactcttt ccactatgaa aggtgcacca	5820
aagaataccc tcaaaatctt tgatggatg aacattggtt tgaatgacat cactgatgcc	5880
agtcgtaaag ctaatgaagc tgtttacact tcttggcagg gtaacctat taagaatggt	5940
tatgaatcat atgctaagtt catgaagaat gtatatttca gcaagctgct cctgaagca	6000
ttggaagcaa ttggtaaatc tgctctgtaa tatgaccaac gtgagaatgc tactgtatg	6060

-continued

---

gatattgcta	acgctgcac	tctgattgaa	cgtaacttac	gtaatattgc	actgggtgta	6120
gatattcgtc	ataaggtgct	ggataaggta	aatctgtcca	ttgaccagat	ggctgctgta	6180
ggctgctcct	atcagaacaa	cggaagatt	gacctcagca	atatgacccc	tgaacaacag	6240
gctgatgaac	tgaataaaact	tttccgtgaa	gagttagaag	cccgtaaaaca	aaaagtcgct	6300
aaggctaggg	ctgaagtcaa	agaagaaact	gtttctgaaa	aagaaccagt	gaatccagac	6360
tttggatagg	taggccgtga	gcataaggca	tctgggtgtc	gtatcctgtc	tgctactgct	6420
attcgtaatc	tggctaagat	tagtaatctg	ccatctactc	aggcagctac	tcttgccggag	6480
attcagaaat	cactggcagc	taaagactat	aagattatct	acggtacacc	tactcagggt	6540
gcagagtatg	ctcgtcagaa	gaatgttact	gaattgactt	ctcaggaaat	ggaagaagct	6600
caggcaggta	atattttatgg	ctggactaac	ttcgatgata	agaccattta	tctggtttagc	6660
ccatctatgg	aaacctcat	tcatgaactg	gttcatgcct	ctaccttcca	ggaagtttat	6720
tccttctatc	agggtaatga	agtaagccct	acttctaagc	aggctattga	gaacctgaa	6780
ggctctgatg	aacagttccg	ttctctggat	atttccaaag	attctccaga	aatgagagaa	6840
gcatatgctg	atgctattgc	aactatcgaa	ggtcatttga	gtaatggatt	tgttgacca	6900
gctatctcta	aagctgctgc	tcttaatgag	tttatggctt	gggggttagc	taaccgtgct	6960
cttgctgcta	aacagaagag	aacatcttca	ctggttcaaa	tggtgaaaga	tgtttatcag	7020
gctattaaga	aattgatattg	gggacgtaaa	caagctcctg	cattgggaga	agatatgttc	7080
tccaatctgc	tgtttaactc	tgcaattctg	atgcgtagcc	aacctacaac	tcaggcagta	7140
gctaaagatg	gcacactggt	ccatagcaaa	gcataatgta	ataatgaacg	tctgtctcag	7200
ttgaaccaga	ctttcgataa	actggtaact	gattaccttc	gtactgaccc	agttacagaa	7260
gtagaacgtc	gtggcaatgt	ggctaatgca	ttaatgagtg	ctactcgact	ggttcgtgat	7320
gttcagtctc	atggcttcaa	tatgactgct	caggaacagt	ctgtattcca	gatggttact	7380
gctgcattag	caactgaagc	tgcgattgac	ccacatgcta	tggctcgtgc	tcaggaactt	7440
tatacccatg	taatgaaaca	ccttacggta	gagcatttca	tggctgaccc	tgatagtact	7500
aacctgctg	accgttacta	tgctcaacag	aaatagaca	ccatctctgg	tgctaactctg	7560
gttgaagtag	atgccaaagg	tagaaccagt	ctgttaccta	cattcctggg	tctggctatg	7620
gttaatgaag	aactacgttc	aatcatmeta	gaaatgcctg	tacctaaagc	agataagaaa	7680
ttaggaatg	atatagatac	tctgcttacc	aatgcaggtg	ctcaggtaat	ggaatctctg	7740
aaccgtcgtg	tggctgggtg	ccagaagact	actaatgttc	aggacagtat	tgatgctttg	7800
tcagaacaaa	tcatggctgc	tgctttgaaa	cgagagtcct	tctatgatgc	tgtagcaacc	7860
cctaccggta	acttcattga	ccgtgcta	cagtacgtaa	cggatagcat	tgaacggtta	7920
tctgaaactg	ttattgagaa	ggcagataag	gtaattgcta	acccttctaa	tatagctgct	7980
aaaggtggtg	ctcatctggc	taaaactgact	gctgctattg	catctgaaaa	acagggtgaa	8040
atagtggtc	agggtgttat	gactgctatg	aaccagggtg	aagtatggca	acctttccat	8100
gacttagtta	atgacattgt	tggccgtact	aagactaatg	ccaatgtcta	tgacttaatc	8160
aaattgggta	agagccagat	ttctcaagac	cgtcagcaat	tccgtgagca	tttacctaca	8220
gtcattgctg	gtaagttctc	tcgtaaatg	actgataccg	aatggctctg	aatgcatact	8280
ggtttaggta	aaacagattt	agctgttcta	cgtgaaacta	tgagcatggc	tgaattaga	8340
gatttactct	cttcatccaa	gaaagtmeta	gatgaaatct	ctactctgga	aaaagagatt	8400

-continued

---

cagaaccaag caggtagaaa ctggaatctg gttcagaaga aatctaagca actggctcaa	8460
tacatgatta tgggggaagt aggtaataac ctccttcgta atgccatgc tattagtcgt	8520
ttgttaggtg aacgtattac taatggctct gtgacagatg tagctgctat tgataagctc	8580
attactttgt actctctgga attgatgaat aagtctgacc gtgacctttt gtcagaattg	8640
gctcaatcag aagtggaagg tatggagttc tccattgctt atatggttgg tcaacgtact	8700
gaagagatgc gtaaagctaa aggtgataac cgtactctgc tgaatcactt taaaggctat	8760
atcctctgag agaaccagca aggtgtgaat ttgattattg ctgacgataa agagtttgc	8820
aagttaaata gccaatcctt tactcgtatt ggtacttata aggggagcac tggtttccgt	8880
actggttcta aaggttatta cttcagccca gtagctgccc gtgcccctta ctctcagggt	8940
attcttcaga acgttcgtaa tactgctggt ggtgtggata ttggtactgg ctttacgtta	9000
ggcactatgg ttgctgggcg tattactgac aaaccaaccg tagagcgtat taccaaagct	9060
ctggctaaag gtgagcgtgg gcgtgaacca ctgatgcaa tttataacag caaaggctcag	9120
gtagttgctt atgaacaatc cggtgacctt aatatgttga agcacctaaa ccaagacaat	9180
cactttgcta agatggttgg tgtatggcgt ggtcgtcagg tgaagaggc taaagcacia	9240
cgttttaatg acattctcat tgagcaatta catgctatgt atgagaaaga cattaagac	9300
tccagtgcta ataaatctca atatgtaaac ctgttaggta aaattgatga cccagtactg	9360
gctgatgcca ttaacctgat gaacattgag actcgtcata aggccgaaga actcttcggt	9420
aaagatgagt tatgggttcg tagggatagc ctgaatgatg cacttggtta tctgtctgca	9480
tctattggtg atgtgtggac cggttaactc ctgttggtcac ctgacacctt tgatactgtt	9540
aagaagatgt tcctcggctc attcggtaat aaggcatatc atgtagtaat gaatgctgaa	9600
aataccattc agaacttagt gaaggacgct aagacagtaa ttgttgttaa atctgttgta	9660
gtaccggcag ttaacttctt tgctaacatc taccagatga ttggacgtgg tgttctctgtt	9720
aaagatattg ctgtgaacat tcctcgtgag acgtcagaga ttaatcagta tattaaatct	9780
cgtttacgtc agattgatgc ggaagcagag ctacgtgctg ctgaaggtta ccctaactctg	9840
gttcgtaaac ttaaaactga gattcaatct attactgata gtcacgtctg tatgagtatc	9900
tggcctttga ttgaagcagg tgagttctct tctattgctg atgctggtat tagtcgtgat	9960
gacctgttag tagctgaagg taagattcat gagtacctgg aaaaacttgc taataaactt	10020
ccagaaaaag tacgtaatgc tggccgttac gctcttattg ctaaggacac tgctctgttc	10080
cagggtatcc agaaaacagt agagtattca gactttattg ctaaagccat catctatgat	10140
gatttagtga aacgtaagaa aaaatcttct tctgaagcat taggtcaggt aactgaagag	10200
tttattaact atgacagatt gcctggctct ttccgtggtt atatgaaag tatgggtctg	10260
atgtggttct acaactttaa aattcgttcc attaaagtgt ctatgagcat gattagaaac	10320
aaccagtac attctctgat tgctacagta gtacctgctc ctaccatggt tggtaacgta	10380
ggtctaccaa ttcaggacaa catgctaacc atgctggctg aaggaagact ggattactca	10440
ttaggcttcg gacaaggatt aagagcacct accctcaatc cttggttcaa ccttactcac	10500
taataa	10506

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 3500

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

---

Peptide

<400> SEQUENCE: 2

Met Ser Val Phe Asp Arg Leu Ala Gly Phe Ala Asp Ser Val Thr Asn  
 1 5 10 15

Ala Lys Gln Val Asp Val Ser Thr Ala Thr Ala Gln Lys Lys Ala Glu  
 20 25 30

Gln Gly Val Thr Thr Pro Leu Val Ser Pro Asp Ala Ala Tyr Gln Met  
 35 40 45

Gln Ala Ala Arg Thr Gly Asn Val Gly Ala Asn Ala Phe Glu Pro Gly  
 50 55 60

Thr Val Gln Ser Asp Phe Met Asn Leu Thr Pro Met Gln Ile Met Asn  
 65 70 75 80

Lys Tyr Gly Val Glu Gln Gly Leu Gln Leu Ile Asn Ala Arg Ala Asp  
 85 90 95

Ala Gly Asn Gln Val Phe Asn Asp Ser Val Thr Thr Arg Thr Pro Gly  
 100 105 110

Glu Glu Leu Gly Asp Ile Ala Thr Gly Val Gly Leu Gly Phe Val Asn  
 115 120 125

Thr Leu Gly Gly Ile Gly Ala Leu Gly Ala Gly Leu Leu Asn Asp Asp  
 130 135 140

Ala Gly Ala Val Val Ala Gln Gln Leu Ser Lys Phe Asn Asp Ala Val  
 145 150 155 160

His Ala Thr Gln Ser Gln Ala Leu Gln Asp Lys Arg Lys Leu Phe Ala  
 165 170 175

Ala Arg Asn Leu Met Asn Glu Val Glu Ser Glu Arg Gln Tyr Gln Thr  
 180 185 190

Asp Lys Lys Glu Gly Thr Asn Asp Ile Val Ala Ser Leu Ser Lys Phe  
 195 200 205

Gly Arg Asp Phe Val Gly Ser Ile Glu Asn Ala Ala Gln Thr Asp Ser  
 210 215 220

Ile Ile Ser Asp Gly Leu Ala Glu Gly Val Gly Ser Leu Leu Gly Ala  
 225 230 235 240

Gly Pro Val Leu Arg Gly Ala Ser Leu Leu Gly Lys Ala Val Val Pro  
 245 250 255

Ala Asn Thr Leu Arg Ser Ala Ala Leu Ala Gly Ala Ile Asp Ala Gly  
 260 265 270

Thr Gly Thr Gln Ser Leu Ala Arg Ile Ala Ser Thr Val Gly Arg Ala  
 275 280 285

Ala Pro Gly Met Val Gly Val Gly Ala Met Glu Ala Gly Gly Ala Tyr  
 290 295 300

Gln Gln Thr Ala Asp Glu Ile Met Lys Met Ser Leu Lys Asp Leu Glu  
 305 310 315 320

Lys Ser Pro Val Tyr Gln Gln His Ile Lys Asp Gly Met Ser Pro Glu  
 325 330 335

Gln Ala Arg Arg Gln Thr Ala Ser Glu Thr Gly Leu Thr Ala Ala Ala  
 340 345 350

Ile Gln Leu Pro Ile Ala Ala Ala Thr Gly Pro Leu Val Ser Arg Phe  
 355 360 365

Glu Met Ala Pro Phe Arg Ala Gly Ser Leu Gly Ala Val Gly Met Asn  
 370 375 380

Leu Ala Arg Glu Thr Val Glu Glu Gly Val Gln Gly Ala Thr Gly Gln  
 385 390 395 400



-continued

820				825				830							
Glu	Trp	Val	Arg	Ser	Arg	Thr	Gly	Leu	Gly	Val	Asn	Pro	Tyr	Asp	Thr
	835						840					845			
Lys	Ser	Val	Lys	Phe	Ala	Gln	Gln	Val	Ala	Leu	Glu	Ala	Lys	Thr	Val
	850					855						860			
Ala	Asp	Ile	Ala	Asn	Ala	Leu	Ala	Ser	Ala	Tyr	Pro	Glu	Leu	Lys	Val
	865				870					875					880
Ser	His	Ile	Lys	Val	Thr	Pro	Leu	Asp	Ser	Arg	Leu	Asn	Ala	Pro	Ala
				885						890				895	
Ala	Glu	Val	Val	Lys	Ala	Phe	Arg	Gln	Gly	Asn	Arg	Asp	Val	Ala	Ser
			900						905				910		
Ser	Gln	Pro	Lys	Ala	Asp	Ser	Val	Asn	Gln	Val	Lys	Glu	Thr	Pro	Val
		915					920					925			
Thr	Lys	Gln	Glu	Pro	Val	Thr	Ser	Thr	Val	Gln	Thr	Lys	Thr	Pro	Val
	930					935					940				
Ser	Glu	Ser	Val	Lys	Thr	Glu	Pro	Thr	Thr	Lys	Glu	Ser	Ser	Pro	Gln
	945				950					955					960
Ala	Ile	Lys	Glu	Pro	Val	Asn	Gln	Ser	Glu	Lys	Gln	Asp	Val	Asn	Leu
				965					970					975	
Thr	Asn	Glu	Asp	Asn	Ile	Lys	Gln	Pro	Thr	Glu	Ser	Val	Lys	Glu	Thr
			980				985						990		
Glu	Thr	Ser	Thr	Lys	Glu	Ser	Thr	Val	Thr	Glu	Glu	Leu	Lys	Glu	Gly
		995					1000					1005			
Ile	Asp	Ala	Val	Tyr	Pro	Ser	Leu	Val	Gly	Thr	Ala	Asp	Ser	Lys	Ala
	1010					1015					1020				
Glu	Gly	Ile	Lys	Asn	Tyr	Phe	Lys	Leu	Ser	Phe	Thr	Leu	Pro	Glu	Glu
	1025				1030					1035				1040	
Gln	Lys	Ser	Arg	Thr	Val	Gly	Ser	Glu	Ala	Pro	Leu	Lys	Asp	Val	Ala
				1045					1050					1055	
Gln	Ala	Leu	Ser	Ser	Arg	Ala	Arg	Tyr	Glu	Leu	Phe	Thr	Glu	Lys	Glu
			1060				1065						1070		
Thr	Ala	Asn	Pro	Ala	Phe	Asn	Gly	Glu	Val	Ile	Lys	Arg	Tyr	Lys	Glu
	1075					1080						1085			
Leu	Met	Glu	His	Gly	Glu	Gly	Ile	Ala	Asp	Ile	Leu	Arg	Ser	Arg	Leu
	1090					1095					1100				
Ala	Lys	Phe	Leu	Asn	Thr	Lys	Asp	Val	Gly	Lys	Arg	Phe	Ala	Gln	Gly
	1105				1110					1115				1120	
Thr	Glu	Ala	Asn	Arg	Trp	Val	Gly	Gly	Lys	Leu	Leu	Asn	Ile	Val	Glu
				1125					1130					1135	
Gln	Asp	Gly	Asp	Thr	Phe	Lys	Tyr	Asn	Glu	Gln	Leu	Leu	Gln	Thr	Ala
		1140					1145						1150		
Val	Leu	Ala	Gly	Leu	Gln	Trp	Arg	Leu	Thr	Ala	Thr	Ser	Asn	Thr	Ala
	1155					1160						1165			
Ile	Lys	Asp	Ala	Lys	Asp	Val	Ala	Ala	Ile	Thr	Gly	Ile	Asp	Gln	Ala
	1170				1175						1180				
Leu	Leu	Pro	Glu	Gly	Leu	Val	Glu	Gln	Phe	Asp	Thr	Gly	Met	Thr	Leu
	1185				1190					1195				1200	
Thr	Glu	Ala	Val	Ser	Ser	Leu	Ala	Gln	Lys	Ile	Glu	Ser	Tyr	Trp	Gly
			1205						1210					1215	
Leu	Ser	Arg	Asn	Pro	Asn	Ala	Pro	Leu	Gly	Tyr	Thr	Lys	Gly	Ile	Pro
			1220				1225						1230		
Thr	Ala	Met	Ala	Ala	Glu	Ile	Leu	Ala	Ala	Phe	Val	Glu	Ser	Thr	Asp
	1235					1240						1245			

-continued

---

Val Val Glu Asn Ile Val Asp Met Ser Glu Ile Asp Pro Asp Asn Lys  
 1250 1255 1260

Lys Thr Ile Gly Leu Tyr Thr Ile Thr Glu Leu Asp Ser Phe Asp Pro  
 1265 1270 1275 1280

Ile Asn Ser Phe Pro Thr Ala Ile Glu Glu Ala Val Leu Val Asn Pro  
 1285 1290 1295

Thr Glu Lys Met Phe Phe Gly Asp Asp Ile Pro Pro Val Ala Asn Thr  
 1300 1305 1310

Gln Leu Arg Asn Pro Ala Val Arg Asn Thr Pro Glu Gln Lys Ala Ala  
 1315 1320 1325

Leu Lys Ala Glu Gln Ala Thr Glu Phe Tyr Val His Thr Pro Met Val  
 1330 1335 1340

Gln Phe Tyr Glu Thr Leu Gly Lys Asp Arg Ile Leu Glu Leu Met Gly  
 1345 1350 1355 1360

Ala Gly Thr Leu Asn Lys Glu Leu Leu Asn Asp Asn His Ala Lys Ser  
 1365 1370 1375

Leu Glu Gly Lys Asn Arg Ser Val Glu Asp Ser Tyr Asn Gln Leu Phe  
 1380 1385 1390

Ser Val Ile Glu Gln Val Arg Ala Gln Ser Glu Asp Ile Ser Thr Val  
 1395 1400 1405

Pro Ile His Tyr Ala Tyr Asn Met Thr Arg Val Gly Arg Met Gln Met  
 1410 1415 1420

Leu Gly Lys Tyr Asn Pro Gln Ser Ala Lys Leu Val Arg Glu Ala Ile  
 1425 1430 1435 1440

Leu Pro Thr Lys Ala Thr Leu Asp Leu Ser Asn Gln Asn Asn Glu Asp  
 1445 1450 1455

Phe Ser Ala Phe Gln Leu Gly Leu Ala Gln Ala Leu Asp Ile Lys Val  
 1460 1465 1470

His Thr Met Thr Arg Glu Val Met Ser Asp Glu Leu Thr Lys Leu Leu  
 1475 1480 1485

Glu Gly Asn Leu Lys Pro Ala Ile Asp Met Met Val Glu Phe Asn Thr  
 1490 1495 1500

Thr Gly Ser Leu Pro Glu Asn Ala Val Asp Val Leu Asn Thr Ala Leu  
 1505 1510 1515 1520

Gly Asp Arg Lys Ser Phe Val Ala Leu Met Ala Leu Met Glu Tyr Ser  
 1525 1530 1535

Arg Tyr Leu Val Ala Glu Asp Lys Ser Ala Phe Val Thr Pro Leu Tyr  
 1540 1545 1550

Val Glu Ala Asp Gly Val Thr Asn Gly Pro Ile Asn Ala Met Met Leu  
 1555 1560 1565

Met Thr Gly Gly Leu Phe Thr Pro Asp Trp Ile Arg Asn Ile Ala Lys  
 1570 1575 1580

Gly Gly Leu Phe Ile Gly Ser Pro Asn Lys Thr Met Asn Glu His Arg  
 1585 1590 1595 1600

Ser Thr Ala Asp Asn Asn Asp Leu Tyr Gln Ala Ser Thr Asn Ala Leu  
 1605 1610 1615

Met Glu Ser Leu Gly Lys Leu Arg Ser Asn Tyr Ala Ser Asn Met Pro  
 1620 1625 1630

Ile Gln Ser Gln Ile Asp Ser Leu Leu Ser Leu Met Asp Leu Phe Leu  
 1635 1640 1645

Pro Asp Ile Asn Leu Gly Glu Asn Gly Ala Leu Glu Leu Lys Arg Gly  
 1650 1655 1660

-continued

---

Ile Ala Lys Asn Pro Leu Thr Ile Thr Ile Tyr Gly Ser Gly Ala Arg  
 1665 1670 1675 1680  
 Gly Ile Ala Gly Lys Leu Val Ser Ser Val Thr Asp Ala Ile Tyr Glu  
 1685 1690 1695  
 Arg Met Ser Asp Val Leu Lys Ala Arg Ala Lys Asp Pro Asn Ile Ser  
 1700 1705 1710  
 Ala Ala Met Ala Met Phe Gly Lys Gln Ala Ala Ser Glu Ala His Ala  
 1715 1720 1725  
 Glu Glu Leu Leu Ala Arg Phe Leu Lys Asp Met Glu Thr Leu Thr Ser  
 1730 1735 1740  
 Thr Val Pro Val Lys Arg Lys Gly Val Leu Glu Leu Gln Ser Thr Gly  
 1745 1750 1755 1760  
 Thr Gly Ala Lys Gly Lys Ile Asn Pro Lys Thr Tyr Thr Ile Lys Gly  
 1765 1770 1775  
 Glu Gln Leu Lys Ala Leu Gln Glu Asn Met Leu His Phe Phe Val Glu  
 1780 1785 1790  
 Pro Leu Arg Asn Gly Ile Thr Gln Thr Val Gly Glu Ser Leu Val Tyr  
 1795 1800 1805  
 Ser Thr Glu Gln Leu Gln Lys Ala Thr Gln Ile Gln Ser Val Val Leu  
 1810 1815 1820  
 Glu Asp Met Phe Lys Gln Arg Val Gln Glu Lys Leu Ala Glu Lys Ala  
 1825 1830 1835 1840  
 Lys Asp Pro Thr Trp Lys Lys Gly Asp Phe Leu Thr Gln Lys Glu Leu  
 1845 1850 1855  
 Asn Asp Ile Gln Ala Ser Leu Asn Asn Leu Ala Pro Met Ile Glu Thr  
 1860 1865 1870  
 Gly Ser Gln Thr Phe Tyr Ile Ala Gly Ser Glu Asn Ala Glu Val Ala  
 1875 1880 1885  
 Asn Gln Val Leu Ala Thr Asn Leu Asp Asp Arg Met Arg Val Pro Met  
 1890 1895 1900  
 Ser Ile Tyr Ala Pro Ala Gln Ala Gly Val Ala Gly Ile Pro Phe Met  
 1905 1910 1915 1920  
 Thr Ile Gly Thr Gly Asp Gly Met Met Met Gln Thr Leu Ser Thr Met  
 1925 1930 1935  
 Lys Gly Ala Pro Lys Asn Thr Leu Lys Ile Phe Asp Gly Met Asn Ile  
 1940 1945 1950  
 Gly Leu Asn Asp Ile Thr Asp Ala Ser Arg Lys Ala Asn Glu Ala Val  
 1955 1960 1965  
 Tyr Thr Ser Trp Gln Gly Asn Pro Ile Lys Asn Val Tyr Glu Ser Tyr  
 1970 1975 1980  
 Ala Lys Phe Met Lys Asn Val Asp Phe Ser Lys Leu Ser Pro Glu Ala  
 1985 1990 1995 2000  
 Leu Glu Ala Ile Gly Lys Ser Ala Leu Glu Tyr Asp Gln Arg Glu Asn  
 2005 2010 2015  
 Ala Thr Val Asp Asp Ile Ala Asn Ala Ala Ser Leu Ile Glu Arg Asn  
 2020 2025 2030  
 Leu Arg Asn Ile Ala Leu Gly Val Asp Ile Arg His Lys Val Leu Asp  
 2035 2040 2045  
 Lys Val Asn Leu Ser Ile Asp Gln Met Ala Ala Val Gly Ala Pro Tyr  
 2050 2055 2060  
 Gln Asn Asn Gly Lys Ile Asp Leu Ser Asn Met Thr Pro Glu Gln Gln  
 2065 2070 2075 2080  
 Ala Asp Glu Leu Asn Lys Leu Phe Arg Glu Glu Leu Glu Ala Arg Lys

-continued

2085					2090					2095					
Gln	Lys	Val	Ala	Lys	Ala	Arg	Ala	Glu	Val	Lys	Glu	Glu	Thr	Val	Ser
			2100					2105						2110	
Glu	Lys	Glu	Pro	Val	Asn	Pro	Asp	Phe	Gly	Met	Val	Gly	Arg	Glu	His
		2115					2120					2125			
Lys	Ala	Ser	Gly	Val	Arg	Ile	Leu	Ser	Ala	Thr	Ala	Ile	Arg	Asn	Leu
		2130				2135					2140				
Ala	Lys	Ile	Ser	Asn	Leu	Pro	Ser	Thr	Gln	Ala	Ala	Thr	Leu	Ala	Glu
		2145				2150					2155				2160
Ile	Gln	Lys	Ser	Leu	Ala	Ala	Lys	Asp	Tyr	Lys	Ile	Ile	Tyr	Gly	Thr
			2165						2170					2175	
Pro	Thr	Gln	Val	Ala	Glu	Tyr	Ala	Arg	Gln	Lys	Asn	Val	Thr	Glu	Leu
			2180					2185					2190		
Thr	Ser	Gln	Glu	Met	Glu	Glu	Ala	Gln	Ala	Gly	Asn	Ile	Tyr	Gly	Trp
		2195					2200					2205			
Thr	Asn	Phe	Asp	Asp	Lys	Thr	Ile	Tyr	Leu	Val	Ser	Pro	Ser	Met	Glu
		2210				2215					2220				
Thr	Leu	Ile	His	Glu	Leu	Val	His	Ala	Ser	Thr	Phe	Glu	Glu	Val	Tyr
		2225				2230					2235				2240
Ser	Phe	Tyr	Gln	Gly	Asn	Glu	Val	Ser	Pro	Thr	Ser	Lys	Gln	Ala	Ile
			2245						2250					2255	
Glu	Asn	Leu	Glu	Gly	Leu	Met	Glu	Gln	Phe	Arg	Ser	Leu	Asp	Ile	Ser
			2260					2265					2270		
Lys	Asp	Ser	Pro	Glu	Met	Arg	Glu	Ala	Tyr	Ala	Asp	Ala	Ile	Ala	Thr
		2275					2280					2285			
Ile	Glu	Gly	His	Leu	Ser	Asn	Gly	Phe	Val	Asp	Pro	Ala	Ile	Ser	Lys
		2290				2295					2300				
Ala	Ala	Ala	Leu	Asn	Glu	Phe	Met	Ala	Trp	Gly	Leu	Ala	Asn	Arg	Ala
			2305			2310					2315				2320
Leu	Ala	Ala	Lys	Gln	Lys	Arg	Thr	Ser	Ser	Leu	Val	Gln	Met	Val	Lys
			2325						2330					2335	
Asp	Val	Tyr	Gln	Ala	Ile	Lys	Lys	Leu	Ile	Trp	Gly	Arg	Lys	Gln	Ala
			2340					2345						2350	
Pro	Ala	Leu	Gly	Glu	Asp	Met	Phe	Ser	Asn	Leu	Leu	Phe	Asn	Ser	Ala
		2355					2360					2365			
Ile	Leu	Met	Arg	Ser	Gln	Pro	Thr	Thr	Gln	Ala	Val	Ala	Lys	Asp	Gly
		2370				2375					2380				
Thr	Leu	Phe	His	Ser	Lys	Ala	Tyr	Gly	Asn	Asn	Glu	Arg	Leu	Ser	Gln
			2385			2390					2395				2400
Leu	Asn	Gln	Thr	Phe	Asp	Lys	Leu	Val	Thr	Asp	Tyr	Leu	Arg	Thr	Asp
			2405						2410					2415	
Pro	Val	Thr	Glu	Val	Glu	Arg	Arg	Gly	Asn	Val	Ala	Asn	Ala	Leu	Met
			2420					2425					2430		
Ser	Ala	Thr	Arg	Leu	Val	Arg	Asp	Val	Gln	Ser	His	Gly	Phe	Asn	Met
		2435					2440					2445			
Thr	Ala	Gln	Glu	Gln	Ser	Val	Phe	Gln	Met	Val	Thr	Ala	Ala	Leu	Ala
		2450				2455					2460				
Thr	Glu	Ala	Ala	Ile	Asp	Pro	His	Ala	Met	Ala	Arg	Ala	Gln	Glu	Leu
			2465			2470					2475				2480
Tyr	Thr	His	Val	Met	Lys	His	Leu	Thr	Val	Glu	His	Phe	Met	Ala	Asp
			2485					2490						2495	
Pro	Asp	Ser	Thr	Asn	Pro	Ala	Asp	Arg	Tyr	Tyr	Ala	Gln	Gln	Lys	Tyr
			2500					2505						2510	

-continued

---

Asp Thr Ile Ser Gly Ala Asn Leu Val Glu Val Asp Ala Lys Gly Arg  
 2515 2520 2525

Thr Ser Leu Leu Pro Thr Phe Leu Gly Leu Ala Met Val Asn Glu Glu  
 2530 2535 2540

Leu Arg Ser Ile Ile Lys Glu Met Pro Val Pro Lys Ala Asp Lys Lys  
 2545 2550 2555 2560

Leu Gly Asn Asp Ile Asp Thr Leu Leu Thr Asn Ala Gly Thr Gln Val  
 2565 2570 2575

Met Glu Ser Leu Asn Arg Arg Met Ala Gly Asp Gln Lys Ala Thr Asn  
 2580 2585 2590

Val Gln Asp Ser Ile Asp Ala Leu Ser Glu Thr Ile Met Ala Ala Ala  
 2595 2600 2605

Leu Lys Arg Glu Ser Phe Tyr Asp Ala Val Ala Thr Pro Thr Gly Asn  
 2610 2615 2620

Phe Ile Asp Arg Ala Asn Gln Tyr Val Thr Asp Ser Ile Glu Arg Leu  
 2625 2630 2635 2640

Ser Glu Thr Val Ile Glu Lys Ala Asp Lys Val Ile Ala Asn Pro Ser  
 2645 2650 2655

Asn Ile Ala Ala Lys Gly Val Ala His Leu Ala Lys Leu Thr Ala Ala  
 2660 2665 2670

Ile Ala Ser Glu Lys Gln Gly Glu Ile Val Ala Gln Gly Val Met Thr  
 2675 2680 2685

Ala Met Asn Gln Gly Lys Val Trp Gln Pro Phe His Asp Leu Val Asn  
 2690 2695 2700

Asp Ile Val Gly Arg Thr Lys Thr Asn Ala Asn Val Tyr Asp Leu Ile  
 2705 2710 2715 2720

Lys Leu Val Lys Ser Gln Ile Ser Gln Asp Arg Gln Gln Phe Arg Glu  
 2725 2730 2735

His Leu Pro Thr Val Ile Ala Gly Lys Phe Ser Arg Lys Leu Thr Asp  
 2740 2745 2750

Thr Glu Trp Ser Ala Met His Thr Gly Leu Gly Lys Thr Asp Leu Ala  
 2755 2760 2765

Val Leu Arg Glu Thr Met Ser Met Ala Glu Ile Arg Asp Leu Leu Ser  
 2770 2775 2780

Ser Ser Lys Lys Val Lys Asp Glu Ile Ser Thr Leu Glu Lys Glu Ile  
 2785 2790 2795 2800

Gln Asn Gln Ala Gly Arg Asn Trp Asn Leu Val Gln Lys Lys Ser Lys  
 2805 2810 2815

Gln Leu Ala Gln Tyr Met Ile Met Gly Glu Val Gly Asn Asn Leu Leu  
 2820 2825 2830

Arg Asn Ala His Ala Ile Ser Arg Leu Leu Gly Glu Arg Ile Thr Asn  
 2835 2840 2845

Gly Pro Val Ala Asp Val Ala Ala Ile Asp Lys Leu Ile Thr Leu Tyr  
 2850 2855 2860

Ser Leu Glu Leu Met Asn Lys Ser Asp Arg Asp Leu Leu Ser Glu Leu  
 2865 2870 2875 2880

Ala Gln Ser Glu Val Glu Gly Met Glu Phe Ser Ile Ala Tyr Met Val  
 2885 2890 2895

Gly Gln Arg Thr Glu Glu Met Arg Lys Ala Lys Gly Asp Asn Arg Thr  
 2900 2905 2910

Leu Leu Asn His Phe Lys Gly Tyr Ile Pro Val Glu Asn Gln Gln Gly  
 2915 2920 2925

-continued

---

Val Asn Leu Ile Ile Ala Asp Asp Lys Glu Phe Ala Lys Leu Asn Ser  
 2930 2935 2940  
 Gln Ser Phe Thr Arg Ile Gly Thr Tyr Gln Gly Ser Thr Gly Phe Arg  
 2945 2950 2955 2960  
 Thr Gly Ser Lys Gly Tyr Tyr Phe Ser Pro Val Ala Ala Arg Ala Pro  
 2965 2970 2975  
 Tyr Ser Gln Gly Ile Leu Gln Asn Val Arg Asn Thr Ala Gly Gly Val  
 2980 2985 2990  
 Asp Ile Gly Thr Gly Phe Thr Leu Gly Thr Met Val Ala Gly Arg Ile  
 2995 3000 3005  
 Thr Asp Lys Pro Thr Val Glu Arg Ile Thr Lys Ala Leu Ala Lys Gly  
 3010 3015 3020  
 Glu Arg Gly Arg Glu Pro Leu Met Pro Ile Tyr Asn Ser Lys Gly Gln  
 3025 3030 3035 3040  
 Val Val Ala Tyr Glu Gln Ser Val Asp Pro Asn Met Leu Lys His Leu  
 3045 3050 3055  
 Asn Gln Asp Asn His Phe Ala Lys Met Val Gly Val Trp Arg Gly Arg  
 3060 3065 3070  
 Gln Val Glu Glu Ala Lys Ala Gln Arg Phe Asn Asp Ile Leu Ile Glu  
 3075 3080 3085  
 Gln Leu His Ala Met Tyr Glu Lys Asp Ile Lys Asp Ser Ser Ala Asn  
 3090 3095 3100  
 Lys Ser Gln Tyr Val Asn Leu Leu Gly Lys Ile Asp Asp Pro Val Leu  
 3105 3110 3115 3120  
 Ala Asp Ala Ile Asn Leu Met Asn Ile Glu Thr Arg His Lys Ala Glu  
 3125 3130 3135  
 Glu Leu Phe Gly Lys Asp Glu Leu Trp Val Arg Arg Asp Met Leu Asn  
 3140 3145 3150  
 Asp Ala Leu Gly Tyr Arg Ala Ala Ser Ile Gly Asp Val Trp Thr Gly  
 3155 3160 3165  
 Asn Ser Arg Trp Ser Pro Ser Thr Leu Asp Thr Val Lys Lys Met Phe  
 3170 3175 3180  
 Leu Gly Ala Phe Gly Asn Lys Ala Tyr His Val Val Met Asn Ala Glu  
 3185 3190 3195 3200  
 Asn Thr Ile Gln Asn Leu Val Lys Asp Ala Lys Thr Val Ile Val Val  
 3205 3210 3215  
 Lys Ser Val Val Val Pro Ala Val Asn Phe Leu Ala Asn Ile Tyr Gln  
 3220 3225 3230  
 Met Ile Gly Arg Gly Val Pro Val Lys Asp Ile Ala Val Asn Ile Pro  
 3235 3240 3245  
 Arg Lys Thr Ser Glu Ile Asn Gln Tyr Ile Lys Ser Arg Leu Arg Gln  
 3250 3255 3260  
 Ile Asp Ala Glu Ala Glu Leu Arg Ala Ala Glu Gly Asn Pro Asn Leu  
 3265 3270 3275 3280  
 Val Arg Lys Leu Lys Thr Glu Ile Gln Ser Ile Thr Asp Ser His Arg  
 3285 3290 3295  
 Arg Met Ser Ile Trp Pro Leu Ile Glu Ala Gly Glu Phe Ser Ser Ile  
 3300 3305 3310  
 Ala Asp Ala Gly Ile Ser Arg Asp Asp Leu Leu Val Ala Glu Gly Lys  
 3315 3320 3325  
 Ile His Glu Tyr Met Glu Lys Leu Ala Asn Lys Leu Pro Glu Lys Val  
 3330 3335 3340  
 Arg Asn Ala Gly Arg Tyr Ala Leu Ile Ala Lys Asp Thr Ala Leu Phe

-continued

3345	3350	3355	3360
Gln Gly Ile Gln Lys Thr Val Glu Tyr Ser Asp Phe Ile Ala Lys Ala	3365	3370	3375
Ile Ile Tyr Asp Asp Leu Val Lys Arg Lys Lys Lys Ser Ser Ser Glu	3380	3385	3390
Ala Leu Gly Gln Val Thr Glu Glu Phe Ile Asn Tyr Asp Arg Leu Pro	3395	3400	3405
Gly Arg Phe Arg Gly Tyr Met Glu Ser Met Gly Leu Met Trp Phe Tyr	3410	3415	3420
Asn Phe Lys Ile Arg Ser Ile Lys Val Ala Met Ser Met Ile Arg Asn	3425	3430	3435
Asn Pro Val His Ser Leu Ile Ala Thr Val Val Pro Ala Pro Thr Met	3445	3450	3455
Phe Gly Asn Val Gly Leu Pro Ile Gln Asp Asn Met Leu Thr Met Leu	3460	3465	3470
Ala Glu Gly Arg Leu Asp Tyr Ser Leu Gly Phe Gly Gln Gly Leu Arg	3475	3480	3485
Ala Pro Thr Leu Asn Pro Trp Phe Asn Leu Thr His	3490	3495	3500

<210> SEQ ID NO 3  
 <211> LENGTH: 3318  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 3

gaaagtacag ttacagaaga attaaaagaa ggtattgatg ctgtttaccc ttcattggta	60
ggtactgctg attctaaagc agagggtatt aagaactatt tcaaattgtc ctttacctta	120
ccagaagaac agaaatcccg tactgttggt tcagaagcac ctctaaaaga tgtagcccaa	180
gctctgtcct ctcggtctcg ttatgaactc tttactgaga aagaaactgc taaccctgct	240
tttaatgggg aagtatttaa gcgatacaaa gaactcatgg aacatgggga aggtattgct	300
gatattcttc gctcccgctc ggctaagttc cttaacacta aggatgttgg taaacgtttt	360
gctcaaggta cagaagccaa cggttgggta ggtggtaagt tacttaacat tgttgagcag	420
gatggggata cctttaagta caacgaacaa ttgctacaga ctgctgtatt agcaggctct	480
caatggagac ttactgctac cagcaatact gctatcaaag atgcaaaaga tgttgctgct	540
attactggta ttgaccaagc tctgctgcca gaaggtttag tagagcaatt tgatactggt	600
atgacactca ctgaagcagt tagttccctg gctcagaaaa ttgagtctta ctggggatta	660
tctcgtaatc caaatgctcc attgggctat accaaaggca tccctacagc aatggctgct	720
gaaattctgg ctgcatttgt agagtctact gatgtttag agaacatcgt ggatattgca	780
gaaattgacc cagataacaa gaagactatt ggtctgtaca ccattactga actggattcc	840
ttcgacccaa ttaatagctt ccctactgct attgaagaag ctgttttagt gaatcctaca	900
gagaagatgt tctttgggta tgacattcct cctgtageta atactcagct tcgtaaccct	960
gctgttcgta atactccaga acagaaggct gcattgaaag cagagcaggc tacagagttc	1020
tatgtacaca ccccaatggt tcaattctat gagacgttag gtaaagaccg tattctcgaa	1080
ctgatgggtg ctggtactct gaataaagag ttacttaatg ataaccatgc taaatctctg	1140
gaaggaaga accgttcagt agaggactct tacaaccaac tgttctcctg cattgagcag	1200

-continued

---

```

gtaagagcac agagcgaaga catctctact gtacctattc actatgcata caatatgacc 1260
cgtggttggtc gtatgcagat gttaggtaaa tacaatcttc aatcagccaa actgggtcgt 1320
gaggccatct tacctactaa agctactttg gatttatcga accagaacaa tgaagacttc 1380
tctgcattcc agttaggctt ggctcaggca ttggacatta aagtcatac tatgactcgt 1440
gaggttatgt ctgacgagtt gactaaatta ctggaagta atctgaaacc agccattgat 1500
atgatggttg agtttaatac cactggttcc ttaccagaaa acgcagtga tgttctgaat 1560
acagcattag gagataggaa gtcattcgta gcattgatgg ctcttatgga gtattcccgt 1620
tacttagtag cagaggataa atctgcattt gtaactccac tgtatgtaga agcagatggt 1680
gttactaatg gtccaatcaa tgccatgatg ctaatgacag gcggtctggt tactcctgac 1740
tggattcgta atattgccaa agggggcttg ttcattggtt ctccaaataa gaccatgaat 1800
gagcatcgct ctactgctga caataatgat ttatatcaag catccactaa tgctttgatg 1860
gaatcgttgg gtaagttacg tagtaactat gcctctaata tgctattca gtctcagata 1920
gacagctctc tttctctgat ggatttgttt ttaccggata ttaatcttgg tgagaatggt 1980
gctttagaac ttaaactggy tattgctaag aaccactga ctattacat ctatggttct 2040
ggtgctcgtg gtattgcagg taagctggtt agttctgtta ctgatgcat ctatgagcgt 2100
atgtctgatg tactgaaagc tctgtctaaa gacccaaata tctctgctgc tatggcaatg 2160
tttggtaagc aagctgcttc agaagcacat gctgaagaac ttcttgcccg tttcctgaaa 2220
gatatggaaa cactgacttc tactgttctc gttaaacgta aaggtgtact ggaactaaa 2280
tccacaggta caggagccaa aggaaaaatc aatcctaaga cctataccat taagggcgag 2340
caactgaagg cacttcagga aaatatgctg cacttctttg tagaaccact acgtaatggt 2400
attactcaga ctgtaggtga aagtctggtg tactctactg aacaattaca gaaagctact 2460
cagattcaat ctgtagtctg ggaagatatg ttcaaacagc gagtacaaga gaagctggca 2520
gagaaggcta aagaccacac atggaagaaa ggtgatttcc ttactcagaa agaactgaat 2580
gatattcagg cttctctgaa taacttagcc cctatgattg agactggttc tcagactttc 2640
tacattgctg gttcagaaaa tgcagaagta gcaaatcagg tattagctac taaccttgat 2700
gaccgtatgc gtgtaccaat gagtatctat gctccagcac aggcgggtgt agcaggtatt 2760
ccatttatga ctattggtac tggatgatgc atgatgatgc aaactctttc cactatgaaa 2820
ggtgcaccaa agaataccct caaaatcttt gatggtatga acattggttt gaatgacatc 2880
actgatgcca gtcgtaaagc taatgaagct gtttacctt cttggcaggg taacctatt 2940
aagaatgttt atgaatcata tgctaagttc atgaagaatg tagatttcag caagctgtcc 3000
cctgaagcat tggaaagcaat tggtaaatct gctctggaat atgaccaacg tgagaatgat 3060
actgtagatg atattgctaa cgctgcatct ctgattgaac gtaacttacg taatattgca 3120
ctgggtgtag atattcgtca taagggtgctg gataaggtaa atctgtccat tgaccagatg 3180
gctgctgtag gtgctcctta tcagaacaac ggtaagattg acctcagcaa tatgaccct 3240
gaacaacagg ctgatgaact gaataaactt ttccgtgaag agttagaagc ccgtaaaaa 3300
aaagtcgcta aggctagg 3318

```

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 1107

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

-continued

---

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic Peptide

<400> SEQUENCE: 4

```

Met Glu Ser Thr Val Thr Glu Glu Leu Lys Glu Gly Ile Asp Ala Val
 1          5          10          15
Tyr Pro Ser Leu Val Gly Thr Ala Asp Ser Lys Ala Glu Gly Ile Lys
 20          25          30
Asn Tyr Phe Lys Leu Ser Phe Thr Leu Pro Glu Glu Gln Lys Ser Arg
 35          40          45
Thr Val Gly Ser Glu Ala Pro Leu Lys Asp Val Ala Gln Ala Leu Ser
 50          55          60
Ser Arg Ala Arg Tyr Glu Leu Phe Thr Glu Lys Glu Thr Ala Asn Pro
 65          70          75          80
Ala Phe Asn Gly Glu Val Ile Lys Arg Tyr Lys Glu Leu Met Glu His
 85          90          95
Gly Glu Gly Ile Ala Asp Ile Leu Arg Ser Arg Leu Ala Lys Phe Leu
100          105          110
Asn Thr Lys Asp Val Gly Lys Arg Phe Ala Gln Gly Thr Glu Ala Asn
115          120          125
Arg Trp Val Gly Gly Lys Leu Leu Asn Ile Val Glu Gln Asp Gly Asp
130          135          140
Thr Phe Lys Tyr Asn Glu Gln Leu Leu Gln Thr Ala Val Leu Ala Gly
145          150          155          160
Leu Gln Trp Arg Leu Thr Ala Thr Ser Asn Thr Ala Ile Lys Asp Ala
165          170          175
Lys Asp Val Ala Ala Ile Thr Gly Ile Asp Gln Ala Leu Leu Pro Glu
180          185          190
Gly Leu Val Glu Gln Phe Asp Thr Gly Met Thr Leu Thr Glu Ala Val
195          200          205
Ser Ser Leu Ala Gln Lys Ile Glu Ser Tyr Trp Gly Leu Ser Arg Asn
210          215          220
Pro Asn Ala Pro Leu Gly Tyr Thr Lys Gly Ile Pro Thr Ala Met Ala
225          230          235          240
Ala Glu Ile Leu Ala Ala Phe Val Glu Ser Thr Asp Val Val Glu Asn
245          250          255
Ile Val Asp Met Ser Glu Ile Asp Pro Asp Asn Lys Lys Thr Ile Gly
260          265          270
Leu Tyr Thr Ile Thr Glu Leu Asp Ser Phe Asp Pro Ile Asn Ser Phe
275          280          285
Pro Thr Ala Ile Glu Glu Ala Val Leu Val Asn Pro Thr Glu Lys Met
290          295          300
Phe Phe Gly Asp Asp Ile Pro Pro Val Ala Asn Thr Gln Leu Arg Asn
305          310          315          320
Pro Ala Val Arg Asn Thr Pro Glu Gln Lys Ala Ala Leu Lys Ala Glu
325          330          335
Gln Ala Thr Glu Phe Tyr Val His Thr Pro Met Val Gln Phe Tyr Glu
340          345          350
Thr Leu Gly Lys Asp Arg Ile Leu Glu Leu Met Gly Ala Gly Thr Leu
355          360          365
Asn Lys Glu Leu Leu Asn Asp Asn His Ala Lys Ser Leu Glu Gly Lys
370          375          380
Asn Arg Ser Val Glu Asp Ser Tyr Asn Gln Leu Phe Ser Val Ile Glu
385          390          395          400

```

-continued

---

Gln Val Arg Ala Gln Ser Glu Asp Ile Ser Thr Val Pro Ile His Tyr  
 405 410 415  
 Ala Tyr Asn Met Thr Arg Val Gly Arg Met Gln Met Leu Gly Lys Tyr  
 420 425 430  
 Asn Pro Gln Ser Ala Lys Leu Val Arg Glu Ala Ile Leu Pro Thr Lys  
 435 440 445  
 Ala Thr Leu Asp Leu Ser Asn Gln Asn Asn Glu Asp Phe Ser Ala Phe  
 450 455 460  
 Gln Leu Gly Leu Ala Gln Ala Leu Asp Ile Lys Val His Thr Met Thr  
 465 470 475 480  
 Arg Glu Val Met Ser Asp Glu Leu Thr Lys Leu Leu Glu Gly Asn Leu  
 485 490 495  
 Lys Pro Ala Ile Asp Met Met Val Glu Phe Asn Thr Thr Gly Ser Leu  
 500 505 510  
 Pro Glu Asn Ala Val Asp Val Leu Asn Thr Ala Leu Gly Asp Arg Lys  
 515 520 525  
 Ser Phe Val Ala Leu Met Ala Leu Met Glu Tyr Ser Arg Tyr Leu Val  
 530 535 540  
 Ala Glu Asp Lys Ser Ala Phe Val Thr Pro Leu Tyr Val Glu Ala Asp  
 545 550 555 560  
 Gly Val Thr Asn Gly Pro Ile Asn Ala Met Met Leu Met Thr Gly Gly  
 565 570 575  
 Leu Phe Thr Pro Asp Trp Ile Arg Asn Ile Ala Lys Gly Gly Leu Phe  
 580 585 590  
 Ile Gly Ser Pro Asn Lys Thr Met Asn Glu His Arg Ser Thr Ala Asp  
 595 600 605  
 Asn Asn Asp Leu Tyr Gln Ala Ser Thr Asn Ala Leu Met Glu Ser Leu  
 610 615 620  
 Gly Lys Leu Arg Ser Asn Tyr Ala Ser Asn Met Pro Ile Gln Ser Gln  
 625 630 635 640  
 Ile Asp Ser Leu Leu Ser Leu Met Asp Leu Phe Leu Pro Asp Ile Asn  
 645 650 655  
 Leu Gly Glu Asn Gly Ala Leu Glu Leu Lys Arg Gly Ile Ala Lys Asn  
 660 665 670  
 Pro Leu Thr Ile Thr Ile Tyr Gly Ser Gly Ala Arg Gly Ile Ala Gly  
 675 680 685  
 Lys Leu Val Ser Ser Val Thr Asp Ala Ile Tyr Glu Arg Met Ser Asp  
 690 695 700  
 Val Leu Lys Ala Arg Ala Lys Asp Pro Asn Ile Ser Ala Ala Met Ala  
 705 710 715 720  
 Met Phe Gly Lys Gln Ala Ala Ser Glu Ala His Ala Glu Glu Leu Leu  
 725 730 735  
 Ala Arg Phe Leu Lys Asp Met Glu Thr Leu Thr Ser Thr Val Pro Val  
 740 745 750  
 Lys Arg Lys Gly Val Leu Glu Leu Gln Ser Thr Gly Thr Gly Ala Lys  
 755 760 765  
 Gly Lys Ile Asn Pro Lys Thr Tyr Thr Ile Lys Gly Glu Gln Leu Lys  
 770 775 780  
 Ala Leu Gln Glu Asn Met Leu His Phe Phe Val Glu Pro Leu Arg Asn  
 785 790 795 800  
 Gly Ile Thr Gln Thr Val Gly Glu Ser Leu Val Tyr Ser Thr Glu Gln  
 805 810 815

-continued

Leu	Gln	Lys	Ala	Thr	Gln	Ile	Gln	Ser	Val	Val	Leu	Glu	Asp	Met	Phe
			820					825					830		
Lys	Gln	Arg	Val	Gln	Glu	Lys	Leu	Ala	Glu	Lys	Ala	Lys	Asp	Pro	Thr
		835					840					845			
Trp	Lys	Lys	Gly	Asp	Phe	Leu	Thr	Gln	Lys	Glu	Leu	Asn	Asp	Ile	Gln
	850					855					860				
Ala	Ser	Leu	Asn	Asn	Leu	Ala	Pro	Met	Ile	Glu	Thr	Gly	Ser	Gln	Thr
865					870					875					880
Phe	Tyr	Ile	Ala	Gly	Ser	Glu	Asn	Ala	Glu	Val	Ala	Asn	Gln	Val	Leu
				885					890					895	
Ala	Thr	Asn	Leu	Asp	Asp	Arg	Met	Arg	Val	Pro	Met	Ser	Ile	Tyr	Ala
			900					905					910		
Pro	Ala	Gln	Ala	Gly	Val	Ala	Gly	Ile	Pro	Phe	Met	Thr	Ile	Gly	Thr
		915					920					925			
Gly	Asp	Gly	Met	Met	Met	Gln	Thr	Leu	Ser	Thr	Met	Lys	Gly	Ala	Pro
	930					935					940				
Lys	Asn	Thr	Leu	Lys	Ile	Phe	Asp	Gly	Met	Asn	Ile	Gly	Leu	Asn	Asp
945					950					955					960
Ile	Thr	Asp	Ala	Ser	Arg	Lys	Ala	Asn	Glu	Ala	Val	Tyr	Thr	Ser	Trp
				965					970					975	
Gln	Gly	Asn	Pro	Ile	Lys	Asn	Val	Tyr	Glu	Ser	Tyr	Ala	Lys	Phe	Met
			980					985					990		
Lys	Asn	Val	Asp	Phe	Ser	Lys	Leu	Ser	Pro	Glu	Ala	Leu	Glu	Ala	Ile
		995					1000					1005			
Gly	Lys	Ser	Ala	Leu	Glu	Tyr	Asp	Gln	Arg	Glu	Asn	Ala	Thr	Val	Asp
	1010					1015					1020				
Asp	Ile	Ala	Asn	Ala	Ala	Ser	Leu	Ile	Glu	Arg	Asn	Leu	Arg	Asn	Ile
1025					1030					1035					1040
Ala	Leu	Gly	Val	Asp	Ile	Arg	His	Lys	Val	Leu	Asp	Lys	Val	Asn	Leu
				1045					1050					1055	
Ser	Ile	Asp	Gln	Met	Ala	Ala	Val	Gly	Ala	Pro	Tyr	Gln	Asn	Asn	Gly
		1060						1065					1070		
Lys	Ile	Asp	Leu	Ser	Asn	Met	Thr	Pro	Glu	Gln	Gln	Ala	Asp	Glu	Leu
	1075						1080					1085			
Asn	Lys	Leu	Phe	Arg	Glu	Glu	Leu	Glu	Ala	Arg	Lys	Gln	Lys	Val	Ala
	1090					1095					1100				

Lys Ala Arg  
1105

<210> SEQ ID NO 5

<211> LENGTH: 3432

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer

<400> SEQUENCE: 5

atgggggggtt	ctcatcatca	tcatcatcat	ggatgggcta	gcatgactgg	tggacagcaa	60
atgggtcggg	atctgtacga	cgatgacgat	aaggatccga	gctcgagatc	tgaagtaca	120
gttacagaag	aattaaaga	aggtattgat	gctgtttacc	cttcattggt	aggtactgct	180
gattctaaag	cagaggggat	taagaactat	ttcaaattgt	cctttacctt	accagaagaa	240
cagaaatccc	gtactgttgg	ttcagaagca	cctctaaaag	atgtagccca	agctctgtct	300
tctcgtgctc	gttatgaact	ctttactgag	aaagaaactg	ctaaccctgc	ttttaatggg	360

-continued

---

gaagttatta agcgatacaa agaactcatg gaacatgggg aaggtattgc tgatattcct	420
cgctcccgtc tggctaagtt cottaacact aaggatgttg gtaaacgttt tgetcaaggt	480
acagaagcca accgttgggt aggtggtaag ttacttaaca ttgttgagca ggatggggat	540
accttaagt acaacgaaca attgtacag actgctgtat tagcaggtct tcaatggaga	600
cttactgcta ccagcaatac tgctatcaaa gatgcaaaag atgttgctgc tattactggt	660
attgaccaag ctctgctgcc agaaggttta gtagagcaat ttgatactgg tatgacactc	720
actgaagcag ttagttccct ggctcagaaa attgagtctt actggggatt atctcgtaat	780
ccaaatgctc cattgggcta taccaaaaggc atccctacag caatggctgc tgaattctg	840
gctgcatttg tagagtctac tgatgttgta gagaacatcg tggatatgtc agaaattgac	900
ccagataaca agaagactat tggctgttac accattactg aactggattc cttcgacca	960
attaataget tccctactgc tattgaagaa gctgttttag tgaatcctac agagaagatg	1020
ttctttggtg atgacattcc tcctgtagct aatactcagc ttcgtaaccg tgctgtcgt	1080
aatactccag aacagaaggc tgcattgaaa gcagagcagg ctacagagtt ctatgtacac	1140
acccaatgg ttcaattcta tgagacgtta ggtaagacc gtattctcga actgatgggt	1200
gctggtaactc tgaataaaga gttacttaat gataaccatg ctaaatctct ggaaggtaa	1260
aaccgttcag tagaggactc ttacaaccaa ctgttctcgg tcattgagca ggtaagagca	1320
cagagcgaag acatctctac tgtacctatt cactatgcat acaatatgac ccgtgttgg	1380
cgtagcaga tgttaggtaa atacaatcct caatcagcca aactggttcg tgaggccatc	1440
ttacctacta aagctacttt ggatttatcg aaccagaaca atgaagactt ctctgcatc	1500
cagttaggtc tggctcagcg attggacatt aaagtccata ctatgactcg tgaggttatg	1560
tctgacgagt tgactaaatt actggaaggt aatctgaaac cagccattga tatgatggt	1620
gagtttaata cactgggttc cttaccagaa aacgcagttg atgttctgaa tacagcatta	1680
ggagatagga agtcattcgt agcattgatg gctcttatgg agtattcccg ttacttagta	1740
gcagaggata aatctgcatt tgtaactcca ctgtatgtag aagcagatgg tgttactaat	1800
ggtccaatca atgccatgat gctaatagaca ggcggtctgt ttactcctga ctggattcgt	1860
aatattgcca aagggggctt gttcattggt tctccaaata agaccatgaa tgagcatcgc	1920
tctactgctg acaataatga tttatatcaa gcatccacta atgctttgat ggaatcgttg	1980
ggtaagttac gtagtaacta tgccctotaat atgctattc agtctcagat agacagctt	2040
ctttctctga tggatttgtt tttaccggat attaactctg gtgagaatgg tgctttagaa	2100
cttaaacgtg gtattgctaa gaaccactg actattacca tctatggttc tggtgctcgt	2160
ggtattgcag gtaagctggt tagttctggt actgatgcca tctatgagcg tatgtctgat	2220
gtactgaaaag ctctgctaa agaccctaat atctctgctg ctatggcaat gtttgtaag	2280
caagctgctt cagaagcaca tgctgaagaa cttcttgccc gtttctgaa agatatggaa	2340
acactgactt ctactgttcc tgttaaacgt aaaggtgtac tggaaactaca atccacaggt	2400
acaggagcca aaggaanaat caatcctaag acctatacca ttaagggcga gcaactgaag	2460
gcacttcagg aaaatatgct gcacttcttt gtagaaccac tacgtaatgg tattactcag	2520
actgtaggty aaagtctggt gtactctact gaacaattac agaaagctac tcagattcaa	2580
tctgtagtgc tggaaagatg gttcaaacag cgagtacaag agaagctggc agagaaggct	2640
aaagacccaa catggaagaa aggtgatttc cttactcaga aagaactgaa tgatattcag	2700

-continued

---

```

gcttctctga ataaacttagc ccctatgatt gagactgggt ctcagacttt ctacattgct 2760
ggttcagaaa atgcagaagt agcaaatcag gtattagcta ctaacctga tgaccgtatg 2820
cgtgtaccaa tgagtatcta tgctccagca caggccgggt tagcaggat tccatttatg 2880
actattggta ctgggtgatgg catgatgatg caaactcttt ccactatgaa aggtgcacca 2940
aagaatcccc tcaaaatctt tgatggatg aacattgggt tgaatgacat cactgatgcc 3000
agtcgtaaag ctaatgaagc tgtttacct tcttggcagg gtaaccctat taagaatgtt 3060
tatgaatcat atgctaagtt catgaagaat gtagatttca gcaagctgtc ccctgaagca 3120
ttggaagcaa ttggtaaate tgctctggaa tatgaccaac gtgagaatgc tactgtagat 3180
gatattgcta acgctgcac tctgattgaa cgtaacttac gtaatattgc actgggtgta 3240
gatattcgtc ataaggtgct ggataaggt aatctgtcca ttgaccagat ggctgctgta 3300
gggtgctcct atcagaacaa cggtaagatt gacctcagca atatgacccc tgaacaacag 3360
gctgatgaac tgaataaaact tttccgtgaa gagttagaag cccgtaaaca aaaagtctgt 3420
aaggctaggt aa 3432

```

```

<210> SEQ ID NO 6
<211> LENGTH: 1143
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Peptide

```

```

<400> SEQUENCE: 6

```

```

Met Gly Gly Ser His His His His His Gly Met Ala Ser Met Thr
 1           5           10          15
Gly Gly Gln Gln Met Gly Arg Asp Leu Tyr Asp Asp Asp Lys Asp
 20          25          30
Pro Ser Ser Arg Ser Glu Ser Thr Val Thr Glu Glu Leu Lys Glu Gly
 35          40          45
Ile Asp Ala Val Tyr Pro Ser Leu Val Gly Thr Ala Asp Ser Lys Ala
 50          55          60
Glu Gly Ile Lys Asn Tyr Phe Lys Leu Ser Phe Thr Leu Pro Glu Glu
 65          70          75          80
Gln Lys Ser Arg Thr Val Gly Ser Glu Ala Pro Leu Lys Asp Val Ala
 85          90          95
Gln Ala Leu Ser Ser Arg Ala Arg Tyr Glu Leu Phe Thr Glu Lys Glu
100         105         110
Thr Ala Asn Pro Ala Phe Asn Gly Glu Val Ile Lys Arg Tyr Lys Glu
115         120         125
Leu Met Glu His Gly Glu Gly Ile Ala Asp Ile Leu Arg Ser Arg Leu
130         135         140
Ala Lys Phe Leu Asn Thr Lys Asp Val Gly Lys Arg Phe Ala Gln Gly
145         150         155         160
Thr Glu Ala Asn Arg Trp Val Gly Gly Lys Leu Leu Asn Ile Val Glu
165         170         175
Gln Asp Gly Asp Thr Phe Lys Tyr Asn Glu Gln Leu Leu Gln Thr Ala
180         185         190
Val Leu Ala Gly Leu Gln Trp Arg Leu Thr Ala Thr Ser Asn Thr Ala
195         200         205
Ile Lys Asp Ala Lys Asp Val Ala Ala Ile Thr Gly Ile Asp Gln Ala
210         215         220

```

-continued

---

Leu Leu Pro Glu Gly Leu Val Glu Gln Phe Asp Thr Gly Met Thr Leu  
 225 230 235 240  
 Thr Glu Ala Val Ser Ser Leu Ala Gln Lys Ile Glu Ser Tyr Trp Gly  
 245 250 255  
 Leu Ser Arg Asn Pro Asn Ala Pro Leu Gly Tyr Thr Lys Gly Ile Pro  
 260 265 270  
 Thr Ala Met Ala Ala Glu Ile Leu Ala Ala Phe Val Glu Ser Thr Asp  
 275 280 285  
 Val Val Glu Asn Ile Val Asp Met Ser Glu Ile Asp Pro Asp Asn Lys  
 290 295 300  
 Lys Thr Ile Gly Leu Tyr Thr Ile Thr Glu Leu Asp Ser Phe Asp Pro  
 305 310 315 320  
 Ile Asn Ser Phe Pro Thr Ala Ile Glu Glu Ala Val Leu Val Asn Pro  
 325 330 335  
 Thr Glu Lys Met Phe Phe Gly Asp Asp Ile Pro Pro Val Ala Asn Thr  
 340 345 350  
 Gln Leu Arg Asn Pro Ala Val Arg Asn Thr Pro Glu Gln Lys Ala Ala  
 355 360 365  
 Leu Lys Ala Glu Gln Ala Thr Glu Phe Tyr Val His Thr Pro Met Val  
 370 375 380  
 Gln Phe Tyr Glu Thr Leu Gly Lys Asp Arg Ile Leu Glu Leu Met Gly  
 385 390 395 400  
 Ala Gly Thr Leu Asn Lys Glu Leu Leu Asn Asp Asn His Ala Lys Ser  
 405 410 415  
 Leu Glu Gly Lys Asn Arg Ser Val Glu Asp Ser Tyr Asn Gln Leu Phe  
 420 425 430  
 Ser Val Ile Glu Gln Val Arg Ala Gln Ser Glu Asp Ile Ser Thr Val  
 435 440 445  
 Pro Ile His Tyr Ala Tyr Asn Met Thr Arg Val Gly Arg Met Gln Met  
 450 455 460  
 Leu Gly Lys Tyr Asn Pro Gln Ser Ala Lys Leu Val Arg Glu Ala Ile  
 465 470 475 480  
 Leu Pro Thr Lys Ala Thr Leu Asp Leu Ser Asn Gln Asn Asn Glu Asp  
 485 490 495  
 Phe Ser Ala Phe Gln Leu Gly Leu Ala Gln Ala Leu Asp Ile Lys Val  
 500 505 510  
 His Thr Met Thr Arg Glu Val Met Ser Asp Glu Leu Thr Lys Leu Leu  
 515 520 525  
 Glu Gly Asn Leu Lys Pro Ala Ile Asp Met Met Val Glu Phe Asn Thr  
 530 535 540  
 Thr Gly Ser Leu Pro Glu Asn Ala Val Asp Val Leu Asn Thr Ala Leu  
 545 550 555 560  
 Gly Asp Arg Lys Ser Phe Val Ala Leu Met Ala Leu Met Glu Tyr Ser  
 565 570 575  
 Arg Tyr Leu Val Ala Glu Asp Lys Ser Ala Phe Val Thr Pro Leu Tyr  
 580 585 590  
 Val Glu Ala Asp Gly Val Thr Asn Gly Pro Ile Asn Ala Met Met Leu  
 595 600 605  
 Met Thr Gly Gly Leu Phe Thr Pro Asp Trp Ile Arg Asn Ile Ala Lys  
 610 615 620  
 Gly Gly Leu Phe Ile Gly Ser Pro Asn Lys Thr Met Asn Glu His Arg  
 625 630 635 640  
 Ser Thr Ala Asp Asn Asn Asp Leu Tyr Gln Ala Ser Thr Asn Ala Leu

-continued

645					650					655					
Met	Glu	Ser	Leu	Gly	Lys	Leu	Arg	Ser	Asn	Tyr	Ala	Ser	Asn	Met	Pro
			660					665					670		
Ile	Gln	Ser	Gln	Ile	Asp	Ser	Leu	Leu	Ser	Leu	Met	Asp	Leu	Phe	Leu
		675					680					685			
Pro	Asp	Ile	Asn	Leu	Gly	Glu	Asn	Gly	Ala	Leu	Glu	Leu	Lys	Arg	Gly
	690					695					700				
Ile	Ala	Lys	Asn	Pro	Leu	Thr	Ile	Thr	Ile	Tyr	Gly	Ser	Gly	Ala	Arg
	705					710					715				720
Gly	Ile	Ala	Gly	Lys	Leu	Val	Ser	Ser	Val	Thr	Asp	Ala	Ile	Tyr	Glu
				725					730					735	
Arg	Met	Ser	Asp	Val	Leu	Lys	Ala	Arg	Ala	Lys	Asp	Pro	Asn	Ile	Ser
			740					745					750		
Ala	Ala	Met	Ala	Met	Phe	Gly	Lys	Gln	Ala	Ala	Ser	Glu	Ala	His	Ala
		755					760					765			
Glu	Glu	Leu	Leu	Ala	Arg	Phe	Leu	Lys	Asp	Met	Glu	Thr	Leu	Thr	Ser
		770				775					780				
Thr	Val	Pro	Val	Lys	Arg	Lys	Gly	Val	Leu	Glu	Leu	Gln	Ser	Thr	Gly
	785					790					795				800
Thr	Gly	Ala	Lys	Gly	Lys	Ile	Asn	Pro	Lys	Thr	Tyr	Thr	Ile	Lys	Gly
				805					810					815	
Glu	Gln	Leu	Lys	Ala	Leu	Gln	Glu	Asn	Met	Leu	His	Phe	Phe	Val	Glu
			820					825					830		
Pro	Leu	Arg	Asn	Gly	Ile	Thr	Gln	Thr	Val	Gly	Glu	Ser	Leu	Val	Tyr
		835					840					845			
Ser	Thr	Glu	Gln	Leu	Gln	Lys	Ala	Thr	Gln	Ile	Gln	Ser	Val	Val	Leu
	850					855					860				
Glu	Asp	Met	Phe	Lys	Gln	Arg	Val	Gln	Glu	Lys	Leu	Ala	Glu	Lys	Ala
	865					870					875				880
Lys	Asp	Pro	Thr	Trp	Lys	Lys	Gly	Asp	Phe	Leu	Thr	Gln	Lys	Glu	Leu
				885					890					895	
Asn	Asp	Ile	Gln	Ala	Ser	Leu	Asn	Asn	Leu	Ala	Pro	Met	Ile	Glu	Thr
			900					905					910		
Gly	Ser	Gln	Thr	Phe	Tyr	Ile	Ala	Gly	Ser	Glu	Asn	Ala	Glu	Val	Ala
		915					920					925			
Asn	Gln	Val	Leu	Ala	Thr	Asn	Leu	Asp	Asp	Arg	Met	Arg	Val	Pro	Met
	930					935					940				
Ser	Ile	Tyr	Ala	Pro	Ala	Gln	Ala	Gly	Val	Ala	Gly	Ile	Pro	Phe	Met
	945					950					955				960
Thr	Ile	Gly	Thr	Gly	Asp	Gly	Met	Met	Met	Gln	Thr	Leu	Ser	Thr	Met
				965					970					975	
Lys	Gly	Ala	Pro	Lys	Asn	Thr	Leu	Lys	Ile	Phe	Asp	Gly	Met	Asn	Ile
			980					985					990		
Gly	Leu	Asn	Asp	Ile	Thr	Asp	Ala	Ser	Arg	Lys	Ala	Asn	Glu	Ala	Val
		995					1000					1005			
Tyr	Thr	Ser	Trp	Gln	Gly	Asn	Pro	Ile	Lys	Asn	Val	Tyr	Glu	Ser	Tyr
	1010					1015					1020				
Ala	Lys	Phe	Met	Lys	Asn	Val	Asp	Phe	Ser	Lys	Leu	Ser	Pro	Glu	Ala
	1025					1030					1035				1040
Leu	Glu	Ala	Ile	Gly	Lys	Ser	Ala	Leu	Glu	Tyr	Asp	Gln	Arg	Glu	Asn
				1045					1050					1055	
Ala	Thr	Val	Asp	Asp	Ile	Ala	Asn	Ala	Ala	Ser	Leu	Ile	Glu	Arg	Asn
			1060				1065						1070		

-continued

Leu Arg Asn Ile Ala Leu Gly Val Asp Ile Arg His Lys Val Leu Asp  
 1075 1080 1085

Lys Val Asn Leu Ser Ile Asp Gln Met Ala Ala Val Gly Ala Pro Tyr  
 1090 1095 1100

Gln Asn Asn Gly Lys Ile Asp Leu Ser Asn Met Thr Pro Glu Gln Gln  
 1105 1110 1115 1120

Ala Asp Glu Leu Asn Lys Leu Phe Arg Glu Glu Leu Glu Ala Arg Lys  
 1125 1130 1135

Gln Lys Val Ala Lys Ala Arg  
 1140

<210> SEQ ID NO 7

<211> LENGTH: 3432

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 7

```

atgggggggtt ctcacatca tcatcatcat ggtatggcta gcatgactgg tggacagcaa    60
atgggtcggg atctgtacga cgatgacgat aaggatccga gctcgagatc tgaagtaca    120
gttacagaag aattaaaga aggtattgat gctgtttacc cttcattggt aggtactgct    180
gattctaaag cagaggggat taagaactat ttcaaattgt cctttacctt accagaagaa    240
cagaaatccc gtactgttgg ttcagaagca cctctaaaag atgtagccca agctctgtct    300
tctcgtgctc gttatgaact ctttactgag aaagaaactg ctaaccctgc ttttaatggg    360
gaagttatta agcgatacaa agaactcatg gaacatgggg aaggtattgc tgaattctt    420
cgctcccgtc tggctaagtt ccttaacct aaggatgttg gtaaactgtt tgctcaaggt    480
acagaagcca accgttgggt aggtggtaag ttacttaaca ttgttgagca ggtggggat    540
acctttaagt acaacgaaca attgctacag actgctgtat tagcaggctt tcaatggaga    600
cttactgcta ccagcaatac tgctatcaaa gatgcaaaag atgttgctgc tattactggt    660
attgaccaag ctctgctgcc agaaggttta gtagagcaat ttgatactgg tatgacactc    720
actgaagcag ttagttccct ggctcagaaa attgagtctt actggggatt atctcgtaat    780
ccaaatgctc cattgggcta taccaaaggc atccctacag caatggctgc tgaattctg    840
gtgcatcttg tagagtctac tgatgttga gagaacatcg tggatatgtc agaaattgac    900
ccagataaca agaagactat tggctgttac accattactg aactggattc cttcgacca    960
attaatagct tcctactgc tattgaagaa gctgttttag tgaatcctac agagaagatg   1020
ttctttgggt atgacattcc tcctgtagct aatactcagc ttcgtaaccg tgctgttcgt   1080
aatactccag aacagaaggc tgcattgaaa gcagagcagg ctacagagtt ctatgtacac   1140
acccaatgg ttcaattcta tgagacgta ggtaaagacc gtattctcga actgatgggt   1200
gctggctact tgaataaaga gttacttaat gataacctg ctaaatctct ggaaggtaag   1260
aacggttcag tagaggactc ttacaaccaa ctgttctccg tcattgagca ggtaagagca   1320
cagagcgaag acatctctac tgtacctatt cactatgcat acaatatgac ccgtgttggt   1380
cgtatgcaga tgttaggtaa atacaatcct caatcagcca aactggttcg tgaggccatc   1440
ttaoctacta aagctacttt ggatttatcg aaccagaaca atgaagactt ctctgcattc   1500
cagttaggtc tggctcaggc attggacatt aaagtccata ctatgactcg tgaggttatg   1560

```

-continued

---

```

tctgacgagt tgactaaatt actggaaggt aatctgaaac cagccattga tatgatggtt 1620
gagtttaata coactggttc cttaccagaa aacgcagttg atgttctgaa tacagcatta 1680
ggagatagga agtcattcgt agcattgatg gctcttatgg agtattcccg ttacttagta 1740
gcagaggata aatctgcatt tgtaactcca ctgtatgtag aagcagatgg tgttactaat 1800
ggccaatca atgccatgat gctaatagaca ggcggctctgt ttactcctga ctggattcgt 1860
aatattgcca aagggggcct gttcattggg tctccaaata agaccatgaa tgagcatcgc 1920
tctactgctg acaataatga tttatatcaa gcattcacta atgctttgat ggaatcgttg 1980
ggtaagttac gtagtaacta tgcctctaat atgcctattc agtctcagat agacagtctt 2040
ctttctctga tggatttgtt tttaccggat attaactcttg gtgagaatgg tgctttagaa 2100
cttaaacgtg gtattgctaa gaaccactg actattacca tcttcggttc tgggtctcgt 2160
ggtattgcag gtaagctggg tagttctgtt actgatgcca tctatgagcg tatgtctgat 2220
gtactgaaag ctcgtgctaa agacccaaat atctctgctg ctatggcaat gtttgtaag 2280
caagctgctt cagaagcaca tgctgaagaa cttcttgccc gtttctgaa agatatggaa 2340
acactgactt ctactgttcc tgttaaactg aaaggtgtac tggaaactaca atccacaggt 2400
acaggagcca aagggaaaat caatcctaag acctatacca ttaagggcga gcaactgaag 2460
gcacttcagg aaaatagct gcacttcttt gtagaaccac tacgtaatgg tattactcag 2520
actgtagggtg aaagtctggt gtactctact gaacaattac agaaagctac tcagattcaa 2580
tctgtagtgc tggaaagatat gttcaaacag cgagtacaag agaagctggc agagaaggt 2640
aaagacccaa catggaagaa aggtgatctt cttactcaga aagaactgaa tgatattcag 2700
gcttctctga ataacttagc ccctatgatt gagactgggt ctcagacttt ctacattgct 2760
ggttcagaaa atgcagaagt agcaaatcag gtattagcta ctaaccttga tgaccgtatg 2820
cgtgtaccaa tgagtatcta tgctccagca caggccggtg tagcaggtat tccatttatg 2880
actattggta ctgggtgatg catgatgatg caaactcttt ccactatgaa aggtgcacca 2940
aagaatacce tcaaaatctt tgatgggatg aacattggtt tgaatgacat cactgatgcc 3000
agtcgtaaag ctaatgaagc tgtttacact tcttggcagg gtaacctat taagaatggt 3060
tatgaatcat atgctaagtt catgaagaat gtagatttca gcaagctgtc ccctgaagca 3120
ttggaagcaa ttggtaaatc tgctctggaa tatgaccaac gtgagaatgc tactgtagat 3180
gatattgcta acgctgcac cctgattgaa cgtaacttac gtaatattgc actgggtgta 3240
gatattcgtc ataaggtgct ggataaggtg aatctgtcca ttgaccagat ggctgctgta 3300
ggtgctcctt atcagaacaa cggtaagatt gacctcagca atatgacccc tgaacaacag 3360
gctgatgaac tgaataaact tttccgtgaa gagttagaag cccgtaaaaca aaaagtcgct 3420
aaggctaggt aa 3432

```

&lt;210&gt; SEQ ID NO 8

&lt;211&gt; LENGTH: 1143

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic Peptide

&lt;400&gt; SEQUENCE: 8

```

Met Gly Gly Ser His His His His His His Gly Met Ala Ser Met Thr
  1             5             10             15

```

```

Gly Gly Gln Gln Met Gly Arg Asp Leu Tyr Asp Asp Asp Asp Lys Asp

```

-continued

20					25					30					
Pro	Ser	Ser	Arg	Ser	Glu	Ser	Thr	Val	Thr	Glu	Glu	Leu	Lys	Glu	Gly
		35					40					45			
Ile	Asp	Ala	Val	Tyr	Pro	Ser	Leu	Val	Gly	Thr	Ala	Asp	Ser	Lys	Ala
	50					55					60				
Glu	Gly	Ile	Lys	Asn	Tyr	Phe	Lys	Leu	Ser	Phe	Thr	Leu	Pro	Glu	Glu
	65				70					75					80
Gln	Lys	Ser	Arg	Thr	Val	Gly	Ser	Glu	Ala	Pro	Leu	Lys	Asp	Val	Ala
				85					90					95	
Gln	Ala	Leu	Ser	Ser	Arg	Ala	Arg	Tyr	Glu	Leu	Phe	Thr	Glu	Lys	Glu
			100					105					110		
Thr	Ala	Asn	Pro	Ala	Phe	Asn	Gly	Glu	Val	Ile	Lys	Arg	Tyr	Lys	Glu
		115					120					125			
Leu	Met	Glu	His	Gly	Glu	Gly	Ile	Ala	Asp	Ile	Leu	Arg	Ser	Arg	Leu
	130					135					140				
Ala	Lys	Phe	Leu	Asn	Thr	Lys	Asp	Val	Gly	Lys	Arg	Phe	Ala	Gln	Gly
	145				150					155					160
Thr	Glu	Ala	Asn	Arg	Trp	Val	Gly	Gly	Lys	Leu	Leu	Asn	Ile	Val	Glu
				165					170					175	
Gln	Asp	Gly	Asp	Thr	Phe	Lys	Tyr	Asn	Glu	Gln	Leu	Leu	Gln	Thr	Ala
			180					185						190	
Val	Leu	Ala	Gly	Leu	Gln	Trp	Arg	Leu	Thr	Ala	Thr	Ser	Asn	Thr	Ala
		195					200						205		
Ile	Lys	Asp	Ala	Lys	Asp	Val	Ala	Ala	Ile	Thr	Gly	Ile	Asp	Gln	Ala
	210					215					220				
Leu	Leu	Pro	Glu	Gly	Leu	Val	Glu	Gln	Phe	Asp	Thr	Gly	Met	Thr	Leu
	225				230					235					240
Thr	Glu	Ala	Val	Ser	Ser	Leu	Ala	Gln	Lys	Ile	Glu	Ser	Tyr	Trp	Gly
				245					250					255	
Leu	Ser	Arg	Asn	Pro	Asn	Ala	Pro	Leu	Gly	Tyr	Thr	Lys	Gly	Ile	Pro
			260					265						270	
Thr	Ala	Met	Ala	Ala	Glu	Ile	Leu	Ala	Ala	Phe	Val	Glu	Ser	Thr	Asp
		275					280					285			
Val	Val	Glu	Asn	Ile	Val	Asp	Met	Ser	Glu	Ile	Asp	Pro	Asp	Asn	Lys
	290					295					300				
Lys	Thr	Ile	Gly	Leu	Tyr	Thr	Ile	Thr	Glu	Leu	Asp	Ser	Phe	Asp	Pro
	305				310					315					320
Ile	Asn	Ser	Phe	Pro	Thr	Ala	Ile	Glu	Glu	Ala	Val	Leu	Val	Asn	Pro
				325					330					335	
Thr	Glu	Lys	Met	Phe	Phe	Gly	Asp	Asp	Ile	Pro	Pro	Val	Ala	Asn	Thr
			340					345						350	
Gln	Leu	Arg	Asn	Pro	Ala	Val	Arg	Asn	Thr	Pro	Glu	Gln	Lys	Ala	Ala
		355					360						365		
Leu	Lys	Ala	Glu	Gln	Ala	Thr	Glu	Phe	Tyr	Val	His	Thr	Pro	Met	Val
	370					375						380			
Gln	Phe	Tyr	Glu	Thr	Leu	Gly	Lys	Asp	Arg	Ile	Leu	Glu	Leu	Met	Gly
	385				390					395					400
Ala	Gly	Thr	Leu	Asn	Lys	Glu	Leu	Leu	Asn	Asp	Asn	His	Ala	Lys	Ser
				405					410					415	
Leu	Glu	Gly	Lys	Asn	Arg	Ser	Val	Glu	Asp	Ser	Tyr	Asn	Gln	Leu	Phe
			420					425					430		
Ser	Val	Ile	Glu	Gln	Val	Arg	Ala	Gln	Ser	Glu	Asp	Ile	Ser	Thr	Val
		435					440						445		

-continued

---

Pro Ile His Tyr Ala Tyr Asn Met Thr Arg Val Gly Arg Met Gln Met  
 450 455 460

Leu Gly Lys Tyr Asn Pro Gln Ser Ala Lys Leu Val Arg Glu Ala Ile  
 465 470 475 480

Leu Pro Thr Lys Ala Thr Leu Asp Leu Ser Asn Gln Asn Asn Glu Asp  
 485 490 495

Phe Ser Ala Phe Gln Leu Gly Leu Ala Gln Ala Leu Asp Ile Lys Val  
 500 505 510

His Thr Met Thr Arg Glu Val Met Ser Asp Glu Leu Thr Lys Leu Leu  
 515 520 525

Glu Gly Asn Leu Lys Pro Ala Ile Asp Met Met Val Glu Phe Asn Thr  
 530 535 540

Thr Gly Ser Leu Pro Glu Asn Ala Val Asp Val Leu Asn Thr Ala Leu  
 545 550 555 560

Gly Asp Arg Lys Ser Phe Val Ala Leu Met Ala Leu Met Glu Tyr Ser  
 565 570 575

Arg Tyr Leu Val Ala Glu Asp Lys Ser Ala Phe Val Thr Pro Leu Tyr  
 580 585 590

Val Glu Ala Asp Gly Val Thr Asn Gly Pro Ile Asn Ala Met Met Leu  
 595 600 605

Met Thr Gly Gly Leu Phe Thr Pro Asp Trp Ile Arg Asn Ile Ala Lys  
 610 615 620

Gly Gly Leu Phe Ile Gly Ser Pro Asn Lys Thr Met Asn Glu His Arg  
 625 630 635 640

Ser Thr Ala Asp Asn Asn Asp Leu Tyr Gln Ala Ser Thr Asn Ala Leu  
 645 650 655

Met Glu Ser Leu Gly Lys Leu Arg Ser Asn Tyr Ala Ser Asn Met Pro  
 660 665 670

Ile Gln Ser Gln Ile Asp Ser Leu Leu Ser Leu Met Asp Leu Phe Leu  
 675 680 685

Pro Asp Ile Asn Leu Gly Glu Asn Gly Ala Leu Glu Leu Lys Arg Gly  
 690 695 700

Ile Ala Lys Asn Pro Leu Thr Ile Thr Ile Phe Gly Ser Gly Ala Arg  
 705 710 715 720

Gly Ile Ala Gly Lys Leu Val Ser Ser Val Thr Asp Ala Ile Tyr Glu  
 725 730 735

Arg Met Ser Asp Val Leu Lys Ala Arg Ala Lys Asp Pro Asn Ile Ser  
 740 745 750

Ala Ala Met Ala Met Phe Gly Lys Gln Ala Ala Ser Glu Ala His Ala  
 755 760 765

Glu Glu Leu Leu Ala Arg Phe Leu Lys Asp Met Glu Thr Leu Thr Ser  
 770 775 780

Thr Val Pro Val Lys Arg Lys Gly Val Leu Glu Leu Gln Ser Thr Gly  
 785 790 795 800

Thr Gly Ala Lys Gly Lys Ile Asn Pro Lys Thr Tyr Thr Ile Lys Gly  
 805 810 815

Glu Gln Leu Lys Ala Leu Gln Glu Asn Met Leu His Phe Phe Val Glu  
 820 825 830

Pro Leu Arg Asn Gly Ile Thr Gln Thr Val Gly Glu Ser Leu Val Tyr  
 835 840 845

Ser Thr Glu Gln Leu Gln Lys Ala Thr Gln Ile Gln Ser Val Val Leu  
 850 855 860

-continued

Glu Asp Met Phe Lys Gln Arg Val Gln Glu Lys Leu Ala Glu Lys Ala  
 865 870 875 880  
 Lys Asp Pro Thr Trp Lys Lys Gly Asp Phe Leu Thr Gln Lys Glu Leu  
 885 890 895  
 Asn Asp Ile Gln Ala Ser Leu Asn Asn Leu Ala Pro Met Ile Glu Thr  
 900 905 910  
 Gly Ser Gln Thr Phe Tyr Ile Ala Gly Ser Glu Asn Ala Glu Val Ala  
 915 920 925  
 Asn Gln Val Leu Ala Thr Asn Leu Asp Asp Arg Met Arg Val Pro Met  
 930 935 940  
 Ser Ile Tyr Ala Pro Ala Gln Ala Gly Val Ala Gly Ile Pro Phe Met  
 945 950 955 960  
 Thr Ile Gly Thr Gly Asp Gly Met Met Met Gln Thr Leu Ser Thr Met  
 965 970 975  
 Lys Gly Ala Pro Lys Asn Thr Leu Lys Ile Phe Asp Gly Met Asn Ile  
 980 985 990  
 Gly Leu Asn Asp Ile Thr Asp Ala Ser Arg Lys Ala Asn Glu Ala Val  
 995 1000 1005  
 Tyr Thr Ser Trp Gln Gly Asn Pro Ile Lys Asn Val Tyr Glu Ser Tyr  
 1010 1015 1020  
 Ala Lys Phe Met Lys Asn Val Asp Phe Ser Lys Leu Ser Pro Glu Ala  
 1025 1030 1035 1040  
 Leu Glu Ala Ile Gly Lys Ser Ala Leu Glu Tyr Asp Gln Arg Glu Asn  
 1045 1050 1055  
 Ala Thr Val Asp Asp Ile Ala Asn Ala Ala Ser Leu Ile Glu Arg Asn  
 1060 1065 1070  
 Leu Arg Asn Ile Ala Leu Gly Val Asp Ile Arg His Lys Val Leu Asp  
 1075 1080 1085  
 Lys Val Asn Leu Ser Ile Asp Gln Met Ala Ala Val Gly Ala Pro Tyr  
 1090 1095 1100  
 Gln Asn Asn Gly Lys Ile Asp Leu Ser Asn Met Thr Pro Glu Gln Gln  
 1105 1110 1115 1120  
 Ala Asp Glu Leu Asn Lys Leu Phe Arg Glu Glu Leu Glu Ala Arg Lys  
 1125 1130 1135  
 Gln Lys Val Ala Lys Ala Arg  
 1140

<210> SEQ ID NO 9  
 <211> LENGTH: 69  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 9

tcccagacaa aaggттаага ttccatacag gattggatgc attacttcat ccaaaagaag 60

cggagcttc 69

<210> SEQ ID NO 10  
 <211> LENGTH: 69  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 10

-continued

---

```
tgggagagaa aaggtaaga ttgatagag gattggatgg attagttgat ggaaaagaag 60
cggagcttc 69
```

```
<210> SEQ ID NO 11
<211> LENGTH: 69
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer
```

```
<400> SEQUENCE: 11
```

```
tcctctctt ttggtttgt tttcttctg gtttggtgc tttcttctt caaaagaag 60
cggagcttc 69
```

```
<210> SEQ ID NO 12
<211> LENGTH: 69
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer
```

```
<400> SEQUENCE: 12
```

```
tcccacaaa aaccttaaca tttcatacac cattccatcc attacttcat caaaagaag 60
cggagcttc 69
```

```
<210> SEQ ID NO 13
<211> LENGTH: 69
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer
```

```
<400> SEQUENCE: 13
```

```
accagacaa aaggaaaaga aaacaaacag gaaaggaagc aaaacaaca caaaagaag 60
cggagcttc 69
```

```
<210> SEQ ID NO 14
<211> LENGTH: 10617
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer
```

```
<400> SEQUENCE: 14
```

```
atggggggtt ctcatcatca tcatcatcat ggtatggcta gcatgactgg tggacagcaa 60
atgggtcggg atctgtacga cgatgacgat aaggatccga gctcgagatc tatgtcagta 120
ttgatagac tggctgggtt cgcagacagc gtaaccaatg caaagcaagt tgacgtctct 180
actgcaaccg cccagaagaa agctgaacaa ggtgtcacta ctctcttgt ttctctgat 240
gctgcttata aatgcaagc tgcccgtact ggtaatgttg gggctaagtc attgaacca 300
gggacagtgc aatcagattt catgaatctg accccaatgc aatcatgaa taagtatggg 360
gttgagcaag gttacaact tatcaatgct cgtgctgatg cagggaacca ggtattcaat 420
gattcagtta ctacaagaac tcctggggaa gaactggggg atattgctac tgggtttggc 480
cttggtttgy ttaataccct tgggggcatt ggtgctcttg gggcaggctt actcaacgat 540
```

-continued

---

gatgcaggtg	ctgttgttgc	tcaacaattg	agtaagttta	atgatgctgt	tcatgctacc	600
caaagccagc	cattacaaga	taaacgtaag	ctctttgctg	ctcgttaact	aatgaatgaa	660
gtagagagtg	aacgtcagta	tcaaacagat	aagaaagaag	gcactaatga	catagtagct	720
tccttatcta	aatttggacg	tgattttgta	ggttcaattg	agaatgctgc	tcaaactgac	780
tctattattt	ctgatgggtt	agcagaagg	gtaggttctc	tattaggtgc	tggtcctgta	840
ttaaggggtg	catctttact	gggtaaagca	gttgttccag	caaatactct	tcgtagtgct	900
gcattggctg	gtgctattga	tcaggtact	ggtactcagt	cactggctcg	tattgcctct	960
actgtaggta	gagctgcacc	gggtatggtt	gggtttggtg	caatggaagc	tggtggtgca	1020
taccaacaaa	ctgctgatga	aattatgaag	atgagtctta	aagacttaga	gaagtctcct	1080
gtttatcagc	aacatattaa	agatggtagt	tcccctgaac	aggctcgtcg	tcagactgca	1140
tctgaaactg	gtcttactgc	tgctgctatt	caattaccta	ttgctgctgc	aaccggctcct	1200
ctggatcccc	gttttgagat	ggctccttcc	cgtgctggct	ctttaggtgc	tgtaggtatg	1260
aaccttgccc	gtgaaacagt	ggaagaaggt	gttcaggggtg	ctacaggcca	actggctcag	1320
aatattgcac	agcaacaaaa	cattgataag	aaccaagacc	tgcttaaggg	tgtcgggtaca	1380
caggctgggt	taggtgctct	ttatggcttt	ggttctgctg	gtgttggtaca	ggctccggct	1440
gggtgctgctc	gtttagcagg	tgctgcaact	gctcctgtat	tgctgaccac	aatggctgggt	1500
gttaagctg	ctggtagtgt	agcaggtaag	gttgtttctc	ctattaagaa	tacttttagta	1560
gctcgtgggt	aacgggttat	gaagcagaat	gaagaagcat	ctcctgttgc	tgatgactat	1620
gttgcacagg	cagcacaaga	agctatggct	caagcaccag	aagcagaagt	tactattcgt	1680
gatgctggtg	aagcaactga	tgctactcca	gaacagaaag	ttgcagcaca	ccagtatggt	1740
tctgacttaa	tgaatgctac	tcgttttaat	cctgaaaatt	atcaggaagc	accagagcat	1800
attcgtaatg	ctgtagctgg	ttctactgac	caagtacagg	ttattcagaa	gtagcagac	1860
ttagttaaca	cattagatga	atctaactct	caagcactga	tggaagctgc	atcttatatg	1920
tatgatgctg	tttcagagtt	tgagcagttc	attaaccgtg	acctgctgc	actggatagc	1980
attcctaagg	attctccggc	tattgagtta	ctcaaccgtt	atacgaatct	gacagctaat	2040
attcagaaca	cacaaaaagt	aattggtgca	ctgaatgtta	ttaatcgaat	gattaatgaa	2100
tctgctcaga	atggttcttt	gaatgtgact	gaagaatcca	gtccacagga	aatgcagaac	2160
gtagcattag	ctgctgaagt	agcccctgaa	aagctcaatc	cagagtctgt	aatgtttggt	2220
cttaaacatg	ctgctgatgg	tcgtatataa	ctgaataatc	gccagattgc	tgccctccag	2280
aatgctgctg	caatcctgaa	gggggcacgg	gaatatgatg	cagaagctgc	ccgtcttggg	2340
ttacgtcctc	aagacattgt	gagtaaacag	atataaacgg	atgagagcag	aactcaggaa	2400
ggacaatact	ctgcgttgca	acatgogaat	aggattcgggt	ctgcgtataa	ctctggtaat	2460
ttcaggttgg	cctccgctta	cctgaacgac	tttatgcagt	tcgccagca	catgcagaat	2520
aaggttggag	cgttgaatga	gcactctggt	acggggaatg	cggtataaga	taagtctgtc	2580
cactaccaag	ctcttactgc	tgacagagaa	tgggttcgta	gccgtaccgg	attgggggtc	2640
aatccctatg	acactaagtc	ggttaaattt	gccagcaag	ttgctcttga	agcgaaaacg	2700
gtagcggata	ttgctaatagc	cctcgtctcg	gcttaccgg	aactgaaggt	cagtcatata	2760
aaagttactc	cattggatcc	acgtcttaac	gctcctgctg	ctgaggtgggt	caaggcattc	2820
cgtcaaggca	atcgagacgt	tgcttcttct	caaccgaaag	ctgactccgt	gaatcagggt	2880
aaagaaactc	ctgttataaa	acaggaacca	gttacatcta	ctgtacagac	taagactcct	2940

-continued

---

gtagtgaat	ctgttaaac	agaacctact	actaaagagt	ctagcccaca	ggctataaaa	3000
gaacctgtga	accagtctga	aaaacaggat	gttaacctta	ctaagagga	caacatcaag	3060
caacctactg	aatctgttaa	agaaactgaa	acttctacaa	aagaaagtac	agttacagaa	3120
gaattaaaag	aaggatttga	tgctgtttac	ccttcattgg	taggtactgc	tgattctaaa	3180
gcagagggta	ttaagaacta	tttcaaatg	tcctttacct	taccagaaga	acagaaatcc	3240
cgtactgttg	gttcagaagc	acctctaaaa	gatgtagccc	aagctctgtc	ttctcgtgct	3300
cgttatgaac	tctttactga	gaaagaaact	gctaacctcg	cttttaatgg	ggaagttatt	3360
aagcgataca	aagaactcat	ggaacatggg	gaaggatttg	ctgatattct	tcgctcccgt	3420
ctggctaagt	tccttaacac	taaggatggt	ggtaaacggt	ttgctcaagg	tacagaagcc	3480
aaccgttggg	taggtggtaa	gttacttaac	attgttgagc	aggatgggga	tacctttaag	3540
tacaacgaac	aattgctaca	gactgctgta	ttagcaggtc	ttcaatggag	acttactgct	3600
accagcaata	ctgctatcaa	agatgcaaaa	gatgttgctg	ctattactgg	tattgaccaa	3660
gctctgctgc	cagaaggttt	agtagagcaa	tttgatactg	gtatgacact	caactgaagca	3720
gttagttccc	tggtcagaa	aattgagtct	tactggggat	tatctcgtaa	tccaaatgct	3780
ccattgggct	ataccaaagg	catccctaca	gcaatggctg	ctgaaattct	ggctgcattt	3840
gtagagtcta	ctgatgttgt	agagaacatc	gtggatatgt	cagaaattga	cccagataac	3900
aagaagacta	ttggtctgta	caccattact	gaactggatt	ccttcgaccc	aattaatagc	3960
ttccctactg	ctattgaaga	agctgtttta	gtgaatccta	cagagaagat	gttctttggt	4020
gatgacattc	ctcctgtagc	taatactcag	cttcgtaacc	ctgctgttcg	taatactcca	4080
gaacagaagg	ctgcattgaa	agcagagcag	gctacagagt	tctatgtaca	cacccaatg	4140
gttcaattct	atgagacggt	aggtaaagac	cgtattctcg	aactgatggg	tgctggtagt	4200
ctgaataaag	agttacttaa	tgataacat	gctaaatctc	tggaaagtaa	gaaccgttca	4260
gtagaggact	cttacaacca	actgttctcc	gtcattgagc	aggtaagagc	acagagcgaa	4320
gacatctcta	ctgtacctat	tcactatgca	tacaatatga	cccgtgttgg	tcgtatgcag	4380
atgttaggta	aatacaatcc	tcaatcagcc	aaactggttc	gtgaggccat	cttacctact	4440
aaagctactt	tggatttate	gaaccagaac	aatgaagact	tctctgcatt	ccagttaggt	4500
ctggctcagg	cattggacat	taaagtccat	actatgactc	gtgaggttat	gtctgacgag	4560
ttgactaaat	tactggaagg	taactgaaa	ccagccattg	atatgatggg	tgagtttaat	4620
accactgggt	ccttaccaga	aaacgcagtt	gatgttctga	atacagcatt	aggagatagg	4680
aagtcattcg	tagcattgat	ggctcttatg	gagtattccc	gttacttagt	agcagaggat	4740
aaatctgcat	ttgtaactcc	actgtatgta	gaagcagatg	gtgttactaa	tggtccaatc	4800
aatgccatga	tgctaagac	aggcggctcg	tttactctcg	actggattcg	taatattgcc	4860
aaagggggct	tgttcattgg	ttctccaaat	aagaccatga	atgagcatcg	ctctactgct	4920
gacaataatg	atztatatca	agcatocact	aatgctttga	tggaaatcgt	gggtaagtta	4980
cgtagtaact	atgcctctaa	tatgcctatt	cagtctcaga	tagacagtct	tctttctctg	5040
atggatttgt	ttttaccgga	tattaatctt	ggtgagaatg	gtgctttaga	acttaaacgt	5100
ggtagtctga	agaaccact	gactattacc	atctatgggt	ctgggtgctcg	tggtattgca	5160
ggtaagctgg	ttagttctgt	tactgatgcc	atctatgagc	gtatgtctga	tgtactgaaa	5220
gctcgtgcta	aagaccctaaa	tatctctgct	gctatggcaa	tgtttggtaa	gcaagctgct	5280

-continued

---

tcagaagcac atgctgaaga acttcttgcc cgttctctga aagatatgga aacctgact	5340
tctactgttc ctgttaaacy taaaggtgta ctggaactac aatccacagg tacaggagcc	5400
aaaggaaaa tcaatcctaa gacctatacc attaagggcg agcaactgaa ggcacttcag	5460
gaaaatatgc tgcacttctt tgtagaacca ctacgtaatg gtattactca gactgtaggt	5520
gaaagtctgg tgtactctac tgaacaatta cagaaagcta ctcagattca atctgtagt	5580
ctggaagata tgttcaaaca gcgagtacaa gagaagctgg cagagaaggc taaagacca	5640
acatggaaga aaggtgattt ccttactcag aaagaactga atgatattca ggcttctctg	5700
aataacttag cccctatgat tgagactggg tctcagactt tctacattgc tggttcagaa	5760
aatgcagaag tagcaaatca ggtattagct actaaccttg atgaccgtat gcgtgtacca	5820
atgagtatct atgctccagc acaggccggt gtatcaggtg tccatttat gactattggg	5880
actggtgatg gcatgatgat gaaaactctt tccactatga aaggtgcacc aaagaatacc	5940
ctcaaaatct ttgatgggat gaacattggg ttgaatgaca tcaactgatgc cagtcgtaa	6000
gctaataagc ctgtttacac ttcttggcag ggtaacccta ttaagaatgt ttatgaatca	6060
tatgctaagt tcatgaagaa tgtagatttc agcaagctgt cccctgaagc attggaagca	6120
attggtaaat ctgctctgga atatgaccaa cgtgagaatg ctactgtaga tgatattgct	6180
aacgctgcat ctctgattga acgtaactta cgtaaatatg cactgggtgt agatattcgt	6240
cataaggtgc tggataaggt aaatctgtcc attgaccaga tggctgctgt aggtgctcct	6300
tatcagaaca acggtaagat tgacctcagc aatatgaccc ctgaacaaca ggctgatgaa	6360
ctgaataaac ttttccgtga agagttagaa gcccgtaaac aaaaagtcgc taaggctagg	6420
gctgaagtca aagaagaaac tgtttctgaa aaagaaccag tgaatccaga ctttggatg	6480
gtaggccggt agcataaggc atctgggtgt cgtatcctgt ctgctactgc tattcgtaat	6540
ctggctaaga ttagtaatct gccatctact caggcagcta ctcttgcgga gattcagaaa	6600
tcaactggcag ctaaagacta taagattatc tacggtacac ctactcaggt tgcagagtat	6660
gctcgtcaga agaattgtac tgaattgact tctcaggaaa tggagaagc tcaggcaggt	6720
aatatttatg gctggactaa cttcgtatgat aagaccattt atctgggttag cccatctatg	6780
gaaaccctca ttcattgaact ggttcattgc tctacctctg aggaagtta ttccttctat	6840
cagggtaatg aagtaagccc tacttctaag caggctattg agaacctga aggtctgatg	6900
gaacagttcc gttctctgga tatttccaaa gattctccag aatgagaga agcatatgct	6960
gatgctattg caactatcga aggtcatttg agtaatggat ttgttgaccc agctatctct	7020
aaagctgctg ctcttaatga gtttatggct tgggggtag ctaaccgtgc tcttgctgct	7080
aaacagaaga gaacatcttc actggttcaa atggtgaaag atgtttatca ggctattaag	7140
aaattgattt ggggacgtaa acaagctcct gcattgggag aagatatggt ctccaatctg	7200
ctgtttaact ctgcaattct gatgcgtagc caacctacaa ctcaggcagt agctaaagat	7260
ggcacactgt tccatagcaa agcatatggt aataatgaac gtctgtctca gttgaaccag	7320
actttcgata aactggtaac tgattacctt cgtactgacc cagttacaga agtagaacgt	7380
cgtggcaatg tggctaagc attaatgagt gctactcgac tggttcgtga tgttcagtct	7440
catggcttca atatgactgc tcaggaacag tctgtattcc agatggttac tgctgcatta	7500
gcaactgaag ctgcgattga cccacatgct atggctcgtg ctcaggaact ttatacccat	7560
gtaatgaaac accttacggt agagcatttc atggctgacc ctgatagtac taaccctgct	7620
gaccgttact atgctcaaca gaaatatgac accatctctg gtgctaactt ggttgaagta	7680

-continued

---

gatgccaag	gtagaacag	tctgttacct	acattcctgg	gtctggctat	ggtaaatgaa	7740
gaactacgtt	caatcattaa	agaaatgcct	gtacctaaag	cagataagaa	attaggggat	7800
gatatagata	ctctgcttac	caatgcaggt	actcaggtaa	tggaaatctct	gaaccgtcgt	7860
atggctgggtg	accagaaagc	tactaatgtt	caggacagta	ttgatgcttt	gtcagaaaca	7920
atcatggctg	ctgctttgaa	acgagagtcc	ttctatgatg	ctgtagcaac	ccctaccggg	7980
aacttcattg	accgtgctaa	tcagtagcgt	acggatagca	ttgaacgggt	atctgaaact	8040
gttattgaga	aggcagataa	ggtaattgct	aaccttcta	atatagctgc	taaagggtgt	8100
gctcatctgg	ctaaactgac	tgctgctatt	gcactgaaa	aacaggggtga	aatagtggct	8160
caggggtgta	tgactgctat	gaaccagggt	aaagtatggc	aacctttcca	tgacttagtt	8220
aatgacattg	ttggcgtgac	taagactaat	gccaatgtct	atgacttaat	caaattgggt	8280
aagagccaga	tttctcaaga	ccgtcagcaa	ttccgtgagc	atttacctac	agtcattgct	8340
ggtaagttct	ctcgtaaatt	gactgatacc	gaatggctcg	caatgcatac	tggtttaggt	8400
aaaaacagatt	tagctgttct	acgtgaaact	atgagcatgg	ctgaaattag	agatttactc	8460
tcttcatcca	agaaagtgaa	agatgaaatc	tctactctgg	aaaaagagat	tcagaaccaa	8520
gcaggtagaa	actggaatct	ggttcagaag	aaatctaagc	aactggctca	atacatgatt	8580
atgggggaag	taggtaataa	cctccttctg	aatgcccctg	ctattagtctg	tttggttaggt	8640
gaacgtatta	ctaatggctc	tgtggcagat	gtagctgcta	ttgataagct	cattactttg	8700
tactctctgg	aattgatgaa	taagtctgac	cgtgaccttt	tgtagaatt	ggctcaatca	8760
gaagtggaag	gtatggaggt	ctccattgct	tatatggttg	gtcaacgtac	tgaagagatg	8820
cgtaaagcta	aaggtgataa	ccgtactctg	ctgaatcact	ttaaaggcta	tatccctgta	8880
gagaaccagc	aaggtgtgaa	tttgattatt	gctgacgata	aagagtttgc	taagttaaat	8940
agccaatcct	ttactcgtat	tggctactat	caggggagca	ctggtttccg	tactggttct	9000
aaaggttatt	acttcagccc	agtagctgcc	cgtgcccctt	actctcaggg	tattcttcag	9060
aacgttcgta	atactgctgg	tgggtgtgat	attggctactg	gctttacggt	aggcactatg	9120
gttgctgggc	gtattactga	caaaccaacc	gtagagcgt	ttaccaaacg	tctggctaaa	9180
ggtagcgtg	ggcgtgaacc	actgatgcca	atataaaca	gcaaaggctca	ggtagttgct	9240
tatgaacaat	ccgttgaccc	taatatgttg	aagcacctaa	accaagacaa	tcactttgct	9300
aagatgggtg	gtgtatggcg	tggctcgtcag	gtggaagagg	ctaaagcaca	acgttttaat	9360
gacattctca	ttgagcaatt	acatgctatg	tatgagaaag	acattaaaga	ctccagtgct	9420
aataaatctc	aatatgtaaa	cctgttaggt	aaaattgatg	accagtagct	ggctgatgctg	9480
attaacctga	tgaacattga	gactcgtcat	aaggccgaag	aactcttcgg	taaagatgag	9540
ttatgggttc	gtagggat	gctgaatgat	gcacttggt	atcgtgctgc	atctattggt	9600
gatgtgtgga	ccggtaacct	tcggttgctca	cctagcacc	ttgatactgt	taagaagatg	9660
ttctcgggtg	cattcggtaa	taaggcatat	catgtagtaa	tgaatgctga	aaataccatt	9720
cagaacttag	tgaaggacgc	taagacagta	attgttgta	aatctgtgtg	agtaccggca	9780
gttaacttcc	ttgctaaccat	ctaccagatg	attggacgtg	gtgttcctgt	taaagatatt	9840
gctgtgaaca	ttctcgtgaa	gacgtcagag	attaatcagt	atattaaatc	tcgtttacgt	9900
cagattgatg	cggaagcaga	gctacgtgct	gctgaaggta	accctaactc	ggttcgtaaa	9960
cttaaaactg	agattcaatc	tattactgat	agtcacgtc	gtatgagtat	ctggcctttg	10020

-continued

---

```

attgaagcag gtgagttctc ttctattgct gatgctggta ttagtctgga tgacctgtta 10080
gtagctgaag gtaagattca tgagtacatg gaaaaacttg ctaataaact tccagaaaaa 10140
gtacgtaatg ctggccggtta cgctcttatt gctaaggaca ctgctctggt ccagggtatc 10200
cagaaaaacag tagagtattc agactttatt gctaaagcca tcacttatga tgatttagtg 10260
aaacgtaaga aaaaatcttc ttctgaagca ttaggtcagg taactgaaga gtttattaac 10320
tatgacagat tgccctggcg tttccgtggc tatatggaaa gtaggggtct gatgtgggtc 10380
tacaacttta aaattcgttc cattaagttt gctatgagca tgattagaaa caaccagta 10440
cattctctga ttgctacagt agtacctgct cctaccatgt ttggtaacgt aggtctacca 10500
attcaggaca acatgctaac catgctggct gaaggaagac tggattactc attaggcttc 10560
ggacaaggat taagagcacc taccctcaat ccttggttca accttactca ctaataa 10617

```

```

<210> SEQ ID NO 15
<211> LENGTH: 3537
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Peptide

```

```

<400> SEQUENCE: 15

```

```

Met Gly Gly Ser His His His His His Gly Met Ala Ser Met Thr
 1          5          10          15
Gly Gly Gln Gln Met Gly Arg Asp Leu Tyr Asp Asp Asp Asp Lys Asp
 20          25          30
Pro Ser Ser Arg Ser Met Ser Val Phe Asp Arg Leu Ala Gly Phe Ala
 35          40          45
Asp Ser Val Thr Asn Ala Lys Gln Val Asp Val Ser Thr Ala Thr Ala
 50          55          60
Gln Lys Lys Ala Glu Gln Gly Val Thr Thr Pro Leu Val Ser Pro Asp
 65          70          75          80
Ala Ala Tyr Gln Met Gln Ala Ala Arg Thr Gly Asn Val Gly Ala Asn
 85          90          95
Ala Phe Glu Pro Gly Thr Val Gln Ser Asp Phe Met Asn Leu Thr Pro
100          105          110
Met Gln Ile Met Asn Lys Tyr Gly Val Glu Gln Gly Leu Gln Leu Ile
115          120          125
Asn Ala Arg Ala Asp Ala Gly Asn Gln Val Phe Asn Asp Ser Val Thr
130          135          140
Thr Arg Thr Pro Gly Glu Leu Gly Asp Ile Ala Thr Gly Val Gly
145          150          155          160
Leu Gly Phe Val Asn Thr Leu Gly Gly Ile Gly Ala Leu Gly Ala Gly
165          170          175
Leu Leu Asn Asp Asp Ala Gly Ala Val Val Ala Gln Gln Leu Ser Lys
180          185          190
Phe Asn Asp Ala Val His Ala Thr Gln Ser Gln Ala Leu Gln Asp Lys
195          200          205
Arg Lys Leu Phe Ala Ala Arg Asn Leu Met Asn Glu Val Glu Ser Glu
210          215          220
Arg Gln Tyr Gln Thr Asp Lys Lys Glu Gly Thr Asn Asp Ile Val Ala
225          230          235          240
Ser Leu Ser Lys Phe Gly Arg Asp Phe Val Gly Ser Ile Glu Asn Ala
245          250          255

```



-continued

675					680					685					
Gly	Ala	Leu	Asn	Val	Ile	Asn	Arg	Met	Ile	Asn	Glu	Ser	Ala	Gln	Asn
690					695					700					
Gly	Ser	Leu	Asn	Val	Thr	Glu	Glu	Ser	Ser	Pro	Gln	Glu	Met	Gln	Asn
705					710					715					720
Val	Ala	Leu	Ala	Ala	Glu	Val	Ala	Pro	Glu	Lys	Leu	Asn	Pro	Glu	Ser
				725					730					735	
Val	Asn	Val	Val	Leu	Lys	His	Ala	Ala	Asp	Gly	Arg	Ile	Lys	Leu	Asn
			740						745				750		
Asn	Arg	Gln	Ile	Ala	Ala	Leu	Gln	Asn	Ala	Ala	Ala	Ile	Leu	Lys	Gly
		755					760					765			
Ala	Arg	Glu	Tyr	Asp	Ala	Glu	Ala	Ala	Arg	Leu	Gly	Leu	Arg	Pro	Gln
		770					775				780				
Asp	Ile	Val	Ser	Lys	Gln	Ile	Lys	Thr	Asp	Glu	Ser	Arg	Thr	Gln	Glu
785					790					795					800
Gly	Gln	Tyr	Ser	Ala	Leu	Gln	His	Ala	Asn	Arg	Ile	Arg	Ser	Ala	Tyr
				805					810					815	
Asn	Ser	Gly	Asn	Phe	Glu	Leu	Ala	Ser	Ala	Tyr	Leu	Asn	Asp	Phe	Met
			820						825					830	
Gln	Phe	Ala	Gln	His	Met	Gln	Asn	Lys	Val	Gly	Ala	Leu	Asn	Glu	His
		835					840						845		
Leu	Val	Thr	Gly	Asn	Ala	Asp	Lys	Asn	Lys	Ser	Val	His	Tyr	Gln	Ala
		850					855					860			
Leu	Thr	Ala	Asp	Arg	Glu	Trp	Val	Arg	Ser	Arg	Thr	Gly	Leu	Gly	Val
865					870					875					880
Asn	Pro	Tyr	Asp	Thr	Lys	Ser	Val	Lys	Phe	Ala	Gln	Gln	Val	Ala	Leu
				885					890					895	
Glu	Ala	Lys	Thr	Val	Ala	Asp	Ile	Ala	Asn	Ala	Leu	Ala	Ser	Ala	Tyr
			900						905					910	
Pro	Glu	Leu	Lys	Val	Ser	His	Ile	Lys	Val	Thr	Pro	Leu	Asp	Ser	Arg
			915					920					925		
Leu	Asn	Ala	Pro	Ala	Ala	Glu	Val	Val	Lys	Ala	Phe	Arg	Gln	Gly	Asn
			930				935					940			
Arg	Asp	Val	Ala	Ser	Ser	Gln	Pro	Lys	Ala	Asp	Ser	Val	Asn	Gln	Val
945					950					955					960
Lys	Glu	Thr	Pro	Val	Thr	Lys	Gln	Glu	Pro	Val	Thr	Ser	Thr	Val	Gln
				965					970					975	
Thr	Lys	Thr	Pro	Val	Ser	Glu	Ser	Val	Lys	Thr	Glu	Pro	Thr	Thr	Lys
			980						985					990	
Glu	Ser	Ser	Pro	Gln	Ala	Ile	Lys	Glu	Pro	Val	Asn	Gln	Ser	Glu	Lys
			995				1000					1005			
Gln	Asp	Val	Asn	Leu	Thr	Asn	Glu	Asp	Asn	Ile	Lys	Gln	Pro	Thr	Glu
1010							1015					1020			
Ser	Val	Lys	Glu	Thr	Glu	Thr	Ser	Thr	Lys	Glu	Ser	Thr	Val	Thr	Glu
1025					1030					1035					1040
Glu	Leu	Lys	Glu	Gly	Ile	Asp	Ala	Val	Tyr	Pro	Ser	Leu	Val	Gly	Thr
				1045					1050					1055	
Ala	Asp	Ser	Lys	Ala	Glu	Gly	Ile	Lys	Asn	Tyr	Phe	Lys	Leu	Ser	Phe
			1060					1065						1070	
Thr	Leu	Pro	Glu	Glu	Gln	Lys	Ser	Arg	Thr	Val	Gly	Ser	Glu	Ala	Pro
			1075				1080					1085			
Leu	Lys	Asp	Val	Ala	Gln	Ala	Leu	Ser	Ser	Arg	Ala	Arg	Tyr	Glu	Leu
1090							1095					1100			

-continued

---

Phe Thr Glu Lys Glu Thr Ala Asn Pro Ala Phe Asn Gly Glu Val Ile  
1105                    1110                    1115                    1120  
Lys Arg Tyr Lys Glu Leu Met Glu His Gly Glu Gly Ile Ala Asp Ile  
                  1125                    1130                    1135  
Leu Arg Ser Arg Leu Ala Lys Phe Leu Asn Thr Lys Asp Val Gly Lys  
                  1140                    1145                    1150  
Arg Phe Ala Gln Gly Thr Glu Ala Asn Arg Trp Val Gly Gly Lys Leu  
                  1155                    1160                    1165  
Leu Asn Ile Val Glu Gln Asp Gly Asp Thr Phe Lys Tyr Asn Glu Gln  
                  1170                    1175                    1180  
Leu Leu Gln Thr Ala Val Leu Ala Gly Leu Gln Trp Arg Leu Thr Ala  
1185                    1190                    1195                    1200  
Thr Ser Asn Thr Ala Ile Lys Asp Ala Lys Asp Val Ala Ala Ile Thr  
                  1205                    1210                    1215  
Gly Ile Asp Gln Ala Leu Leu Pro Glu Gly Leu Val Glu Gln Phe Asp  
                  1220                    1225                    1230  
Thr Gly Met Thr Leu Thr Glu Ala Val Ser Ser Leu Ala Gln Lys Ile  
                  1235                    1240                    1245  
Glu Ser Tyr Trp Gly Leu Ser Arg Asn Pro Asn Ala Pro Leu Gly Tyr  
                  1250                    1255                    1260  
Thr Lys Gly Ile Pro Thr Ala Met Ala Ala Glu Ile Leu Ala Ala Phe  
1265                    1270                    1275                    1280  
Val Glu Ser Thr Asp Val Val Glu Asn Ile Val Asp Met Ser Glu Ile  
                  1285                    1290                    1295  
Asp Pro Asp Asn Lys Lys Thr Ile Gly Leu Tyr Thr Ile Thr Glu Leu  
                  1300                    1305                    1310  
Asp Ser Phe Asp Pro Ile Asn Ser Phe Pro Thr Ala Ile Glu Glu Ala  
                  1315                    1320                    1325  
Val Leu Val Asn Pro Thr Glu Lys Met Phe Phe Gly Asp Asp Ile Pro  
                  1330                    1335                    1340  
Pro Val Ala Asn Thr Gln Leu Arg Asn Pro Ala Val Arg Asn Thr Pro  
1345                    1350                    1355                    1360  
Glu Gln Lys Ala Ala Leu Lys Ala Glu Gln Ala Thr Glu Phe Tyr Val  
                  1365                    1370                    1375  
His Thr Pro Met Val Gln Phe Tyr Glu Thr Leu Gly Lys Asp Arg Ile  
                  1380                    1385                    1390  
Leu Glu Leu Met Gly Ala Gly Thr Leu Asn Lys Glu Leu Leu Asn Asp  
                  1395                    1400                    1405  
Asn His Ala Lys Ser Leu Glu Gly Lys Asn Arg Ser Val Glu Asp Ser  
                  1410                    1415                    1420  
Tyr Asn Gln Leu Phe Ser Val Ile Glu Gln Val Arg Ala Gln Ser Glu  
1425                    1430                    1435                    1440  
Asp Ile Ser Thr Val Pro Ile His Tyr Ala Tyr Asn Met Thr Arg Val  
                  1445                    1450                    1455  
Gly Arg Met Gln Met Leu Gly Lys Tyr Asn Pro Gln Ser Ala Lys Leu  
                  1460                    1465                    1470  
Val Arg Glu Ala Ile Leu Pro Thr Lys Ala Thr Leu Asp Leu Ser Asn  
                  1475                    1480                    1485  
Gln Asn Asn Glu Asp Phe Ser Ala Phe Gln Leu Gly Leu Ala Gln Ala  
                  1490                    1495                    1500  
Leu Asp Ile Lys Val His Thr Met Thr Arg Glu Val Met Ser Asp Glu  
1505                    1510                    1515                    1520



-continued

1940					1945					1950				
Gly	Ile	Pro	Phe	Met	Thr	Ile	Gly	Thr	Gly	Asp	Gly	Met	Met	Gln
	1955					1960					1965			
Thr	Leu	Ser	Thr	Met	Lys	Gly	Ala	Pro	Lys	Asn	Thr	Leu	Lys	Ile
	1970					1975					1980			
Asp	Gly	Met	Asn	Ile	Gly	Leu	Asn	Asp	Ile	Thr	Asp	Ala	Ser	Arg
	1985					1990					1995			2000
Ala	Asn	Glu	Ala	Val	Tyr	Thr	Ser	Trp	Gln	Gly	Asn	Pro	Ile	Lys
			2005						2010					2015
Val	Tyr	Glu	Ser	Tyr	Ala	Lys	Phe	Met	Lys	Asn	Val	Asp	Phe	Ser
			2020					2025					2030	
Leu	Ser	Pro	Glu	Ala	Leu	Glu	Ala	Ile	Gly	Lys	Ser	Ala	Leu	Glu
		2035					2040					2045		
Asp	Gln	Arg	Glu	Asn	Ala	Thr	Val	Asp	Asp	Ile	Ala	Asn	Ala	Ala
	2050					2055					2060			
Leu	Ile	Glu	Arg	Asn	Leu	Arg	Asn	Ile	Ala	Leu	Gly	Val	Asp	Ile
	2065					2070					2075			2080
His	Lys	Val	Leu	Asp	Lys	Val	Asn	Leu	Ser	Ile	Asp	Gln	Met	Ala
			2085						2090					2095
Val	Gly	Ala	Pro	Tyr	Gln	Asn	Asn	Gly	Lys	Ile	Asp	Leu	Ser	Asn
			2100					2105						2110
Thr	Pro	Glu	Gln	Gln	Ala	Asp	Glu	Leu	Asn	Lys	Leu	Phe	Arg	Glu
			2115				2120					2125		
Leu	Glu	Ala	Arg	Lys	Gln	Lys	Val	Ala	Lys	Ala	Arg	Ala	Glu	Val
	2130					2135					2140			
Glu	Glu	Thr	Val	Ser	Glu	Lys	Glu	Pro	Val	Asn	Pro	Asp	Phe	Gly
	2145					2150					2155			2160
Val	Gly	Arg	Glu	His	Lys	Ala	Ser	Gly	Val	Arg	Ile	Leu	Ser	Ala
			2165						2170					2175
Ala	Ile	Arg	Asn	Leu	Ala	Lys	Ile	Ser	Asn	Leu	Pro	Ser	Thr	Gln
			2180						2185					2190
Ala	Thr	Leu	Ala	Glu	Ile	Gln	Lys	Ser	Leu	Ala	Ala	Lys	Asp	Tyr
		2195					2200						2205	
Ile	Ile	Tyr	Gly	Thr	Pro	Thr	Gln	Val	Ala	Glu	Tyr	Ala	Arg	Gln
	2210					2215					2220			
Asn	Val	Thr	Glu	Leu	Thr	Ser	Gln	Glu	Met	Glu	Glu	Ala	Gln	Ala
	2225					2230					2235			2240
Asn	Ile	Tyr	Gly	Trp	Thr	Asn	Phe	Asp	Asp	Lys	Thr	Ile	Tyr	Leu
			2245						2250					2255
Ser	Pro	Ser	Met	Glu	Thr	Leu	Ile	His	Glu	Leu	Val	His	Ala	Ser
			2260						2265					2270
Phe	Glu	Glu	Val	Tyr	Ser	Phe	Tyr	Gln	Gly	Asn	Glu	Val	Ser	Pro
			2275					2280					2285	
Ser	Lys	Gln	Ala	Ile	Glu	Asn	Leu	Glu	Gly	Leu	Met	Glu	Gln	Phe
	2290					2295					2300			
Ser	Leu	Asp	Ile	Ser	Lys	Asp	Ser	Pro	Glu	Met	Arg	Glu	Ala	Tyr
	2305					2310					2315			2320
Asp	Ala	Ile	Ala	Thr	Ile	Glu	Gly	His	Leu	Ser	Asn	Gly	Phe	Val
			2325						2330					2335
Pro	Ala	Ile	Ser	Lys	Ala	Ala	Ala	Leu	Asn	Glu	Phe	Met	Ala	Trp
			2340					2345					2350	
Leu	Ala	Asn	Arg	Ala	Leu	Ala	Ala	Lys	Gln	Lys	Arg	Thr	Ser	Ser
	2355							2360						2365

-continued

---

Val Gln Met Val Lys Asp Val Tyr Gln Ala Ile Lys Lys Leu Ile Trp  
 2370 2375 2380

Gly Arg Lys Gln Ala Pro Ala Leu Gly Glu Asp Met Phe Ser Asn Leu  
 2385 2390 2395 2400

Leu Phe Asn Ser Ala Ile Leu Met Arg Ser Gln Pro Thr Thr Gln Ala  
 2405 2410 2415

Val Ala Lys Asp Gly Thr Leu Phe His Ser Lys Ala Tyr Gly Asn Asn  
 2420 2425 2430

Glu Arg Leu Ser Gln Leu Asn Gln Thr Phe Asp Lys Leu Val Thr Asp  
 2435 2440 2445

Tyr Leu Arg Thr Asp Pro Val Thr Glu Val Glu Arg Gly Asn Val  
 2450 2455 2460

Ala Asn Ala Leu Met Ser Ala Thr Arg Leu Val Arg Asp Val Gln Ser  
 2465 2470 2475 2480

His Gly Phe Asn Met Thr Ala Gln Glu Gln Ser Val Phe Gln Met Val  
 2485 2490 2495

Thr Ala Ala Leu Ala Thr Glu Ala Ala Ile Asp Pro His Ala Met Ala  
 2500 2505 2510

Arg Ala Gln Glu Leu Tyr Thr His Val Met Lys His Leu Thr Val Glu  
 2515 2520 2525

His Phe Met Ala Asp Pro Asp Ser Thr Asn Pro Ala Asp Arg Tyr Tyr  
 2530 2535 2540

Ala Gln Gln Lys Tyr Asp Thr Ile Ser Gly Ala Asn Leu Val Glu Val  
 2545 2550 2555 2560

Asp Ala Lys Gly Arg Thr Ser Leu Leu Pro Thr Phe Leu Gly Leu Ala  
 2565 2570 2575

Met Val Asn Glu Glu Leu Arg Ser Ile Ile Lys Glu Met Pro Val Pro  
 2580 2585 2590

Lys Ala Asp Lys Lys Leu Gly Asn Asp Ile Asp Thr Leu Leu Thr Asn  
 2595 2600 2605

Ala Gly Thr Gln Val Met Glu Ser Leu Asn Arg Arg Met Ala Gly Asp  
 2610 2615 2620

Gln Lys Ala Thr Asn Val Gln Asp Ser Ile Asp Ala Leu Ser Glu Thr  
 2625 2630 2635 2640

Ile Met Ala Ala Ala Leu Lys Arg Glu Ser Phe Tyr Asp Ala Val Ala  
 2645 2650 2655

Thr Pro Thr Gly Asn Phe Ile Asp Arg Ala Asn Gln Tyr Val Thr Asp  
 2660 2665 2670

Ser Ile Glu Arg Leu Ser Glu Thr Val Ile Glu Lys Ala Asp Lys Val  
 2675 2680 2685

Ile Ala Asn Pro Ser Asn Ile Ala Ala Lys Gly Val Ala His Leu Ala  
 2690 2695 2700

Lys Leu Thr Ala Ala Ile Ala Ser Glu Lys Gln Gly Glu Ile Val Ala  
 2705 2710 2715 2720

Gln Gly Val Met Thr Ala Met Asn Gln Gly Lys Val Trp Gln Pro Phe  
 2725 2730 2735

His Asp Leu Val Asn Asp Ile Val Gly Arg Thr Lys Thr Asn Ala Asn  
 2740 2745 2750

Val Tyr Asp Leu Ile Lys Leu Val Lys Ser Gln Ile Ser Gln Asp Arg  
 2755 2760 2765

Gln Gln Phe Arg Glu His Leu Pro Thr Val Ile Ala Gly Lys Phe Ser  
 2770 2775 2780

-continued

Arg	Lys	Leu	Thr	Asp	Thr	Glu	Trp	Ser	Ala	Met	His	Thr	Gly	Leu	Gly	2785	2790	2795	2800
Lys	Thr	Asp	Leu	Ala	Val	Leu	Arg	Glu	Thr	Met	Ser	Met	Ala	Glu	Ile	2805	2810	2815	
Arg	Asp	Leu	Leu	Ser	Ser	Ser	Lys	Lys	Val	Lys	Asp	Glu	Ile	Ser	Thr	2820	2825	2830	
Leu	Glu	Lys	Glu	Ile	Gln	Asn	Gln	Ala	Gly	Arg	Asn	Trp	Asn	Leu	Val	2835	2840	2845	
Gln	Lys	Lys	Ser	Lys	Gln	Leu	Ala	Gln	Tyr	Met	Ile	Met	Gly	Glu	Val	2850	2855	2860	
Gly	Asn	Asn	Leu	Leu	Arg	Asn	Ala	His	Ala	Ile	Ser	Arg	Leu	Leu	Gly	2865	2870	2875	2880
Glu	Arg	Ile	Thr	Asn	Gly	Pro	Val	Ala	Asp	Val	Ala	Ala	Ile	Asp	Lys	2885	2890	2895	
Leu	Ile	Thr	Leu	Tyr	Ser	Leu	Glu	Leu	Met	Asn	Lys	Ser	Asp	Arg	Asp	2900	2905	2910	
Leu	Leu	Ser	Glu	Leu	Ala	Gln	Ser	Glu	Val	Glu	Gly	Met	Glu	Phe	Ser	2915	2920	2925	
Ile	Ala	Tyr	Met	Val	Gly	Gln	Arg	Thr	Glu	Glu	Met	Arg	Lys	Ala	Lys	2930	2935	2940	
Gly	Asp	Asn	Arg	Thr	Leu	Leu	Asn	His	Phe	Lys	Gly	Tyr	Ile	Pro	Val	2945	2950	2955	2960
Glu	Asn	Gln	Gln	Gly	Val	Asn	Leu	Ile	Ile	Ala	Asp	Asp	Lys	Glu	Phe	2965	2970	2975	
Ala	Lys	Leu	Asn	Ser	Gln	Ser	Phe	Thr	Arg	Ile	Gly	Thr	Tyr	Gln	Gly	2980	2985	2990	
Ser	Thr	Gly	Phe	Arg	Thr	Gly	Ser	Lys	Gly	Tyr	Tyr	Phe	Ser	Pro	Val	2995	3000	3005	
Ala	Ala	Arg	Ala	Pro	Tyr	Ser	Gln	Gly	Ile	Leu	Gln	Asn	Val	Arg	Asn	3010	3015	3020	
Thr	Ala	Gly	Gly	Val	Asp	Ile	Gly	Thr	Gly	Phe	Thr	Leu	Gly	Thr	Met	3025	3030	3035	3040
Val	Ala	Gly	Arg	Ile	Thr	Asp	Lys	Pro	Thr	Val	Glu	Arg	Ile	Thr	Lys	3045	3050	3055	
Ala	Leu	Ala	Lys	Gly	Glu	Arg	Gly	Arg	Glu	Pro	Leu	Met	Pro	Ile	Tyr	3060	3065	3070	
Asn	Ser	Lys	Gly	Gln	Val	Val	Ala	Tyr	Glu	Gln	Ser	Val	Asp	Pro	Asn	3075	3080	3085	
Met	Leu	Lys	His	Leu	Asn	Gln	Asp	Asn	His	Phe	Ala	Lys	Met	Val	Gly	3090	3095	3100	
Val	Trp	Arg	Gly	Arg	Gln	Val	Glu	Glu	Ala	Lys	Ala	Gln	Arg	Phe	Asn	3105	3110	3115	3120
Asp	Ile	Leu	Ile	Glu	Gln	Leu	His	Ala	Met	Tyr	Glu	Lys	Asp	Ile	Lys	3125	3130	3135	
Asp	Ser	Ser	Ala	Asn	Lys	Ser	Gln	Tyr	Val	Asn	Leu	Leu	Gly	Lys	Ile	3140	3145	3150	
Asp	Asp	Pro	Val	Leu	Ala	Asp	Ala	Ile	Asn	Leu	Met	Asn	Ile	Glu	Thr	3155	3160	3165	
Arg	His	Lys	Ala	Glu	Glu	Leu	Phe	Gly	Lys	Asp	Glu	Leu	Trp	Val	Arg	3170	3175	3180	
Arg	Asp	Met	Leu	Asn	Asp	Ala	Leu	Gly	Tyr	Arg	Ala	Ala	Ser	Ile	Gly	3185	3190	3195	3200
Asp	Val	Trp	Thr	Gly	Asn	Ser	Arg	Trp	Ser	Pro	Ser	Thr	Leu	Asp	Thr				

-continued

3205				3210				3215							
Val	Lys	Lys	Met	Phe	Leu	Gly	Ala	Phe	Gly	Asn	Lys	Ala	Tyr	His	Val
			3220								3225				3230
Val	Met	Asn	Ala	Glu	Asn	Thr	Ile	Gln	Asn	Leu	Val	Lys	Asp	Ala	Lys
			3235				3240								3245
Thr	Val	Ile	Val	Val	Lys	Ser	Val	Val	Val	Val	Pro	Ala	Val	Asn	Phe
			3250				3255								3260
Ala	Asn	Ile	Tyr	Gln	Met	Ile	Gly	Arg	Gly	Val	Pro	Val	Lys	Asp	Ile
			3265				3270				3275				3280
Ala	Val	Asn	Ile	Pro	Arg	Lys	Thr	Ser	Glu	Ile	Asn	Gln	Tyr	Ile	Lys
			3285								3290				3295
Ser	Arg	Leu	Arg	Gln	Ile	Asp	Ala	Glu	Ala	Glu	Leu	Arg	Ala	Ala	Glu
			3300												3310
Gly	Asn	Pro	Asn	Leu	Val	Arg	Lys	Leu	Lys	Thr	Glu	Ile	Gln	Ser	Ile
			3315				3320								3325
Thr	Asp	Ser	His	Arg	Arg	Met	Ser	Ile	Trp	Pro	Leu	Ile	Glu	Ala	Gly
			3330				3335								3340
Glu	Phe	Ser	Ser	Ile	Ala	Asp	Ala	Gly	Ile	Ser	Arg	Asp	Asp	Leu	Leu
			3345				3350				3355				3360
Val	Ala	Glu	Gly	Lys	Ile	His	Glu	Tyr	Met	Glu	Lys	Leu	Ala	Asn	Lys
			3365								3370				3375
Leu	Pro	Glu	Lys	Val	Arg	Asn	Ala	Gly	Arg	Tyr	Ala	Leu	Ile	Ala	Lys
			3380												3390
Asp	Thr	Ala	Leu	Phe	Gln	Gly	Ile	Gln	Lys	Thr	Val	Glu	Tyr	Ser	Asp
			3395				3400								3405
Phe	Ile	Ala	Lys	Ala	Ile	Ile	Tyr	Asp	Asp	Leu	Val	Lys	Arg	Lys	Lys
			3410				3415								3420
Lys	Ser	Ser	Ser	Glu	Ala	Leu	Gly	Gln	Val	Thr	Glu	Glu	Phe	Ile	Asn
			3425				3430				3435				3440
Tyr	Asp	Arg	Leu	Pro	Gly	Arg	Phe	Arg	Gly	Tyr	Met	Glu	Ser	Met	Gly
			3445								3450				3455
Leu	Met	Trp	Phe	Tyr	Asn	Phe	Lys	Ile	Arg	Ser	Ile	Lys	Val	Ala	Met
			3460				3465								3470
Ser	Met	Ile	Arg	Asn	Asn	Pro	Val	His	Ser	Leu	Ile	Ala	Thr	Val	Val
			3475				3480								3485
Pro	Ala	Pro	Thr	Met	Phe	Gly	Asn	Val	Gly	Leu	Pro	Ile	Gln	Asp	Asn
			3490				3495								3500
Met	Leu	Thr	Met	Leu	Ala	Glu	Gly	Arg	Leu	Asp	Tyr	Ser	Leu	Gly	Phe
			3505				3510				3515				3520
Gly	Gln	Gly	Leu	Arg	Ala	Pro	Thr	Leu	Asn	Pro	Trp	Phe	Asn	Leu	Thr
			3525								3530				3535

His

<210> SEQ ID NO 16  
 <211> LENGTH: 32  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

&lt;400&gt; SEQUENCE: 16

ggcattactt catccaaaag aagcggagct tc

32

&lt;210&gt; SEQ ID NO 17

-continued

---

<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer  
  
<400> SEQUENCE: 17  
  
ggccatccat tacttcatcc aaaagaagcg gagcttc 37

<210> SEQ ID NO 18  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer  
  
<400> SEQUENCE: 18  
  
ggatccaaaa gaagcggagc ttc 23

<210> SEQ ID NO 19  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer  
  
<400> SEQUENCE: 19  
  
ggcattactt catccaaaag aagctgagct tc 32

<210> SEQ ID NO 20  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer  
  
<400> SEQUENCE: 20  
  
ggcattactt catccaaaag aagcggagc 29

<210> SEQ ID NO 21  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer  
  
<400> SEQUENCE: 21  
  
ggaggctcct cggagtctcc tttt 24

<210> SEQ ID NO 22  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer  
  
<400> SEQUENCE: 22  
  
ggactacctt cgggtagtcc ttttt 25

<210> SEQ ID NO 23  
<211> LENGTH: 33

-continued

---

<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer

<400> SEQUENCE: 23

agaagggggc tactaagccc tcttcttatt ttt 33

<210> SEQ ID NO 24  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer

<400> SEQUENCE: 24

aagctgctcc gcagctttt 19

<210> SEQ ID NO 25  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer

<400> SEQUENCE: 25

aaggctatcc ctacgggggt agcctttatt ttttt 35

<210> SEQ ID NO 26  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer

<400> SEQUENCE: 26

gccctccttg tgaggccttt tt 22

<210> SEQ ID NO 27  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer

<400> SEQUENCE: 27

caacgaagcg ttgaatacct 20

<210> SEQ ID NO 28  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
Primer

<400> SEQUENCE: 28

ttcttcgagg cgaagaaaac ct 22

<210> SEQ ID NO 29  
<211> LENGTH: 20  
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 29

cgacgaggcg tcgaaaacca

20

What is claimed is:

1. An isolated nucleic acid comprising a region encoding a polypeptide having an amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:15.

2. The nucleic acid of claim 1, wherein said nucleic acid comprises the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:14.

3. The nucleic acid of claim 1, wherein said nucleic acid is operatively linked to a promoter.

4. The nucleic acid of claim 3, wherein said promoter is an N4 vRNAP promoter set forth in SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:27, SEQ ID NO:28 or SEQ ID NO:29.

5. The nucleic acid of claim 3, wherein said promoter is a P2 sequence set forth in SEQ ID NO:16 or SEQ ID NO:28.

6. An isolated recombinant host cell comprising a DNA segment comprising a DNA segment comprising the nucleic acid sequence of claim 1.

7. The recombinant host cell of claim 6, wherein said DNA segment is a single-stranded DNA segment.

8. The recombinant host cell of claim 6, wherein said DNA segment is a double-stranded DNA segment.

9. The recombinant host cell of claim 6, wherein said DNA segment encodes a polypeptide having an amino acid sequence set forth in SEQ ID NO:4.

10. The recombinant host cell of claim 6, wherein said DNA segment encodes a polypeptide having an amino acid sequence set forth in SEQ ID NO:6.

11. The recombinant host cell of claim 6, wherein said cell is an *E. coli* cell.

12. A recombinant vector comprising a DNA segment encoding a N4 virion RNA polymerase polypeptide under the control of a promoter, wherein the DNA segment comprises the nucleic acid sequence of claim 1.

13. An isolated polynucleotide comprising a sequence identical or completely complementary to the full length of SEQ ID NO:1.

14. An isolated polynucleotide comprising a sequence identical or completely complementary to the full length of SEQ ID NO:3.

15. The nucleic acid of claim 1, wherein said polypeptide comprises the amino acid sequence of SEQ ID NO:4.

16. An isolated nucleic acid encoding the polypeptide of SEQ ID NO:4 wherein said polypeptide has a mutation at position Y678.

17. A method of making a full-length N4 vRNAP or mini-vRNAP comprising:

(a) expressing an isolated nucleic acid sequence encoding vRNAP, wherein said vRNAP has the amino sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or, SEQ ID NO:15; and

(b) purifying said vRNAP.

18. The method of claim 17, wherein said expressing occurs in a bacteria, yeast, CHO, Cos, HeLa, NIH3T3, Jurkat, 293, Saos, or PC12 host cell.

19. The method of claim 17, further comprising using a promoter appropriate for expression in the host cell line being used.

20. The method of claim 19, wherein said promoter is pBAD.

21. The method of claim 19, wherein said promoter is a promoter recognized by T7 RNA polymerase, T3 RNA polymerase or SP6 RNA polymerase.

22. The method of claim 19, wherein said promoter is a promoter recognized by T7-like RNA polymerase.

23. The method of claim 17, wherein said vRNAP has a specific recombinant sequence for use in purification.

24. The method of claim 23, wherein said vRNAP has at least one histidine, FLAG, hemagglutinin or c-myc tag.

25. The method of claim 23, wherein said vRNAP has at least one histidine tag.

26. The method of claim 24, wherein said purifying occurs in one step.

27. The method of claim 17, wherein said vRNAP does not have a tag.

\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 7,452,705 B2  
APPLICATION NO. : 10/153219  
DATED : November 18, 2008  
INVENTOR(S) : Krystyna M. Kazmierczak et al.

Page 1 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the title page, item (56) References Cited - Other Publications, please insert:

--Paule and White, "Transcription by RNA polymerases I and III," Nuc. Acids Res., 28:1283-1298, 2000.  
Roeder, "The role of general transcription factors in transcription by RNA polymerase II," Trends Biochem. Sci., 21: 327-335, 1996.  
Rong et al., "Promoter specificity determinants of T7 RNA polymerase," Proc. Natl. Acad. Sci. USA, 95:515-519, 1998.  
Rothman-Denes et al., "Transcriptional Regulation by DNA Structural Transitions and Single-Stranded DNA Binding Proteins," 63rd Cold Spring Harbor Symp. Quant. Biol., 63: 63-73, 1998.  
Sanders et al., "Dual targets of a transcriptional activator that tracks on DNA," EMBO Journal, 16: 3124-3132, 1997.  
Shadel and Clayton, "Mitochondrial transcription initiation, variation and conservation," J. Biol. Chem., 268:16083-16086, 1993.  
Sousa, "Structural and mechanistic relationships between nucleic acid polymerases," Trends in Biochem. Sci., 21: 186-190, 1996.  
Sousa et al., "Crystal structure of bacteriophage T7 RNA polymerase at 3.3 Å resolution," Nature, 364:593-599, 1993.  
Zhang et al., "Crystal structure of Thermus aquaticus core RNA polymerase at 3.3.Å resolution," Cell, 98:811-824, 1999.--.

In column 1, lines 9-11, delete "The government may own rights in the present invention pursuant to grant number R01 A1 12575 from the National Institute of Health" and insert --This invention was made with government support under grant number R01 A1 12575 awarded by the National Institutes of Health. The government has certain rights in the invention.-- therefor.

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 7,452,705 B2  
APPLICATION NO. : 10/153219  
DATED : November 18, 2008  
INVENTOR(S) : Krystyna M. Kazmierczak et al.

Page 2 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In claim 1, column 149, line 15, delete "an" and insert --the-- therefor.

In claim 2, column 149, lines 18-19, insert --sequence-- between "acid" and "comprises".


In claim 6, column 149, line 29, delete "a DNA segment comprising".

In claim 9, column 149, line 37, delete "an" and insert --the-- therefor.

In claim 10, column 149, line 40, delete "an" and insert --the-- therefor.

Signed and Sealed this

Third Day of February, 2009



JOHN DOLL  
*Acting Director of the United States Patent and Trademark Office*