



US 20080288264A1

(19) **United States**

(12) **Patent Application Publication**  
**Mach et al.**

(10) **Pub. No.: US 2008/0288264 A1**

(43) **Pub. Date: Nov. 20, 2008**

(54) **METHODS FOR GENERATION OR INCREASING REVENUES FROM CROPS**

(22) Filed: **Feb. 2, 2007**

**Related U.S. Application Data**

(75) Inventors: **Jennifer Mach**, Chicago, IL (US);  
**Helge Zieler**, Chicago, IL (US);  
**RongGuan Jin**, Chicago, IL (US);  
**Kevin Keith**, Chicago, IL (US);  
**Gregory P. Copenhaver**, Chapel Hill, NC (US);  
**Daphne Preuss**, Chicago, IL (US)

(63) Continuation of application No. 10/170,944, filed on Jun. 12, 2002, now Pat. No. 7,193,128, which is a continuation-in-part of application No. 09/531,120, filed on Mar. 17, 2000, now Pat. No. 6,972,197.

**Publication Classification**

(51) **Int. Cl.**  
**G06Q 50/00** (2006.01)  
**G06Q 30/00** (2006.01)  
(52) **U.S. Cl.** ..... **705/1**


Correspondence Address:  
**MARSHALL, GERSTEIN & BORUN LLP**  
**233 S. WACKER DRIVE, SUITE 6300, SEARS TOWER**  
**CHICAGO, IL 60606 (US)**

(57) **ABSTRACT**

The present invention provides methods of doing business and providing services. For example, methods of increasing the revenue of crops are provided. To this end, the method includes the use of a nucleic acid sequences of plant centromeres. This will permit construction of stably inherited recombinant DNA constructs and mini chromosomes which can serve as vectors for the construction of transgenic plant and animal cells.

(73) Assignees: **CHROMATIN INC.**, Chicago, IL (US); **THE UNIVERSITY OF CHICAGO**, CHICAGO, IL (US)

(21) Appl. No.: **11/701,900**



AGCTTGATTTGGATACATAAAGTGGTGGAGAATCACCAGGAAGT  
TGAATAAATCTCATAGGAGTTGGCATGAAGAAGTTATCCCCTT  
TCAAATCAGGTGATTCCAGTTTCCCAGTTTGGGAATAGCACAGC  
TTCTTCGTCGTTCCAATCAAACCAGGATGAATCWCTTTGTRARA  
AGCT

FIG. 1A

AGCTTGATTTTGATACATAAAGTAGTGGAGAATCAYTWGGAAGT  
GGAATAAATCTCATAGGAGTTAGGATGAAGAAGCTATCMCACTT  
TCAAATCAGGTGATCCCARTTTTCTGTTTGGGAATATGACAAC  
TTMTTTGTCATTCTAATCAAACCAGGAWGAATCGCBATGTAARA  
AGCT

FIG. 1B

Chrbo1 AGCTTGATTTG GATAACATAAAGTGGTGGAGAATCACCAGGAAGTGAATAAATCTCATAG  
ChrBo2 AGCTTGATTTG GATAACATAAAGTAGTGGAGAATCAYTWGGAAGTGAATAAATCTCATAG  
position 1.....10.....20.....30.....40.....50.....60

Chrbo1 GAGTTGGCATGAAGAAGTATCCCACTTTCAAATCAGGTGATCCAGTTTCCCAGTTTGG  
ChrBo2 GAGTTAGGATGAAGAAGCTATCMCACTTTCAAATCAGGTGATCCCARTTTTCTGTTTGG  
position 61.....70.....80.....90.....100.....110.....120

Chrbo1 GAATAGCACAGCTTCTTCGTCGTTCCAATCAAACCAGGATGAATCWCTTTGTRARAAGCT  
ChrBo2 GAATATGACAACCTTMTTGTCAATTCTAATCAAACCAGGAWGAATCGCBATGTAARAAGCT  
position 121.....130.....140.....150.....160.....170.....180

FIG. 1C

AGCTTGATTTGGATACATAAAGTGGTGGAGAATCACCAGGAAGT  
TGAATAAATCTCATAGGAGTTGGSATGAAGAAGTTATCCCCTT  
TCAAATCAGGTGATTCCAGTTTCCCAGTTTGGGAATAGCACAGC  
TTCTTCGTCGTTCCAATCAAACCAGGATGAATCACTTTGTRAGA  
AGCT

FIG. 1D

AGCTTGATTTTGGATACATAAAGTARTGGAGAATCAYYAGGAAGT  
KGAATAAATCTCATAGGAGTTAGGATGAAGAAGCTATCCCCTT  
TCAAATCAGGTGATCCCARTTTTCTGTTTGGGAATAKGACARC  
TTCTTTGTCATTCTAATCAAACCAGGAWGAATCGCKATGTAARA  
AGCT

FIG. 1E

<u>ChrBo1</u>	AGCTTGATTTGGATAACATAAAGTGGTGGAGAATCACAGGAAGTGAATAAATCTCATAG
<u>ChrBo2</u>	AGCTTGATTTTGATAACATAAAGTARTGGAGAATCAYYAGGAAGTKGAATAAATCTCATAG
<u>position</u>	1.....10.....20.....30.....40.....50.....60
<u>ChrBo1</u>	GAGTTGSATGAAGAAGTTATCCCACTTTCAAATCAGGTGATCCAGTTTCCAGTTTGG
<u>ChrBo2</u>	GAGTTAGGATGAAGAAGCTATCCCACTTTCAAATCAGGTGATCCARTTTTCCTGTTTGG
<u>position</u>	.....70.....80.....90.....100.....110.....120
<u>ChrBo1</u>	GAATAGCACAGCTTCTTCGTCGTTCCAATCAAACCAGGATGAATCACTTTGTRAGAAGCT
<u>ChrBo2</u>	GAATAKGACARCTTCTTTGTCATTCTAATCAAACCAGGAWGAATCGCKATGTAARAAGCT
<u>position</u>	.....130.....140.....150.....160.....170.....180

FIG. 1F

AAATTCAAATGGTCATAACTTTTMACWCGGAKGTCCGATTCAGG  
CGCATAATATATCGAGACGCTCGAAATTGAACAAYGGAAGCTCT  
CGAG

FIG. 2A

AAATTCAAACGACAATAACTTTTTACTCGGATGTCYGATTGAGT  
CCCGTAATATATCGAGACGCTCGAAATTGAATRYTGAAGCTCTG  
AGC

FIG. 2B

ChrGm1 AAATTCAAATGGTCATAACTTTTMACWCGGAKGTCCGATTCAGGCGCATAATATATCGAG  
ChrGm2 AAATTCAAACGACAATAACTTTTACTCGGATGTCYGATTGAGTCCCGTAATATATCGAG  
position 1.....10.....20.....30.....40.....50.....60

ChrGm1 ACGCTCGAAATTGAACAAYGGAAGCTCTCGAG 92  
ChrGm2 ACGCTCGAAATTGAATRYT-GAAGCTCT-GAGC 91  
position .....70.....80.....90...

FIG. 2C

AAATTCAAATGGTCATAACTTTTMACWCGGAKGTCCGATTCAGG  
CGCATAATATATCGAGACGCTCGAAATTGAACAAYGGAAGCTCT  
CGAG

FIG. 2D

AAATTCAAACGACAATAACTTTTTACTCGGATGTCYGATTGAGT  
CCCGTAATATATCGAGACGCTCGAAATTGAATRYTGAAGCTCTG  
AGC

FIG. 2E

ChrGm1	AAATTCAAATGGTCATAACTTTTTMACWCGGAKGTCGATTGAGCCGCATAATATATCGAG
ChrGm2	AAATTCAAACGACAATAACTTTTTACTCGGATGTCYGATTGAGTCCCGTAATATATCGAG
position	1.....10.....20.....30.....40.....50.....60

ChrGm1	ACGCTCGAAATTGAACAAYGGAAGCTCTCGAG	92
ChrGm2	ACGCTCGAAATTGAATRYT-GAAGCTCT-GAGC	91
position	.....70.....80.....90...	

FIG. 2F



CCATCACGGGTTTTCTGGGCCRRTTTGGAAGGTCAAACGAGCCCC  
GGAGCGAGCATACGCCTCATTTTGACGATTTTCGTGTGCTATTG  
CACACCATTTTTTTGGGTGATCGGGATTCCGACGTCAAAAATGCC  
AAATTTGTTTCGTGGACGTCCGTCAAGACGTTGTCTATGCATACG  
GTTGG

FIG. 3A

CCATCACGGGTTTTCTGGGCCRRTTTGGAAGGTCAAACGAGCCCC  
GRAGCGAGCATACGCCTCATTTTGACGATTTTCGTGTGCTATTG  
CACACCATTTTTTTGGGTGATCGGGATTCCGACGTCAAAAATGCC  
AAATTTGTTTCGTGGACGTCCGTCAAGACGTTGTCTATGCATACG  
GTTGG

FIG. 3B

GGCCACACAACCCCCATTTTTGTCGAAAATAGCCATGAACGACC  
ATTTTCAATAATACYRAAGGCTAACACCTACGGATTTTTTRACCA  
AGAAATGGTCTCCACCAGAAATCCAAGAATGTGATCTATGGCAA  
GGAAACATATGTGGGGTGAGGTGTAYGAGCCTCTGGTCGAYGAT  
CAAT

FIG. 4A

GGCCACACAACCCCCATTTTTGTCGAAAATAGCCATGAAYGACC  
ATTTTCAATAATACCGAAGGCTAACACCTACGGATTTTTTGACCA  
AGAAATGGTCTCCACCAGAAATCCAAGAATGTGATCTATGGCAA  
GGAAACATATGTGGGGTGAGGTGTAYGAGCCTCTGGTCGATGAT  
CAAT


FIG. 4B

GGTTCCGGTGGCAAAAACCTCGTAGCTTTGTATGCACCCMGACAC  
CCGTTTTTCGGAATGGGTGACGTGYGACAACAGAAATTGCGMGAA  
ACCACCCCAAACATGAGTTTTGKACCTAAAGTAGTGGATTGGGC  
ATGTTTCGTTGYGAAAAACGAAGAAAT

FIG. 4C

FIG. 5

Mini-chromosome construction:  
BAC retrofitting by Cre-Lox recombination



- plasmid backbone sequences
- BAC insert (centromere sequences)
- ▷ LoxP site
- vector sequences from mini-chromosome vector




FIG. 6




FIG. 7A




FIG. 7B




FIG. 7C






FIG. 7D




FIG. 7E




FIG. 7F




FIG. 7G




FIG. 7H




FIG. 7I




FIG. 7J




FIG. 7K




FIG. 7L






FIG. 7M




FIG. 7N




FIG. 8A



FIG. 8B




FIG. 9




FIG. 10A




FIG. 10B




FIG. 10C






FIG. 10D




FIG. 10E




FIG. 10F




FIG. 10G




FIG. 11A-T




FIG. 12




FIG. 13A,B

Chromosome 2

Sequenced clone	marker name	marker or primer pair	Marker location	marker properties, position	Forward Primer	Reverse Primer
T13E11	T13E11.01	primer	1755-2385	Lan & Col	AGCGCTGGGATGGGT TGGTTG	TTAATGGGCAATGGCTGA ACAAG
T13E11	T13E11.30	primer	30628-31270	Lan & Col	CAGGTTGCGTTACTA CATGGGTTTCAG	TATGCATGCGAGTTGGTGG AGGTAAG
T13E11	T13E11.48	primer	48187-48969	Lan & Col	AACCCCGTAAATTA AACCACAC	CGATACGGGCATGACTCCA G
T13E11	T13E11.63	primer	63886-64530	Lan & Col	ACGGCGTTGAGAGG AGAAGC	CCCCAACCGCAGCAAGAC AATC
T13E11	T13E11.78	primer	78190-78878	Lan & Col	AACAAAACAAATGCC AGGTCAGG	CTCCGGTCGCAAGTTACA TACAG
T13E11	T13E11.93	primer	93907-94579	Lan & Col	GTTACCCCGTCTCTGA GATTGAG	TTGGGGAGCAGGATTTGAT GTG
F27C21	F27C21.18	primer	18383-19057	Lan & Col	CAGGCGATTGCTTTT ATAGGCTGTAAG	TTTTGCTGAAACGGAGGGA GTAC
F27C21	F27C21.02	primer	2570-3293	Lan & Col	ACAAAAGCCGAATC GTGGAAG	TGCTGGTTGATTATTGCT GAAAG
F9A16	F9A16.71	primer	71978-72592	Lan & Col	AATGCTTTTCCGACTC TTTTGAC	TTGTTATTTGGGTTTGGG TTGG
F9A16	F9A16.53	primer	53253-53921	Dom	CGCAGGCGGCTACTT GTTTG	GATGAATTGATCCGTTGTT TTATGCT
F9A16	F9A16.38	primer	37116-37733	Dom	CAAGGCGGGAAAAACA ACTC	AATTATTTTCAAACGGCTCT TTACC
F9A16	F9A16.22	primer	22166-22889	Dom	GTATTTAGCATTATGT TAGTCTGTTAGTGG	GCTCCTTGCGTATTCTTCAC C
F9A16	F9A16.03		3308-4091	Lan & Col	GATCCAGCAACCTTA GCCTCCTC	GATCCCTCAGTTGAAAATC AATCTTC
TSM2	mito border	marker	17685	Dom	AAGGCATCAACGTTT GTGTG	CTACCAGGTAGGTGAAAACG AGGGCCGGAGCTCGTATGG A
TSE7	TSE7.23	primer	109092-109688	Lan & Col	ATTATTGGCTGCTGCA CTTCTGTCAC	CGGTCGCCCTCGTTCGTGA TCTG
TSE7	TSE7.73	primer	73460-74120	Lan & Col	TCTCGGGAGTAGGGG CITTGTTCTG	TCGCCAATGAAAGAGGGT AG
TSE7	TSE7.58	primer	57942-58583	Lan & Col	AGGGGCTTACAAGA ATGAAC	CTAAATCCCGAAAAACCAA ACCAC
TSE7	TSE7.40	primer	40913-41537	Lan & Col	GTAGCCGGCTCAGTCT CATAACATC	CGAATTOCTTCAGATG ATGC
TSE7	mito border	marker	13507	Dom	AAATGGCAGAAGCAG AAGCAGGAATAG	TTGCTTAOCTATAACCGAC TAATGCAAGGGTCTGTAA TGGAAATG
TSE7	TSE7.02	primer	2919-3585	Lan & Col	GTGTGGCTCGTGTGA CCTGAC	AGTTTGCTTGGTTGCTTGGT TATTATG
T12J2	T12J2.01	primer	1373-1998	Lan & Col	CTGGCCATCCCTTAT CGGTTTAC	AGCGCAATCAAGCTATCC TACATA
T12J2	T12J2.19	primer	19369-20038	Lan & Col	ACCTCCCCACACTTA AAGACACTG	CTCAGAATCCAAAACAGA GCCACAC
T12J2	T12J2.37	primer	37750-38359	Lan & Col	TGAATGCTATGAAAG ATGGATGAAAC	AGACGGCTAGTATTTGGT GG
T12J2	ATEDA59	marker	50592	Lan & Col	AATGGGCTCGGTTGT GTAGAAAC	ATGGCCCAATCAAAAAGCA ATCC

FIG. 14A



Chromosome 2

T12J2	T12J2.73	primer	73911-74556	Lan & Col	ACTTGTAGGCCCTTTG ATGTCTG	TGCTTTGTGTTGCTTTGATT ATTCTATTAG
T14C8	T14C8.1	primer	8862-9544	Lan & Col	AOGAACCOGACGACC ACTG	ACGCCCTTGATTCCATTTC TACC
T14C8	T14C8.6	primer	10837-11485	Lan & Col	GACGGGTGAAAGAA AGCACAG	AGAAGATGATGGCAAGTT ACGAAGAG
T14C8	T14C8.7	primer	45334-46016	Lan & Col	CGACCATTCACGACCC ATAC	AAGCCCATTCAAAAGAGTT AGGAGAG
T14C8	T14C8.3	primer	46672-47283	Lan & Col	ATAGGGTCAGCOCTCA TTTCAG	ACCCCTTTGCTTGTATTTTC GTG
T14C8	T14C8.8	primer	48833-49538	Lan & Col	CAAATGGGCGGAGGG GG	TGTGGCAAGTCATGGGTAA GGAG
T14C8	T14C8.9	primer	71115-71878	Lan & Col	GGTGCGGGAGAACGA TGAC	COGGTTTCTGCGATATTTG GTTAG
T14C8	T14C8.5	primer	73841-74456	Lan & Col	TTCCGCGOCCAAAAG GTG	AGAGTCAAGCCAAGCAAT AACAGG
T14C8	T14C8.4	primer	75442-76122	Lan & Col	AAGAAGGCTGGAAT TGGTTGAG	GAGCGGAAGTAGATGCAG AATGTC
F7B19	F7B19.1	primer	108-757	Lan & Col	AAGGTCCGCGGTGG TGAG	GGGTCCGAGTGAATGTGATTG AGTG
F7B19	F7B19.12	primer	13831	Lan & Col	TAACTGCATCAGCGGT AGGAAAC	TTACAAGCGAGAAAAGAT GAGAAGC
F7B19	F7B19.27	primer	27033	Lan & Col	CCCCGCTGAAGTACT GACTACGAG	TCCGCCACCGATAAGATA CGAC
F7B19	F7B19.2	primer	30189-30791	Lan & Col	GCTGGTTGCGGTTGCT GTTC	CCGCGGTGGCTGCTTTTAG TGCCCTTCCGGAACTGGT G
F7B19	F7B19.43	primer	43142	Lan & Col	CAGGAAAGTGGTTG GATTGATG	TGCTCTTCCGGAACTGGT G
F7B19	F7B19.3	primer	55446-56209	Lan & Col	ATTTGTTGCCATCGT CCTTC	TTATCAATGTATTTCCCTG TGTATC
F7B19	F7B19.59	primer	59771	Lan & Col	TTGAAGAATGCGCAT AGCCGTAG	TCTGGGATGAAGAGAAAG AGAAGTCTC
F7B19	F7B19.4	primer	70859-71492	Lan & Col	AAAAACCGTGAGACC CATAAATG	TCCAAAATCGGAAAGTGAC AG
F7B19	F7B19.77	primer	77633	Lan & Col	ATAACCGAAGAAGCC GAGAAATC	ATCCGGAGACGAAAATGA ACTTAG
F7B19	F7B19.95	primer	95351	Lan & Col	TCGATTGCCAGCAGA GTCAGAAC	TGGGGCTTGTAAAGGAGG AGTAAC
F7B19	F7B19.5	primer	98977-99658	Dom?	AGATGGGGTCTATT TTGTATG	CCGGTCCGAGTGAATTCCTG TAG
F7B19	F7B19.6	primer	112337-113039	Lan & Col	AGGGGAAAACCTTGA GAGCAC	TATCGGGTTTTGAAGAGGG AAGG
T15D9	T15D9.3	primer	2985	Lan & Col	AGGGTGGCGGTGTG GAG	TCCTGGCAAATGCTTCT CGTTG
T15D9	T15D9.1	primer	12299-12914	Lan & Col	GCTCCGCCATCTCCTC GTC	GAAGTCCATGCTATGCC TG
T15D9	T15D9.19	MARKER	18991	LAST RECOMBINANT Col	GAGOOCTTCTATGAGC CTAACCTGTTTC	AGAGATCCOCTGTACTAA AGCCTATTCTG
T15D9	T15D9.2	primer	37103-37728	Lan & Col	ATGGGGTAATCGAAT AGTGTGGTC	CCCTAGGGCATCCGTTTTT ATCTC
T15D9	T15D9.3a	primer	52189-52811	Col	CGGAGAAAGTTGGGG GTTAGTTG	GAGAGGTTTGGGTGGGCT TGTAG
T15D9	T15D9.55	primer	55134	Lan & Col	GCTGCGAACCCACAC TTTGCTC	ATGTTATCGTCCCGGTT TTATG

FIG. 14A (cont'd)

Chromosome 2

T15D9	T15D9.73	primer	72993	Lan & Col	AACCGGTTGATAGTA GAOGAGATG	TCCGGGGTTGGGATAGAG
T15D9	T15D9.4	primer	73930-74552	Lan & Col	GTAAGACGGAGOOCC TGAAG	AACATGTAAAGCCAATAC OCTCTC
T15D9	T15D9.5	primer	86724-87494	Lan & Col	TCGGAAAGGCTAGAG ATGGGTAAC TG	ATTGGACTATATGGGCCTC GTGAC
T15D9	T15D9.93	primer	93763	Lan & Col	TTTGGGGATATTCTAA AGGTGATG	TACTATTGGCCTGCTGTTG AGG
F7K9	F7K9.3	primer	21647-22276	Lan & Col	GGATGCAATGCCCGTT ATGATG	TCGAGGGAGGATGCTGAGT ATG
F7K9	F7K9.2	primer	12216-12843	Lan & Col	CAAAGCGGOCATCTC CTTC	GCAATTCATACCGCCACAT CTG
F7K9	F7K9.1	primer	3590-4226	Lan & Col	ACTATGCGTGGGTGG CTTTGIG	CAGGGGCATGGGAATCTC
F12P23	F12P23.3	primer	61772-62430	Lan & Col	AGCGAGGTTATCTATC AGGGTIG	GATTAGGTCGGCTTCTTCC AGTTAG
F12P23	F12P23.5	primer	44870-45511	Lan & Col	CTTCATTGTCATCATC GTATTAG	GGTGTGAAAGTCTGAGGCTC CC
F12P23	F12P23.4	primer	40880-41507	Lan & Col	TACCCATGOCCTGACT GCTG	TTCTGAAOGTGTGTGTCT ATTTG
F12P23	F12P23.2	primer	22431-23107	Lan & Col	TGTGCGAACTAATTGG TGGGAAC	TCGGGAAGAGTGCTAAG AG
F12P23	F12P23.1	primer	3352-4026	Lan & Col	ACAATGGCAACAATG GGCTGATAG	TTCCGGTGGTTGTCTCTAA AG
T4D8	T4D8.5	primer	81647-82250	Lan & Col	CTCGGTCGTGTAATGT GAAGTGGT	ATACGTCGGGGAGTTGAG
T4D8	T4D8.3	primer	47146-47883	Lan & Col	GCCCGTCTGCCATCTC TATC	CGCCTCCTCACAGCCACA A
T4D8	T4D8.2	primer	21848-22453	Lan & Col	AAACTGGCCGCTCGT GTAAC	AGGATAAAOCCATAGCTTG ACCAG
T4D8	T4D8.1	primer	18915-19589	Lan & Col	CTCGTCTCATCCAAAT COGTCC	CAATATAACCCCGTCCCGT GAAG

FIG. 14A (cont'd)

Chromosome IV

Sequenced clone	marker name	marker or primer pair	Marker location	marker properties, position	Forward Primer	Reverse primer
T5H22	T5H22.00	primer	248-2654	Lan & Col SSLP	TTTGTTACCCCTTTGGC TCGGACTGG	AAGGGGACACGCAC AAAAACGCTCTC
T5H22	T5H22.21	primer	21508-22868	Col Dom	GTCGCCCTGGTCTAG TAAATGG	CTGTTCGTCCCTTC TGCTG
T5H22	T5H22.41.3	primer	35072-35719	Lan & Col	TTGCGAGAACTTGGC AGGAACATC	TTAGAAAAGCATCG GGCACCAAAC
T5H22	T5H22.41.4	primer	64404-65591	Col Dom	CTCCCTCGCATATTTG TGA CTG	GTTGCCAAAAGTTCTC TACGATTC
T5H22	T5H22.65	primer	65036-66470	Lan & Col	TCAACCTAAGGCAAAT TTTCTAAG	TTTAATGAAGGCC AACACC
T7M24	T7M24.04	primer	4816-8214	Lan & Col	GTGCATGGCCTAAACA ACAG	GTTCCTATAACGGGT CAGTCC
T7M24	T7M24.46	primer	46240-47868	Col Dom	ATGTTATGTTTACGTC GGGGTTGTGTTG	TCTGGCTCCGGATG CTATTTGTATTTTC
T25H8	T25H8.01	primer	1889-2953	Lan & Col	TGACGAAGAAGGGGG AAAAGTTG	TGACGTGGTGAAAG TAGGCTGTGAAG
T25H8	T25H8.17	primer	16846-17990	Col dom	ACTAAAGCCCAACTG AAGAGGAAG	AAACCCCACTACC GCCATAA
T25H8	T25H8.22.9	primer	22482-25074	Ler & Col	AATCGATCCGTCTTTC ACCAAC	CTTCCTGCAGCCGTT CTTC
T24M8	T24M8.65	primer	65402-66309	Lan & Col	CGGCATGACCAAACCC TAAACTC	AGGGGAAAGATGAA AGATGAAATAAG
T24M8	T24M8.54	primer	53851-54655	Col Dom	TAATAAACCGCTCAGC CACC ACTCTAAG	GGGCTGCTCAATCT CGCTACAC
T24M8	T24M8.43	primer	42439-43274	Lan & Col	CTFAAATTGCCCGTGA TGATGGTTG	GATGGAGTCGGCAA AAGATAGGATG
T24M8	T24M8.22	primer	22640-23386	Lan & Col	ACGAGAAGCGAAAAAC CGAAGATAG	CGAACCTAAACCAA ACCTAAACTGAATC
T24M8	T24M8.09	primer	5961-8374	Lan & Col	GAGTTCGGGGTAATT TCCTCTCG	ATTCTTGGGTGCC TGGTGTAAAC
T24H24	T24H24.82	primer	82814-82890	Ler & Col	AGACAGCCGGAAGCA ATGGTGG	TCTCGTGTGGACA TACTCACTCAC
T24H24	T24H24.66	primer	66082-66765	Ler & Col	GTTGAAGGACCGGAGT TGTTAGAC	TGTGGATCGGTTATT GGAGGG
T24H24	T24H24.48	primer	47836-48636	Ler & Col	CCCCAGCCCATTGAG TGAGTAG	AGCGGGGCGCTTGA GAGTATC
T24H24	T24H24.11	primer	11212-11867	Ler & Col	GGCGGGTAGTTATGT TGATTGAG	TACCAAGGCCCGA GATACTAAC
T27D20	T27D20.77	primer	77681-78420	Lan & Col	TCGCGCAAATGGGACA CG	CGGGGAGGCTCGGG AATC
T27D20	T27D20.64	primer	64198-66686	Ler & Col	ACCTGCGATAGAGTTG TGAGTTC	CTGCCTTTGOOGATA ATAGTC
T27D20	T27D20.51	primer	51084-51783	Lan & Col	CCGCCGGCTTATGCTG AG	GAAGAGAAATGCC TGTGAGTCC
T27D20	T27D20.41	primer	41203-42770	Lan & Col	GTGATTGCGAGGACAT TGAGTG	TACATTTTTCAGCC ATTTTGTG
T27D20	T27D20.06	primer	6107-6887	Col Dom	TCGAATGGCTGAAAGA AAAGAATAAGAG	AAAAACGGTGGCGG AGAATG
T19B17	T19B17.96	primer	96402-97060	Lan & Col	CGTCTCCGTGAGGTG GC	ATTTTCATAATTATT TGGCGTGTGC
T19B17	T19B17.77	primer	77318-78093	Lan & Col	GGAACCCATCCGAAC TAAC	TGCCACAACAACT CCACTATG

FIG. 14B

Chromosome IV

F14G16	F14G16.81	primer	22905-23604	Lan & Col	CGTCTTCATCGGCTTC GTTCAG	TGGGGAGCGGAGGA TTCTTG
F14G16	F14G16.66	primer	37689-38299	Lan & Col	AGCGATTGTACCCCA CCATTC	GCTCCGGCAATCTTC TTCTCTC
F14G16	F14G16.49	primer	54150-54777	Lan & Col	ACTTTGGGCAATGAAG CGTATG	AACCCCTTAGGATT ATTCTAGTGTTC
F28D6	F14G16.32	primer	8172-8825	Lan & Col	TCTCGCAGTTGCAGAG ATGGTG	TCCGGAAGAGAAG AGTGATGG
F28D6	F14G16.66	primer	9445-10055	Lan & Col	AGCGATTGTACCCCA CCATTC	GCTCCGGCAATCTTC TTCTCTC
F28D6	F14G16.16	primer	24251-24873	Lan & Col	TGGTGTATTTTTCCTT GTTTCTCAGG	GTGTTCGCTATGG GGCTAAGG
F28D6	F14G16.01	primer	39801-40577	Lan & Col	GTGCGGAAATGTCTGG GCTC	AATCACTCAACCGC GAAACTCTATC
F28D6	F28D6.42	primer	42565-43225	Lan & Col	ATCAAACCCCAATCA CCAGAAAC	AATCGGGTTAGCC ACTTCATC
F28D6	F28D6.50k	MARKER	50323	Col Dom	CGGCTGGCTTTATTAT CTGAGTTG	TTCGGGAAGCCTGT GGAAG
F28D6	F28D6.58	primer	58994-59869	Col Dom	ACCCCGAGCTCAACTT CTTAGG	GGACGGGAGATGGG ATTACC
F28D6	F28D6.76	primer	76571-77289	Lan & Col	AGAATAGGAGCTGGG AGGTCAAAC	ATACTTAGATGCAA TGGGTGTGGTG
F28D6	F28D6.93	primer	93823-94512	Lan & Col	CCCCATCCTGCCGACA TAAAG	TACTCCGCATCATCT TCCATCTCTTC
F28D6	F28D6.120	primer	7985-8702	Lan & Col	GAGGGCGAGTAGTTG AATCTGC	CCTAAGCCCGAAAC CAAGTGAG

FIG. 14B (cont'd)

Chromosome IV

T1J1	T1J1.50	primer	50248-50937	Lan & Col	AAGTCTGGGAAGAGG ATGAGAAACCC	ATAAAGTACGCCGC CCATCAATAG
T32N4	T32N4.09	primer	10175-11108	Lan & Col	GGCAGATACGGCGGGT CCATAC	TCTGAATGCGATCTC CTCGTGTAAG
T32N4	T32N4.24	primer	24917-25724	Lan & Col	CGTGGGAGCTGCGGTA GAAG	GCGGTTGATGATGA AAA'AGGGTG
T32N4	T32N4.45	primer	45840-46451	Lan & Col	CGCCCCCTCAGGTTAG TCC	GTTTGCTCCCCCTCC AGTG
T32N4	T32N4.46	primer	46637-47558	Col Dom	CTGGCGTAOGAGAGTG CTGTG	ATGACCCGTGTGCTTT TGCTCCTC
T32N4	T32N4.60	primer	60777-61645	Lan & Col	CTCTCGGCGTTGCTTCT GG	GCCCGGCTGGTGCT ATTG
T32N4	T32N4.66	primer	66497-67374	Col Dom	AAAGAAGCGAAACAA CATAACCATAG	GGAGACAAAGAAAT CGGCAGAGTAG
T1J24	T1J24.114	primer	114825-115648	Lan & Col	CATGCCCGAATTACGA CACTC	GCGCCAAATCTCTA AACAACACTC
T1J24	T1J24.90	primer	90665-91646	Lan & Col	AATGAATGGGACGAA AAGGAAACT	GCATCCCCGGTACT GGTGAG
T1J24	T1J24.81	primer	80921-81638	Col Dom	AATCGGACTTTGCGT TCC	TAAACTACTATCCCA CCACCCTACC
T1J24	T1J24.79	primer	79569-80351	Lan & Col	GTGTATCGGGGGCCAT CTCAG	GCTCAACATCGCCG CAATCT
T1J24	T1J24.61	primer	60440-61245	Lan & Col	CCCAAAGTATAAGCGC CCACCTA	TAAAGCGCTCACTTC ACCATTG
T1J24	T1J24.51	primer	51061-51798	Lan & Col	TCCGGAAGGAGCCACA TAAG	TCCCAGACCTCTCG TTGAC
T1J24	T1J24.27	primer	27855-28895	Lan & Col	GGCCGGGAGTTGGTCA TAAGG	TCAATTTCAATCCCC GCTGGTC
T1J24	T1J24.23	primer	23943-24800	Lan & Col	TGGTGGGCATATTGT TTTTCTGTG	CGGCGCTGCCCTG GTTCC
T1J24	T1J24.01	primer	683-1663	Lan & Col	TTCCCCAAAATCGTT CAGC	ACATCGCTCTTCAA CCCCTC
F6H8	F6H8.70	primer	unknown-unique seq	Lan & Col -	ACCCGAGAAGCCGATG ACC	AAATTTGGGGGAGT TGATAAGTGTG
F6H8	F6H8.51	primer	unknown	Lan & Col -	GCTAAGCCATCCAAGT TCTGAG	GTTTGAGTCTTTGGC TTTGATGTTC
F6H8	F6H8.94	primer	unknown	Lan & Col -	CGTGCAGGGGAGTGTG GTG	CAATTTCAATCCCOG CTGGTC
F6H8	F6H8.114	primer	unknown	Lan & Col -	CGGGCTGCTTCATG TATCTATC	GOCCATTTGTGCGCT TATTCTATTG
F2I12	F2I12.82	primer	82463-83233	Lan & Col	TTTTGGGGATAGGGA TTGAGTGTG	TAAGCGGAAGGAGA GGTTTGAAGTGTG
F2I12	F2I12.70	primer	70415-71220	Lan & Col	TGCTGGCGCTTTGTGATC TATTGTG	COGCGGGGACTGOC TACTC
F2I12	F2I12.68	primer	68874-69938	Lan & Col	CCAGAGCGGGGAAA GCAATAC	TAGCCGGGGTGGTC TCGTGG
F2I12	F2I12.50	primer	50288-50891	Lan & Col	TGACTATAGGGGCGGT TGTGGTAAG	TTGCGTTGGAGTTTG CGTGGTC
F2I12	F2I12.48	primer	48960-50345	Col	ACCTTCTTCTCAACG CACTCAACC	AAOCCCTTGGCATAT AACTCOGACTC
F2I12	F2I12.29	primer	29895-30702	Lan & Col	GTGGGGTCGAGTGGTG TGGTAG	GGATCOCCGTGTTACT TAAGCCTATTG
F2I12	F2I12.02	primer	2313-3098	Lan & Col	AAAATCCTCCCGCGTC AACATC	CATCATCCCAATCCC AAATACAAGTC
F14G16	F14G16.100	primer	3496-4174	Lan & Col	AAACTTGGCACTCT CCTCTATTATG	ATTTGGTAAGGGG TTGATGACTC

FIG. 14B (cont'd)

Chromosome IV

T19B17	T19B17.59	primer	59092-59808	Lan & Col	AGCGGTCAATGTTCTT CAATGTCGTAG	TATCGCGCGGAGT CAGGAG
T19B17	T19B17.44	primer	44057-44788	Col Dom	CTGCCCCGACCACCTT TCAAC	TTGCGGATTCGTTAT GCTGTTCTC
T19B17	T19B17.30	primer	30680-31352	Lan & Col	ATCGCCCGCGTCTTCTT CAG	CAGGTTCAAGCCGGT CAACTATAATC
T19B17	T19B17.11	primer	11260-12044	Lan & Col	TCATTTGCGTCTAGAG GTGGAGTGC	GGGGTAGAAAGAAG CGAGAGGGATAG
T26N6	T26N6.12	primer	12724-13462	Lan & Col	CAGGCATCATTTCATC AAACGAG	GTAGGATCCGGCTG AATAGTGGTGG
T26N6	T26N6.27	primer	27839-28536	Lan & Col	TCTTCCGATGACGACA ACGACAC	ATTCTGCTGCTGCTG ATTCTCTG
T26N6	T26N6.42	primer	43996-44639	Col only	GACGGCCTTTTCATTCT CACACAG	TTTCATATTTGCTCA TCTAACCCCTC
T26N6	T26N6.59	primer	59333-59938	Col only	GCCTCGAACCCCTACAC CTCCAC	AGTCGCGTAGCAA ATGAAAACC
T26N6	T26N6.74	primer	74460-75083	Lan & Col	ATGGGGCCCTTTGACT ACTGAG	TCCGGAGACGATTTT GATGAC
T26N6	T26N6.93	primer	93352-93986	Lan & Col	TTCCCGCATGCATTAG TTCTTGIG	TTGOCATCATCTTTC TGTGTTTGTCTATC
F4H6	F4H6.44	primer	42567-43173	Lan & Col	GCAGACGCGAGGACA CAGACAG	CAGCCTAAGCCCAT TTGTTTTGAAG
F4H6	F4H6.60	primer	60209-60835	Lan & Col	GTTC AACGCTAGCAA GGTCTG	AGGGCCAACATGC ACTACAAG
F4H6	F4H6.82	primer	82859-83642	Lan & Col	ACAAATCAGAGGOCOA AAGTCAATG	TGGGCGAATAACA GCAAGTOC
F4H6	F4H6.100	primer	100331-101001	Lan & Col	ATCCAAACGCCCAAAT GTCAAC	TTAAGTCCGGTGG GTTCAAATAC
T19J18	T19J18.12	primer	12781-13435	Lan & Col	ATGCCATAAAGAAAG CCAGTC	CGCCTATCTTCGGTG TCTOGTC
T19J18	T19J18.27	primer	28093-29954	Lan & Col	CAGCGCTGTACAGTGG TCAAATG	CGTGGTTCAGGTGG GTCAGG
T4B21	T19J18.71	primer	6380-7009	Lan & Col	CATTACTTACCGCTTC CGTCTTATC	AATGTTAGTGGAG TTTATGGTTGTGTC
T4B21	T19J18.42	primer	20045-20648	Lan & Col	TGTCGCCTTACTCCATT CGTTCAAC	CGGCCGCTTCATGT ATCTATCTC
T4B21	T4B21.20	primer	21757-22522	SSLP polymorphic	AATAGGCCTTTCCGGT GCTTCTC	AATTGATTTTGGGGT TTCCTGTTC
T4B21	T4B21.35	primer	37346-38074	Lan & Col	GTGAAAGGAGCAGCA GGAACAGTG	ATTTATAGGCCAAT GAOCCAATCG
T4B21	T19J18.57	primer	38498-39157	Lan & Col	CTATCAAAACGAGTCA AAGAAAGG	AGAAAGTGAGOCOA AGAGATTAGTG
T4B21	T4B21.52	primer	54320-55077	Lan & Col	ATAGACAAAATTGGCA ACACATAAC	CAGCCACTCTTCAT CTCCTTTC
T4B21	T4B21.68	primer	69927-70543	Lan & Col	TTGTCATTGGCGCTGC TCTATC	GCTTCCCAOCCAAT ATCCTTTC
T4B21	T4B21.83	primer	85772-86299	Lan & Col	AAGCCCGGATTGGT TC	CGCTACGCATGGGT CTATTTG
T1J1	T1J1.08	primer	8862-9483	Lan & Col	TAGAGGGTAACITAA CGAATGTGC	ATGTGGGGCCAAAT AAATCAAAAC
T1J1	T1J1.23	primer	23155-23843	Lan & Col	TGGAGGGCTTGCATGT GAGAGTG	CAGAGCCGGATGAG AAAACAGAGC
T1J1	T15D16	marker	38027	CAPS	AATCAATTGGTTTCTA CTTTTTAG	AACTCCGACTGAAG GTATAAC
T1J1	T1J1.39	primer	39177-40174	Lan & Col	ACCGGCTCATTGGCTA AAAAGTTC	TTAAGGGTTGGGGT TCATCTGTAC

FIG. 14B (cont'd)




FIG. 15

## METHODS FOR GENERATION OR INCREASING REVENUES FROM CROPS

[0001] This application is a continuation-in-part of U.S. patent application Ser. No. 09/553,231, filed Mar. 13, 2002, which is a continuation of U.S. patent application Ser. No. 09/090,051, filed Jun. 3, 1998, now U.S. Pat. No. 6,156,953 which claims the priority of U.S. Provisional Patent Application Ser. No. 60/048,451, filed Jun. 3, 1997; and U.S. Provisional Patent Application Ser. No. 60/073,741, filed Feb. 5, 1998, both of the disclosures of which are specifically incorporated herein by reference in their entirety. This application is also a continuation-in-part of U.S. patent application Ser. No. 09/531,120, filed Mar. 17, 2000, which claims the priority of U.S. Provisional Application Ser. No. 60/125,219, filed Mar. 18, 1999; U.S. Provisional Application Ser. No. 60/127,409, filed Apr. 1, 1999; U.S. Provisional Application Ser. No. 60/134,770, filed May 18, 1999; U.S. Provisional Application Ser. No. 60/153,584, filed Sep. 13, 1999, U.S. Provisional Application Ser. No. 60/154,603, filed Sep. 17, 1999 and U.S. Provisional Application Ser. No. 60/172,493, filed Dec. 16, 1999, each of which disclosures is specifically incorporated herein by reference in its entirety.

[0002] The government owns rights in the present invention pursuant to U.S. Department of Agriculture Grant No. 96-35304-3491 and Grant No. DE-FC05-92OR22072 from the Consortium for Plant Biotechnology Research, National Science Foundation Grant No. 9872641, and Department of Energy Small Business Innovation Research Grants DE-FG02-01 ER83163, DE-FG02-01 ER83165, and DE-FG02-01 ER83166.

### BACKGROUND OF THE INVENTION

[0003] The present invention relates generally to methods of doing business. More specifically, the present invention relates to methods of generating and/or increasing the revenue derived from crops.

[0004] To improve agricultural traits or produce certain products, it has been known to add a gene to an organism such as a plant. Although genes, for a vast array of products, have been proposed or identified, the technologies in the industry, heretofore, have severely curtailed delivery of several genes into a plant.

[0005] Two general approaches are used for introduction of new genetic information ("transformation") into cells. One approach is to introduce the new genetic information as part of another DNA molecule, referred to as an "episomal vector," or "mini chromosome", which can be maintained as an independent unit (an episome) apart from the host chromosomal DNA molecule(s). Episomal vectors contain all the necessary DNA sequence elements required for DNA replication and maintenance of the vector within the cell. Many episomal vectors are available for use in bacterial cells (for example, see Maniatis et al., 1982). However, only a few episomal vectors that function in higher eukaryotic cells have been developed. Higher eukaryotic episomal vectors were primarily based on naturally occurring viruses. In higher plant systems gemini viruses are double-stranded DNA viruses that replicate through a double-stranded intermediate upon which an episomal vector could be based, although the gemini virus is limited to an approximately 800 bp insert. Although an episomal plant vector based on the Cauliflower

Mosaic Virus has been developed, its capacity to carry new genetic information also is limited (Brisson et al., 1984).

[0006] The other general method of genetic transformation involves integration of introduced DNA sequences into the recipient cell's chromosomes, permitting the new information to be replicated and partitioned to the cellos progeny as a part of the natural chromosomes. The introduced DNA usually is broken and joined together in various combinations before it is integrated at random sites into the cell's chromosome (see, for example Wigler et al., 1977). Common problems with this procedure are the rearrangement of introduced DNA sequences and unpredictable levels of expression due to the location of the transgene in the genome or so called "position effect variation" (Shingo et al., 1986). Further, unlike episomal DNA, integrated DNA cannot normally be precisely removed. A more refined form of integrative transformation can be achieved by exploiting naturally occurring viruses that integrate into the host's chromosomes as part of their life cycle, such as retroviruses (see Cepko et al., 1984). In mouse, homologous integration has recently become common, although it is significantly more difficult to use in plants (Lam et al. 1996).

[0007] The most common genetic transformation method used in higher plants is based on the transfer of bacterial DNA into plant chromosomes that occurs during infection by the phytopathogenic soil bacterium *Agrobacterium* (see Nester et al., 1984). By substituting genes of interest for the naturally transferred bacterial sequences (called T-DNA), investigators have been able to introduce new DNA into plant cells. However, even this more "refined" integrative transformation system is limited in three major ways. First, DNA sequences introduced into plant cells using the *Agrobacterium* T-DNA system are frequently rearranged (see Jones et al., 1987). Second, the expression of the introduced DNA sequences varies between individual transformants (see Jones et al., 1985). This variability is presumably caused by rearranged sequences and the influence of surrounding sequences in the plant chromosome (i.e., position effects), as well as methylation of the transgene. A third drawback of the *Agrobacterium* T-DNA system is the reliance on a "gene addition" mechanism: the new genetic information is added to the genome (i.e., all the genetic information a cell possesses) but does not replace information already present in the genome.

[0008] One attractive alternative to commonly used methods of transformation is the use of an artificial chromosome. Artificial chromosomes are man-made linear or circular DNA molecules constructed from cis-acting DNA sequence elements that provide replication and partitioning of the constructed chromosomes (see Murray et al., 1983). Desired elements include: (1) Autonomous Replication Sequences (ARS) (these have properties of replication origins, which are the sites for initiation of DNA replication), (2) Centromeres (site of kinetochore assembly and responsible for proper distribution of replicated chromosomes at mitosis or meiosis), and (3) if the chromosome is linear, telomeres (specialized DNA structures at the ends of linear chromosomes that function to stabilize the ends and facilitate the complete replication of the extreme termini of the DNA molecule).

[0009] The essential chromosomal elements for construction of artificial chromosomes have been precisely characterized in lower eukaryotic species, and more recently in mouse and human. ARSs have been isolated from unicellular fungi, including *Saccharomyces cerevisiae* (brewer's yeast) and *Schizosaccharomyces pombe* (see Stinchcomb et al., 1979



and Hsiao et al., 1979). An ARS behaves like a replication origin allowing DNA molecules that contain the ARS to be replicated as an episome after introduction into the cell nuclei of these fungi. DNA molecules containing these sequences replicate, but in the absence of a centromere they are partitioned randomly into daughter cells.

**[0010]** Artificial chromosomes have been constructed in yeast using the three cloned essential chromosomal elements. Murray et al., 1983, disclose a cloning system based on the in vitro construction of linear DNA molecules that can be transformed into yeast, where they are maintained as artificial chromosomes. These yeast artificial chromosomes (YACs) contain cloned genes, origins of replication, centromeres and telomeres and are segregated in daughter cells with high fidelity when the YAC is at least 100 kB in length. Smaller CEN-containing vectors may be stably segregated, however, when in circular form.

**[0011]** None of the essential components identified in unicellular organisms, however, function in higher eukaryotic systems. For example, a yeast CEN sequence will not confer stable inheritance upon vectors transformed into higher eukaryotes. While such DNA fragments can be readily introduced, they do not stably exist as episomes in the host cell. This has seriously hampered efforts to produce artificial chromosomes in higher organisms.

**[0012]** In one case, a plant artificial chromosome was discussed (Richards et al., U.S. Pat. No. 5,270,201). However, this vector was based on plant telomeres, as a functional plant centromere was not disclosed. While telomeres are important in maintaining the stability of chromosomal termini, they do not encode the information needed to ensure stable inheritance of an artificial chromosome. It is well documented that centromere function is crucial for stable chromosomal inheritance in almost all eukaryotic organisms (reviewed in Nicklas 1988). For example, broken chromosomes that lack a centromere (acentric chromosomes) are rapidly lost from cell lines, while fragments that have a centromere are faithfully segregated. The centromere accomplishes this by attaching, via centromere binding proteins, to the spindle fibers during mitosis and meiosis, thus ensuring proper gene segregation during cell divisions.

**[0013]** In contrast to the detailed studies done in *S. cerevisiae* and *S. pombe*, less is known about the molecular structure of functional centromere DNA of higher eukaryotes. Ultrastructural studies indicate that higher eukaryotic kinetochores, which are specialized complexes of proteins that form on the chromosome during late prophase, are large structures (mammalian kinetochore plates are approximately 0.3  $\mu\text{m}$  in diameter) which possess multiple microtubule attachment sites (reviewed in Rieder, 1982). It is therefore possible that the centromeric DNA regions of these organisms will be correspondingly large, although the minimal amount of DNA necessary for centromere function may be much smaller.

**[0014]** The above studies have been useful in elucidating the structure and function of centromeres. The extensive literature indicating both the necessity of centromeres for stable inheritance of chromosomes, and the non-functionality of yeast centromeres in higher organisms, demonstrate that cloning of a functional centromere from a higher eukaryote is a necessary first step in the production of artificial chromosomes suitable for use in higher plants and animals. The production of artificial chromosomes with centromeres which function in higher eukaryotes would overcome many

of the problems associated with the prior art and represent a significant breakthrough in biotechnology research.

#### SUMMARY OF THE INVENTION

**[0015]** The present invention provides methods for improving crops as well as reducing the time necessary to produce new crops. Pursuant to the present invention, methods are provided that allow one to increase revenues associated with crops, develop new crops, develop new avenues for generating revenues from crops, and provide new services to a third party.

**[0016]** By allowing for the isolation and identification of plant centromere DNA sequences from the total genomic DNA of an organism or fractions thereof it is possible to construct chromosomes having functional centromeres and carrying large number of genes. As noted above, genes for producing a vast set of products have been identified, but technologies used within the industry severely limit the delivery of these genes to plant cells. One or at most a few genes are typically inserted into random locations in the host chromosomes, which can irreversibly disrupt host gene functions while causing variable and uncontrolled expression of the introduced genes. The present invention makes it possible to overcome the technical limitations associated with gene delivery in crop species, thereby allowing for the ability to shorten the time required for crop development.

**[0017]** To this end, in an embodiment, the present invention provides a method for providing a service. The method comprises the steps of: identifying a sequence associated with a centromere of a crop; using a gene or a number of genes and the sequence to create a minichromosome; and introducing the minichromosome into a cell of the commercial plant to create a transgenic plant having a desirable characteristic not present in the crop.

**[0018]** In an embodiment, the sequence is isolated after it is identified.

**[0019]** In an embodiment, the commercial plant is chosen from the group consisting of vegetable crop, fruit and vine crops, and field crop plants.

**[0020]** In an embodiment, the desirable characteristic is chosen from the group consisting of insect resistances, disease resistances, herbicide resistances, environmental resistances, agronomic characteristics, nutrient utilization, nutrient content, production of bioproducts, production of pharmaceutical products, production of chemical products, chemical assimilation, and reduction of length to maturity.

**[0021]** In an embodiment, the gene is selected from the group consisting of a selectable marker gene, a screenable marker gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene, a gene encoding an enzyme, a gene encoding an antibody, a gene encoding an antigen for a vaccine, a transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones.

**[0022]** In an embodiment, the new commercial plant has at least two desirable characteristics not present in the commercial plant.

**[0023]** In an embodiment, the minichromosome includes at least two genes each producing a different desirable characteristic.

**[0024]** In an embodiment, the method includes the step of producing seeds that generate the new commercial plant.

**[0025]** In an embodiment, the method includes the step of extracting the desired product from the new commercial plant.

**[0026]** In a further embodiment, the present invention provides a method of increasing revenues from a species of plant. The method comprises the steps of: identifying a sequence associated with a centromere of the species of plant; using a gene or a number of genes and the sequence to create a minichromosome; and introducing the minichromosome into the species of plant to create a plant having a desirable characteristic not naturally present in the species plant or to increase or decrease the level of a naturally occurring characteristic.

**[0027]** In yet another embodiment of the present invention, a method for increasing the value of a specific crop is provided. The method comprises the steps of: identifying a sequence associated with a centromere of the crop; using a gene or a number of genes and the sequence to create a minichromosome; and introducing the minichromosome into the crop to create a transgenic plant having a characteristic that causes the modified crop to be more valuable.

**[0028]** Still further, in an embodiment, a method for increasing crop yield is provided. The method comprises the steps of: identifying a sequence associated with a centromere of the crop; using a gene or a number of genes and the sequence to create a minichromosome; and introducing the minichromosome into the crop to create a transgenic plant having a desirable characteristic not naturally present in the crop or to increase or decrease the level of a naturally occurring characteristic.

**[0029]** Moreover, in an embodiment, the present invention provides a method for producing a new seed. The method includes the steps of: identifying a sequence associated with a centromere of the commercial plant; using a gene or a number of genes and the sequence to create a minichromosome; introducing the minichromosome into the commercial plant to create a transgenic plant having a desirable characteristic not present in the commercial plant or to increase or decrease the level of a naturally occurring characteristic; allowing the transgenic plant to mature; and collecting seeds from the mature plant.

**[0030]** Further, the present invention provides a method for generating revenue based on agricultural products. The method comprises the steps of: developing a new crop based on a minichromosome comprising a sequence of a centromere; and receiving a fee for the new crop.

**[0031]** In an embodiment, the fee is received as a royalty payment.

**[0032]** In an embodiment, the fee is received through sales of seed generated by the new crop.

**[0033]** In an embodiment, the fee is based on sales of a product made from the new crop.

**[0034]** In an embodiment, the fee is based on licensing of the new crop or product.

**[0035]** In an embodiment, the fee is based on the receipt of products or services.

**[0036]** In addition, in an embodiment, the present invention provides a method for performing a service comprising the steps of: developing a modified plant having a desired characteristic that does not naturally occur in a non-modified plant by inserting into a cell of the non-modified plant a minichromosome comprising a sequence of a centromere and a gene expressing the desired characteristic.

**[0037]** In a still further embodiment, the present invention provides a method for performing a service for a third party including the step of: isolating from a total genomic DNA of a plant a nucleotide sequence relating to the plant's centromere and using the nucleotide sequence to make a minichromosome for the third party.

**[0038]** Yet another embodiment of the present invention provides methods for establishing strategic relationships with a third party. The method can comprise the steps of initiating a research and development program with a third party to develop a crop that includes a mini chromosome comprising a gene and a centromere sequence.

**[0039]** If desired, the method can include the step of working with the third party to develop the crop.

**[0040]** In an embodiment, the method includes the step of licensing to the third party at least a portion of the technology necessary to develop the crop.

**[0041]** In order to obtain a centromere DNA sequence from a selected organism, in an embodiment, the method that is used comprises the steps of preparing a sample of genomic DNA from a selected organism, obtaining a plurality of nucleic acid segments from the genomic DNA and screening the nucleic acid segments to identify one or more centromere nucleic acid sequences. In an embodiment, the method of obtaining the plurality of nucleic acid segments comprises contacting said genomic DNA with a restriction endonuclease and selecting nucleic acid segments containing repetitive DNA to obtain said plurality of nucleic acid segments. In another embodiment, the method of obtaining the plurality of nucleic acid segments comprises contacting said genomic DNA with a methylation sensitive restriction endonuclease and selecting nucleic acid segments exhibiting resistance to cleavage with said methylation sensitive restriction endonuclease to obtain said plurality of nucleic acid segments. In yet another embodiment, the method of obtaining the plurality of nucleic acid segments comprises contacting said genomic DNA with a restriction endonuclease or physically shearing said genomic DNA and selecting nucleic acid segments that anneal rapidly after denaturation to obtain said plurality of nucleic acid segments.

**[0042]** In another aspect, the invention provides a method for identifying a centromere nucleic acid sequence from a dataset of the genomic sequences of an organism. The method comprises the steps of (1) providing a first dataset consisting of the genomic sequences, or a representative fraction of genomic sequence, of the organism; (2) identifying and eliminating known non-centromeric repeat sequences from the first dataset by using the BLAST sequence-comparison algorithm to create a second dataset; (3) comparing each sequence in the second dataset to itself by using the BLAST sequence comparison algorithm, obtaining a BLAST score for each pair of sequence compared, and collecting high score pairs to create a third dataset; (4) examining the BLAST score of each high score pair in the third dataset and eliminating the pairs having a score greater than  $10^{-20}$  to create a fourth dataset; (5) eliminating the high score pairs in the fourth dataset having less than 80 bp or more than 250 bp to create a fifth dataset; (6)

examining the nucleotide position of each high score pair in the fifth dataset and eliminating pairs having 100% identity as well as identical nucleotide positions to create a sixth dataset; (7) examining the nucleotide position of each high score pair in the sixth dataset and eliminating pairs having opposite orientation of the nucleotides to create a seventh dataset; (8) examining the nucleotide position of both sequences for each high score pair in the seventh dataset and eliminating sequences that are overlapping to create an eighth dataset; and (9) examining the nucleotide position of each sequence in the eighth dataset and eliminating sequences not having at least one neighboring sequence within 250 bp to create a ninth dataset; and (10) comparing each sequence in the ninth dataset to all other sequences in the ninth dataset by using the BLAST sequence comparison algorithm and selecting the most common sequence as a centromere sequence of the organism. In one embodiment, the known non-centromeric repeat sequence in the second step is a ribosomal DNA.

**[0043]** A number of centromeres can be used pursuant to the present invention. In an aspect of the present invention, a *Brassica oleracea* centromere comprising *Brassica oleracea* centromere DNA is used. In one embodiment, the *Brassica oleracea* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Brassica oleracea* given by SEQ ID NO:1, 2, 3, or 4.

**[0044]** In yet another aspect, a *Glycine max* centromere comprising *glycine max* centromere DNA is used. In an embodiment, the *Glycine max* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100; 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Glycine max* given by SEQ ID NO:5, 6, 7, or 8.

**[0045]** In yet another aspect, a *Lycopersicon esculentum* centromere comprising *Lycopersicon esculentum* centromere DNA is used. In an embodiment, the *Lycopersicon esculentum* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Lycopersicon esculentum* given by SEQ ID NO:9 or 10.

**[0046]** In yet another aspect, a *Zea mays* centromere comprising *Zea mays* centromere DNA is used. In an embodiment, the centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially

any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Zea mays* given by SEQ ID NO:11, 12 or 13.

**[0047]** In yet another aspect, recombinant DNA construct comprising a plant centromere sequence is used. The recombinant DNA construct may additionally comprise any other desired sequences, for example, a telomere. Still further, one may wish to include a structural gene on the construct, or multiple genes. Examples of structural genes one may wish to use include a selectable or screenable marker gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, a gene encoding an enzyme, a gene encoding an antibody, a gene encoding an antigen for a vaccine, a transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones.

**[0048]** In still yet another aspect, a recombinant DNA construct comprising a plant centromere sequence and which is capable of being maintained as a chromosome, wherein the chromosome is transmitted in dividing cells is used. The plant centromere may be from any plant or may be from any other source of DNA or may be partially or entirely synthetic in origin.

**[0049]** In yet another aspect, a recombinant DNA construct comprising a plant centromere sequence and which is a plasmid is used. The plasmid may contain any desired sequences, such as an origin of replication. The plasmid may also comprise a selection marker.

**[0050]** In still yet another aspect, a mini chromosome comprising a plant centromere sequence and may also contain a telomere sequence is used. Any additional desired sequences may be added to the mini chromosome, such as an autonomous replicating sequence and a structural gene such as those described above. The mini chromosome may comprise any of the centromere compositions disclosed herein.

**[0051]** The mini chromosome also may contain "negative" selectable markers which confer susceptibility to an antibiotic, herbicide or other agent, thereby allowing for selection against plants, plant cells or cells of any other organism of interest containing a mini chromosome. The mini chromosome also may include genes or other sequences which control the copy number of the mini chromosome within a cell. One or more structural genes also may be included in the mini chromosome. Specifically contemplated as being useful will be as many structural genes as may be inserted into the mini chromosome.

**[0052]** In still yet another aspect, a cell transformed with a recombinant DNA construct comprising a plant centromere sequence is used. The cell may be of any type, including a prokaryotic cell or eukaryotic cell. Where the cell is a eukaryotic

otic cell, the cell may be, for example, a yeast cell or a higher eukaryotic cell, such as plant cell. The plant cell may be from a dicotyledonous plant, such as tobacco, tomato, potato, soybean, canola, sunflower, alfalfa, cotton and *Arabidopsis*, or may be a monocotyledonous plant cell, such as wheat, maize, rye, rice, turfgrass, oat, barley, sorghum, millet, and sugarcane. In one embodiment of the invention, the plant centromere is a centromere chosen from the group consisting of *Brassica oleracea*, *Glycine max*, *Lycopersicon esculentum*, and *Zea mays* and the cell may be a cell chosen from one of the above species or any other species. The recombinant DNA construct may comprise additional sequences, such as a telomere, an autonomous replicating sequence (ARS), a structural gene or genes, or a selectable or screenable marker gene or genes, including as many of such sequences as may physically be placed on said recombinant DNA construct. In one embodiment of the invention, the cell is further defined as capable of expressing said structural gene. In another embodiment of the invention, a plant is provided comprising the aforementioned cells.

**[0053]** In still yet another aspect, a method for preparing a transgenic plant cell is used. The method comprises the steps of contacting a starting plant cell with a recombinant DNA construct comprising a plant centromere sequence of the present invention, whereby the starting plant cell is transformed with the recombinant DNA construct.

**[0054]** In still yet another aspect, a transgenic crop comprising a mini chromosome, wherein the mini chromosome comprises a plant centromere sequence of the present invention is used. The mini chromosome may further comprise a telomere sequence, an autonomous replicating sequence or a structural gene, such as a selectable or screenable marker gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, a gene encoding an enzyme, a gene encoding an antibody, a gene encoding an antigen for a vaccine, a transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones. The transgenic crop may be any type of crop, such as a dicotyledonous plant, for example, tobacco, tomato, potato, pea, carrot, cauliflower, broccoli, soybean, canola, sunflower, alfalfa, cotton and *Arabidopsis*, or may be a monocotyledonous plant, such as wheat, maize, rye, rice, turfgrass, oat, barley, sorghum, millet, and sugarcane.

**[0055]** In still yet another aspect, the method includes preparing a transgenic crop tissue. The method comprises the steps of contacting a starting crop tissue with a recombinant DNA construct comprising a plant centromere sequence of the present invention, whereby the starting crop tissue is transformed with the recombinant DNA construct.

**[0056]** In still yet another aspect, the method includes preparing a transgenic crop seed. The method comprises the steps of contacting a starting crop, crop tissue, or crop cell, with a recombinant DNA construct comprising a plant centromere sequence of the present invention, whereby the starting crop, crop tissue, or crop cell is transformed with the recombinant DNA construct. These transformed crops, crop

tissues, or crop cells are allowed to develop into mature crops, using standard agricultural techniques. Transgenic seed is then collected from these crops.

**[0057]** In still yet another aspect, the method includes preparing an extract of a transgenic crop, crop tissue, crop seed, or crop cell. The method comprises the steps of contacting a starting crop, crop tissue, or crop cell with a recombinant DNA construct comprising a plant centromere sequence of the present invention, whereby the starting crop cell is transformed with the recombinant DNA construct. The resulting transgenic crop, crop tissue, crop seed, or crop cell is then extracted and processed to yield the desirable product. One preferred desirable product is a food product. Another preferred desirable product is a pharmaceutical product. Yet another preferred desirable product is a chemical product.

**[0058]** Additional features and advantages of the present invention are described in, and will be apparent from, the following Detailed Description of the Invention and the figures.

#### BRIEF DESCRIPTION OF THE FIGURES

**[0059]** The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

**[0060]** FIG. 1A-F Consensus sequences of repeats from *Brassica oleracea*. FIG. 1A is the consensus sequence of ChrBo1. This consensus was assembled from 33 sequences collected by the inventors. The length of this repeat is  $180 \pm 0$ . 86 base pairs and A and T compose 60% of the consensus. FIG. 1B is the consensus sequence of ChrBo2. This consensus was assembled from 7 sequences collected by the inventors. The length of this repeat is  $180 \pm 0.45$  base pairs and A and T compose 63% of the consensus. FIG. 1C is a comparison of the consensus sequences of ChrBo1 and ChrBo2. The two repeats (ChrBo1 and ChrBo2) were aligned to each other using the ClustalX program (ClustalX is a free multiple sequence alignment program for Windows. Those sites with significant differences between the two sequences (Chi-squared,  $P < 0.05$ ) are highlighted. FIG. 1D is a revised consensus sequence of ChrBo1. This consensus was assembled from 33 DNA sequences collected by the inventors and 18 sequences from Genbank, identified by the accession numbers:

M30962	M30963	M31436	M31435
M31438	M31434	M31439	M31437
X68786	X12736	X07519	X16589
X15291	X68783	X68784	X61583
AJ228348	Z22947		

**[0061]** FIG. 1E is a revised consensus sequence of ChrBo2. This consensus was assembled from 7 DNA sequences collected by the inventors and 5 sequences from Genbank, identified by the accession numbers AJ228347, M30962, X12736, X61583, and X68785. FIG. 1F is a comparison of the revised consensus sequences of ChrBo1 and ChrBo2, aligned as for FIG. 1C.

**[0062]** FIG. 2A-F Consensus sequences of repeats from *Glycine max*. FIG. 2A is a consensus sequence of ChrGm1. This consensus was assembled from 32 sequences collected

by the inventors. The length of this repeat is  $92 \pm 0.79$  base pairs and A and T compose 63% of the consensus. FIG. 2B is a consensus sequence of ChrGm2. This consensus was assembled from 21 sequences collected by the inventors. The length of this repeat is  $91 \pm 0.48$  base pairs and A and T compose 62% of the consensus. FIG. 2C is a comparison of the consensus-sequences of ChrGm1 and ChrGm2. The two repeats (ChrGm1 and ChrGm2) were aligned to each other using the ClustalX program (ClustalX is a free multiple sequence alignment program for Windows. Those sites with significant differences between the two sequences (Chi-squared,  $P < 0.05$ ) are highlighted. FIG. 2D is a revised consensus sequence of ChrGm1. This consensus was assembled from 32 DNA sequences collected by the inventors and 1 sequence from Genbank, identified by the accession number Z26334. FIG. 2E is a revised consensus sequence of ChrGm2. This consensus was assembled from 21 DNA sequences collected by the inventors and 13 sequences from Genbank, identified by the accession numbers AF297983, AF297984, and AF297985. FIG. 2F is a comparison of the revised consensus sequences of ChrGm1 and ChrGm2, aligned as for FIG. 2C.

**[0063]** FIG. 3A-B Consensus sequences of repeats from *Lycopersicon esculentum*. FIG. 3A is a consensus sequence of ChrLe1. This consensus was assembled from 42 sequences collected by the inventors. The length of this repeat is  $181 \pm 0.61$  base pairs and A and T compose 50% of the consensus. FIG. 3B is a revised consensus sequence of ChrLe1. This consensus was assembled from 32 sequences collected by the inventors and 2 Genbank sequences identified by the accession numbers X87233 and AY007367.

**[0064]** FIG. 4A-C Consensus sequences of repeats from *Zea mays*. FIG. 4A is a consensus sequence of ChrZm1. This consensus was assembled from 38 sequences collected by the inventors. The length of this repeat is  $180 \pm 1.15$  base pairs and A and T compose 56% of the consensus. FIG. 4B is a revised consensus sequence of ChrZm1. This consensus was assembled from 38 sequences collected by the inventors and 26 sequences from Genbank, identified by the accession numbers:

M32521	M32522	M32523	M32524	M32525	M32526
M32527	M32528	M32529	M32530	M32531	M32532
M32533	M32534	M325375	M32536	M32537	M32538
M35408	AF030934	AF030935	AF030936	AF030937	AF030938
AF030939	AF030940				

**[0065]** FIG. 4C is a consensus sequence of ChrZm2. This consensus was assembled from 6 sequences collected from Genbank identified by the accession numbers:

AF078918	AF078919	AF078920
AF0789121	AF078922	AF078923

**[0066]** The length of this repeat is  $158 \pm 1.6$  base pairs and A and T compose 53% of the consensus.

**[0067]** FIG. 5 Mini chromosome containing centromere sequences as well as mini chromosome vector sequences

**[0068]** FIG. 6 Mini chromosome construct formed by mini chromosome vector tailing method.

**[0069]** FIG. 7A-7N. Exemplary Mini chromosome vectors: The vectors shown in FIG. 7A, FIG. 7B, FIG. 7E, FIG. 7F, FIG. 7I and FIG. 7J have an *E. coli* origin of replication which can be high copy number, low copy number or single copy. In FIGS. 7A-7N, the vectors include a multiple cloning site which can contain recognition sequences for conventional restriction endonucleases with 4-8 bp specificity as well as recognition sequences for very rare cutting enzymes such as, for example, I-Ppo I, I-Cue I, PI-Tli, PI-Psp I, Not I, and PI Sce I. In FIG. 7A-7N, the centromere is flanked by *Lox* sites which can act as targets for the site specific recombinase Cre. FIG. 7A. Shows an *E. coli* plant circular shuttle vector with a plant ARS. FIG. 7B. Shows a plant circular vector without a plant ARS. The vector relies on a plant origin of replication function found in other DNA sequences such as selectable or screenable markers. FIG. 7C. Shows a yeast-plant circular shuttle vector with a plant ARS. The yeast ARS is included twice, once on either side of multiple cloning site to ensure that large inserts are stable. FIG. 7D. Shows a yeast-plant circular shuttle vector without a plant ARS. The vector relies on a plant origin of replication function found in other plant DNA sequences such as selectable markers. The yeast ARS is included twice, once on either side of the multiple cloning site to ensure that large inserts are stable. FIG. 7E. Shows an *E. coli*-*Agrobacterium*-plant circular shuttle vector with a plant ARS. Vir functions for T-DNA transfer would be provided in trans by a using the appropriate *Agrobacterium* strain. FIG. 7F. Shows an *E. coli*-*Agrobacterium*-plant circular shuttle vector without a plant ARS. The vector relies on a plant origin of replication function found in other plant DNA sequences such as selectable markers. Vir functions for T-DNA transfer would be provided in trans by a using the appropriate *Agrobacterium* strain. FIG. 7G. Shows a linear plant vector with a plant ARS. The linear vector could be assembled in vitro and then transferred into the plant by, for example, mechanical means such as micro projectile bombardment, electroporation, or PEG-mediated transformation. FIG. 7H. Shows a linear plant vector without a plant ARS. The linear vector could be assembled in vitro and then transferred into the plant by, for example, mechanical means such as micro projectile bombardment, electroporation, or PEG-mediated transformation. FIGS. 7I-7N. The figures are identical to FIGS. 7A-7F, respectively, with the exception that they do not contain plant telomeres. These vectors will remain circular once delivered into the plant cell and therefore do not require telomeres to stabilize their ends.

**[0070]** FIG. 8. Sequence features at *Arabidopsis* CEN2 (A) and CEN4 (B). Central bars depict annotated genomic sequence of indicated BAC clones; black, genetically-defined centromeres; white, regions flanking the centromeres. Sequences corresponding to genes and repetitive features, filled boxes (above and below the bars, respectively), are defined as in FIG. 11A-T; predicted nonmobile genes, red; genes carried by mobile elements, black; nonmobile pseudogenes, pink; pseudogenes carried by mobile elements, gray; retroelements, yellow; transposons, green; previously defined centromeric repeats, dark blue; 180 bp repeats, pale blue. Chromosome-specific centromere features include a large mitochondrial DNA insertion (orange; CEN2), and a novel array of tandem repeats (purple; CEN4). Gaps in the physical maps (//), unannotated regions (hatched boxes), and expressed genes (filled circles) are shown.

**[0071]** FIG. 9. Method for converting a BAC clone (or any other bacterial clone) into a mini chromosome. A portion of

the conversion vector will integrate into the BAC clone (or other bacterial clone of interest) either through non-homologous recombination (transposable element mediated) or by the action of a site specific recombinase system, such as Cre-Lox or FLP-FRT.

**[0072]** FIG. 10A-G. Method for converting a BAC clone (or any other bacterial clone) into a mini chromosome. The necessary selectable markers and origins of replication for propagation of genetic material in *E. coli*, *Agrobacterium* and *Arabidopsis* as well as the necessary genetic loci for *Agrobacterium* mediated transformation into *Arabidopsis* are cloned into a conversion vector. Using Cre/loxP recombination, the conversion vectors are recombined into BACs containing centromere fragments to form mini chromosomes.

**[0073]** FIG. 11A-T. Properties of centromeric regions on chromosomes II and IV of *Arabidopsis*. (Top) Drawing of genetically-defined centromeres (gray shading, CEN2, left; CEN4, right), adjacent pericentromeric DNA, and a distal segment of each chromosome, scaled in Mb as determined by DNA sequencing (gaps in the grey shading correspond to gaps in the physical maps). Positions in cM on the R1 map ([http://nasc.nott.ac.uk/new\\_ri\\_map.html](http://nasc.nott.ac.uk/new_ri_map.html)) and physical distances in Mb, beginning at the northern telomere and at the centromeric gap, are shown. (Bottom) The density of each feature (FIGS. 11A-11T) is plotted relative to the position on the chromosome in Mb. (FIG. 11A, 11K) cM positions for markers on the R1 map (solid squares) and a curve representing the genomic average of 1 cM/221 kb (dashed line). A single crossover within CEN4 in the R1 mapping population ([http://nasc.nott.ac.uk/new\\_ri\\_map.html](http://nasc.nott.ac.uk/new_ri_map.html); Somerville and Somerville, 1999) may reflect a difference between male meiotic recombination monitored here and recombination in female meiosis. (FIGS. 11B-11E and FIGS. 11L-11O) The % of DNA occupied by repetitive elements was calculated for a 100 kb window with a sliding interval of 10 kb. (FIGS. 11B, 11L) 180 bp repeats; (FIGS. 12C, 12M) sequences with similarity to retroelements, including del, Ta1, Ta11, copia, Athila, LINE, Ty3, TSCL, 106B (Athila-like), Tat1, LTRs and Cinfu; (FIGS. 11D, 11N) sequences with similarity to transposons, including Tag1, En/Spm, Ac/Ds, Tam1 MuDR, Limpet, MITES and Mariner; (FIGS. 10E, 11O) previously described centromeric repeats including 163A, 164A, 164B, 278A, 11B7RE, mi167, pAT27, 160-, 180- and 500-bp repeats, and telomeric sequences. (FIGS. 11F, 11P) % adenosine+thymidine was calculated for a 50 kb window with a sliding interval of 25 kb (FIGS. 11G-11J, 11Q-11T). The number of predicted genes or pseudogenes was plotted over a window of 100 kb with a sliding interval of 10 kb. (FIGS. 11G, 11I, 11Q, 11S) predicted genes (FIGS. 11G, 11Q) and pseudogenes (FIGS. 11I, 11S) typically not found on mobile DNA elements; (FIGS. 11H, 11J, 11R, 11T) predicted genes (FIGS. 11H, 11R) and pseudogenes (FIGS. 11J, 11T) often carried on mobile DNA, including reverse transcriptase, transposase, and retroviral polyproteins. Dashed lines indicate regions in which sequencing or annotation is in progress, annotation was obtained from GenBank records (<http://www.ncbi.nlm.nih.gov/Entrez/nucleotide.html>), from the AGAD database (<http://www.tigr.org/tdb/at/agad/>), and by BLAST comparisons to the database of repetitive *Arabidopsis* sequences (<http://nucleus.cshl.org/protarab/AtRepBase.htm>); though updates to annotation records may change individual entries, the overall structure of the region will not be significantly altered.

**[0074]** FIG. 12. Methods for converting a BAC clone containing centromere DNA into a mini chromosome for introduction into plant cells. The specific elements described are provided for exemplary purposes and are not limiting. A) diagram of the BAC clone, noting the position of the centromere DNA, a site-specific recombination site (for example, lox P), and the F origin of replication. B) Conversion vector containing selectable and color markers (for example, 35S-Bar, nptII, LAT52-GUS, Scarecrow-GFP), telomeres, a site-specific recombination site (for example, lox P), antibiotic resistance markers (for example, amp or spc/str), *Agrobacterium* T-DNA borders (Agro Left and Right) and origin of replication (RiA4). C) The product of site specific recombination with the Cre recombinase at the lox P sites yields a circular product with centromeric DNA and markers flanked by telomeres. D) Mini chromosome immediately after transformation into plants; subsequently, the left and right borders will likely be removed by the plant cell and additional telomeric sequence added by the plant telomerase.

**[0075]** FIG. 13A-B. Conservation of *Arabidopsis* centromere DNA. BAC clones (bars) used to sequence CEN2 (FIG. 13A) and CEN4 (FIG. 13B) are indicated; arrows denote the boundaries of the genetically-defined centromeres. PCR primer pairs yielding products from only Columbia (filled circles) or from both Landsberg and Columbia (open circles); BACs encoding DNA with homology to the mitochondrial genome (gray bars); 180 bp repeats (gray boxes); unsequenced DNA (dashed lines); and gaps in the physical map (double slashes) are shown.

**[0076]** FIG. 14A-B. Primers used to analyze conservation of centromere sequences in the *A. thaliana* Columbia and Landsberg ecotypes. FIG. 14A: Primers used for amplification of chromosome 2 sequences. FIG. 14B: Primers used for amplification of chromosome 4 sequences.

**[0077]** FIG. 15. Sequences common to *Arabidopsis* CEN2 and CEN4. Genetically-defined centromeres (bold lines), sequenced (thin lines), and unannotated (dashed lines) BAC clones are displayed as in FIG. 14A, B. Repeats AtCCS1 (*A. thaliana* centromere conserved sequence) and AtCCS2 (closed and open circles, respectively), AtCCS3 (triangles), and AtCCS4-7 (4-7, respectively) are indicated (GenBank Accession numbers AF204874 to AF204880), and were identified using BLAST 2.0 (<http://blast.wustl.edu>).

#### DETAILED DESCRIPTION OF THE INVENTION

**[0078]** The present invention provides numerous methods for doing business or providing a service. These methods include increasing revenues generated by crops, developing new crops, and improving existing crops. As noted above, although it has been known to add a gene to improve the agricultural traits of a plant, this technology has been severely limited. One, or at most a few genes, are typically inserted into random locations in the host chromosome. This technology thereby can irreversibly disrupt host gene functions, can cause variable and uncontrolled expression of the introduced genes, and is increasingly time consuming and expensive as the number of genes introduced increases.

**[0079]** The inventors have identified the centromeres of a number of crops allowing them to overcome technical limitations associated with gene delivery in crops. This technology allows the inventors to develop methods for improving crops and developing new crops. Through these methods, the inventors are able to increase revenues associated with a crop, generate additional revenues from crops, provide a multitude

of services associated with crops as well as produce new products from crops. This technology also affords the ability to perform a number of services for third parties (entities that request a service from another party) based on crops.

**[0080]** For example, in an embodiment, the present invention allows the inventors to provide a service to a third party. The service includes identifying a sequence associated with the centromere of a commercial plant. A gene or set of genes to be added to the commercial plant is then chosen. The gene or set of genes and the sequence is then utilized to create a mini chromosome. This mini chromosome is then introduced into a cell of the commercial plant to create a transgenic plant having desirable characteristics not present in the commercial plant. A variety of different genes can be utilized.

**[0081]** This method can be used to increase the revenues generated by commercial plants. In this regard, the desirable characteristic can be chosen so that the resultant plant is more valuable. As used herein, the term "more valuable" can have a variety of meanings. In its most basic form, it means more revenue is generated by the crop. However, the term can also mean that the costs associated with the crop are reduced. In this regard, the gene or set of genes can add characteristics that make the commercial plant more hearty (including but not limited to characteristics such as standility, draught tolerance, improved stature, cold tolerance, and pest resistance). On the other hand, the gene or set of genes can cause the plant to mature quicker. A variety of characteristics can be added that can increase the value of the commercial plant.

**[0082]** In an embodiment, a method of generating revenues based on agricultural products is provided. The method includes the step of developing a new crop based on a mini chromosome including a sequence of a centromere and receiving a fee for the new crop. The term "crop" is defined herein in the definitions section. As used herein, the term "new" as modifying crop means a crop, that includes a gene or set of genes or expresses a trait, that is in nature not typically part of the genetic sequence of or expressed by the crop. Additionally, the term "new" as modifying crop can also mean a crop that includes a gene or set of genes naturally found in the crop but modified so that their copy number is changed. Additionally, the term "new" as modifying crop can also mean a crop that is modified to increase or decrease a trait that in nature is typically found at undesirable levels. Thus, developing a "new soybean" means that a soybean is provided that includes a trait or a gene in its sequence that is not typically found in soybeans in nature or a soybean that has been modified to increase or decrease an existing trait or gene product.

**[0083]** A variety of mechanisms can be utilized as a method of receiving a fee for the new crop. Such methods can include a fee that is received as a royalty payment. The royalty payment may take a variety of forms. The royalty can be charged either as a one-time payment, fixed fee paid on a regular basis, or a portion of the sales, or any combination thereof. For example, the technology for creating a new crop or the new crop itself may be licensed to a third party. The third party may sell seeds to a farmer or other commercial entity for a set price and the fee is set as a portion of the sales price. For example, a fee of 1% of the sales of the seeds can be charged as the fee. If desired, the fee can be based on sales of a product made from the new crop, e.g., foods, pharmaceuticals, or chemicals. If desired, the fee can comprise, in whole or in part, a cross-license of technology from a third party. A vari-

ety of mechanisms can be used to meet the step receiving a fee. Essentially, all that is required is that a valve is received.

**[0084]** The present invention affords a variety of different business methods. In part, these methods are due to the ability to introduce a number and variety of different genes into a crop. These genes can be introduced into the plants in a single step. Because the genes can be introduced independent of the host genome, this reduces the damaging effects that occur in transformation technologies. This independence allows the additional advantage that the chromosome can be eliminated or modified when desired. Further, because the DNA sequences surrounding the genes are defined, copy number and expression levels of the genes can be controlled precisely. This technology allows one to shorten both the cost and the time required for crop development.

**[0085]** Pursuant to the present invention, strategic relationships can be forged with third parties, e.g., agricultural companies that can benefit from the use of the technology in crops. These relationships can be used to initiate and drive a research and development program using crop centromeres to introduce engineered chromosomes into crops. Such programs can be based on the ability to use existing genes, or genes to be developed, to produce a unique variety of seeds. It should be noted that these programs can be used to not only generate improved crops, but also to produce pharmaceuticals, chemicals, and compounds based, at least in part, on crops.

**[0086]** In addition to the methods set forth above, it is also envisioned that methods of generating revenue can include, without limitation: selling centromeres, possibly for specific fields of use or exclusive use; developing chromosomes per customer specifications; designing plants carrying custom chromosomes to enhance crop values; and producing high value products synthesized by genes carried on engineered chromosomes.

**[0087]** The methods of the present invention are premised, in part, on the fact that the inventors have overcome the deficiencies in the prior art by providing the nucleic acid sequences of plant centromeres. The significance of this achievement relative to the prior art is exemplified by the general lack of detailed information in the art regarding the centromeres of multicellular organisms in general. To date, the most extensive and reliable characterization of centromere sequences has come from studies of lower eukaryotes such as *S. cerevisiae* and *S. pombe*, where the ability to analyze centromere functions has provided a clear picture of the desired DNA sequences. The *S. cerevisiae* centromere consists of three essential regions, CDEI, CDEII, and CDEIII, totaling only 125 bp, or approximately 0.006 to 0.06% of each yeast chromosome (Carbon et al., 1990; Bloom 1993). *S. pombe* centromeres are between 40 and 100 kB in length and consist of repetitive elements that comprise 1 to 3% of each chromosome (Baum et al., 1994). Subsequent studies, using tetrad analysis to follow the segregation of artificial chromosomes, demonstrated that less than 1/3 of the naturally occurring *S. pombe* centromere is sufficient for centromere function (Baum et al., 1994).

**[0088]** In contrast, the centromeres of mammals and other higher eukaryotes are less understood. Although DNA fragments that hybridize to centromeric regions in higher eukaryotes have been identified, in many cases, little is known regarding the functionality of these sequences (see Tyler-Smith et al., 1993). Centromere repeats often correlate with centromere location, with probes to the repeats mapping both cytologically and genetically to centromere regions. Many of



these sequences are tandemly-repeated satellite elements and dispersed repeated sequences in arrays ranging from 300 kB to 5000 kB in length (Willard 1990). To date, only one of these repeats, a 171 bp element known as the alphoid satellite, has been shown by in situ hybridization to be present at each human centromere (Tyler-Smith et al., 1993). Whether repeats themselves represent functional centromeres remains controversial, as other genomic DNA can be required to confer efficient inheritance upon a region of DNA (Willard, 1997). Alternatively, the positions of some higher eukaryotic centromeres have been estimated by analyzing the segregation of chromosome fragments. This approach is imprecise, however, because a limited set of fragments can be obtained, and because normal centromere function is influenced by surrounding chromosomal sequences (for example, see Koornneef, 1983; FIG. 2).

**[0089]** A more precise method for mapping centromeres that can be used in intact chromosomes is tetrad analysis (Mortimer et al., 1981), which provides a functional definition of a centromere in its native chromosomal context. Centromeres that have been mapped in this manner include those from the yeasts *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Kluyveromyces lactis* (Carbon et al., 1990; Hegemann et al., 1993). In many of these systems, accurate mapping of the centromeres made it possible to clone centromeric DNA, using a chromosome walking strategy (Clarke et al., 1980). Subsequently, artificial chromosome assays were used to define more precisely the centromere sequences (Hegemann et al., 1993; Baum et al., 1994).

**[0090]** Attempts to develop a reliable centromeric assay in mammals have yielded ambiguous results. For example, Hadlaczkzy et al., (1991) identified a 14 kB human fragment that can, at low frequency, result in de novo centromere formation in a mouse cell line. In situ hybridization studies, however, have shown that this fragment is absent from naturally occurring centromeres, calling into question the reliability of this approach for testing centromere function (Tyler-Smith et al., 1993). Similarly, transfection of alphoid satellites into cell lines results in the formation of new chromosomes, yet some of these chromosomes also required host sequences that could contribute centromere activity (Haaf et al., 1992; Willard, 1997). Further, the novel chromosomes can have alphoid DNA spread throughout their length yet have only a single centromeric constriction, indicating that a block of alphoid DNA alone may be insufficient for centromere function (Tyler-Smith et al., 1993).

**[0091]** Although plant centromeres can be visualized easily in condensed chromosomes, they have not been characterized as extensively as centromeres from yeast or mammals. Genetic characterization has relied on segregation analysis of chromosome fragments, and in particular on analysis of trisomic strains that carry a genetically marked, telocentric fragment (for example, see Koornneef 1983). In addition, repetitive elements have been identified that are either genetically (Richards et al., 1991) or physically (Alfenito et al., 1993; Maluszynska et al., 1991) linked to a centromere. In no case, however, has the functional significance of these sequences been tested.

**[0092]** Cytology in *Arabidopsis thaliana* has served to correlate centromere structure with repeat sequences. A fluorescent dye, DAPI, allows visualization of centromeric chromatin domains in metaphase chromosomes. A fluorescence in situ hybridization (FISH) probe based on 180 bp pAL1 repeat sequences colocalized with the DAPI signature near the cen-

tromeres of all five *Arabidopsis* chromosomes (Maluszynska et al., 1991; Martinez-Zapater et al., 1986). Although a functional role for pAL1 has been proposed, more recent studies have failed to detect this sequence near the centromeres in species closely related to *Arabidopsis thaliana* (Maluszynska et al., 1993). These results are particularly troubling because one of the species tested, *A. pumila*, is thought to be an amphidiploid, derived from a cross between *A. thaliana* and another close relative (Maluszynska et al., 1991; Price et al., 1995). Another repetitive sequence, pAtT12, has been genetically mapped to within 5 cM of the centromere on chromosome 1 and to the central region of chromosome 5 (Richards et al., 1991), although its presence on other chromosomes has not been established. Like pAL1, a role for pAtT12 in centromere function remains to be demonstrated.

**[0093]** Due to the fact that kinetochores constitute a necessary link between centromeric DNA and the spindle apparatus, the proteins that are associated with these structures recently have been the focus of intense investigation (Bloom 1993; Earnshaw 1991). Human autoantibodies that bind specifically in the vicinity of the centromere have facilitated the cloning of centromere-associated proteins (CENPs, Rattner 1991), and at least one of these proteins belongs to the kinesin superfamily of microtubule-based motors (Yen 1991). Yeast centromere-binding proteins also have been identified, both through genetic and biochemical studies (Bloom 1993; Lechner et al., 1991).

**[0094]** The centromeres of *Arabidopsis thaliana* have been mapped using trisomic strains, where the segregation of chromosome fragments (Koornneef 1983) or whole chromosomes (Sears et al., 1970) was used to localize four of the centromeres to within 5, 12, 17 and 38 cM, respectively. These positions have not been refined by more recent studies because the method is limited the difficulty of obtaining viable trisomic strains (Koornneef 1983). These factors introduce significant error into the calculated position of the centromere, and in *Arabidopsis*, where 1 cM corresponds roughly to 200 kB (Koornneef 1987; Hwang et al., 1991), this method did not map any of the centromeres with sufficient precision to make chromosome walking strategies practical. Mapping of the *Arabidopsis* genome was also discussed by (Hauge et al., 1991).

## I. ISOLATION OF CENTROMERE CLONES

**[0095]** The present invention relates to methods of isolating and identifying centromere DNA sequences from total genomic DNA of an organism without genetic mapping of the organism. Centromere DNA can be purified from total genomic DNA using several methods which include: 1) digesting genomic DNA with restriction enzymes and separating the fragments on agarose gels, to reveal major classes of repetitive DNA; 2) digesting genomic DNA with restriction enzymes sensitive to DNA methylation and separating the fragments on agarose gels to reveal the heavily methylated fraction of the genome; and 3) collecting the rapidly annealing fraction of denatured genomic DNA. These three methods isolate centromere DNA; therefore, these methods are expected to independently isolate the same sequences, thus validating the sequences' centromere origin. It is anticipated that each of these methods can be applied to genomic DNA from any organism, including some lower organisms such as yeasts, as well as higher organisms such as plants and animals. Each of these methods is described in detail below.

### 1. Isolation of repetitive DNA



[0096] Centromere regions often contain many copies of the same DNA sequence (repetitive DNA); such repeats can range in size from a few nucleotides long to hundreds or thousands of bases. Such repetitive DNA can be identified following digestion of genomic DNA with restriction endonucleases. Digestion of non-repetitive genomic DNA with a particular restriction enzyme produces a distribution of size fragments; in contrast, digestion of repeats with a restriction enzyme that cuts within each repeat produces a fragment of a typical size. Thus, genomic DNA that has been cut with a restriction enzyme can be size fractionated by agarose gel electrophoresis to reveal repetitive DNA elements; after staining the gel to reveal the DNA, the repetitive fragment can be excised and purified using conventional techniques or commercial kits. Such repeats can be introduced into cloning vectors and characterized as described below. By using this method with a variety of restriction enzymes, different repetitive elements can be purified from genomic DNA.

## 2. Purification of Methylated DNA

[0097] This method is disclosed in detail in co-pending U.S. patent application Ser. No. 09/888,220, filed Jun. 22, 2001, the disclosure of which is incorporated herein by reference in its entirety and made a part hereof. Plant centromere DNA is often extensively modified by methylation; the presence of this methylation can be used to purify centromere fragments. Digestion of genomic DNA with a methylation-sensitive restriction endonuclease (for example *Sau3A* or *HpaII*) yields a range of fragment sizes; endonuclease sites that are methylated are protected from digestion. Heavily methylated DNA molecules, such as centromere DNA, yield large fragments after digestion and can therefore be separated from the lightly or non-methylated fraction by virtue of their size. For example agarose gel electrophoresis, acrylamide gel electrophoresis, sucrose gradient fractionation, or other size fractionation techniques can be used to separate these fragments into pools of "large" (7-12 kb) and "smaller" fragments (3-7 kb and 0-3 kb).

## 3. Isolation of Rapidly Annealing DNA.

[0098] The rapidity with which denatured single stranded DNA can reanneal with another single stranded DNA molecule of complementary sequence upon renaturation is dependant upon its abundance. Therefore when genomic DNA is denatured and allowed to renature, the repetitive fraction of the genome, including centromere DNA, will renature before the unique and low copy fractions of the genome. Thus by fragmenting purified genomic DNA, denaturing it, collecting fractions at specific time points (such as 2, 4, 6, 8, and 10 minutes) during renaturation and treating those fractions to remove unannealed DNA it is possible to purify repetitive DNA from total genomic DNA. Several methods can be used to remove unannealed from annealed DNA including treatment of the sample with an enzyme, such as *S1* nuclease, that degrades single-stranded DNA or exposure to an agent that binds single-stranded DNA such as hydroxylapatite. By varying the time at which fractions are collected during renaturation it is possible to separate DNA fragments into highly repetitive, moderately repetitive, and non-repetitive fractions.

## II. CLONING AND SEQUENCING SMALL FRAGMENTS OF CENTROMERE DNA

[0099] Repetitive or methylated DNA fragments isolated using the methods described above can be ligated (using *T4*

DNA ligase, for example) to a plasmid vector and cloned by transformation into *E. coli*. These clones can then be propagated, sequenced, used to assemble mini chromosomes, or used to identify larger centromere clones, generate molecular markers that facilitate genetic mapping of centromeres, or create probes for chromosome mapping experiments such as fluorescent in situ hybridization (FISH).

## III. IDENTIFYING CENTROMERE CLONES IN GENOMIC LIBRARIES

[0100] A genomic library can be screened for clones carrying centromere DNA by arraying the clones onto solid supports, such as membrane filters, and probing with labeled fragments of purified centromere DNA, including cloned repetitive or methylated DNA fragments described above, or alternatively, the entire set of rapidly annealing genomic DNA or highly methylated genomic DNA fragments. Probes can be used singly or in combination. Typically these probes are labeled by incorporation of radionucleotides, fluorescent nucleotides, or other chemical or enzymatic ligands that enable easy detection. The labeled probe DNA is denatured and hybridized to the arrayed library using standard molecular biology techniques. Hybridization is performed at a temperature that will discourage non-specific DNA annealing while promoting the hybridization of the labeled probe to complementary sequences. After incubation, the arrayed library is washed to remove unannealed probe, and a detection method appropriate to the label incorporated in the probe is used. For example, if the probe is radiolabeled, the labeled filter is exposed to X-ray film.

[0101] To identify centromere clones, the results of several hybridization experiments are quantitated and compared. In some cases, centromere clones may hybridize to only one probe; in other cases, the clones will hybridize to multiple probes. The hybridization intensity of each clone to each probe can be measured and stored in a database. A preferred method for this analysis is to use software that digitizes the hybridization signals, assigns each signal to its corresponding clone address, ensures that duplicate copies of the clones successfully hybridized, and enters the resulting information into a relational database (MySQL for example). Another possible method for this analysis is to examine the hybridization results visually, estimate the hybridization intensity, and tabulate the resulting information.

[0102] The results of each hybridization experiment can be classified by grouping clones that show hybridization to each probe above a threshold value. For example, a computerized relational database can be queried for clones giving hybridization signals above a certain threshold for individual probes or for multiple probes. Based on these hybridization patterns, clones can be grouped into categories, and representative members of each category can be tested in mini chromosomes.

## IV. IDENTIFYING CENTROMERE SEQUENCES OF AN ORGANISM FROM GENOMIC SEQUENCE DATASETS

[0103] It is possible to devise computational algorithms to search databases of genomic sequences and select centromere sequences by identifying those with the characteristics of centromeres. For example, by selecting the most abundant tandem repeat of a particular size will yield centromere sequences. Other sets of characteristics could also be useful.

The following is a computational algorithm designed to extract centromere sequences from genomic sequence datasets. It is important to note that this algorithm examines primary sequence data and does not rely on prior annotation of the sequence. The algorithm consists of steps 1 through 10. However, not all the steps must occur in the listed order without altering the output. Other rearrangements are easily recognizable by one skilled in the art. The following terms are used in describing the algorithm. BLAST is Basic Local Alignment Search Tool, a family of freely available algorithms for sequence database searches. BLAST aligns two sequences and yields an estimate of the probability that this alignment is significant, i.e. that it did not occur by chance. The two sequences compared by BLAST are called the 'query', usually a single sequence of interest, and the 'subject', often part of a large database of sequences that are compared to the query. The query sequence (query) can also be part of a database of sequences. The outputs of BLAST are High Scoring Pairs (HSPs) that are alignments of subject and query sequences. Nucleotide position describes the position of a given nucleotide within the sequence, relative to the first nucleotide of the sequence. BLAST score (e value) is the likelihood that a given sequence alignment is significant (the lower the value the higher the significance). The algorithm is as follows:

**[0104]** (1) provide a first dataset consisting of the genomic sequences, or a representative fraction of genomic sequence, of the organism of interest;

**[0105]** (2) identify and eliminate known non-centromeric repeat sequences from the first dataset by using the BLAST sequence comparison algorithm to create a second dataset;

**[0106]** (3) compare each sequence in the second dataset to itself by using the BLAST sequence comparison algorithm, obtain a BLAST score for each pair of sequence compared, and collect high score pairs to create a third dataset;

**[0107]** (4) examine the BLAST score of each high score pair in the third dataset and eliminate the pairs having a score greater than  $10^{-20}$  to create a fourth dataset;

**[0108]** (5) eliminate the high score pairs in the fourth dataset having less than 80 bp or more than 250 bp to create a fifth dataset;

**[0109]** (6) examine the nucleotide position of each high score pair in the fifth dataset and eliminate pairs having 100% identity and identical nucleotide positions (i.e. self matches) to create a sixth dataset;

**[0110]** (7) examine the nucleotide position of each high score pair in the sixth dataset and eliminate pairs having opposite orientation of the nucleotides to create a seventh dataset;

**[0111]** (8) examine the nucleotide position of both sequences for each high score pair in the seventh dataset and eliminate sequences that are overlapping to create an eighth dataset; and

**[0112]** (9) examine the nucleotide position of each sequence in the eighth dataset and eliminate sequences not having at least one neighboring sequence within 250 bp to create a ninth dataset; and

**[0113]** (10) compare each sequence in the ninth dataset to all other sequences in the ninth dataset by using the BLAST sequence comparison algorithm and select the most common sequence as a centromere sequence of the organism.

**[0114]** Optimally, the dataset used in step (1) in the above algorithm would be the whole genome dataset such as the *Arabidopsis* genome which was derived by methodical

sequencing of mapped clones or the rice genome dataset which was derived by shotgun sequencing. Alternatively, the algorithm would also work well on representative genome datasets. By the term "representative genome datasets", it is meant that the genomic sequences in the dataset is a subset of the sequences of the whole genome collected from the whole genome without bias, such as bias toward coding sequences. These sequences would be representative of the genome as a whole. For example, the use of a 0.5× or even a 0.1× library of *Arabidopsis* with representative genome datasets would return a true positive result. On the contrary, the use of a subset of genomic sequences of the whole genome which are not representative of the whole genome and biased toward certain sequences, such as the coding sequence, would return false positive results.

## V. CENTROMERE COMPOSITIONS

**[0115]** The present invention concerns nucleic acid segments, isolatable from various plant cells, that are enriched relative to total genomic DNA, or isolated from other sources or chemically synthesized with a novel sequence, or other nucleic acids that are capable of conferring centromere activity to a recombinant molecule when incorporated into the host cell. As used herein, the term "nucleic acid segment" refers to a nucleic acid molecule that has been purified from total genomic nucleic acids of a particular species. Therefore, a nucleic acid segment conferring centromere function refers to a nucleic acid segment that contains centromere sequences yet is isolated away from, or purified free from, total genomic nucleic acids. Included within the term "nucleic acid segment", are nucleic acid segments and smaller fragments of such segments, and also recombinant vectors, including, for example, mini chromosomes, artificial chromosomes, BACs, YACs, plasmids, cosmids, phage, viruses, and the like.

**[0116]** Similarly, a nucleic acid segment comprising an isolated or purified centromeric sequence refers to a nucleic acid segment including centromere sequences and, in certain aspects, regulatory sequences, isolated substantially away from other naturally occurring sequences, or other nucleic acid sequences. In this respect, the term "gene" is used for simplicity to refer to a protein, polypeptide or peptide-encoding unit. As will be understood by those in the art, this functional term includes both genomic sequences, cDNA sequences and smaller engineered gene segments that may express, or may be adapted to express, proteins, polypeptides or peptides.

**[0117]** "Isolated substantially away from other sequences" means that the sequences of interest, in this case centromere sequences, are included within the genomic nucleic acid clones provided herein. Of course, this refers to the nucleic acid segment as originally isolated, and does not exclude all genes or coding regions.

**[0118]** In particular embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a centromere functional sequence that includes a contiguous sequence from the centromeres of the current invention. Again, nucleic acid segments that exhibit centromere function activity will be most preferred.

**[0119]** In still yet another aspect, the invention provides a plant centromere which is further defined as an *Arabidopsis thaliana* centromere. In yet another embodiment of the invention, the plant centromere comprises an *Arabidopsis thaliana* chromosome 2 centromere. The chromosome 2 centromere

may comprise, for example, from about 100 to about 611,000, about 500 to about 611,000, about 1,000 to about 611,000, about 10,000 to about 611,000, about 20,000 to about 611,000, about 40,000 to about 611,000, about 80,000 to about 611,000, about 150,000 to about 611,000, or about 300,000 to about 611,000 contiguous nucleotides of a first nucleic acid sequence flanking a first series of 180 bp repeats in centromere 2 of *A. thaliana*. The centromere may also be defined as comprising from about 100 to about 50,959, about 500 to about 50,959, about 1,000 to about 50,959, about 5,000 to about 50,959, about 10,000 to about 50,959, 20,000 to about 50,959, about 30,000 to about 50,959, or about 40,000 to about 50,959 contiguous nucleotides of a second nucleic acid sequence flanking a second series of 180 bp repeats in centromere 2 of *A. thaliana*. The centromere may comprise sequences from both of the third and the fourth sequences, including the aforementioned fragments, or the entirety of these sequences. In particular embodiments, the inventors contemplate a 3' fragment of the first sequence can be fused to a 5' fragment of the second sequence, optionally including one or more 180 bp repeat sequence disposed therebetween.

**[0120]** In still yet another aspect, the invention provides an *Arabidopsis thaliana* chromosome 4 centromere. In certain embodiments of the invention, the centromere may comprise from about 100 to about 1,082,000, about 500 to about 1,082,000, about 1,000 to about 1,082,000, about 5,000 to about 1,082,000, about 10,000 to about 1,082,000, about 50,000 to about 1,082,000, about 100,000 to about 1,082,000, about 200,000 to about 1,082,000, about 400,000 to about 1,082,000, or about 800,000 to about 1,082,000 contiguous nucleotides of a third nucleic acid sequence flanking a third series of repeated sequences, including comprising the nucleic acid sequence of the third sequence. The centromere may also be defined as comprising from about 100 to about 163,317, about 500 to about 163,317, about 1,000 to about 163,317, about 5,000 to about 163,317, about 10,000 to about 163,317, about 30,000 to about 163,317, about 50,000 to about 163,317, about 80,000 to about 163,317, or about 120,000 to about 163,317 contiguous nucleotides of the nucleic acid sequence of a fourth sequence flanking a fourth series of repeated sequences, and may be defined as comprising the nucleic acid sequence of the fourth sequence. The centromere may comprise sequences from both the third and the fourth sequences, including the aforementioned fragments, or the entirety of the third and the fourth sequences. In particular embodiments, the inventors contemplate a 3' fragment of the third sequence can be fused to a 5' fragment of the fourth sequence, optionally including one or more 180 bp repeat sequence disposed therebetween.

**[0121]** In yet another embodiment, there is provided a *Arabidopsis thaliana* chromosome 1, 3 or 5 centromere selected from the nucleic acid sequence given by one of the repeated sequences in these chromosomes, or fragments thereof. The length of the repeat used may vary, but will preferably range from about 20 bp to about 250 bp, from about 50 bp to about 225 bp, from about 75 bp to about 210 bp, from about 100 bp to about 205 bp, from about 125 bp to about 200 bp, from about 150 bp to about 195 bp, from about 160 bp to about 190 bp and from about 170 bp to about 185 bp including about 180 bp. In one embodiment, the construct comprises at least 100 base pairs, up to an including the full length, of one of the preceding sequences. In addition, the construct may include 1 or more 180 base pair repeats.

**[0122]** In one embodiment, the centromere n copies of a repeated nucleotide sequence obtained by the method disclosed herein, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. Moreover, the copies, while largely identical, can vary from each other. Such repeat variation is commonly observed in naturally occurring centromeres.

**[0123]** In another embodiment, the centromere is a *Brassica oleracea* centromere comprising *Brassica oleracea* centromere DNA. In one embodiment, the *Brassica oleracea* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Brassica oleracea* given by SEQ ID NO:1, 2, 3, or 4.

**[0124]** In yet another embodiment, the centromere is a *Glycine max* centromere comprising *glycine max* centromere DNA. In an embodiment, the *Glycine max* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Glycine max* given by SEQ ID NO:5, 6, 7, or 8.

**[0125]** In yet another embodiment, the centromere is a *Lycopersicon esculentum* centromere comprising *Lycopersicon esculentum* centromere DNA. In an embodiment, the *Lycopersicon esculentum* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Lycopersicon esculentum* given by SEQ ID NO:9 or 10.

**[0126]** In yet another embodiment, the centromere is a *Zea mays* centromere comprising *Zea mays* centromere DNA. In an embodiment, the centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000,

60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Zea mays* given by SEQ ID NO:11, 12 or 13.

[0127] The centromere can additionally be defined as the region of the chromosome where the sister chromatids pair during cell division. The centromere is also the chromosomal region where the kinetochore (the chromosomal attachment structure for the spindle) and the spindle (the cellular machinery that provides the motive force for chromosome segregation) attach to the chromosome during mitosis and meiosis. The centromere is also defined as the region of the primary constriction in a condensed chromosome. The DNA of the centromere is characteristically heavily methylated, repetitive, and condensed (heterochromatic).

## VI. MINI CHROMOSOME CONSTRUCTION

[0128] Mini chromosomes are constructed by combining fragments of centromere DNA with other DNA sequences useful for propagation of the resultant recombinant DNA molecule in *E. coli*, other bacteria, yeast or plants. Recombinant plasmids containing large fragments of centromere DNA are referred to as centromere clones. Centromere sequences removed from centromere clones, or centromere sequences derived directly from genomic DNA, are referred to as centromere fragments. Recombinant constructs containing DNA sequences necessary for the propagation, delivery, selection, and detection of mini chromosomes will be referred to as mini chromosome vector sequences or mini chromosome vectors; these sequences can include but are not limited to selectable marker genes, visible marker genes, origins of replication, restriction endonuclease recognition sites, homing endonuclease recognition sites, sequences recognized by site specific recombinase enzymes, telomere sequences, and sequences required for delivery of mini chromosomes into bacteria, yeast or plant cells. Recombinant constructs containing both large centromere fragments as well as mini chromosome vector sequences are referred to as mini chromosomes. The process of assembling mini chromosomes from centromere clones/fragments and mini chromosome vector sequences can be done in several ways, and involves techniques that are common practice among those trained in molecular biology:

1) Joining Centromere Fragments to Mini Chromosome Vector Sequences:

[0129] Centromere DNA fragments and mini chromosome vector DNA fragments are generated and purified using conventional techniques, some of which include restriction enzyme digestion, agarose gel electrophoresis, gel purification of specific fragments, anion-exchange purification and ethanol precipitation. The resulting purified centromere and vector fragments are enzymatically joined *in vitro*, using for example T4 DNA ligase. The ends of the fragments can be cohesive, as the result of digestion with compatible restriction endonucleases or from the addition of compatible oligonucleotide linkers; alternatively the ends of the fragments can be blunt and can be directly joined. Following ligation, the resulting mini chromosomes are introduced into *E. coli*, other bacteria, yeast, or plant cells using chemical or physical transformation methods. The structure of the resulting mini chromosomes can be determined by recovering them from the host organism and assessing DNA fragment size and composition.

2) Transfer of Mini Chromosome Vector Sequences into Centromere Clones by Site-Specific Recombination:

[0130] The mini chromosome vector sequences can be constructed to include site-specific recombination sequences (for example those recognized by the bacteriophage P1 Cre recombinase, or the bacteriophage lambda integrase, or similar recombination enzymes). A compatible recombination site, or a pair of such sites, can also be included in the centromere clones. Incubation of the mini chromosome vector and the centromere clone in the presence of the recombinase enzyme causes strand exchange to occur between the recombination sites in the two plasmids; the resulting mini chromosomes contain centromere sequences as well as mini chromosome vector sequences (FIG. 5). Introducing the DNA molecules formed in such recombination reactions into *E. coli*, other bacteria, yeast or plant cells can be followed by selection for marker genes present on both parental plasmids, allowing the isolation of mini chromosomes.

3) Mini Chromosome Vector Tailing Method for Mini Chromosome Construction:

[0131] Centromere DNA fragments isolated from genomic DNA or from centromere clones can be modified on their ends by treatment with restriction endonucleases, or by ligation with DNA molecules including, but not limited to, oligonucleotide linkers, or by the addition of nucleotides, to produce a desired cohesive or blunt end. These fragments are size-fractionated by, agarose gel electrophoresis or other methods, and the centromere fragments purified using conventional techniques. Mini chromosome vector fragments are generated and purified in a similar manner, resulting in linear mini chromosome vector sequences with DNA ends compatible with those on the centromere fragments. Compatible ends in this case are defined by ends that can be joined *in vitro* by the action of a ligase enzyme. As shown in FIG. 6, the two fragments are then mixed so that the mini chromosome vector molecules are present in at least two-fold molar excess over the centromere fragments. The fragments are joined by the addition of a ligase enzyme (for example bacteriophage T4 DNA ligase), resulting in the formation of DNA molecules in which mini chromosome vector molecules have been joined to both ends of the same centromere fragment. Digestion of the ligation mixture with a rare-cutting restriction or homing endonuclease (for example endonucleases with recognition sequences of 8 or more bases) results in linear mini chromosome precursors consisting of a fragment of the original mini chromosome vector attached to each end of the centromere fragment. The ends of this hybrid molecule are compatible because they were created by the same restriction enzyme. This linear mini chromosome precursor is purified, for example, by agarose gel electrophoresis followed by gel purification of the DNA fragments of the expected length. The purified DNA molecules are circularized by joining the ends, for example by treatment with a DNA ligase enzyme. The resulting mini chromosome molecules can be introduced into *E. coli*, other bacteria, yeast or plant cells, followed by purification and characterization using conventional methods.

## VII. USE OF MINI CHROMOSOMES FOR PLANT TRANSFORMATION

[0132] 1) Delivery of Mini Chromosomes into Plant Cells:

[0133] Mini chromosomes are purified and delivered into plant cells, either individually or as a mixture. The mini

chromosomes can be either circular or linear or mixtures thereof. The plant cells used for mini chromosome delivery can be either intact seedlings, immature or mature plants, parts of seedlings or plants, specific plant tissues (for example leaves, stems, roots, flowers, fruits), differentiated tissues cultured in vitro (for example roots), or undifferentiated cells (for example callus) cultured in vitro. The mini chromosome DNA can be delivered into plant cells by a variety of methods including but not limited to the following: electroporation; *Agrobacterium*-mediated DNA delivery; virus-mediated DNA delivery; delivery mediated by salts or lipids that facilitate the cellular uptake of DNA; microinjection of DNA, manipulation into a cell of DNA-coated or DNA-containing particles, droplets, micelles, microspheres, or chemical complexes using a variety of techniques, including biolistic particle bombardment, optical tweezers, particle beams, and electrospray apparatus; manipulation of DNA-coated magnetic particles into the cells by magnetic fields; DNA delivery into cells by cell wounding using micro-needles (for example silicon carbide needles); sonication or other acoustic treatment of the cells to facilitate DNA uptake; fusion of plant cells with other cell types carrying a mini chromosome, including bacterial, yeast, or other plant cells; any other electrical, chemical, physical, or biological mechanism that results in the introduction of mini chromosome DNA into the plant cell

## 2) Isolating Plant Cells Containing Mini Chromosomes:

**[0134]** Following mini chromosome delivery, plant cells, plant tissues, or complete plants carrying the mini chromosome can be isolated by a variety of selection methods. Selection involves subjecting the plant cells, tissues or plants to chemical, environmental, or mechanical treatments that enrich for those cells, tissue or plants that contain a mini chromosome. The selection methods include but are not limited to: fluorescence-activated cell sorting of cells, cell clumps, or cell protoplasts based on expression of a marker protein encoded by the mini chromosome. (for example, a fluorescent protein such as DsRed); affinity purification of cells, cell clumps, or protoplasts based on expression of a cell wall protein, membrane protein, or membrane-associated protein encoded by the mini chromosome; any cell fractionation method capable of separating cells based on their density, size or shape to enrich for cells with a property that differs from that of the starting population and is conferred by the mini chromosome; selection of cells for resistance to an antibiotic conferred by the mini chromosome; selection of cells for resistance to an herbicide conferred by the mini chromosome; selection of cells for resistance to a toxic metal, salt, mineral or other substance conferred by the mini chromosome; selection of cells for resistance to abiotic stress (for example heat, cold, acid, base, osmotic stress) conferred by the mini chromosome; selection of cells capable of utilizing a carbon source or other nutrient source not normally utilized by plant cells, this utilization function being conferred by the mini chromosome. As a result of the treatment, a population of plant cells can be obtained that contain mini chromosomes. Individual clones or sub-populations of these cells can be expanded in culture for further characterization.

**[0135]** Alternatively, plant cells, plant tissues, or complete plants that carry mini chromosomes can be identified by direct screening. Such methods involve subjecting each cell, plant, or tissue to diagnostic tests indicative of the presence of the mini chromosome. These tests can include direct assays

for the presence of mini chromosome DNA, or indirect assays for properties conferred by the mini chromosome. Direct assays for the presence of the mini chromosome DNA include but are not limited to: staining of cells with DNA-binding molecules to allow detection of an additional chromosome; in situ hybridization with labeled DNA probes corresponding to sequences present on the mini chromosome; southern blots or dot blots of DNA extracted from the cells, plant or tissue and probed with labeled DNA sequences corresponding to sequences present on the mini chromosome; electrophoresis of genomic DNA extracted from the cells, plant or tissue under conditions that allow identification of the mini chromosome; amplification of specific sequences present on the mini chromosome from genomic DNA extracted from the cells, plant or tissue using the polymerase chain reaction. Indirect assays for properties conferred by the mini chromosome include but are not limited to: detection of the expression of a fluorescent marker encoded by the mini chromosome by fluorescence microscopy, flow cytometry or fluorimetry; detection of the expression of a protein encoded by the mini chromosome by use of specific antibodies, or any other reagent capable of specifically binding to the protein; use of cell fractionation methods capable of detecting a specific density, size or shape of the cells or tissues, that is conferred by the mini chromosome; growth of cells, seedlings, plants or tissues on an antibiotic-containing medium to determine the presence of an antibiotic-resistance gene encoded by the mini chromosome; growth of cells, seedlings, plants or tissues on an herbicide-containing medium to determine the presence of an herbicide-resistance gene encoded by the mini chromosome; growth of cells, seedlings, plants or tissues on a medium containing a toxic metal, salt, mineral or other substance to determine the presence of a gene conferring resistance to this substance encoded by the mini chromosome; growth of cells, tissues or plants under conditions of abiotic stress (for example heat, cold, acid, base, osmotic stress) to determine the presence of a gene conferring resistance to this stress encoded by the mini chromosome; growth of cells on a medium containing a carbon source or other nutrient source normally not utilized by plant cells, to determine the presence of a utilization function conferred by the mini chromosome.

## 3) Characterization of Plant Cell Clones Containing Mini Chromosomes

**[0136]** Plant cells, tissues, or entire plants containing mini chromosomes can be further characterized to determine whether the mini chromosome is an autonomous DNA molecule, or whether it is associated with one of the plant cell's chromosomes by integration. The methods used for this analysis include, but are not limited to, the following:

1) Detection of marker protein expression by microscopy, flow cytometry, fluorimetry, enzymatic assays, cell staining or any other technique that allows the detection of a marker protein having a specific enzymatic activity, or conferring a specific color, or fluorescence property onto the cells. For example, if a cell line has been selected for containing a mini chromosome by selecting for the function of a resistance gene encoded by the mini chromosome, and if a marker protein is also encoded by the mini chromosome, then expression of this marker protein in the selected cells is an indication of the presence of the entire mini chromosome, and could indicate autonomy of this mini chromosome from the cell's other chromosomes.

2) Use of gel electrophoresis to detect a mini chromosome in genomic DNA isolated from the plant cells, tissue or entire plants. For example, genomic DNA isolated from the cells, tissues or plants can be fractionated by gel electrophoresis, either intact or following digestion with restriction endonucleases or homing endonucleases, allowing the detection of a mini chromosome or a fragment of a mini chromosome.

3) Use of southern blots or dot blots of DNA extracted from the cells, tissue or plants to detect the presence of specific sequences contained on the mini chromosome. For example, digestion of genomic DNA extracted from the cells, tissues or plants can be fractionated by agarose gel electrophoresis, blotted onto a DNA-binding membrane, and probed with labeled DNA sequences corresponding to sequences present on the mini chromosome to detect specific fragments of mini chromosome DNA, and thus allowing the determination of the autonomous, or integrated structure of the mini chromosome.

4) Cytological techniques for directly visualizing the mini chromosome in the transformed cells, such as staining of cells with DNA-binding dyes or in situ hybridization with labeled DNA probes corresponding to sequences present on the mini chromosome.

5) Genetic analysis of marker segregation by scoring marker inheritance in progeny of a plant containing a mini chromosome. For example, markers present on an autonomous mini chromosome will segregate independently from markers on the arms of the host chromosomes in a population of F2 progeny generated from a cross between a line carrying a mini chromosome and a second marked line that doesn't carry the mini chromosome. Markers include but are not limited to: visible markers conferring a visible characteristic to the plant; selectable markers, conferring resistance to an antibiotic, herbicide, or other toxic compound; enzymatic markers, conferring an enzymatic activity that can be assayed in the plant or in extracts made from the plant; protein markers, allowing the specific detection of a protein expressed in the plant; molecular markers, such as restriction fragment length polymorphisms, amplified fragment length polymorphisms, short sequence repeat (microsatellite) markers, presence of certain sequences in the DNA of the plant as detected by the polymerase chain reaction, single nucleotide polymorphisms or cleavable amplified polymorphic sites.

4) Plant Regeneration from Transformed Cell Clones:

[0137] Plant cells or tissues that harbor mini chromosomes can be used to regenerate entire plants. This will be accomplished with standard techniques of plant regeneration from differentiated tissues or undifferentiated cells. Typically, transformed tissues or callus are subjected to a series of treatments with media containing various mixtures of plant hormones and growth regulators that promote the formation of a plant embryo, specific plant tissues or organs, or a complete plant (roots and shoot) from the starting cells or tissues. Following plant regeneration, the plant can be grown either in sterile media or in soil.

#### VIII. TESTING MINI CHROMOSOME INHERITANCE IN PLANT CELLS

[0138] The inheritance of mini chromosomes can be measured through one or more cell divisions. After isolating cells, tissues, or entire plants that contain the mini chromosome, the population of cells is allowed to grow (either with or without selection), and the presence of the mini chromosome is monitored as the cells divide. Mini chromosomes can be detected

in cells by a variety of methods, including but not limited to: detection of fluorescence or any other visual characteristic arising from a marker protein gene present on the mini chromosome; resistance to an antibiotic, herbicide, toxic metal, salt, mineral or other substance, or abiotic stress as outlined above (isolating plant cells containing mini chromosomes); staining of cells with DNA-binding molecules to allow detection of an additional chromosome; in situ hybridization with labeled DNA probes corresponding to sequences present on the mini chromosome; southern blots or dot blots of DNA extracted from the cell population and probed with labeled DNA sequences corresponding to sequences present on the mini chromosome; expression of a marker enzyme encoded by a gene present on the mini chromosome (i.e. luciferase, alkaline phosphatase, beta-galactosidase, etc.) that can be assayed in the cells or in an extract made from the cells.

[0139] The percentage of cells containing the chromosome is determined at regular intervals during this growth phase. The change in the fraction of cells harboring the mini chromosome, divided by the number of cell divisions, represents the average mini chromosome loss rate. Mini chromosomes with the lowest loss rates have the highest level of inheritance.

#### IX. RECOVERY OF MINI CHROMOSOMES FROM PLANT CELLS

[0140] Recovery of mini chromosomes from plant cells can be achieved by a variety of techniques, including, but not limited to, the following:

1) Extracting the genomic DNA of transformed plant cells and introducing that DNA into *E. coli*, other bacteria or yeast and selecting for the antibiotic resistance genes present on the mini chromosome.

2) Isolation of chromosomes from cells, tissues or plants containing mini chromosomes, and sorting these by flow cytometry to allow the separation of chromosomes of different size;

3) Isolation of individual chromosomes from a cell harboring mini chromosomes by micro-manipulation involving mechanical devices such as needles made of glass, metal or other suitable substances, or other techniques such as optical tweezers, or micro-suction devices.

4) Combinations of the above, for example chromosome isolation by flow cytometry or micromanipulation followed by introduction into *E. coli*, other bacteria, yeast or plant cells.

[0141] The resulting mini chromosomes "rescued" in this fashion may differ from their parental molecules in total size, size of the centromere, presence or absence of additional sequences, and overall arrangement of the sequences. These procedures allow the isolation of DNA molecules capable of replicating and segregating in plant cells without having to test mini chromosomes individually. For example, after delivery of pools of mini chromosomes, or pools of centromere clones into plant cells, tissues or whole plants, and recovering them by the methods listed above, facilitates the selection of specific mini chromosomes or centromere clones that remain autonomous in plant cells. Whereas plant transformation with mini chromosomes relies on the sequences contributed by mini chromosome vectors, the recovery methods do not necessarily require mini chromosome vector sequences; as a

result, pools of centromere clones can be delivered into plant cells followed by recovery of the ones that replicated and persist.

#### X. EXOGENOUS GENES FOR EXPRESSION IN PLANTS

**[0142]** One particularly important advance of the present invention is that it provides methods and compositions for expression of exogenous genes in plant cells. One advance of the constructs of the current invention is that they enable the introduction of multiple genes (often referred to as gene “stacking”), potentially representing an entire biochemical pathway, or any combination of genes encoding different biochemical processes or pathways. Significantly, the current invention allows for the transformation of plant cells with a mini chromosome comprising a number of structural genes. Another advantage is that more than one mini chromosome could be introduced, allowing combinations of genes to be moved and shuffled. Moreover, the ability to eliminate a mini chromosome from a plant would provide additional flexibility, making it possible to alter the set of genes contained within a plant. Further, by using site-specific recombinases, it should be possible to add genes to an existing mini chromosome once it is in a plant.

**[0143]** Added genes often will be genes that direct the expression of a particular protein or polypeptide product, but they also may be non-expressible DNA segments, e.g., transposons such as Ds that do not direct their own transposition. As used herein, an “expressible gene” is any gene that is capable of being transcribed into RNA (e.g., mRNA, antisense RNA, etc.) or translated into a protein, expressed as a trait of interest, or the like, etc., and is not limited to selectable, screenable or non-selectable marker-genes. The inventors also contemplate that, where both an expressible gene that is not necessarily a marker gene is employed in combination with a marker gene, one may employ the separate genes on either the same or different DNA segments for transformation. In the latter case, the different vectors may be delivered concurrently to recipient cells to maximize cotransformation or may be delivered sequentially.

**[0144]** The choice of the particular DNA segments to be delivered to the recipient cells often will depend on the purpose of the transformation. One of the major purposes of transformation of crop plants is to add some commercially desirable, agronomically important traits to the plant. Such traits include, but are not limited to, herbicide resistance or tolerance; insect resistance or tolerance; disease resistance or tolerance (viral, bacterial, fungal, nematode); stress tolerance and/or resistance, as exemplified by resistance or tolerance to drought, heat, chilling, freezing, excessive moisture, salt stress; oxidative stress; increased yields; food content and makeup; physical appearance; male sterility; drydown; standability; prolificacy; starch quantity and quality; oil quantity and quality; protein quality and quantity; amino acid composition; the production of a pharmaceutically active protein; the production of a small molecule with medicinal properties; the production of a chemical including those with industrial utility; the production of nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, hormones, and the like. Additionally one could create a library of an entire genome from any organism or organelle including mammals, plants, microbes, fungi, bacteria, represented on mini chromosomes. Furthermore one could incorporate a desired genomic segment such

as one that includes a quantitative trait onto a mini chromosome. One may desire to incorporate one or more genes conferring any such desirable trait or traits, such as, for example, a gene or genes encoding herbicide resistance.

**[0145]** In certain embodiments, the present invention contemplates the transformation of a recipient cell with mini chromosomes comprising more than one exogenous gene. An “exogenous gene,” can be a gene not normally found in the host genome in an identical context, or alternatively, the mini chromosome could be used to introduce extra copies of host genes into a cell. The gene may be isolated from a different species than that of the host genome, or alternatively, isolated from the host genome but operably linked to one or more regulatory regions which differ from those found in the unaltered, native gene. Two or more exogenous genes also can be supplied in a single transformation event using either distinct transgene-encoding vectors, or using a single vector incorporating two or more gene coding sequences. For example, plasmids bearing the bar and aroA expression units in either convergent, divergent, or colinear orientation, are considered to be particularly useful. Further preferred combinations are those of an insect resistance gene, such as a Bt gene, along with a protease inhibitor gene such as pinII, or the use of bar in combination with either of the above genes. Of course, any two or more transgenes of any description, such as those conferring herbicide, insect, disease (viral, bacterial, fungal, nematode) or drought resistance, male sterility, drydown, standability, prolificacy, starch properties, oil quantity and quality, modified chemical production, pharmaceutical or nutraceutical properties, bioremediation properties, increased biomass, altered growth rate, altered fitness, altered salinity tolerance, altered thermal tolerance, altered growth form, altered composition, altered metabolism, altered biodegradability, altered CO<sub>2</sub> fixation, altered stress tolerance, presence of bioindicator activity, altered digestibility by humans or animals, altered allergenicity, altered mating characteristics, altered pollen dispersal, altered appearance, improved environmental impact, nitrogen fixation capability, or those increasing yield or nutritional quality may be employed as desired.

#### **[0146]** (i) Herbicide Resistance

**[0147]** The genes encoding phosphinothricin acetyltransferase (bar and pat), glyphosate tolerant EPSP synthase genes, the glyphosate degradative enzyme gene gox encoding glyphosate oxidoreductase, deh (encoding a dehalogenase enzyme that inactivates dalapon), herbicide resistant (e.g., sulfonyleurea and imidazolinone) acetolactate synthase, and bxn genes (encoding a nitrilase enzyme that degrades bromoxynil) are good examples of herbicide resistant genes for use in transformation. The bar and pat genes code for an enzyme, phosphinothricin acetyltransferase (PAT), which inactivates the herbicide phosphinothricin and prevents this compound from inhibiting glutamine synthetase enzymes. The enzyme 5-enolpyruvylshikimate 3-phosphate synthase (EPSP Synthase), is normally inhibited by the herbicide N-(phosphonomethyl)glycine (glyphosate). However, genes are known that encode glyphosate-resistant EPSP synthase enzymes. These genes are particularly contemplated for use in plant transformation. The deh gene encodes the enzyme dalapon dehalogenase and confers resistance to the herbicide dalapon. The bxn gene codes for a specific nitrilase enzyme that converts bromoxynil to a non-herbicidal degradation product.

[0148] (ii) Insect Resistance

[0149] Potential insect resistance genes that can be introduced include *Bacillus thuringiensis* crystal toxin genes or Bt genes (Watrud et al., 1985). Bt genes may provide resistance to lepidopteran or coleopteran pests such as European Corn Borer (ECB). Preferred Bt toxin genes for use in such embodiments include the CryIA(b) and CryIA(c) genes. Endotoxin genes from other species of *B. thuringiensis* which affect insect growth or development also may be employed in this regard.

[0150] It is contemplated that preferred Bt genes for use in the transformation protocols disclosed herein will be those in which the coding sequence has been modified to effect increased expression in plants, and more particularly, in monocot plants. Means for preparing synthetic genes are well known in the art and are disclosed in, for example, U.S. Pat. No. 5,500,365 and U.S. Pat. No. 5,689,052, each of the disclosures of which are specifically incorporated herein by reference in their entirety. Examples of such modified Bt toxin genes include a synthetic Bt CryIA(b) gene (Perlak et al., 1991), and the synthetic CryIA(c) gene termed 1800b (PCT Application WO 95/06128). Some examples of other Bt toxin genes known to those of skill in the art are given in Table 1 below.

TABLE 1

<i>Bacillus thuringiensis</i> $\delta$ -Endotoxin Genes <sup>a</sup>		
New Nomenclature	Old Nomenclature	GenBank Accession
Cry1Aa	CryIA(a)	M11250
Cry1Ab	CryIA(b)	M13898
Cry1Ac	CryIA(c)	M11068
Cry1Ad	CryIA(d)	M73250
Cry1Ae	CryIA(e)	M65252
Cry1Ba	CryIB	X06711
Cry1Bb	ET5	L32020
Cry1Bc	PEG5	Z46442
Cry1Bd	CryE1	U70726
Cry1Ca	Cry1C	X07518
Cry1Cb	Cry1C(b)	M97880
Cry1Da	Cry1D	X54160
Cry1Db	PrtB	Z22511
Cry1Ea	Cry1E	X53985
Cry1Eb	Cry1E(b)	M73253
Cry1Fa	Cry1F	M63897
Cry1Fb	PrtD	Z22512
Cry1Ga	PrtA	Z22510
Cry1Gb	CryH2	U70725
Cry1Ha	PrtC	Z22513
Cry1Hb		U35780
Cry1Ia	CryV	X62821
Cry1Ib	CryV	U07642
Cry1Ja	ET4	L32019
Cry1Jb	ET1	U31527
Cry1K		U28801
Cry2Aa	CryIIA	M31738
Cry2Ab	CryIIB	M23724
Cry2Ac	CryIIC	X57252
Cry3A	CryIIIA	M22472
Cry3Ba	CryIIIB	X17123
Cry3Bb	CryIIIB2	M89794
Cry3C	CryIIID	X59797
Cry4A	CryIVA	Y00423
Cry4B	CryIVB	X07423
Cry5Aa	CryVA(a)	L07025
Cry5Ab	CryVA(b)	L07026
Cry6A	CryVIA	L07022
Cry6B	CryVIB	L07024
Cry7Aa	CryIIIC	M64478
Cry7Ab	CryIIICb	U04367

TABLE 1-continued

<i>Bacillus thuringiensis</i> $\delta$ -Endotoxin Genes <sup>a</sup>		
New Nomenclature	Old Nomenclature	GenBank Accession
Cry8A	CryIIIE	U04364
Cry8B	CryIIIG	U04365
Cry8C	CryIIIF	U04366
Cry9A	CryIG	X58120
Cry9B	CryIX	X75019
Cry9C	CryIH	Z37527
Cry10A	CryIVC	M12662
Cry11A	CryIVD	M31737
Cry11B	Jeg80	X86902
Cry12A	CryVB	L07027
Cry13A	CryVC	L07023
Cry14A	CryVD	U13955
Cry15A	34 kDa	M76442
Cry16A	cbm71	X94146
Cry17A	cbm71	X99478
Cry18A	CryBP1	X99049
Cry19A	Jeg65	Y08920
Cyt1Aa	CytA	X03182
Cyt1Ab	CytM	X98793
Cyt2A	CytB	Z14147
Cyt2B	CytB	U52043

<sup>a</sup>Adapted from: [http://epunix.biols.susx.ac.uk/Home/Neil\\_Crickmore/Bt/index.html](http://epunix.biols.susx.ac.uk/Home/Neil_Crickmore/Bt/index.html)

[0151] Protease inhibitors also may provide insect resistance (Johnson et al., 1989), and will thus have utility in plant transformation. The use of a protease inhibitor II gene, pinII, from tomato or potato is envisioned to be particularly useful. Even more advantageous is the use of a pinII gene in combination with a Bt toxin gene, the combined effect of which has been discovered to produce synergistic insecticidal activity. Other genes which encode inhibitors of the insect's digestive system, or those that encode enzymes or co-factors that facilitate the production of inhibitors, also may be useful. This group may be exemplified by oryzacystatin and amylase inhibitors such as those from wheat and barley.

[0152] Also, genes encoding lectins may confer additional or alternative insecticide properties. Lectins (originally termed phytohemagglutinins) are multivalent carbohydrate-binding proteins which have the ability to agglutinate red blood cells from a range of species. Lectins have been identified recently as insecticidal agents with activity against weevils, ECB and rootworm (Murdock et al., 1990; Czaplá & Lang, 1990). Lectin genes contemplated to be useful include, for example, barley and wheat germ agglutinin (WGA) and rice lectins (Gatehouse et al., 1984), with WGA being preferred.

[0153] Genes controlling the production of large or small polypeptides active against insects when introduced into the insect pests, such as, e.g., lytic peptides, peptide hormones and toxins and venoms, form another aspect of the invention. For example, it is contemplated that the expression of juvenile hormone esterase, directed towards specific insect pests, also may result in insecticidal activity, or perhaps cause cessation of metamorphosis (Hammock et al., 1990).

[0154] Transgenic plants expressing genes which encode enzymes that affect the integrity of the insect cuticle form yet another aspect of the invention. Such genes include those encoding, e.g., chitinase, proteases, lipases and also genes for the production of nikkomycin, a compound that inhibits chitin synthesis, the introduction of any of which is contemplated to produce insect resistant plants. Genes that code for



activities that affect insect molting, such as those affecting the production of ecdysteroid UDP-glucosyl transferase, also fall within the scope of the useful transgenes of the present invention.

**[0155]** Genes that code for enzymes that facilitate the production of compounds that reduce the nutritional quality of the host plant to insect pests also are encompassed by the present invention. It may be possible, for instance, to confer insecticidal activity on a plant by altering its sterol composition. Sterols are obtained by insects from their diet and are used for hormone synthesis and membrane stability. Therefore alterations in plant sterol composition by expression of novel genes, e.g., those that directly promote the production of undesirable sterols or those that convert desirable sterols into undesirable forms, could have a negative effect on insect growth and/or development and hence endow the plant with insecticidal activity. Lipoxygenases are naturally occurring plant enzymes that have been shown to exhibit anti-nutritional effects on insects and to reduce the nutritional quality of their diet. Therefore, further embodiments of the invention concern transgenic plants with enhanced lipoxygenase activity which may be resistant to insect feeding.

**[0156]** *Tripsacum dactyloides* is a species of grass that is resistant to certain insects, including corn root worm. It is anticipated that genes encoding proteins that are toxic to insects or are involved in the biosynthesis of compounds toxic to insects will be isolated from *Tripsacum* and that these novel genes will be useful in conferring resistance to insects. It is known that the basis of insect resistance in *Tripsacum* is genetic, because said resistance has been transferred to *Zea mays* via sexual crosses (Branson and Guss, 1972). It is further anticipated that other cereal, monocot or dicot plant species may have genes encoding proteins that are toxic to insects which would be useful for producing insect resistant plants.

**[0157]** Further genes encoding proteins characterized as having potential insecticidal activity also may be used as transgenes in accordance herewith. Such genes include, for example, the cowpea trypsin inhibitor (CpTI; Hilder et al., 1987) which may be used as a rootworm deterrent; genes encoding avermectin (*Avermectin and Abamectin.*, Campbell, W. C., Ed., 1989; Ikeda et al., 1987) which may prove particularly useful as a corn rootworm deterrent; ribosome inactivating protein genes; and even genes that regulate plant structures. Transgenic plants including anti-insect antibody genes and genes that code for enzymes that can convert a non-toxic insecticide (pro-insecticide) applied to the outside of the plant into an insecticide inside the plant also are contemplated.

**[0158]** (iii) Environment or Stress Resistance

**[0159]** Improvement of a plants ability to tolerate various environmental stresses such as, but not limited to, drought, excess moisture, chilling, freezing, high temperature, salt, and oxidative stress, also can be effected through expression of novel genes. It is proposed that benefits may be realized in terms of increased resistance to freezing temperatures through the introduction of an "antifreeze" protein such as that of the Winter Flounder (Cutler et al., 1989) or synthetic gene derivatives thereof. Improved chilling tolerance also may be conferred through increased expression of glycerol-3-phosphate acetyltransferase in chloroplasts (Wolter et al., 1992). Resistance to oxidative stress (often exacerbated by conditions such as chilling temperatures in combination with high light intensities) can be conferred by expression of

superoxide dismutase (Gupta et al., 1993), and may be improved by glutathione reductase (Bowler et al., 1992). Such strategies may allow for tolerance to freezing in newly emerged fields as well as extending later maturity higher yielding varieties to earlier relative maturity zones.

**[0160]** It is contemplated that the expression of novel genes that favorably effect plant water content, total water potential, osmotic potential, and turgor will enhance the ability of the plant to tolerate drought. As used herein, the terms "drought resistance" and "drought tolerance" are used to refer to a plants increased resistance or tolerance to stress induced by a reduction in water availability, as compared to normal circumstances, and the ability of the plant to function and survive in lower-water environments. In this aspect of the invention it is proposed, for example, that the expression of genes encoding for the biosynthesis of osmotically-active solutes, such as polyol compounds, may impart protection against drought. Within this class are genes encoding for mannitol-L-phosphate dehydrogenase (Lee and Saier, 1982) and trehalose-6-phosphate synthase (Kaasen et al., 1992). Through the subsequent action of native phosphatases in the cell or by the introduction and coexpression of a specific phosphatase, these introduced genes will result in the accumulation of either mannitol or trehalose, respectively, both of which have been well documented as protective compounds able to mitigate the effects of stress. Mannitol accumulation in transgenic tobacco has been verified and preliminary results indicate that plants expressing high levels of this metabolite are able to tolerate an applied osmotic stress (Tarczynski et al., 1992, 1993).

**[0161]** Similarly, the efficacy of other metabolites in protecting either enzyme function (e.g., alanopine or propionic acid) or membrane integrity (e.g., alanopine) has been documented (Loomis et al., 1989), and therefore expression of genes encoding for the biosynthesis of these compounds might confer drought resistance in a manner similar to or complimentary to mannitol. Other examples of naturally occurring metabolites that are osmotically active and/or provide some direct protective effect during drought and/or desiccation include fructose, erythritol (Coxson et al., 1992), sorbitol, dulcitol (Karsten et al., 1992), glucosylglycerol (Reed et al., 1984; Erdmann et al., 1992), sucrose, stachyose (Koster and Leopold, 1988; Blackman et al., 1992), raffinose (Bernal-Lugo and Leopold, 1992), proline (Rensburg et al., 1993), glycine betaine, ononitol and pinitol (Vernon and Bohnert, 1992). Continued canopy growth and increased reproductive fitness during times of stress will be augmented by introduction and expression of genes such as those controlling the osmotically active compounds discussed above and other such compounds. Currently preferred genes which promote the synthesis of an osmotically active polyol compound are genes which encode the enzymes mannitol-1-phosphate dehydrogenase, trehalose-6-phosphate synthase and myoinositol 0-methyltransferase.

**[0162]** It is contemplated that the expression of specific proteins also may increase drought tolerance. Three classes of Late Embryogenic Proteins have been assigned based on structural similarities (see Dure et al., 1989). All three classes of LEAs have been demonstrated in maturing (i.e. desiccating) seeds. Within these 3 types of LEA proteins, the Type-II (dehydrin-type) have generally been implicated in drought and/or desiccation tolerance in vegetative plant parts (i.e. Mundy and Chua, 1988; Piatkowski et al., 1990; Yamaguchi-Shinozaki et al., 1992). Recently, expression of a Type-III

LEA (HVA-1) in tobacco was found to influence plant height, maturity and drought tolerance (Fitzpatrick, 1993). In rice, expression of the HVA-1 gene influenced tolerance to water deficit and salinity (Xu et al., 1996). Expression of structural genes from all three LEA groups may therefore confer drought tolerance. Other types of proteins induced during water stress include thiol proteases, aldolases and transmembrane transporters (Guerrero et al., 1990), which may confer various protective and/or repair-type functions during drought stress. It also is contemplated that genes that effect lipid biosynthesis and hence membrane composition might also be useful in conferring drought resistance on the plant.

**[0163]** Many of these genes for improving drought resistance have complementary modes of action. Thus, it is envisaged that combinations of these genes might have additive and/or synergistic effects in improving drought resistance in plants. Many of these genes also improve freezing tolerance (or resistance); the physical stresses incurred during freezing and drought are similar in nature and may be mitigated in similar fashion. Benefit may be conferred via constitutive expression of these genes, but the preferred means of expressing these novel genes may be through the use of a turgor-induced promoter (such as the promoters for the turgor-induced genes described in Guerrero et al., 1990 and Shagan et al., 1993 which are incorporated herein by reference). Spatial and temporal expression patterns of these genes may enable plants to better withstand stress.

**[0164]** It is proposed that expression of genes that are involved with specific morphological traits that allow for increased water extractions from drying soil would be of benefit. For example, introduction and expression of genes that alter root characteristics may enhance water uptake. It also is contemplated that expression of genes that enhance reproductive fitness during times of stress would be of significant value. For example, expression of genes that improve the synchrony of pollen shed and receptiveness of the female flower parts, i.e., silks, would be of benefit. In addition it is proposed that expression of genes that minimize kernel abortion during times of stress would increase the amount of grain to be harvested and hence be of value.

**[0165]** Given the overall role of water in determining yield, it is contemplated that enabling plants to utilize water more efficiently, through the introduction and expression of novel genes, will improve overall performance even when soil water availability is not limiting. By introducing genes that improve the ability of plants to maximize water usage across a full range of stresses relating to water availability, yield stability or consistency of yield performance may be realized.

**[0166]** (iv) Disease Resistance

**[0167]** It is proposed that increased resistance to diseases may be realized through introduction of genes into plants, for example, into monocotyledonous plants such as maize. It is possible to produce resistance to diseases caused by viruses, bacteria, fungi and nematodes. It also is contemplated that control of mycotoxin producing organisms may be realized through expression of introduced genes.

**[0168]** Resistance to viruses may be produced through expression of novel genes. For example, it has been demonstrated that expression of a viral coat protein in a transgenic plant can impart resistance to infection of the plant by that virus and perhaps other closely related viruses (Cuozzo et al., 1988, Hemenway et al., 1988, Abel et al., 1986). It is contemplated that expression of antisense genes targeted at essential viral functions may also impart resistance to viruses. For

example, an antisense gene targeted at the gene responsible for replication of viral nucleic acid may inhibit replication and lead to resistance to the virus. It is believed that interference with other viral functions through the use of antisense genes also may increase resistance to viruses. Further, it is proposed that it may be possible to achieve resistance to viruses through other approaches, including, but not limited to the use of satellite viruses.

**[0169]** It is proposed that increased resistance to diseases caused by bacteria and fungi may be realized through introduction of novel genes. It is contemplated that genes encoding so-called "peptide antibiotics," pathogenesis related (PR) proteins, toxin resistance, and proteins affecting host-pathogen interactions such as morphological characteristics will be useful. Peptide antibiotics are polypeptide sequences which are inhibitory to growth of bacteria and other microorganisms. For example, the classes of peptides referred to as cecropins and magainins inhibit growth of many species of bacteria and fungi. It is proposed that expression of PR proteins in monocotyledonous plants such as maize may be useful in conferring resistance to bacterial disease. These genes are induced following pathogen attack on a host plant and have been divided into at least five classes of proteins (Bol, Linthorst, and Cornelissen, 1990). Included amongst the PR proteins are  $\beta$ -1,3-glucanases, chitinases, and osmotin and other proteins that are believed to function in plant resistance to disease organisms. Other genes have been identified that have antifungal properties, e.g., UDA (stinging nettle lectin) and hevein (Brookaert et al., 1989; Barkai-Golan et al., 1978). It is known that certain plant diseases are caused by the production of phytotoxins. It is proposed that resistance to these diseases would be achieved through expression of a novel gene that encodes an enzyme capable of degrading or otherwise inactivating the phytotoxin. It also is contemplated that expression of novel genes that alter the interactions between the host plant and pathogen may be useful in reducing the ability of the disease organism to invade the tissues of the host plant, e.g., an increase in the waxiness of the leaf cuticle or other morphological characteristics.

**[0170]** (v) Plant Agronomic Characteristics

**[0171]** Two of the factors determining where crop plants can be grown are the average daily temperature during the growing season and the length of time between frosts. Within the areas where it is possible to grow a particular crop, there are varying limitations on the maximal time it is allowed to grow to maturity and be harvested. For example, a variety to be grown in a particular area is selected for its ability to mature and dry down to harvestable moisture content within the required period of time with maximum possible yield. Therefore, crops of varying maturities is developed for different growing locations. Apart from the need to dry down sufficiently to permit harvest, it is desirable to have maximal drying take place in the field to minimize the amount of energy required for additional drying post-harvest. Also, the more readily a product such as grain can dry down, the more time there is available for growth and kernel fill. It is considered that genes that influence maturity and/or dry down can be identified and introduced into plant lines using transformation techniques to create new varieties adapted to different growing locations or the same growing location, but having improved yield to moisture ratio at harvest. Expression of genes that are involved in regulation of plant development may be especially useful.

**[0172]** It is contemplated that genes may be introduced into plants that would improve standability and other plant growth characteristics. Expression of novel genes in plants which confer stronger stalks, improved root systems, or prevent or reduce ear droppage would be of great value to the farmer. It is proposed that introduction and expression of genes that increase the total amount of photoassimilate available by, for example, increasing light distribution and/or interception would be advantageous. In addition, the expression of genes that increase the efficiency of photosynthesis and/or the leaf canopy would further increase gains in productivity. It is contemplated that expression of a phytochrome gene in crop plants may be advantageous. Expression of such a gene may reduce apical dominance, confer semidwarfism on a plant, and increase shade tolerance (U.S. Pat. No. 5,268,526). Such approaches would allow for increased plant populations in the field.

**[0173]** (vi) Nutrient Utilization

**[0174]** The ability to utilize available nutrients may be a limiting factor in growth of crop plants. It is proposed that it would be possible to alter nutrient uptake, tolerate pH extremes, mobilization through the plant, storage pools, and availability for metabolic activities by the introduction of novel genes. These modifications would allow a plant such as maize to more efficiently utilize available nutrients. It is contemplated that an increase in the activity of, for example, an enzyme that is normally present in the plant and involved in nutrient utilization would increase the availability of a nutrient. An example of such an enzyme would be phytase. It is further contemplated that enhanced nitrogen utilization by a plant is desirable. Expression of a glutamate dehydrogenase gene in plants, e.g., *E. coli* *gdhA* genes, may lead to increased fixation of nitrogen in organic compounds. Furthermore, expression of *gdhA* in plants may lead to enhanced resistance to the herbicide glufosinate by incorporation of excess ammonia into glutamate, thereby detoxifying the ammonia. It also is contemplated that expression of a novel gene may make a nutrient source available that was previously not accessible, e.g., an enzyme that releases a component of nutrient value from a more complex molecule, perhaps a macromolecule.

**[0175]** (vii) Male Sterility

**[0176]** Male sterility is useful in the production of hybrid seed. It is proposed that male sterility may be produced through expression of novel genes. For example, it has been shown that expression of genes that encode proteins that interfere with development of the male inflorescence and/or gametophyte result in male sterility. Chimeric ribonuclease genes that express in the anthers of transgenic tobacco and oilseed rape have been demonstrated to lead to male sterility (Mariani et al., 1990).

**[0177]** A number of mutations were discovered in maize that confer cytoplasmic male sterility. One mutation in particular, referred to as T cytoplasm, also correlates with sensitivity to Southern corn leaf blight. A DNA sequence, designated TURF-13 (Levings, 1990), was identified that correlates with T cytoplasm. It is proposed that it would be possible through the introduction of TURF-13 via transformation, to separate male sterility from disease sensitivity. As it is necessary to be able to restore male fertility for breeding purposes and for grain production, it is proposed that genes encoding restoration of male fertility also may be introduced.

**[0178]** (viii) Improved Nutritional Content

**[0179]** Genes may be introduced into plants to improve the nutrient quality or content of a particular crop. Introduction of genes that alter the nutrient composition of a crop may greatly enhance the feed or food value. For example, the protein of many grains is suboptimal for feed and food purposes, especially when fed to pigs, poultry, and humans. The protein is deficient in several amino acids that are essential in the diet of these species, requiring the addition of supplements to the grain. Limiting essential amino acids may include lysine, methionine, tryptophan, threonine, valine, arginine, and histidine. Some amino acids become limiting only after corn is supplemented with other inputs for feed formulations. The levels of these essential amino acids in seeds and grain may be elevated by mechanisms which include, but are not limited to, the introduction of genes to increase the biosynthesis of the amino acids, decrease the degradation of the amino acids, increase the storage of the amino acids in proteins, or increase transport of the amino acids to the seeds or grain.

**[0180]** The protein composition of a crop may be altered to improve the balance of amino acids in a variety of ways including elevating expression of native proteins, decreasing expression of those with poor composition, changing the composition of native proteins, or introducing genes encoding entirely new proteins possessing superior composition.

**[0181]** The introduction of genes that alter the oil content of a crop plant may also be of value. Increases in oil content may result in increases in metabolizable-energy-content and density of the seeds for use in feed and food. The introduced genes may encode enzymes that remove or reduce rate-limitations or regulated steps in fatty acid or lipid biosynthesis. Such genes may include, but are not limited to, those that encode acetyl-CoA carboxylase, ACP-acyltransferase,  $\beta$ -ketoacyl-ACP synthase, plus other well known fatty acid biosynthetic activities. Other possibilities are genes that encode proteins that do not possess enzymatic activity such as acyl carrier protein. Genes may be introduced that alter the balance of fatty acids present in the oil providing a more healthful or nutritive feedstuff. The introduced DNA also may encode sequences that block expression of enzymes involved in fatty acid biosynthesis, altering the proportions of fatty acids present in crops.

**[0182]** Genes may be introduced that enhance the nutritive value of the starch component of crops, for example by increasing the degree of branching, resulting in improved utilization of the starch in livestock by delaying its metabolism. Additionally, other major constituents of a crop may be altered, including genes that affect a variety of other nutritive, processing, or other quality aspects. For example, pigmentation may be increased or decreased.

**[0183]** Feed or food crops may also possess sub-optimal quantities of vitamins, antioxidants or other nutraceuticals, requiring supplementation to provide adequate nutritive value and ideal health value. Introduction of genes that enhance vitamin biosynthesis may be envisioned including, for example, vitamins A, E, B<sub>12</sub>, choline, and the like. Mineral content may also be sub-optimal. Thus genes that affect the accumulation or availability of compounds containing phosphorus, sulfur, calcium, manganese, zinc, and iron among others would be valuable.

**[0184]** Numerous other examples of improvements of crops may be used with the invention. The improvements may not necessarily involve grain, but may, for example, improve the value of a crop for silage. Introduction of DNA to accom-

plish this might include sequences that alter lignin production such as those that result in the "brown midrib" phenotype associated with superior feed value for cattle.

**[0185]** In addition to direct improvements in feed or food value, genes also may be introduced which improve the processing of crops and improve the value of the products resulting from the processing. One use of crops is via wetmilling. Thus novel genes that increase the efficiency and reduce the cost of such processing, for example by decreasing steeping time, may also find use. Improving the value of wetmilling products may include altering the quantity or quality of starch, oil, corn gluten meal, or the components of gluten feed. Elevation of starch may be achieved through the identification and elimination of rate limiting steps in starch biosynthesis or by decreasing levels of the other components of crops resulting in proportional increases in starch.

**[0186]** Oil is another product of wetmilling, the value of which may be improved by introduction and expression of genes. Oil properties may be altered to improve its performance in the production and use of cooking oil, shortenings, lubricants or other oil-derived products or improvement of its health attributes when used in the food-related applications. Novel fatty acids also may be synthesized which upon extraction can serve as starting materials for chemical syntheses. The changes in oil properties may be achieved by altering the type, level, or lipid arrangement of the fatty acids present in the oil. This in turn may be accomplished by the addition of genes that encode enzymes that catalyze the synthesis of novel fatty acids and the lipids possessing them or by increasing levels of native fatty acids while possibly reducing levels of precursors. Alternatively, DNA sequences may be introduced which slow or block steps in fatty acid biosynthesis resulting in the increase in precursor fatty acid intermediates. Genes that might be added include desaturases, epoxidases, hydratases, dehydratases, and other enzymes that catalyze reactions involving fatty acid intermediates. Representative examples of catalytic steps that might be blocked include the desaturations from stearic to oleic acid and oleic to linolenic acid resulting in the respective accumulations of stearic and oleic acids. Another example is the blockage of elongation steps resulting in the accumulation of  $C_8$  to  $C_{12}$  saturated fatty acids.

**[0187]** (ix) Production or Assimilation of Chemicals or Biologicals

**[0188]** It may further be considered that a transgenic plant prepared in accordance with the invention may be used for the production or manufacturing of useful biological compounds that were either not produced at all, or not produced at the same level, in the corn plant previously. Alternatively, plants produced in accordance with the invention may be made to metabolize or absorb and concentrate certain compounds, such as hazardous wastes, thereby allowing bioremediation of these compounds.

**[0189]** The novel plants producing these compounds are made possible by the introduction and expression of one or potentially many genes with the constructs provided by the invention. The vast array of possibilities include but are not limited to any biological compound which is presently produced by any organism such as proteins, nucleic acids, primary and intermediary metabolites, carbohydrate polymers, enzymes for uses in bioremediation, enzymes for modifying pathways that produce secondary plant metabolites such as flavonoids or vitamins, enzymes that could produce pharmaceuticals, and for introducing enzymes that could produce

compounds of interest to the manufacturing industry such as specialty chemicals and plastics. The compounds may be produced by the plant, extracted upon harvest and/or processing, and used for any presently recognized useful purpose such as pharmaceuticals, fragrances, and industrial enzymes to name a few.

**[0190]** (x) Non-Protein-Expressing Sequences

**[0191]** DNA may be introduced into plants for the purpose of expressing RNA transcripts that function to affect plant phenotype yet are not translated into protein. Two examples are antisense RNA and RNA with ribozyme activity. Both may serve possible functions in reducing or eliminating expression of native or introduced plant genes. However, as detailed below, DNA need not be expressed to effect the phenotype of a plant.

**[0192]** 1. Antisense RNA

**[0193]** Genes may be constructed or isolated, which when transcribed, produce antisense RNA that is complementary to all or part(s) of a targeted messenger RNA(s). The antisense RNA reduces production of the polypeptide product of the messenger RNA. Genes may also be constructed to produce double-stranded RNA molecules complementary to all or part of the targeted messenger RNA(s). Genes designed in this manner will be referred to as RNAi constructs; the double-stranded RNA or RNAi constructs can trigger the sequence-specific degradation of the target messenger RNA. The polypeptide product of the target messenger RNA may be any protein. The aforementioned genes will be referred to as antisense genes and RNAi constructs, respectively. An antisense gene or RNAi construct may thus be introduced into a plant by transformation methods to produce a novel transgenic plant with reduced expression of a selected protein of interest. For example, the protein may be an enzyme that catalyzes a reaction in the plant. Reduction of the enzyme activity may reduce or eliminate products of the reaction which include any enzymatically synthesized compound in the plant such as fatty acids, amino acids, carbohydrates, nucleic acids and the like. Alternatively, the protein may be a storage protein, such as a zein, or a structural protein, the decreased expression of which may lead to changes in seed amino acid composition or plant morphological changes respectively. The possibilities cited above are provided only by way of example and do not represent the full range of applications.

**[0194]** 2. Ribozymes

**[0195]** Genes also may be constructed or isolated, which when transcribed, produce RNA enzymes (ribozymes) which can act as endoribonucleases and catalyze the cleavage of RNA molecules with selected sequences. The cleavage of selected messenger RNAs can result in the reduced production of their encoded polypeptide products. These genes may be used to prepare novel transgenic plants which possess them. The transgenic plants may possess reduced levels of polypeptides including, but not limited to, the polypeptides cited above.

**[0196]** Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, 1987; Gerlach et al., 1987; Forster and Symons, 1987). For example, a large number of ribozymes accelerate phosphoester transfer reactions with a high degree of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech et al., 1981; Michel and Westhof, 1990; Reinhold-Hurek and Shub, 1992).

This specificity has been attributed to the requirement that the substrate bind via specific base-pairing interactions to the internal guide sequence (“IGS”) of the ribozyme prior to chemical reaction.

**[0197]** Ribozyme catalysis has primarily been observed as part of sequence-specific cleavage/ligation reactions involving nucleic acids (Joyce, 1989; Cech et al., 1981). For example, U.S. Pat. No. 5,354,855 reports that certain ribozymes can act as endonucleases with a sequence specificity greater than that of known ribonucleases and approaching that of the DNA restriction enzymes.

**[0198]** Several different ribozyme motifs have been described with RNA cleavage activity (Symons, 1992). Examples include sequences from the Group I self splicing introns including Tobacco Ringspot Virus (Prody et al., 1986), Avocado Sunblotch Viroid (Palukaitis et al., 1979; Symons, 1981), and Lucerne Transient Streak Virus (Forster and Symons, 1987). Sequences from these and related viruses are referred to as hammerhead ribozyme based on a predicted folded secondary structure.

**[0199]** Other suitable ribozymes include sequences from RNase P with RNA cleavage activity (Yuan et al., 1992; Yuan and Altman, 1994, U.S. Pat. Nos. 5,168,053 and 5,624,824), hairpin ribozyme structures (Berzal-Herranz et al., 1992; Chowrira et al., 1993) and Hepatitis Delta virus based ribozymes (U.S. Pat. No. 5,625,047). The general design and optimization of ribozyme directed RNA cleavage activity has been discussed in detail (Haseloff and Gerlach, 1988; Symons, 1992; Chowrira et al., 1994; Thompson et al., 1995).

**[0200]** The other variable on ribozyme design is the selection of a cleavage site on a given target RNA. Ribozymes are targeted to a given sequence by virtue of annealing to a site by complimentary base pair interactions. Two stretches of homology are required for this targeting. These stretches of homologous sequences flank the catalytic ribozyme structure defined above. Each stretch of homologous sequence can vary in length from 7 to 15 nucleotides. The only requirement for defining the homologous sequences is that, on the target RNA, they are separated by a specific sequence which is the cleavage site. For hammerhead ribozyme, the cleavage site is a dinucleotide sequence on the target RNA is a uracil (U) followed by either an adenine, cytosine or uracil (A, C or U) (Perriman et al., 1992; Thompson et al., 1995). The frequency of this dinucleotide occurring in any given RNA is statistically 3 out of 16. Therefore, for a given target messenger RNA of 1,000 bases, 187 dinucleotide cleavage sites are statistically possible.

**[0201]** Designing and testing ribozymes for efficient cleavage of a target RNA is a process well known to those skilled in the art. Examples of scientific methods for designing and testing ribozymes are described by Chowrira et al., (1994) and Lieber and Strauss (1995), each incorporated by reference. The identification of operative and preferred sequences for use in down regulating a given gene is simply a matter of preparing and testing a given sequence, and is a routinely practiced “screening” method known to those of skill in the art.

**[0202]** 3. Induction of Gene Silencing

**[0203]** It also is possible that genes may be introduced to produce novel transgenic plants which have reduced expression of a native gene product by the mechanism of co-suppression. It has been demonstrated in tobacco, tomato, and petunia (Goring et al., 1991; Smith et al., 1990; Napoli et al., 1990; van der Krol et al., 1990) that expression of the sense

transcript of a native gene will reduce or eliminate expression of the native gene in a manner similar to that observed for antisense genes. The introduced gene may encode all or part of the targeted native protein but its translation may not be required for reduction of levels of that native protein.

**[0204]** 4. Non-RNA-Expressing Sequences

**[0205]** DNA elements including those of transposable elements such as Ds, Ac, or Mu, may be inserted into a gene to cause mutations. These DNA elements may be inserted in order to inactivate (or activate) a gene and thereby “tag” a particular trait. In this instance the transposable element does not cause instability of the tagged mutation, because the utility of the element does not depend on its ability to move in the genome. Once a desired trait is tagged, the introduced DNA sequence may be used to clone the corresponding gene, e.g., using the introduced DNA sequence as a PCR primer together with PCR gene cloning techniques (Shapiro, 1983; Dellaporta et al., 1988). Once identified, the entire gene(s) for the particular trait, including control or regulatory regions where desired, may be isolated, cloned and manipulated as desired. The utility of DNA elements introduced into an organism for purposes of gene tagging is independent of the DNA sequence and does not depend on any biological activity of the DNA sequence, i.e., transcription into RNA or translation into protein. The sole function of the DNA element is to disrupt the DNA sequence of a gene.

**[0206]** It is contemplated that unexpressed DNA sequences, including novel synthetic sequences, could be introduced into cells as proprietary “labels” of those cells and plants and seeds thereof. It would not be necessary for a label DNA element to disrupt the function of a gene endogenous to the host organism, as the sole function of this DNA would be to identify the origin of the organism. For example, one could introduce a unique DNA sequence into a plant and this DNA element would identify all cells, plants, and progeny of these cells as having arisen from that labeled source. It is proposed that inclusion of label DNAs would enable one to distinguish proprietary germplasm or germplasm derived from such, from unlabelled germplasm.

**[0207]** Another possible element which may be introduced is a matrix attachment region element (MAR), such as the chicken lysozyme A element (Stief, 1989), which can be positioned around an expressible gene of interest to effect an increase in overall expression of the gene and diminish position dependent effects upon incorporation into the plant genome (Stief et al., 1989; Phi-Van et al., 1990).

**[0208]** 5. Other

**[0209]** Other examples of non-protein expressing sequences specifically envisioned for use with the invention include tRNA sequences, for example, to alter codon usage, and rRNA variants, for example, which may confer resistance to various agents such as antibiotics.

## XI. BIOLOGICAL FUNCTIONAL EQUIVALENTS

**[0210]** Modification and changes may be made in the centromeric DNA segments of the current invention and still obtain a functional molecule with desirable characteristics. The following is a discussion based upon changing the nucleic acids of a centromere to create an equivalent, or even an improved, second-generation molecule.

**[0211]** In particular embodiments of the invention, mutated centromeric sequences are contemplated to be useful for increasing the utility of the centromere. It is specifically contemplated that the function of the centromeres of the current

invention may be based upon the secondary structure of the DNA sequences of the centromere, modification of the DNA with methyl groups or other adducts, and/or the proteins which interact with the centromere. By changing the DNA sequence of the centromere, one may alter the affinity of one or more centromere-associated protein(s) for the centromere and/or the secondary structure or modification of the centromeric sequences, thereby changing the activity of the centromere. Alternatively, changes may be made in the centromeres of the invention which do not affect the activity of the centromere. Changes in the centromeric sequences which reduce the size of the DNA segment needed to confer centromere activity are contemplated to be particularly useful in the current invention, as would changes which increased the fidelity with which the centromere was transmitted during mitosis and meiosis.

## XII. PLANTS

[0212] The term “plant,” as used herein, refers to any type of plant. The inventors have provided below an exemplary description of some plants that may be used with the invention. However, the list is not in any way limiting, as other types of plants will be known to those of skill in the art and could be used with the invention.

[0213] A common class of plants exploited in agriculture are vegetable crops, including artichokes, kohlrabi, arugula, leeks, asparagus, lettuce (e.g., head, leaf, romaine), bok choy, malanga, broccoli, melons (e.g., muskmelon, watermelon, crenshaw, honeydew, cantaloupe), brussels sprouts, cabbage, cardoni, carrots, napa, cauliflower, okra, onions, celery, parsley, chick peas, parsnips, chicory, chinese cabbage, peppers, collards, potatoes, cucumber plants (marrows, cucumbers), pumpkins, cucurbits, radishes, dry bulb onions, rutabaga, eggplant, salsify, escarole, shallots, endive, garlic, spinach, green onions, squash, greens, beet (sugar beet and fodder beet), sweet potatoes, swiss chard, horseradish, tomatoes, kale, turnips, and spices.

[0214] Other types of plants frequently finding commercial use include fruit and vine crops such as apples, apricots, cherries, nectarines, peaches, pears, plums, prunes, quince almonds, chestnuts, filberts, pecans, pistachios, walnuts, citrus, blueberries, boysenberries, cranberries, currants, loganberries, raspberries, strawberries, blackberries, grapes, avocados, bananas, kiwi, persimmons, pomegranate, pineapple, tropical fruits, pomes, melon, mango, papaya, and lychee.

[0215] Many of the most widely grown plants are field crop plants such as evening primrose, meadow foam, corn (field, sweet, popcorn), hops, jojoba, peanuts, rice, safflower, small grains (barley, oats, rye, wheat, etc.), sorghum, tobacco, kapok, leguminous plants (beans, lentils, peas, soybeans), oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts), fibre plants (cotton, flax, hemp, jute), lauraceae (cinnamon, camphor), or plants such as coffee, sugarcane, tea, and natural rubber plants.

[0216] Still other examples of plants include bedding plants such as flowers, cactus, succulents and ornamental plants, as well as trees such as forest (broad-leaved trees and evergreens, such as conifers), fruit, ornamental, and nut-bearing trees, as well as shrubs and other nursery stock.

## XIII. DEFINITIONS

[0217] As used herein, the terms “autonomous replicating sequence” or “ARS” or “origin of replication” refer to an origin of DNA replication recognized by proteins that initiate DNA replication.

[0218] As used herein, the terms “binary BAC” or “binary bacterial artificial chromosome” refer to a bacterial vector that contains the T-DNA border sequences necessary for *Agrobacterium* mediated transformation (see, for example, Hamilton et al., 1996; Hamilton, 1997; and Liu et al., 1999).

[0219] As used herein, the term “candidate centromere sequence” refers to a nucleic acid sequence which one wishes to assay for potential centromere function.

[0220] As used herein, a “centromere” is any DNA sequence that confers an ability to segregate to daughter cells through cell division. In one context, this sequence may produce a segregation efficiency to daughter cells ranging from about 1% to about 100%, including to about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or about 95% of daughter cells. Variations in such a segregation efficiency may find important applications within the scope of the invention; for example, mini chromosomes carrying centromeres that confer 100% stability could be maintained in all daughter cells without selection, while those that confer 1% stability could be temporarily introduced into a transgenic organism, but be eliminated when desired. In particular embodiments of the invention, the centromere may confer stable segregation of a nucleic acid sequence, including a recombinant construct comprising the centromere, through mitotic or meiotic divisions, including through both meiotic and mitotic divisions. A plant centromere is not necessarily derived from plants, but has the ability to promote DNA segregation in plant cells.

[0221] As used herein, the term “centromere-associated protein” refers to a protein encoded by a sequence of the centromere or a protein which is encoded by host DNA and binds with relatively high affinity to the centromere.

[0222] As used herein, the term “circular permutations” refer to variants of a sequence that begin at base n within the sequence, proceed to the end of the sequence, resume with base number one of the sequence, and proceed to base n-1. For this analysis, n may be any number less than or equal to the length of the sequence. For example, circular permutations of the sequence ABCD are: ABCD, BCDA, CDAB, and DABC.

[0223] As used herein, the term “crop” includes any plant or portion of a plant grown or harvested for commercial or beneficial purposes.

[0224] As used herein, “eukaryote” refers to living organisms whose cells contain nuclei. A eukaryote may be distinguished from a “prokaryote” which is an organism which lacks nuclei. Prokaryotes and eukaryotes differ fundamentally in the way their genetic information is organized, as well as their patterns of RNA and protein synthesis.

[0225] As used herein, the term “expression” refers to the process by which a structural gene produces an RNA molecule, typically termed messenger RNA (mRNA). The mRNA is typically, but not always, translated into polypeptide(s).

[0226] As used herein, the term “genome” refers to all of the genes and DNA sequences that comprise the genetic information within a given cell of an organism. Usually, this is taken to mean the information contained within the nucleus, but also includes the organelles.

[0227] As used herein, the term “higher eukaryote” means a multicellular eukaryote, typically characterized by its greater complex physiological mechanisms and relatively large size. Generally, complex organisms such as plants and animals are included in this category. Preferred higher eukaryotes to be transformed by the present invention include, for example, monocot and dicot angiosperm species, gymnosperm species, fern species, plant tissue culture cells of these species, animal cells and algal cells. It will of course be understood that prokaryotes and eukaryotes alike may be transformed by the methods of this invention.

[0228] As used herein, the term “host” refers to any organism that contains a plasmid, expression vector, or integrated construct comprising a plant centromere. Preferred examples of host cells for cloning, useful in the present invention, are bacteria such as *Escherichia coli*, *Bacillus subtilis*, *Pseudomonas*, *Streptomyces*, *Salmonella*, and yeast cells such as *S. cerevisiae*. Host cells which can be targeted for expression of a mini chromosome may be plant cells of any source and specifically include *Arabidopsis*, maize, rice, sugarcane, sorghum, barley, soybeans, tobacco, wheat, tomato, potato, citrus, or any other agronomically or scientifically important species.

[0229] As used herein, the term “hybridization” refers to the pairing of complementary RNA and DNA strands to produce an RNA-DNA hybrid, or alternatively, the pairing of two DNA single strands from genetically different or the same sources to produce a double stranded DNA molecule.

[0230] As used herein, the term “linker” refers to a DNA molecule, generally up to 50 or 60 nucleotides long and synthesized chemically, or cloned from other vectors. In a preferred embodiment, this fragment contains one, or preferably more than one, restriction enzyme site for a blunt-cutting enzyme and a staggered-cutting enzyme, such as BamHI. One end of the linker fragment is adapted to be ligatable to one end of the linear molecule and the other end is adapted to be ligatable to the other end of the linear molecule.

[0231] As used herein, a “library” is a pool of random DNA fragments which are cloned. In principle, any gene can be isolated by screening the library with a specific hybridization probe (see, for example, Young et al., 1977). Each library may contain the DNA of a given organism inserted as discrete restriction enzyme-generated fragments or as randomly sheered fragments into many thousands of plasmid vectors. For purposes of the present invention, *E. coli*, yeast, and *Salmonella* plasmids are particularly useful when the genome inserts come from other organisms.

[0232] As used herein, the term “lower eukaryote” refers to a eukaryote characterized by a comparatively simple physiology and composition, and most often unicellularity. Examples of lower eukaryotes include flagellates, ciliates, and yeast.

[0233] As used herein, a “mini chromosome” is a recombinant DNA construct including a centromere and capable of transmission to daughter cells. Mini chromosome may remain separate from the host genome (as episomes) or may integrate into host chromosomes. The stability of this construct through cell division could range between from about 1% to about 100%, including about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and about 95%. The mini chromosome construct may be a circular or linear molecule. It may include elements such as one or more telomeres, ARS sequences, and genes. The number of such sequences included is only limited by the physical size limitations of the

construct itself. It could contain DNA derived from a natural centromere, although it may be preferable to limit the amount of DNA to the minimal amount required to obtain a segregation efficiency in the range of 1-100%. The mini chromosome could also contain a synthetic centromere composed of tandem arrays of repeats of any sequence, either derived from a natural centromere, or of synthetic DNA. The mini chromosome could also contain DNA derived from multiple natural centromeres. The mini chromosome may be inherited through mitosis or meiosis, or through both meiosis and mitosis. As used herein, the term mini chromosome specifically encompasses and includes the terms “plant artificial chromosome” or “PLAC,” or engineered chromosomes or microchromosomes and all teachings relevant to a PLAC or plant artificial chromosome specifically apply to constructs within the meaning of the term mini chromosome.

[0234] As used herein, by “mini chromosome-encoded protein” it is meant a polypeptide which is encoded by a sequence of a mini chromosome of the current invention. This includes sequences such as selectable markers, telomeres, etc., as well as those proteins encoded by any other selected functional genes on the mini chromosome.

[0235] As used herein, the term “plant” includes plant cells, plant protoplasts, plant calli, and the like, as well as whole plants regenerated therefrom.

[0236] As used herein, the term “plasmid” or “cloning vector” refers to a closed covalently circular extrachromosomal DNA or linear DNA which is able to replicate in a host cell and which is normally nonessential to the survival of the cell. A wide variety of plasmids and other vectors are known and commonly used in the art (see, for example, Cohen et al., U.S. Pat. No. 4,468,464, which discloses examples of DNA plasmids, and which is specifically incorporated herein by reference).

[0237] As used herein, a “probe” is any biochemical reagent (usually tagged in some way for ease of identification), used to identify or isolate a gene, a gene product, a DNA segment or a protein.

[0238] As used herein, the term “recombination” refers to any genetic exchange that involves breaking and rejoining of DNA strands.

[0239] As used herein the term “regulatory sequence” refers to any DNA sequence that influences the efficiency of transcription or translation of any gene. The term includes, but is not limited to, sequences comprising promoters, enhancers and terminators.

[0240] As used herein, a “selectable marker” is a gene whose presence results in a clear phenotype, and most often a growth advantage for cells that contain the marker. This growth advantage may be present under standard conditions, altered conditions such as elevated temperature, or in the presence of certain chemicals such as herbicides or antibiotics. Use of selectable markers is described, for example, in Broach et al. (1979). Examples of selectable markers include the thymidine kinase gene, the cellular adenine-phosphoribosyltransferase gene and the dihydrofolate reductase gene, hygromycin phosphotransferase genes, the bar gene and neomycin phosphotransferase genes, among others. Preferred selectable markers in the present invention include genes whose expression confer antibiotic or herbicide resistance to the host cell, sufficient to enable the maintenance of a vector within the host cell, and which facilitate the manipulation of the plasmid into new host cells. Of particular interest in the



present invention are proteins conferring cellular resistance to ampicillin, chloramphenicol, tetracycline, G-418, bialaphos, and glyphosate for example.

[0241] As used herein, a “screenable marker” is a gene whose presence results in an identifiable phenotype. This phenotype may be observable under standard conditions, altered conditions such as elevated temperature, or in the presence of certain chemicals used to detect the phenotype.

[0242] As used herein, the term “site-specific recombination” refers to any genetic exchange that involves breaking and rejoining of DNA strands at a specific DNA sequence.

[0243] As used herein, a “structural gene” is a sequence which codes for a polypeptide or RNA and includes 5' and 3' ends. The structural gene may be from the host into which the structural gene is transformed or from another species. A structural gene will preferably, but not necessarily, include one or more regulatory sequences which modulate the expression of the structural gene, such as a promoter, terminator or enhancer. A structural gene will preferably, but not necessarily, confer some useful phenotype upon an organism comprising the structural gene, for example, herbicide resistance. In one embodiment of the invention, a structural gene may encode an RNA sequence which is not translated into a protein, for example a tRNA or rRNA gene.

[0244] As used herein, the term “telomere” refers to a sequence capable of capping the ends of a chromosome, thereby preventing degradation of the chromosome end, ensuring replication and preventing fusion to other chromosome sequences. Telomeres can include naturally occurring telomere sequences or synthetic sequences. Telomeres from one species may confer telomere activity in another species.

[0245] As used herein, the terms “transformation” or “transfection” refer to the acquisition in cells of new DNA sequences through the chromosomal or extra-chromosomal addition of DNA. This is the process by which naked DNA, DNA coated with protein, or whole mini chromosomes are introduced into a cell, resulting in a potentially heritable change.

[0246] As used herein the term “consensus” refers to a nucleic acid sequence derived by comparing two or more related sequences. A consensus sequence defines both the conserved and variable sites between the sequences being compared. Any one of the sequences used to derive the consensus or any permutation defined by the consensus may be useful in construction mini chromosomes.

[0247] As used herein the term “repeated nucleotide sequence” refers to any nucleic acid sequence of at least 25 bp present in a genome or a recombinant molecule that occurs at least two or more times and that are preferably at least 80% identical either in head to tail or head to head orientation either with or without intervening sequence between repeat units.

#### XIV. EXAMPLES

[0248] The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skilled the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventors to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without

departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

#### Example 1

##### Isolation of Genomic DNA

[0249] Tissue from various plants are harvested for DNA extraction. For DNA extraction, leaf tissue is cooled in liquid nitrogen, ground to a fine powder and transferred to an organic solvent-resistant test tube or beaker. Warm CTAB extraction solution (2% (w/v) CTAB, 100 mM Tris-Cl, pH 9.5, 20 mM EDTA, pH 8.0, 1.4 M NaCl, 1% polyethylene glycol) is added in a ratio of 20 ml per gram of tissue and mixed thoroughly. For each 20 ml extraction buffer, 50 microliters of  $\beta$ -mercaptoethanol and 30 microliters of 30 mg/ml RNase A are added and the mixture is incubated for 10-60 min. at 65° C. with occasional mixing. The homogenate is extracted with an equal volume of chloroform, and is then centrifuged 5 min at 7500 $\times$ g (8000 rpm in JA20; 10,000 rpm in a microcentrifuge, for smaller samples), 4° C. The top (aqueous) phase is recovered and nucleic acids are precipitated by adding 1 volume isopropanol. After mixing, the precipitate is pelleted at 15 min at 7500 $\times$ g, 4° C. The pellet is washed with 70% ethanol, dried and resuspended in a minimal volume of TE (10 mM Tris-Cl, pH 8.0, 0.1 mM EDTA, pH 8.0).

#### Example 2

##### *Brassica oleracea* Centromere Repeat Sequences

[0250] We purified repetitive sequences from *Brassica oleracea* (*Brassica oleracea* fast plants, obtained from the Wisconsin Crucifer Cooperative). We set forth herein two centromere repeats, termed ChrBo1 and ChrBo2. We determined the consensus of each repeat as described in Example 6.

[0251] The consensus sequence of ChrBo1 is shown in FIG. 1A (SEQ ID NO:1). This consensus was assembled from DNA sequences collected by the inventors. Twenty-four of these sequences completely spanned the repeat, and nine others partially covered the repeat. The length of this repeat is 180 $\pm$ 0.86 base pairs, and A and T comprise of 60% of the consensus.

[0252] The consensus sequence of ChrBo2 is shown in FIG. 1B (SEQ ID NO:2). This consensus was assembled from DNA sequences collected by the inventors. Five of these sequences completely spanned the repeat, and two others partially covered the repeat. The length of this repeat is 180 $\pm$ 0.45 base pairs, and A and T comprise 63% of the consensus.

[0253] The two repeats (ChrBo1 and ChrBo2) were aligned to each other using the ClustalX program (ClustalX is a free multiple sequence alignment program for Windows. Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. and Higgins, D. G. (1997) The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*, 24:4876-4882.). The two consensus sequences differ significantly from each



other at several bases. Those sites with significant differences (chi-squared,  $P < 0.05$ ) are highlighted as shown in FIG. 1C.

**[0254]** The GenBank nt database (<ftp://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on Apr. 7, 2002) and the plant satellite DNA database (<http://w3lamc.umbr.cas.cz/PlantSat/>, downloaded on Apr. 14, 2002) were compared to the inventors' consensus sequences using the blastn program and an Expect value threshold score of  $-3$ . Consensus sequences were assembled using all inventors' and GenBank sequences that matched with an Expect (E) value of less than  $-45$ .

**[0255]** The revised consensus sequence of ChrBo1 is shown in FIG. 1D (SEQ ID NO:3). This consensus was assembled from thirty-three DNA sequences collected by the inventors and eighteen GenBank sequences (Table 10). Thirty of these sequences completely spanned the repeat, and twenty-one others partially covered the repeat. The length of this repeat is  $180 \pm 0.81$  base pairs, and A and T comprise of 59% of the consensus.

TABLE 1

GenBank sequences (accession numbers) that match inventors' ChrBo1 consensus			
M30962	M30963	M31436	M31435
M31438	M31434	M31439	M31437
X68786	X12736	X07519	X16589
X15291	X68783	X68784	X61583
AJ228348	Z22947		

**[0256]** The revised consensus sequence of ChrBo2 is shown in FIG. 1E (SEQ ID NO:4). This consensus was assembled from seven DNA sequences collected by the inventors and five GenBank sequences (Table 2). Seven of these sequences completely spanned the repeat, and five others partially covered the repeat. The length of this repeat is  $180 \pm 0.44$  base pairs, and A and T comprise of 63% of the consensus.

TABLE 2

GenBank sequences (accession numbers) that match inventors' ChrBo2 consensus		
AJ228347	M30962	X12736
X61583	X68785	

**[0257]** The two revised consensus sequences (ChrBo1 and ChrBo2) were aligned to each other using the ClustalX program. The two consensus sequences differ significantly (chi-squared,  $P < 0.05$ ) from each other at several bases (highlighted as shown in FIG. 1F).

**[0258]** A total of 20 GenBank entries match the *Brassica oleracea* centromere sequences defined by the inventors. These are annotated as follows:

- [0259]** X1e7-2EB gene
- [0260]** X1e4-7B gene
- [0261]** X1e6-14H gene
- [0262]** Satellite tandem repeat monomer
- [0263]** HindIII satellite repeat
- [0264]** Satellite DNA inverted direct repeat
- [0265]** Tandem repeated DNA
- [0266]** Highly repetitive DNA

**[0267]** They are not annotated as centromere repeats in GenBank. A completed list of these sequences are shown in Table 3.

TABLE 3

GenBank entries match the <i>Brassica oleracea</i> centromere sequences defined by the inventors					
GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
X68786	<i>B. juncea</i> X1e7-2EB gene	Complete	472-651	180	97
X68786	<i>B. juncea</i> X1e7-2EB gene	Complete	763-942	180	94
X68786	<i>B. juncea</i> X1e7-2EB gene	Partial	648-761	115	96
X12736	<i>B. campestris</i> DNA for satellite tandem repeat monomer (consensus sequence)	Complete	181-2	180	97
X07519	Wild cabbage satellite DNA	Complete	179-1	179	97
X61583	<i>B. napus</i> Canrep highly repetitive DNA	Complete	2-173	176	98
X68783	<i>B. juncea</i> repetitive DNA sequence canrep subfamily A	Partial	2-173	172	97
X68784	<i>B. juncea</i> X1e4-7B gene	Complete	983-1162	180	95
X68784	<i>B. juncea</i> X1e4-7B gene	Partial	815-986	172	94
AJ228348	<i>B. carinata</i> DNA, HindIII satellite repeat (clone pBcar3)	Partial	2-173	172	96
M31438	<i>B. oleracea</i> satellite DNA inverted direct repeat	Partial	176-1	176	94
X16589	<i>B. nigra</i> tandem repeat DNA (clone BN1G 9, BN1G 23, BG1G 14)	Partial	177-1	177	94
M31434	<i>B. oleracea</i> satellite DNA inverted direct repeat	Partial	176-8	169	95
M31437	<i>B. oleracea</i> satellite DNA inverted direct repeat	Partial	175-1	175	94
M30963	<i>B. juncea</i> tandemly repeated DNA	Complete	181-2	180	93
M31435	<i>B. oleracea</i> satellite DNA inverted direct repeat	Partial	174-8	169	94
M31436	<i>B. oleracea</i> satellite DNA inverted direct repeat	Partial	175-1	177	94
X15291	<i>B. juncea</i> satellite DNA	Partial	1-161	161	95
M31439	<i>B. oleracea</i> satellite DNA inverted direct repeat	Partial	176-1	177	90
Z22947	<i>B. campestris</i> satellite DNA	Partial	181-347	170	90
Z22947	<i>B. campestris</i> satellite DNA	Partial	2-179	178	89
M30962	<i>B. campestris</i> tandemly repeated DNA	Complete	181-2	180	87
X68785	<i>B. juncea</i> X1e6-14H gene	Complete	580-758	180	92

TABLE 3-continued

GenBank entries match the <i>Brassica oleracea</i> centromere sequences defined by the inventors					
GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
X68785	<i>B. juncea</i> Xle6-14H gene	Partial	404-568	165	90
AJ228347	<i>B. carinata</i> DNA, HindIII satellite repeat (clone pBcar5)	Partial	177-2	176	90

## Example 3

*Glycine max* Centromere Repeat Sequences

[0268] We purified repetitive sequences from soybean (*Glycine max*, variety Williams 82), and set forth herein two centromere repeats, termed ChrGm1 and ChrGm2. We determined the consensus of each repeat as shown in Example 6.

[0269] The consensus sequence for ChrGm1 is shown in FIG. 2A (SEQ ID NO:5). This consensus was assembled from DNA sequences collected by the inventors. Seven of these sequences completely spanned the repeat, and twenty-five others partially covered the repeat. It is  $92 \pm 0.79$  base pairs in length, and A and T comprise of 63% of the consensus.

[0270] The consensus sequence for ChrGm2 is shown in FIG. 2B (SEQ ID NO:6). This consensus was assembled from DNA sequences collected by the inventors. Ten of these sequences completely spanned the repeat, and eleven others partially covered the repeat. It is  $91 \pm 0.48$  base pairs in length, and A and T comprise of 62% of the consensus.

[0271] The two repeats (ChrGm1 and ChrGm2) were aligned to each other using the ClustalX program Those sites which differ significantly from each other (chi-squared,  $P < 0.05$ ) are highlighted in FIG. 2C.

[0272] The GenBank nt database (<ftp://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on Apr. 7, 2002) and the plant satellite DNA database (<http://w3lanc. umbr.cas.cz/PlantSat/> downloaded on Apr. 14, 2002) were compared to the inventors' consensus sequences using the blastn program and an Expect value threshold of  $-3$ . Consensus sequences were built using all inventors' and GenBank sequences that matched with an Expect (E) value of less than  $-25$ .

[0273] The revised consensus sequence for ChrGm1 is shown in FIG. 2D (SEQ ID NO:7). This consensus was assembled from thirty-two DNA sequences collected by the inventors and one matching sequence from GenBank (accession number Z26334). Eight of these sequences completely spanned the repeat, and twenty-five others partially covered the repeat. It is  $92 \pm 0.74$  base pairs in length, and A and T comprise of 56% of the consensus.

[0274] The revised consensus sequence for ChrGm2 is shown in FIG. 2E (SEQ ID NO:8). This consensus was assembled from twenty-one DNA sequences collected by the inventors and three matching sequences from GenBank (accession numbers AF297983, AF297984, AF297985). Ten of these sequences completely spanned the repeat, and fourteen others partially covered the repeat. It is  $91 \pm 0.53$  base pairs in length, and A and T comprise of 61% of the consensus.

[0275] The two repeats (ChrGm1 and ChrGm2) were aligned to each other using the ClustalX program Those sites with significant differences (chi-squared,  $P < 0.05$ ) are highlighted in FIG. 2F.

[0276] A total of 4 GenBank entries match the *Glycine max* centromere sequences defined by the inventors. These are annotated as follows:

[0277] Satellite DNA

[0278] Tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes

[0279] They are not annotated as centromere repeats in GenBank. A complete list of these sequences is shown in Table 4:

TABLE 4

GenBank entries match the <i>Glycine max</i> centromere sequences defined by the inventors					
GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
Z26334	<i>G. max</i> satellite DNA	Complete	92-1	92	95
AF297985	<i>G. max</i> clone TRS3 tandem repetitive repeat region	Partial	259-173	87	93
AF297985	<i>G. max</i> clone TRS3 tandem repetitive repeat region	Partial	78-3	76	94
AF297985	<i>G. max</i> clone TRS3 tandem repetitive repeat region	Partial	168-83	86	90
AF297984	<i>G. max</i> clone TRS2 tandem repetitive repeat region	Partial	170-84	87	91
AF297984	<i>G. max</i> clone TRS2 tandem repetitive repeat region	Partial	260-175	86	88
AF297984	<i>G. max</i> clone TRS2 tandem repetitive repeat region	Partial	79-3	77	89
AF297983	<i>G. max</i> clone TRS1 tandem repetitive repeat region	Partial	77-3	75	94

## Example 4

*Lycopersicon esculentum* Centromere Repeat Sequences

[0280] We purified repetitive sequences from tomato (*Lycopersicon esculentum*, variety Microtom) and set forth herein one centromere repeat. We determined the consensus of this repeat as shown in Example 6.

[0281] The consensus sequence of ChrLe1 is shown in FIG. 3A (SEQ ID NO:9). This consensus was assembled from forty-two DNA sequences collected by the inventors. Eighteen of these sequences completely spanned the repeat, and twenty-four others partially covered the repeat. The repeat is  $181 \pm 0.61$  base pairs in length, and A and T comprise of 50% of the consensus.

[0282] The GenBank nt database (<http://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on Apr. 7, 2002) and the plant satellite DNA database (<http://w3lamc.umbr.cas.cz/PlantSat/>, downloaded on Apr. 14, 2002) were compared to the inventors' consensus sequences using the blastn program and an Expect value threshold value of  $-3$ . Consensus sequences were built using all inventors' and GenBank sequences matched with an Expect (E) value of less than  $-40$ .

[0283] We determined the consensus of this repeat. The repeat is  $181 \pm 0.61$  base pairs in length, and A and T comprise of 50% of the consensus.

[0284] The revised consensus sequence of ChrLe1 is shown in FIG. 3B (SEQ ID NO:10). This consensus was assembled from forty-two sequences collected by the inventors and two GenBank sequence (nt database at <http://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on Apr. 7, 2002). Eighteen of these sequences completely spanned the repeat, and twenty-six others partially covered the repeat. The GenBank sequences are accession numbers X87233 and AY007367.

[0285] Neither of the 2 GenBank entries that match the *Lycopersicon esculentum* centromere sequences defined by the inventors are complete repeats; they match only a portion of the sequence identified by the company. These are annotated as follows:

[0286] Satellite DNA

[0287] Tandem Repetitive Repeat Region

[0288] They are not annotated as centromere repeats in GenBank. A complete list of these sequences is shown in Table 5.

TABLE 5

GenBank entries match the <i>Lycopersicon esculentum</i> centromere sequences defined by the inventors					
GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
X87233	<i>L. esculentum</i> satellite DNA	Partial	163-1	161	93
AY007367	<i>L. esculentum</i> tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes	Partial	12003-12156	154	93
AY007367	<i>L. esculentum</i> tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes	Partial	12184-12344	161	90
AY007367	<i>L. esculentum</i> tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes	Partial	12546-12700	155	90

TABLE 5-continued

GenBank entries match the <i>Lycopersicon esculentum</i> centromere sequences defined by the inventors					
GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
AY007367	tospovirus resistance protein E (Sw5-e) genes	Partial	12365-12526	157	89
	tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes				

Example 5

*Zea mays* Centromere Repeat Sequences

[0289] We purified repetitive sequences from corn (*Zea mays*, variety B73), and set forth herein one centromere repeat, termed ChrZm1. We determined the consensus of the repeat as shown in Example 5. The repeat is  $180 \pm 1.15$  base pairs in length, and A and T comprise of 56% of the consensus.

[0290] The consensus sequence of ChrZm1 is shown in FIG. 4A (SEQ ID NO:11). This consensus was assembled from thirty-eight DNA sequences collected by the inventors. Three of these sequences completely spanned the repeat, and thirty-five others partially covered the repeat.

[0291] The GenBank nt database (<http://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on Apr. 7, 2002) and the plant satellite DNA database (<http://w3lamc.umbr.cas.cz/PlanSat/>, downloaded on Apr. 14, 2002) were compared to the inventors' consensus sequences using the blastn program and an Expect value threshold score of  $-3$ . Consensus sequences were built using all inventors' and GenBank sequences matched with an Expect (E) value of  $-50$ .

[0292] The revised consensus sequence of ChrZm1 is shown in FIG. 4B (SEQ ID NO:12). This consensus was assembled from thirty-eight DNA sequences collected by the inventors and twenty-six matching GenBank sequences (Table 6). Twenty of these sequences completely spanned the repeat, and forty-four others partially covered the repeat. The length of the repeat is  $180 \pm 0.51$  base pairs, and A and T comprise the consensus.

TABLE 6

GenBank sequences that match the inventors' ChrZm1 consensus					
M32521	M32522	M32523	M32524	M32525	M32526
M32527	M32528	M32529	M32530	M32531	M32532
M32533	M32534	M32535	M32536	M32537	M32538
M35408	AF030934	AF030935	AF030936	AF030937	AF030938
AF030939	AF030940				

[0293] A total of 26 GenBank entries match the *Zea mays* centromere sequences defined by the inventors. These are annotated as follows:

[0294] 180-bp knob-specific repeat region

[0295] heterochromatin repetitive DNA

[0296] They are not annotated as centromere repeats in GenBank. A complete list of these sequences is shown in Table 7.

TABLE 7

GenBank entries match the <i>Lycopersicon esculentum</i> centromere sequences defined by the inventors					
GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
M32522	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M32521	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M32533	<i>Z. mays</i> subsp. <i>mexicana</i> 180-bp knob-specific repeat region	Complete	1-180	180	96
M32525	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M32524	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M32523	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M35408	Corn heterochromatin repetitive DNA	Complete	1-180	180	96
M32526	Maize 180-bp knob-specific repeat region	Complete	1-180	180	95
AF030939	<i>Z. mays</i> 180-bp knob-associated tandem repeat 15-T3-2	Complete	1-180	180	95
M32528	Maize 180-bp knob-specific repeat region	Complete	1-180	180	95
M32534	<i>Z. mays</i> subsp. <i>mexicana</i> 180-bp knob-specific repeat region	Complete	1-180	180	94
M32527	Maize 180-bp knob-specific repeat region	Partial	8-179	172	95
M32538	<i>T. dactyloides</i> ( <i>Tripsacum dactyloides</i> , gama grass) 180-bp knob-specific repeat region	Complete	1-179	179	94
M32529	Maize 180-bp knob-specific repeat region	Complete	1-180	180	93
AF030938	<i>Z. mays</i> 180-bp knob-associated tandem repeat 15-T3-1	Partial	4-180	177	93
M32532	Maize 180-bp knob-specific repeat region	Complete	1-180	180	93
AF030937	<i>Z. mays</i> 180-bp knob-associated tandem repeat 1-T7-2	Complete	1-180	180	92

TABLE 7-continued

GenBank entries match the <i>Lycopersicon esculentum</i> centromere sequences defined by the inventors					
GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
AF030940	<i>Z. mays</i> 180-bp knob-associated tandem repeat 15-T7-1	Complete	1-180	180	92
AF030936	<i>Z. mays</i> 180-bp knob-associated tandem repeat 1-T7-1	Partial	10-180	172	93
M32537	<i>T. dactyloides</i> 180-bp knob-specific repeat region	Complete	1-180	180	92
M32530	Maize 180-bp knob-specific repeat region	Complete	1-180	180	92
M32531	Maize 180-bp knob-specific repeat region	Complete	1-179, introduced one gap	180	91
AF030935	<i>Z. mays</i> 180-bp knob-associated tandem repeat 1-T3-2	Partial	1-175	175	90
AF030934	<i>Z. mays</i> 180-bp knob-associated tandem repeat 1-T3-1	Partial	47-201	155	92
M32536	<i>T. dactyloides</i> 180-bp knob-specific repeat region	Complete	1-180	180	94
M32535	<i>T. dactyloides</i> 180-bp knob-specific repeat region	Complete	1-177, 2% gaps	177	91

Six GenBank sequences of *Zea mays* centromeric repeat CentC were collected (Table 13) and assigned the identifier ChrZm2. The consensus of the repeat was determined as shown in Example 6. The repeat is 158 ± 1.6 base pairs in length. A and T comprises of 53% of the bases. All 6 sequences are of unit length.

[0297] The consensus sequence of ChrZm2 (SEQ ID NO:13) is shown in FIG. 4C.

TABLE 8

GenBank sequences of <i>Zea mays</i> centromeric repeat ChrZm2		
AF078918	AF078919	AF078920
AF0789121	AF078922	AF078923

Example 6

Determining Consensus Sequences

[0298] Sequences were first aligned and edited in Vector NTI suite7 (InforMax, 7600 Wisconsin Ave., Suite 1100, Bethesda, Md. 20814) and exported as a fasta file. A perl program, consensus.pl, was written and used to determine the consensus for each position within the repeats based on the following rules:

[0299] The most common base is designated as the consensus if it occurs three times more frequently than the second most common base.

[0300] If the occurrence of the most common base is not three times more frequent than the second most common base, but the combined frequency of the two most common

bases is three times that of the third most common base, and the frequency of the second most common base is greater than the frequency of the third most common base, then the second and first bases are together considered as a consensus polymorphism, and designated using the IUPAC codes (M=A or C, R=A or G, W=A or T, S=C or G, Y=C or T, k=G or T, V=A or C or G, H=A or C or T, D=A or G or T, B=C or G or T, N=G or T or C or A).

**[0301]** If the combined frequency of the two most common bases is not three times greater than that of the third most common base, but the combined frequency of the three most common bases is three times that of the fourth most common base, and the third most common base is more common than the fourth most common base, and the frequency of occurrence of the fourth base is less than or equal to 22%, the consensus is assigned according to the IUPAC ambiguity codes for the three most common bases. If the four bases occur approximately equally (23-27%), the consensus is assigned as N.

#### Example 7

##### Constructing BAC Vectors for Testing Centromere Function

**[0302]** A BAC clone may be retrofitted with one or more plant telomeres and selectable markers together with the DNA elements necessary for *Agrobacterium* transformation (FIG. 9). This method will provide a means to deliver any BAC clone into plant cells and to test it for centromere function.

**[0303]** The method works in the following way. The conversion vector contains a retrofitting cassette. The retrofitting cassette is flanked by Tn10, Tn5, Tn7, Mu or other transposable elements and contains an origin of replication and a selectable marker for *Agrobacterium*, a plant telomere array followed by T-DNA right and left borders followed by a second plant telomere array and a plant selectable marker (FIG. 9). The conversion vector is transformed into an *E. coli* strain carrying the target BAC. The transposable elements flanking the retrofitting cassette then mediate transposition of the cassette randomly into the BAC clone. The retrofitted BAC clone can now be transformed into an appropriate strain of *Agrobacterium* and then into plant cells where it can be tested for high fidelity meiotic and mitotic transmission which would indicate that the clone contained a complete functional plant centromere.

#### Example 8

##### Sequence Analysis of *Arabidopsis* Centromeres

**[0304]** A. Abundance of genes in the centromeric regions

**[0305]** Expressed genes are located within 1 kb of essential centromere sequences in *S. cerevisiae*, and multiple copies of tRNA genes reside within an 80 kb fragment necessary for centromere function in *S. pombe* (Kuhn et al., 1991). In contrast, genes are thought to be relatively rare in the centromeres of higher eukaryotes, though there are notable exceptions. The *Drosophila* light, concertina, responder, and rolled loci all map to the centromeric region of chromosome 2, and translocations that remove light from its native heterochromatic context inhibit gene expression. In contrast, many *Drosophila* and human genes that normally reside in euchromatin become inactive when they are inserted near a centromere. Thus, genes that reside near centromeres likely have

special control elements that allow expression (Karpen, 1994; Lohe and Hilliker, 1995). The sequences of *Arabidopsis* CEN2 and CEN4, provided herein, provide a powerful resource for understanding how gene density and expression correlate with centromere position and associated chromatin.

**[0306]** Annotation of chromosome II and IV (<http://www.ncbi.nlm.nih.gov/Entrez/nucleotide.html>) identified many genes within and adjacent to CEN2 and CEN4 (FIG. 8, FIGS. 11A-11T). The density of predicted genes on *Arabidopsis* chromosome arms averages 25 per 100 kb, and in the repeat-rich regions flanking CEN2 and CEN4 this decreases to 9 and 7 genes per 100 kb, respectively (Bevan et al., 1999). Many predicted genes also reside within the recombination-deficient, genetically-defined centromeres. Within CEN2, there were 5 predicted genes per 100 kb; while CEN4 was strikingly different, with 12 genes per 100 kb.

**[0307]** There was strong evidence that several of the predicted centromeric genes are transcribed. The phosphoenolpyruvate gene (CUE1) defines one CEN5 border; mutations in this gene cause defects in light-regulated gene expression (Li et al., 1995). Within the sequenced portions of CEN2 and CEN4, 17% (27/160) of the predicted genes shared >95% identity with cloned cDNAs (ESTs), with three-fold more matches in CEN4 than in CEN2 (<http://www.tigr.org/tdb/at/agad/>). Twenty-four of these genes have multiple exons, and four correspond to single-copy genes with known functions. A list of the predicted genes identified is given in Table 9, below. A list of additional genes encoded within the boundaries of CEN4 are listed in Table 10. The identification of these genes is significant in that the genes may themselves contain unique regulatory elements or may reside in genomic locations flanking unique control or regulatory elements involved in centromere function or gene-expression. In particular, the current inventors contemplate use of these genes, or DNA sequences 0 to 5 kb upstream or downstream of these sequences, for insertion into a gene of choice in a mini chromosome. It is expected that such elements could potentially yield beneficial regulatory controls of the expression of these genes, even when in the unique environment of a centromere.

**[0308]** To investigate whether the remaining 23 genes were uniquely encoded at the centromere, a search was made in the database of annotated *Arabidopsis* sequences. With the exception of two genes, no homologs with >95% identity were found elsewhere in the 80% of the genome that has been sequenced. The number of independent cDNA clones that correspond to a single-copy gene provides an estimate of the level of gene expression. On chromosome II, predicted genes with high quality matches to the cDNA database (>95% identity) match an average of four independent cDNA clones (range 1-78). Within CEN2 and CEN4, 11/27 genes exceed this average (Table 9). Finally, genes encoded at CEN2 and CEN4 are not members of a single gene family, nor do they correspond to genes predicted to play a role in centromere functions, but instead have diverse roles.

**[0309]** Many genes in the *Arabidopsis* centromeric regions are nonfunctional due to early stop codons or disrupted open reading frames, but few pseudogenes were found on the chromosome arms. Though a large fraction of these pseudogenes have homology to mobile elements, many correspond to genes that are typically not mobile (FIGS. 11I-J and FIGS. 11S-T). Within the genetically-defined centromeres there were 1.0 (CEN2) and 0.7 (CEN4) of these nonmobile pseudogenes per 100 kb; the repeat-rich regions bordering the centromeres have 1.5 and 0.9 per 100 kb respectively. The dis-

tributions of pseudogenes and transposable elements are overlapping, indicating that DNA insertions in these regions contributed to gene disruptions.

TABLE 9

Predicted genes within CEN2 and CEN4 that correspond to the cDNA database.		
Putative function	GenBank protein accession	# of EST matches*
<b>CEN2</b>		
Unknown	AAC69124	1
SH3 domain protein	AAD15528	5
Unknown	AAD15529	1
unknown†	AAD37022	1
RNA helicase‡	AAC26676	2
40S ribosomal protein S16	AAD22696	9
<b>CEN4</b>		
Unknown	AAD36948	1
Unknown	AAD36947	4
leucyl tRNA synthetase	AAD36946	4
aspartic protease	AAD29758	6
Peroxisomal membrane protein (PPM2)§	AAD29759	5
5'-adenylylsulfate reductase§	AAD29775	14
symbiosis-related protein	AAD29776	3
ATP synthase gamma chain 1 (APC1)§	AAD48955	3
protein kinase and EF hand	AAD03453	3
ABC transporter	AAD03441	1
Transcriptional regulator	AAD03444	14
Unknown	AAD03446	12
human PCF11p homolog	AAD03447	6
NSF protein	AAD17345	2
1,3-beta-glucan synthase	AAD48971	2
pyridine nucleotide-disulphide oxidoreductase	AAD48975	4
Polyubiquitin (UBQ11)§	AAD48980	72
wound induced protein	AAD48981	6
short chain dehydrogenase/reductase	AAD48959	7
SL15†	AAD48939	2
WD40-repeat protein	AAD48948	2

\*Independent cDNAs with >95% identity,

†related gene present in non-centromeric DNA,

‡potentially associated with a mobile DNA element,

§characterized gene (B. Tugal, 1999; J. F. Gutierrez-Marcos, 1996; N. Inohara, 1991; J. Callis, 1995).

TABLE 10

List of additional genes encoded within the boundaries of CEN4.		
Putative Function	GenBank accession	Nucleotide Position
3'(2'),5'-Bisphosphate Nucleotidase	AC012392	71298-73681
Transcriptional regulator	AC012392	80611-81844
Equilibrative nucleoside transporter 1	AC012392	88570-90739
Equilibrative nucleoside transporter 1	AC012392	94940-96878
Equilibrative nucleoside transporter 1	AC012392	98929-101019
Equilibrative nucleoside transporter 1	AC012392	113069-115262
unknown	AC012392	122486-124729
4-coumarate--CoA ligase	AC012392	126505-128601
ethylene responsive protein	AC012392	130044-131421
Oxygen-evolving enhancer protein precursor	AC012392	134147-135224
Kinesin	AC012392	137630-141536
receptor-like protein kinase	AC012392	141847-144363
LpxD-like protein	AC012392	144921-146953
hypersensitivity induced protein	AC012392	147158-147838
ubiquitin	AC012392	149057-149677
unknown	AC012392	150254-151072
ubiquitin-like protein	AC012392	153514-154470
ubiquitin-like protein	AC012392	155734-156513
ubiquitin-like protein	AC012392	156993-157382

TABLE 10-continued

List of additional genes encoded within the boundaries of CEN4.		
Putative Function	GenBank accession	Nucleotide Position
unknown	AC012392	159635-165559
unknown	AC012392	166279-166920
unknown	AC012392	167724-170212
ubiquitin-like protein	AC012392	176819-178066
polyubiquitin (UBQ10)§	AC012392	180613-182007
phosphatidylinositol-3,4,5-triphosphate binding protein	AC012477	89384-91291
Mitochondrial ATPase	AC012477	94302-94677
RING-H2 finger protein	AC012477	95522-96142
unknown	AC012477	104747-105196
Mitochondrial ATPase	AC012477	105758-106595
ferredoxin--NADP+ reductase	AC012477	107451-109095
unknown	AC012477	109868-110620
U3 snoRNP-associated protein	AC012477	111841-114133
UV-damaged DNA binding factor	AC012477	114900-121275
Glucan endo-1,3-Beta-Glucosidase precursor	AC012477	122194-122895
D123-like protein	AC012477	125886-126887
Adrenodoxin Precursor	AC012477	127660-129246
N7 like-protein	AC012477	129718-131012
N7 like-protein	AC012477	131868-133963
N7 like-protein	AC012477	134215-136569
N7 like-protein	AC012477	139656-140864

§characterized gene (J. Callis, 1995).

### [0310] B. Conservation of Centromeric DNA

[0311] To investigate the conservation of CEN2 and CEN4 sequences, PCR primer pairs were designed that correspond to unique regions in the Columbia sequence and used to survey the centromeric regions of Landsberg and Columbia at ~20 kb intervals (FIGS. 13A, B). The primers used for the analysis are listed in FIGS. 14A, B. Amplification products of the appropriate length were obtained in both ecotypes for most primer pairs. (85%), indicating that the amplified regions were highly similar. In the remaining cases, primer pairs amplified Columbia, but not Landsberg DNA, even at very low stringencies. In these regions, additional primers were designed to determine the extent of nonhomology. In addition to a large insertion of mitochondrial DNA in CEN2, two other non-conserved regions were identified (FIGS. 13A, B). Because this DNA is absent from Landsberg centromeres, it is unlikely to be required for centromere function; consequently, the relevant portion of the centromeric sequence is reduced to 577 kb (CEN2) and 1250 kb (CEN4). The high degree of sequence conservation between Landsberg and Columbia centromeres indicated that the inhibition of recombination frequencies was not due to large regions of nonhomology, but instead was a property of the centromeres themselves.

### [0312] C. Sequence similarity between CEN2 and CEN4

[0313] In order to discern centromere function, a search was conducted for novel sequence motifs shared between CEN2 and CEN4, excluding from the comparison retroelements, transposons, characterized centromeric repeats, and coding sequences resembling mobile genes. After masking simple repetitive sequences, including homopolymer tracts and microsatellites, contigs of unique sequence measuring 417 kb and 851 kb for CEN2 and CEN4, respectively, were compared with BLAST (<http://blast.wustl.edu>).

[0314] The comparison showed that the complex DNA within the centromere regions was not homologous over the entire sequence length. However, 16 DNA segments in CEN2 matched 11 regions in CEN4 with >60% identity (FIG. 15).

The sequences were grouped into families of related sequences, and were designated AtCCS1-7 (*Arabidopsis thaliana* centromere conserved sequences 1-7). These sequences were not previously known to be repeated in the *Arabidopsis* genome. The sequences comprised a total of 17 kb (4%) of CEN2 DNA, had an average length of 1017 bp, and had an A+T content of 65%. Based on similarity, the matching sequences were sorted into groups, including two families containing 8 sequences each, 3 sequences from a small family encoding a putative open reading frame, and 4 sequences found once within the centromeres, one of which corresponds to predicted CEN2 and CEN4 proteins with similarity throughout their exons and introns (FIG. 15).

[0315] Searches of the *Arabidopsis* genomic sequence database demonstrated that AtCCS1-AtCCS5 were moderately repeated sequences that appear in centromeric and pericentromeric regions. The remaining sequences were present only in the genetically-defined centromeres. Similar comparisons of all 16 *S. cerevisiae* centromeres defined a consensus consisting of a conserved 8 bp CDEI motif, an AT-rich 85 bp CDEII element, and a 26 bp CDEII region with 7 highly conserved nucleotides (Fleig et al., 1995). In contrast, surveys of the three *S. pombe* centromeres revealed conservation of overall centromere structure, but no universally conserved motifs (Clark, 1998).

#### Example 9

##### Construction of Plant Mini Chromosomes

[0316] Mini chromosomes are constructed by combining the previously isolated essential chromosomal elements. Exemplary mini chromosome vectors include those designed to be "shuttle vectors"; i.e., they can be maintained in a convenient host (such as *E. coli*, *Agrobacterium* or yeast) as well as plant cells.

[0317] A. General Techniques for Mini Chromosome Construction

[0318] A mini chromosome can be maintained in *E. coli* or other bacterial cells as a circular molecule by placing a removable stuffer fragment between the telomeric sequence blocks. The stuffer fragment is a dispensable DNA sequence, bordered by unique restriction sites, which can be removed by restriction digestion of the circular DNAs to create linear molecules with telomeric ends. The linear mini chromosome can then be isolated by, for example, gel electrophoresis. In addition to the stuffer fragment and the plant telomeres, the mini chromosome contains a replication origin and selectable marker that can function in plants to allow the circular molecules to be maintained in bacterial cells. The mini chromosomes also include a plant selectable marker, a plant centromere, and a plant ARS to allow replication and maintenance of the DNA molecules in plant cells. Finally, the mini chromosome includes several unique restriction sites where additional DNA sequence inserts can be cloned. The most expeditious method of physically constructing such a mini chromosome, i.e., ligating the various essential elements together for example, will be apparent to those of ordinary skill in this art.

[0319] A number of mini chromosome vectors have been designed by the current inventors and are disclosed herein for the purpose of illustration (FIGS. 7A-7H). These vectors are not limiting however, as it will be apparent to those of skill in the art that many changes and alterations may be made and still obtain a functional vector.

[0320] B. Modified Technique for Mini Chromosome Construction

[0321] A two step method was developed for construction of mini chromosomes, which allows adding essential elements to BAC clones containing centromeric DNA. These procedures can take place in vivo, eliminating problems of chromosome breakage that often happen in the test tube. The details and advantages of the techniques are as follows:

[0322] 1.) One plasmid can be created that contains markers, origins and border sequences for *Agrobacterium* transfer, markers for selection and screening in plants, plant telomeres, and a loxP site or other site useful for site-specific recombination in vivo or in vitro. The second plasmid can be an existing BAC clone, isolated from the available genomic libraries (FIG. 10A).

[0323] 2.) The two plasmids are mixed, either within a single *E. coli* cell, or in a test tube, and the site-specific recombinase cre is introduced. This will cause the two plasmids to fuse at the loxP sites (FIG. 10B).

[0324] 3.) If deemed necessary, useful restriction sites (AseI/PacI or Not I) are included to remove excess material. (for example other selectable markers or replication origins)

[0325]<sub>r</sub> 4.) Variations include vectors with or without a Kan<sup>r</sup> gene (FIGS. 10B, 10C), with or without a LAT52 GUS gene, with a LAT52 GFP gene, and with a GUS gene under the control of other plant promoters. (FIGS. 10C, 10D and 10E).

[0326] C. Method for Preparation of Stable Non-Integrated Mini chromosomes

[0327] A technique has been developed to ensure that mini chromosomes do not integrate into the host genome (FIG. 10F). In particular, mini chromosomes must be maintained as distinct elements separate from the host chromosomes. In one method for ensuring that the introduced mini chromosome does not integrate, the inventors envision a variety that would encode a lethal plant gene (such as diphtheria toxin or any other gene product that, when expressed, causes lethality in plants). This gene could be located between the right *Agrobacterium* border and the telomere. Mini chromosomes that enter a plant nucleus and integrate into a host chromosome would result in lethality. However, if the mini chromosome remains separate, and further, if the ends of this construct are degraded up to the telomeres, then the lethal gene would be removed and the cells would survive.

[0328] It should be understood that various changes and modifications to the presently preferred embodiments described herein will be apparent to those skilled in the art. Such changes and modifications can be made without departing from the spirit and scope of the present invention and without diminishing its intended advantages. It is therefore intended that such changes and modifications be covered by the appended claims.

#### REFERENCES

- [0329] The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.
- [0330] Abdullah et al., *Biotechnology*, 4:1087, 1986.
- [0331] Abel et al., *Science*, 232:738-743, 1986.
- [0332] Alfenito et al., "Molecular characterization of a maize B chromosome centric sequence," *Genetics*, 135: 589-597, 1993.

- [0333] Alfenito and Birchler, "Molecular characterization of a maize B chromosome centric sequence," *Genetics* 135: 589-597, 1993.
- [0334] Ananiev et al., "A knob-associated tandem repeat in maize capable of forming fold-back DNA segments: are chromosome knobs megatransposons?" *Proc. Natl. Acad. Sci. U.S.A.* 95 (18), 10785-10790, 1998.
- [0335] Ananiev et al., "Chromosome-specific molecular organization of maize (*Zea mays* L.) centromeric regions," *Proc. Natl. Acad. Sci. U.S.A.* 95 (22), 13073-13078, 1998.
- [0336] Ananiev et al., "Complex structure of knob DNA on maize chromosome 9. Retrotransposon invasion into heterochromatin," *Genetics* 149 (4), 2025-2037, 1998
- [0337] Araki et al., "Site-specific recombinase, R, encoded by yeast plasmid pSR1," *J. Mol. Biol.* 225:25-37, 1992.
- [0338] Armstrong et al., "Physical mapping of DNA repetitive sequences to mitotic and meiotic chromosomes of *Brassica oleracea* var. *alboglabra* by fluorescence in situ hybridization," *Heredity* 81: 666-673, 1998.
- [0339] Barkai-Golan et al., *Arch. Microbiol.*, 116:119-124, 1978.
- [0340] Baum et al., "The centromeric K-type repeat and the central core are together sufficient to establish a functional *Schizosaccharomyces pombe* centromere," *Mol. Bio. Cell.*, 5:747-761, 1994.
- [0341] Bell et al., "Assignment of 30 microsatellite loci to the linkage map of *Arabidopsis*," *Genomics*, 19:137-144, 1994.
- [0342] Bernal-Lugo and Leopold, *Plant Physiol.*, 98:1-207-1210, 1992.
- [0343] Berzal-Herranz et al., *Genes and Devel.*, 6:129-134, 1992.
- [0344] Bevan et al., *Nucleic Acids Research*, 11(2):369-385, 1983.
- [0345] Bevan et al., *Biossays* 21:110, 1999.
- [0346] Blackman et al., *Plant Physiol.*, 100:225-230, 1992.
- [0347] Bloom, "The centromere frontier: Kinetochores components, microtubule-based motility, and the CEN-value paradox," *Cell*, 73:621-624, 1993.
- [0348] Bol et al., *Annu. Rev. Phytopath.*, 28:113-138, 1990.
- [0349] Bowler et al., *Ann Rev. Plant Physiol.*, 43:83-116, 1992.
- [0350] Brandes et al., *Chrom. Res.*, 5:238, 1997.
- [0351] Branson and Guss, *Proceedings North Central Branch Entomological Society of America*, 27:91-95, 1972.
- [0352] Brisson et al., *Nature*, 310:511, 1984.
- [0353] Broach et al., *Gene*, 8:121-133, 1979.
- [0354] Broakaert et al., *Science*, 245:1100-1102, 1989.
- [0355] Burke et al., *Science*, 236:806-812, 1987.
- [0356] Bytebier et al., *Proc. Natl Acad. Sci. USA*, 84:5345, 1987.
- [0357] Callis et al., *Genes and Development*, 1: 1183, 1987.
- [0358] Cambareri et al., *Mol. Cell. Biol.*, 18:5465, 1998.
- [0359] Campbell (ed.), *In: Avermectin and Abamectin*, 1989.
- [0360] Campbell, "Monoclonal Antibody Technology, Laboratory Techniques in Biochemistry and Molecular Biology," Vol. 13, Burden and Von Knippenberg, Eds. pp. 75-83, Elsevier, Amsterdam, 1984.
- [0361] Capecchi, "High efficiency transformation by direct microinjection of DNA into cultured mammalian cells," *Cell* 22(2):479-488, 1980.
- [0362] Carbon et al., *In: Recombinant Molecules: Impact on Science and Society* (Raven Press), 335-378, 1977.
- [0363] Carbon et al., "Centromere structure and function in budding and fission yeasts," *New Biologist*, 2:10-19, 1990.
- [0364] Carpenter et al., "The control of the distribution of meiotic exchange in *Drosophilla melanogaster*," *Genetics*, 101:81-90, 1982.
- [0365] Cech et al., "In vitro splicing of the ribosomal RNA precursor of Tetrahymena: involvement of a guanosine nucleotide in the excision of the intervening sequence," *Cell*, 27:487-496, 1981.
- [0366] Chandler et al., *The Plant Cell*, 1:1175-1183, 1989.
- [0367] Chang et al., "Restriction fragment length polymorphism linkage map for *Arabidopsis thaliana*," *Proc. Natl. Acad. Sci., USA*, 85:6856-6860, 1988.
- [0368] Charlesworth et al., *Nature*, 371:215, 1994.
- [0369] Charlesworth, C. H. Langley, W. Stephan, 112:947, 1986
- [0370] Chepko, *Cell*, 37:1053, 1984.
- [0371] Choi et al., *Plant Mol Biol Rep*, 13:124-29, 1995.
- [0372] Choo, K. H. A. *Genome Res.* 8:81, 1998.
- [0373] Chowrira et al., "In vitro and in vivo comparison of hammerhead, hairpin, and hepatitis delta virus self-processing ribozyme cassettes," *J. Biol. Chem.*, 269:25856-25864, 1994.
- [0374] Chu et al., "Separation of large DNA molecules by contour-clamped homogeneous electric fields" *Science*, 234, 1582-1585, 1986.
- [0375] Chye et al., *Plant Mol. Biol.*, 35:893, 1997.
- [0376] Clapp, "Somatic gene therapy into hematopoietic cells. Current status and future implications," *Clin. Perinatol.* 20(1):155-168, 1993.
- [0377] Clark, L. *Curr. Op. Gen. & Dev.*, 8:212, 1998
- [0378] Clarke et al., "Isolation of a yeast centromere and construction of functional small circular chromosomes," *Nature*, 287:504-509, 1980.
- [0379] Cohen et al., *Proc. Nat'l Acad. Sci. USA*, 70:3240, 1973.
- [0380] Conkling et al., *Plant Physiol.*, 93:1203-1211, 1990.
- [0381] Copenhaver and Pikaard, "RFLP and physical mapping with an rDNA-specific endonuclease reveals that nucleolus organizer regions of *Arabidopsis thaliana* adjoin the telomeres on chromosomes 2 and 4," *Plant J.*, 9:259-276, 1996.
- [0382] Copenhaver et al., "Use of RFLPs larger than 100 kbp to map position and internal organization of the nucleolus organizer region on chromosome 2 in *Arabidopsis thaliana*," *Plant J.* 7, 273-286, 1995.
- [0383] Copenhaver et al., *Proc. Natl. Acad. Sci.* 95:247, 1998.
- [0384] Copenhaver et al., *Science*. 286:2468-2474, 1999.
- [0385] Copenhaver and Preuss, *Plant Biology*, 2:104-108, 1999.
- [0386] Coxson et al., *Biotropica*, 24:121-133, 1992.
- [0387] Creusot et al., *Plant Journal*, 8:763-70, 1995
- [0388] Cristou et al., *Plant Physiol*, 87:671-674, 1988.
- [0389] Cuzzo et al., *Bio/Technology*, 6:549-553, 1988.
- [0390] Curiel et al., "Adenovirus enhancement of transferin-polylysine-mediated gene delivery," *Proc. Natl. Acad. Sci. USA* 88(19):8850-8854, 1991.
- [0391] Curiel et al., high-efficiency gene transfer mediated by adenovirus coupled to DNA-polylysine complexes," *Hum. Gen. Ther.* 3(2):147-154, 1992.



- [0392] Cutler et al., *J. Plant Physiol.*, 135:351-354, 1989.
- [0393] Czapla and Lang, *J. Econ. Entomol.*, 83:2480-2485, 1990.
- [0394] Davies et al., *Plant Physiol.*, 93:588-595, 1990.
- [0395] Dellaporta et al., *In: Chromosome Structure and Function: Impact of New Concepts*, 18th Stadler Genetics Symposium, 11:263-282, 1988.
- [0396] Dennis and Peacock, "Knob heterochromatin homology in maize and its relatives," *J. Mol. Evol.* 20, 341-350, 1984.
- [0397] Depicker et al., *Plant Cell Reports*, 7:63-66, 1988.
- [0398] DiLaurenzio et al., *Cell*, 86:423-33, 1996
- [0399] Dillon et al., *Recombinant DNA Methodology*, 1985.
- [0400] Donahue et al., "The nucleotide sequence of the HIS4 region of yeast," *Gene April*; 18(1):47-59, 1982.
- [0401] Dure et al., *Plant Molecular Biology*, 12:475-486, 1989.
- [0402] Earnshaw et al., "Proteins of the inner and outer centromere of mitotic chromosomes," *Genome*, 31:541-552, 1989.
- [0403] Earnshaw, "When is a centromere not a kinetochore?," *J. Cell Sci.*, 99:1-4, 1991.
- [0404] Ebert et al., 84:5745-5749, *Proc. Nat'l Acad. Sci. USA*, 1987
- [0405] Ecker, J R, *Genomics*, 19:137-144
- [0406] Ecker, *Methods*, 1:186-94, 1990.
- [0407] Eglitis et al., "Retroviral vectors for introduction of genes into mammalian cells," *Biotechniques* 6(7):608-614, 1988.
- [0408] Eglitis et al., "Retroviral-mediated gene transfer into hemopoietic cells," *Adv. Exp. Med. Biol.* 241:19-27, 1988.
- [0409] Enomoto et al., "Mapping of the pin locus coding for a site-specific recombinase that causes flagellar-phase variation in *Escherichia coli* K-12," *J. Bacteriol.*, 156:663-668, 1983.
- [0410] Erdmann et al., *J. Gen. Microbiology*, 138:363-368, 1992.
- [0411] Ferrin et al., "Selective cleavage of human DNA: RecA-Assited Restriction Endonuclease (RARE) cleavage," *Science*, 254:1494-1497, 1991.
- [0412] Fitzpatrick, *Gen. Engineering News*, 22:7, 1993.
- [0413] Fleig, U. et al., "Functional selection for the centromere DNA from yeast chromosome VIII," *Nuc. Acids. Res.* 23:922-924, 1995.
- [0414] Forster and Symons, "Self-cleavage of plus and minus RNAs of a virusoid and a structural model for the active sites," *Cell*, 49:211-220, 1987.
- [0415] Fraley et al., *Biotechnology*, 3:629, 1985.
- [0416] Franz et al., *Plant J.*, 13:867, 1998.
- [0417] Fromm et al., *Nature*, 312:791-793, 1986.
- [0418] Fromm et al., "Expression of genes transferred into monocot and dicot plant cells by electroporation," *Proc. Natl Acad. Sci. USA* 82(17):5824-5828, 1985.
- [0419] Fujimura et al., *Plant Tissue Culture Letters*, 2:74, 1985.
- [0420] Fynan et al., "DNA vaccines: protective immunizations by parenteral, mucosal, and gene gun inoculations," *Proc. Nat'l Acad. Sci. USA* 90(24):11478-11482, 1993.
- [0421] Gatehouse et al., *J. Sci. Food. Agric.*, 35:373-380, 1984.
- [0422] Geftter et al., *Somatic Cell Genet.* 3:231-236, 1977.
- [0423] Gerlach et al., "Construction of a plant disease resistance gene from the satellite RNA of tobacco ringspot virus," *Nature (London)*, 328:802-805, 1987.
- [0424] Goding, "Monoclonal Antibodies: Principles and Practice," pp. 60-74. 2nd Edition, Academic Press, Orlando, Fla., 1986.
- [0425] Golic and Lindquist, "The FLP recombinase of yeast catalyses site-specific recombination in the *Drosophila* genome," *Cell*, 59:499-509, 1989.
- [0426] Goring et al., *Proc. Natl. Acad. Sci. USA*, 88:1770-1774, 1991.
- [0427] Graham et al., "Transformation of rat cells by DNA of human adenovirus 5," *Virology* 54(2):536-539, 1973.
- [0428] Grellet et al., "Organization and evolution of a higher plant alphoid-like satellite DNA sequence," *J. Mol. Biol.* 187:495-507, 1986.
- [0429] Grill and Somerville, *Mol Gen Genet*, 226:484-90, 1991
- [0430] Guerrero et al., *Plant Molecular Biology*, 15:11-26, 1990.
- [0431] Gupta et al., *Proc. Natl. Acad. Sci. USA*, 90:1629-1633, 1993.
- [0432] Gutierrez-Marcos et al., *Proc. Natl. Acad. Sci.*, USA, 93:13377, 1996.
- [0433] Haaf et al., "Integration of human  $\alpha$ -satellite DNA into simian chromosomes: centromere protein binding and disruption of normal chromosome segregation," *Cell*, 70:681-696, 1992.
- [0434] Hadlaczy et al., "Centromere formation in mouse cells cotransformed with human DNA and a dominant marker gene," *Proc. Natl. Acad. Sci. USA*, 88:8106-8110, 1991.
- [0435] Hamilton et al., "Stable transfer of intact high molecular weight DNA into plant chromosomes," *Proc Natl Acad Sci USA* 93(18):9975-9, 1996
- [0436] Hamilton, "A binary-BAC system for plant transformation with high-molecular-weight DNA," *Gene*, 4; 200 (1-2):107-16, 1997.
- [0437] Hammock et al., *Nature*, 344:458-461, 1990.
- [0438] Harrison and Heslop-Harrison, "Centromeric repetitive DNA sequences in the genus *Brassica*," *Theor. Appl. Genet.* 90:157-165, 1995.
- [0439] Haseloff et al., *Proc. Nat'l Acad. Sci. USA* 94(6): 2122-2127, 1997.
- [0440] Hauge et al., *Symp Soc Exp Biol*, 45:45-56, 1991
- [0441] Hegemann et al., "The centromere of budding yeast," *Bioassays*, 15(7):451-460, 1993.
- [0442] Hemenway et al., *The EMBO J.*, 7:1273-1280, 1988.
- [0443] Heslop-Harrison et al., *Plant Cell*, 11:31, 1999.
- [0444] Hilder et al., *Nature*, 330:160-163, 1987.
- [0445] Hinchee et al., *Bio/technol.*, 6:915-922, 1988.
- [0446] Hoess et al., *Proc Natl Acad Sci*, 79:3398-402, 1982
- [0447] Hsiao et al., *J. Proc. Nat'l Acad. Sci. USA*, 76:3829-3833, 1979.
- [0448] Hudspeth and Grula, *Plant Mol. Biol.*, 12:579-589, 1989.
- [0449] Hwang et al., "Identification and map position of YAC clones comprising one-third of the *Arabidopsis* genome," *The Plant Journal*, 1:367-374, 1991.
- [0450] Ikeda et al., *J. Bacteriol.*, 169:5615-5621, 1987.
- [0451] Ikuta et al., *Bio/technol.*, 8:241-242, 1990.
- [0452] Inohara et al., *J. Biol. Chem.*, 266, 7333, 1991.

- [0453] Johnston et al., "Gene gun transfection of animal cells and genetic immunization," *Methods Cell. Biol.* 43(A):353-365, 1994.
- [0454] Jones, *Embo J.*, 4:2411-2418, 1985.
- [0455] Jones, *Mol. Gen. Genet.*, 207:478, 1987.
- [0456] Jorgensen et al., *Mol. Gen. Genet.*, 207:471, 1987.
- [0457] Jouanin et al., *Mol Gene Genet.*, 201:370-4, 1985
- [0458] Joyce, "RNA evolution and the origins of life," *Nature*, 338:217-244, 1989.
- [0459] Kaasen et al., *J. Bacteriology*, 174:889-898, 1992.
- [0460] Karpen, *Curr. Op. Gen. & Dev.*, 4:281, 1994.
- [0461] Karsten et al., *Botanica Marina*, 35:11-19, 1992.
- [0462] Katz et al., *J. Gen. Microbiol.*, 129:2703-2714, 1983.
- [0463] Kim and Cech, "Three dimensional model of the active site of the self-splicing rRNA precursor of Tetrahymena," *Proc. Natl. Acad. Sci. USA*, 84:8788-8792, 1987.
- [0464] Klee et al., *Bio/Technology* 3:637-642, 1985.
- [0465] Klein et al., *Nature*, 327:70-73, 1987.
- [0466] Klein et al., *Proc. Nat'l Acad. Sci. USA*, 85:8502-8505, 1988.
- [0467] Kohler et al., *Eur. J. Immunol.* 6:511-519, 1976.
- [0468] Kohler et al., *Nature* 256:495-497, 1975.
- [0469] Kolchinski and Gresshoff, "A major satellite DNA of soybean is a 92-base pairs tandem repeat," *Theor. Appl. Genet.* 90(5): 621-626, 1995.
- [0470] Konieczny et al., "A procedure for mapping *Arabidopsis* mutations using codominant ecotype-specific PCR-based markers," *The Plant Journal*, 4:403-410, 1993.
- [0471] Konieczny et al., *Genetics*, 127:801, 1991.
- [0472] Koorneef et al., *Genetica*, 61:41-46, 1983.
- [0473] Koorneef, "Linkage map of *Arabidopsis thaliana* (2n=10)," In S J O'Brien, ed. *Genetic Maps 1987: A compilation of linkage and restriction maps of genetically studied organisms*, 724-745, 1987.
- [0474] Koorneef, "The use of telotrisomics for centromere mapping in *Arabidopsis thaliana* (L.) Heynh," *Genetica*, 62:33-40, 1983.
- [0475] Koster and Leopold, *Plant Physiol.*, 88:829-832, 1988.
- [0476] Kubly, J., *Immunology 2nd Edition*, W.H. Freeman & Company, NY, 1994
- [0477] Kuhn et al., *Proc. Natl. Acad. Sci.*, 88:1306, 1991.
- [0478] Kyte et al., "A simple method for displaying the hydropathic character of a protein," *J. Mol. Biol.* 157(1): 105-132, 1982.
- [0479] Lawton et al., *Plant Mol. Biol.* 9:315-324, 1987.
- [0480] Lakshmikumaran and Ranade, "Isolation and characterization of a highly repetitive DNA of *Brassica campestris*," *Plant Mol. Biol.* 14:447-448, 1990.
- [0481] Lechner et al., "A 240 kd multisubunit protein complex, CBF3 is a major component of the budding yeast centromere," *Cell*, 64:717-725, 1991.
- [0482] Lee and Saier, *J. of Bacteriol.*, 153-685, 1983.
- [0483] Levings, *Science*, 250:942-947, 1990.
- [0484] Lewin, *Genes II*, John Wiley & Sons, Publishers, N.Y., 1985.
- [0485] Li et al., *Plant Cell*, 7:1599, 1995.
- [0486] Li et al., *Proc. Natl. Acad. Sci.*, 87:4580-4584, 1990.
- [0487] Lieber and Strauss, "Selection of efficient cleavage sites in target RNAs by using a ribozyme expression library," *Mol. Cell. Biol.*, 15: 540-551, 1995.
- [0488] Lin, S., Kaul, S. Rounsley, T. P. Shea, M-I. Benito, C. D. Town, C. Y. Fujii, T. Mason, C. L. Bowman, M. Barnstead, T. Feldblyum, C. R. Buell, K. A. Ketchum, C. M. Ronning, H. Koo, K. Moffat, L. Cronin, M. Shen, G. Pai, S. Van Aken, L., Umayam, L. Tallon, J. Gill, M. D. Adams, A. J. Carrera, T. H. Creasy, H. M. Goodman, C. R. Somerville, G. P. Copenhaver, D. Preuss, W. C. Nierman, O. White, J. A. Eisen, S. Salzberg, C. M. Fraser, and J. C. Venter, "Sequence and Analysis of Chromosome 2 of *Arabidopsis thaliana*," *Nature* 402: 761-768, 1999.
- [0489] Liu, Y G., Shirano, Y., Fukaki, H., Yanai, Y., Tasaka, M., Tabata, S., Shibata, D, *Proc. Natl Acad Sci USA* 96:6535-40, 1999.
- [0490] Lohe and Hilliker, *Curr. Op. Gen. & Dev.*, 5:746, 1995.
- [0491] Loomis et al., *J. Expt. Zoology*, 252:9-15, 1989.
- [0492] Lorz et al., *Mol. Gen. Genet.*, 199:178, 1985.
- [0493] Louis, E J, "Corrected sequence for the right telomere of *Saccharomyces cerevisiae* chromosome III," *Yeast*, 10(2):271-4, 1994.
- [0494] Lu et al., "High efficiency retroviral mediated gene transduction into single isolated immature and replatable CD34(3+) hematopoietic stem/progenitor cells from human umbilical cord blood," *J. Exp. Med.* 178(6):2089-2096, 1993.
- [0495] Maeser and Kahmann, "The GIN recombinase of phage Mu can catalyze site-specific recombination in plant protoplasts," *Mol. Gen. Genet.*, 230:170-176, 1991.
- [0496] Mahtani, M. M. and Willard, H. F. *Genome Res.* 8:100, 1998.
- [0497] Maloy, S. R., "Experimental Techniques in Bacterial Genetics" Jones and Bartlett Prokop, A., and Bajpai, R. K. "Recombinant DNA Technology I" *Ann. N.Y.Acad. Sci. vol.* 646, 1991.
- [0498] Maluszynaska et al., "Molecular cytogenetics of the genus *Arabidopsis*: In situ localization of rDNA sites, chromosome numbers and diversity in centromeric heterochromatin," *Annals Botany*, 71:479-484, 1993.
- [0499] Maluszynaska et al., "Localization of tandemly repeated DNA sequences in *Arabidopsis thaliana*," *Plant Jour.*, 1(2):159-166, 1991.
- [0500] Maniatis et al., "Molecular Cloning: a Laboratory Manual," Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1982.
- [0501] Marcotte et al., *Nature*, 335:454, 1988.
- [0502] Mariani et al., *Nature*, 347:737-741, 1990.
- [0503] Marra et al., *Nature Genet.* 22:265, 1999.
- [0504] Martinez-Zapater et al., *Mol. Gen. Genet.*, 204:417-423, 1986.
- [0505] Matsuura et al., *Journal of Bacteriology*, 178:3374-6, 1996
- [0506] McCabe et al., *Biotechnology*, 6:923, 1988.
- [0507] Michel and Westhof, "Modeling of the three-dimensional architecture of group I catalytic introns based on comparative sequence analysis," *J. Mol. Biol.*, 216:585-610, 1990.
- [0508] Mortimer et al., "Genetic mapping in *Saccharomyces cerevisiae*," *Life Cycle and Inheritance, In: The Molecular Biology of the Yeast Saccharomyces*, 11-26, 1981.
- [0509] Mozo et al., *Mol Gen Genet*, 258:562-70, 1998.
- [0510] Mozo et al., *Nature Genet.* 22:271, 1999.
- [0511] Mundy and Chua, *The EMBO J.*, 7:2279-2286, 1988.
- [0512] Murakami et al., *Mol. Gen. Genet.*, 205:42-50, 1986.

- [0513] Murata et al., *Plant J.*, 12:31, 1997.
- [0514] Murdock et al., *Phytochemistry*, 29:85-89, 1990.
- [0515] Murray et al., *Nature*, 305:189-193, 1983.
- [0516] Mysore et al., "An *Arabidopsis* histone H2A mutant is deficient in *Agrobacterium* T-DNA integration," *Proc Natl Acad Sci USA* 18; 97(2):948-53, 2000a.
- [0517] Mysore et al., "*Arabidopsis* ecotypes and mutants that are recalcitrant to *Agrobacterium* root transformation are susceptible to germ-line transformation." *Plant J* 21(1): 9-16, 2000b.
- [0518] Napoli, Lemieux, Jorgensen, "Introduction of a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes in trans," *Plant Cell*, 2:279-289, 1990.
- [0519] Negrutiu, I., Hinnisdaels, S., Cammaerts, D., Cherdshewasart, W., Gharti-Chhetri, G., and Jacobs, M. "Plant protoplasts as genetic tool: selectable markers for developmental studies," *Int. J. Dev. Biol.* 36: 73-84, 1992.
- [0520] Nester, *Ann. Rev. Plant Phys.*, 35:387-413, 1984.
- [0521] Nicklas, "The forces that move chromosomes in mitosis," *Annu. Rev. Biophys. Biophys. Chem.*, 17:431-39, 1988.
- [0522] Nussbaum et al., *Proc. Nat'l Acad. Sci. USA*, 73:1068, 1976.
- [0523] Odell et al., *Nature*, 313:810-812, 1985.
- [0524] Ohmori and Tomizawa, "Nucleotide sequence of the region required for maintenance of colicin E1 plasmid," *Mol Gen Genet*, October 3; 176(2):161-70, 1979.
- [0525] Omirulleh et al., *Plant Molecular Biology*, 21:415-428, 1993.
- [0526] Ow et al., *Science*, 234:856-859, 1986.
- [0527] Page et al., "Characterization of a maize chromosome 4 centromeric sequence: evidence for an evolutionary relationship with the B chromosome centromere," *Genetics* 159: 291-302, 2001.
- [0528] Palukaitis et al., "Characterization of a viroid associated with avacado sunblotch disease," *Virology*, 99:145-151, 1979.
- [0529] Peacock et al., "Highly repeated DNA sequence limited to knob heterochromatin in maize," *Proc. Natl. Acad. Sci. U.S.A.* 78, 4490-4494, 1981.
- [0530] Pelissier et al., *Genetica*, 97:141, 1996.
- [0531] Pelissier et al., *Plant Mol. Biol.*, 26:441, 1995.
- [0532] Perkins, "The detection of linkage in tetrad analysis," *Genetics*, 38, 187-197, 1953.
- [0533] Perlak et al., *Proc. Natl. Acad. Sci. USA*, 88:3324-3328, 1991.
- [0534] Perriman et al., "Extended target-site specificity for a hammerhead ribozyme," *Gene*, 113:157-163, 1992.
- [0535] Peterson et al., "Production of transgenic mice with yeast artificial chromosomes," *Trends Genet.* 13: 61-66, 1997.
- [0536] Phi-Van et al., *Mol. Cell. Biol.*, 10:2302-2307, 1990.
- [0537] Piatkowski et al., *Plant Physiol.*, 94:1682-1688, 1990.
- [0538] Potrykus et al., *Mol. Gen. Genet.*, 199:183-188, 1985.
- [0539] Prasher et al., *Biochem. Biophys. Res. Commun.*, 126(3):1259-1268, 1985.
- [0540] Pruss et al., "Tetrad analysis possible in *Arabidopsis* with mutation of the QUARTET (QRT) genes," *Science*, 264:1458, 1994.
- [0541] Price et al., "Systematic relationships of *Arabidopsis*: a molecular and morphological perspective", in: Somerville, C. and Meyerowitz, E. (eds.) *Arabidopsis*, Cold Spring Harbor Press, NY, 1995.
- [0542] Prody et al., "Autolytic processing of dimeric plant virus satellite RNA." *Science*, 231:1577-1580, 1986.
- [0543] Prokop et al., *Ann. N.Y. Acad. Sci.* 646, 1991 Puechberty, J. *Genomics* 56:247, 1999
- [0544] Rathore et al., *Plant Mol Biol*, 21:871-84, 1993
- [0545] Rattner, "The structure of the mammalian centromere," *Bioassays*, 13(2):51-56, 1991.
- [0546] Ravatn et al., *Journal of Bacteriology*, 180:5505-14, 1998.
- [0547] Reed et al., *J. Gen. Microbiology*, 130:1-4, 1984.
- [0548] Reichel et al., *Proc. Nat'l Acad. Sci. USA*, 93 (12) p. 5888-5893. 1996
- [0549] Reinhold-Hurek and Shub, "Self-splicing introns in tRNA genes of widely divergent bacteria," *Nature*, 357: 173-176, 1992.
- [0550] Rensburg et al., *J. Plant Physiol.*, 141:188-194, 1993.
- [0551] Richards and Ausubel, "Isolation of a higher eukaryotic telomere from *Arabidopsis thaliana*," *Cell*, 8:53(1):127-36, 1988.
- [0552] Richards et al., "The centromere region of *Arabidopsis thaliana* chromosome 1 contains telomere-similar sequences," *Nucleic Acids Research*, 19(12):3351-3357, 1991.
- [0553] Rieder, "The formation, structure and composition of the mammalian kinetochore and kinetochore fiber," *Int. Rev. Cytol*, 79:1-58, 1982.
- [0554] Rogers et al., *Meth. in Enzymol.*, 153:253-277, 1987.
- [0555] Rosenberg et al, "RFLP subtraction: A method for making libraries of polymorphic markers," *Proc. Nat'l Acad. Sci. USA*, 91:6113-6117, 1994.
- [0556] Round et al., *Genome Res*, 7, 1053, 1997.
- [0557] Sauer, "Functional expression of the cre-lox site-specific recombination system in the yeast *Saccharomyces cerevisiae*," *Mol. and Cell. Biol.*, 7: 2087-2096, 1987.
- [0558] Schmidt et al., *Plant Journal*, 5:735-44, 1994
- [0559] Schwartz et al., Cold Spring Harbor Symp. *Quant. Biol.*, 47, 195-198, 1982.
- [0560] Schweizer et al., "Species specific sequences for the identification of somatic hybrids between *Lycopersicon esculentum* and *Solanum acaule*," *Theor. Appl. Genet.* 75, 679-684, 1998
- [0561] Sears et al., "Cytogenetic studies in *Arabidopsis thaliana*," *Can. J. Genet. Cytol.*, 12:217-233, 1970.
- [0562] Segal, "Biochemical Calculations" 2nd Edition. John Wiley & Sons, New York, 1976.
- [0563] Setlow et al., *Genetic Engineering: Principles and Methods*, 1979.
- [0564] Shagan and Bar-Zvi, *Plant Physiol.*, 101:1397-1398, 1993.
- [0565] Shapiro, *In: Mobile Genetic Elements*, 1983.
- [0566] Sheen et al., *Plant Journal*, 8(5):777-784, 1995.
- [0567] Shingo et al., *Mol. Cell. Biol.*, 6:1787, 1986.
- [0568] Simoens et al., *Nuc. Acids Res.*, 16:6753, 1988.
- [0569] Smith, Watson, Bird, Ray, Schuch, Grierson, "Expression of a truncated tomato polygalacturonase gene inhibits expression of the endogenous gene in transgenic plants," *Mol. Gen. Genet.*, 224:447-481, 1990.
- [0570] Smithies et al., *Nature*, 317:230-234, 1985.
- [0571] Smythe, "Pollen clusters," *Current Biology*, 4:851-853, 1994.

- [0572] Somerville, C. and Somerville, S., *Science* 285:380, 1999.
- [0573] Spielmann et al., *Mol. Gen. Genet.*, 205:34, 1986.
- [0574] Stalker et al., *Science*, 242:419-422, 1988.
- [0575] Stiefel et al., *Nature*, 341:343, 1989.
- [0576] Stinchcomb et al., *Nature*, 282:39-43, 1979.
- [0577] Stougaard, *The Plant Journal*, 3:755-761, 1993.
- [0578] Sullivan, Christensen, Quail, *Mol. Gen. Genet.*, 215 (3):431-440, 1989.
- [0579] Sun et al., *Cell*, 91:1007, 1997.
- [0580] Sutcliffe, *Proc. Nat'l Acad. Sci. USA*, 75:3737-3741, 1978.
- [0581] Symington et al., *Cell*, 52:237-240, 1988.
- [0582] Symons, "Avacado sunblotch viroid: primary sequence and proposed secondary structure." *Nucl. Acids Res.*, 9:6527-6537, 1981.
- [0583] Symons, "Small catalytic RNAs." *Annu. Rev. Biochem.*, 61:641-671, 1992.
- [0584] Tarczynski et al., "Expression of a bacterial mtlD gene in transgenic tobacco leads to production and accumulation of mannitol," *Proc. Natl. Acad. Sci. USA*, 89:1-5, 1992.
- [0585] Tarczynski et al., "Stress Protection of Transgenic Tobacco by Production of the Osmolyte Mannitol," *Science*, 259:508-510, 1993.
- [0586] Thillet et al., *J. Biol. Chem.*, 263:12500-12508, 1988.
- [0587] Thomas et al., *Cell*, 44:419-428, 1986.
- [0588] Thomas et al., *Proc. Natl. Acad. Sci. USA*, 71:4579, 1974.
- [0589] Thompson et al., "Decreased expression of BRCA1 accelerates growth and is often present during sporadic breast cancer progression," *Nature Genet.*, 9:444-450, 1995.
- [0590] Thompson et al., *Nuc. Acids Res.*, 24:3017, 1996.
- [0591] Tian, Sequin, Charest, *Plant Cell Rep.*, 16:267-271, 1997.
- [0592] Tominaga, *Microbiology*, 143:2057-63, 1997
- [0593] Toriyama et al., *Theor. Appl. Genet.*, 73:16, 1986.
- [0594] Tsay et al., *Science*, 260:342, 1993.
- [0595] Tugal et al., *Plant Physiol.*, 120:309, 1999
- [0596] Twell et al., *Genes Dev* 5:496-507, 1991
- [0597] Twell et al., *Plant Physiol* 91:1270-1274, 1989.
- [0598] Tyler-Smith et al., "Mammalian chromosome structure," *Current Biology*, 3:390-397, 1993.
- [0599] Uchimiya et al., *Mol. Gen. Genet.*, 204:204, 1986.
- [0600] Vahedian et al., "Genomic organization and evolution of the soybean SB92 satellite sequence," *Plant Mol. Biol.* 29: 857-862, 1997.
- [0601] Van der Krol, Mur, Beld, Mol, Stuitje, "Flavonoid genes in petunia: addition of a limiting number of copies may lead to a suppression of gene expression," *Plant Cell*, 2:291-99, 1990.
- [0602] Van't Hof, Kuniyuki, Bjerken, "The size and number of replicon families of chromosomal DNA of *Arabidopsis thaliana*," *Chromosoma*, 68: 269-285, 1978.
- [0603] Vasil et al., "Herbicide-resistant fertile transgenic wheat plants obtained by microprojectile bombardment of regenerable embryogenic callus," *Biotechnology*, 10:667-674, 1992.
- [0604] Vasil, *Biotechnology*, 6:397, 1988.
- [0605] Vernon and Bohnert, *The EMBO J.*, 11:2077-2085, 1992.
- [0606] Voytas and Ausubel, *Nature*, 336:242, 1988.
- [0607] Wagner et al., "Coupling of adenovirus to transferin-polylysine/DNA complexes greatly enhances receptor-mediated gene delivery and expression of transfected genes," *Proc. Nat'l Acad. Sci. USA* 89 (13):6099-6103, 1992.
- [0608] Walker et al., *Proc. Nat'l Acad. Sci. USA*, 84:6624-6628, 1987.
- [0609] Wang et al., *Molecular and Cellular Biology*, 12(8): 3399-3406, 1992.
- [0610] Watrud et al., *In: Engineered Organisms and the Environment*, 1985.
- [0611] Watson et al., *Recombinant DNA: A Short Course*, 1983.
- [0612] Weide et al., "Paracentromeric sequences on tomato chromosome 6 show homology to human satellite III and to the mammalian CENP-B binding box," *Mol. Gen. Genet.* 259 (2): 190-197, 1998.
- [0613] Weinsink et al., *Cell*, 3:315-325, 1974.
- [0614] Wevrick et al., "Partial deletion of alpha satellite DNA association with reduced amounts of the centromere protein CENP-B in a mitotically stable human chromosome rearrangement," *Mol Cell Biol.*, 10:6374-6380, 1990.
- [0615] Whitehouse, *Nature*, No. 4205: 893, 1950.
- [0616] Wigler et al., *Cell*, 11:223, 1977.
- [0617] Willard, H., *Nature Genetics* 15:345-354, 1997
- [0618] Willard, H., "Centromeres of mammalian chromosomes" *Trends Genet.*, 6:410-416, 1990.
- [0619] Wolter et al., *The EMBO J.*, 4685-4692, 1992.
- [0620] Wong et al., "Electric field mediated gene transfer," *Biochim. Biophys. Res. Commun.* 107(2):584-587, 1982.
- [0621] Wright et al., *Genetics*, 142:569, 1996.
- [0622] Xia, X. et al., "Structure and evolution of a highly repetitive DNA sequence from *Brassica napus*," *Plant Mol. Biol.* 21:213-224, 1993.
- [0623] Xia, X. et al., "Genomic organization of the canrep repetitive DNA in *Brassica juncea*," *Plant Mol. Biol.* 26:817-832, 1994.
- [0624] Xiang and Guerra, *Plant Physiol.*, 102:287-293, 1993.
- [0625] Xu et al., *Plant Physiol.*, 110:249-257, 1996.
- [0626] Yamada et al., *Plant Cell Rep.*, 4:85, 1986.
- [0627] Yamaguchi-Shinozaki et al., *Plant Cell Physiol.*, 33:217-224, 1992.
- [0628] Yang and Russell, *Proc. Nat'l Acad. Sci. USA*, 87:4144-4148, 1990.
- [0629] Yen, *Embo J.* 10(5), 1245-1254, 1991.
- [0630] Young et al., *In: Eukaryotic Genetic Systems ICN-UCLA Symposia on Molecular and Cellular Biology, VII*, 315-331, 1977.
- [0631] Yuan and Altman, "Selection of guide sequences that direct efficient cleavage of mRNA by human ribonuclease P," *Science*, 263:1269-1273, 1994.
- [0632] Yuan et al., "Targeted cleavage of mRNA by human RNase P," *Proc. Natl. Acad. Sci. USA*, 89:8006-8010, 1992.
- [0633] Zatloukal et al., "Transferrinfection: a highly efficient way to express gene constructs in eukaryotic cells," *Ann. N.Y. Acad. Sci.*, 660:136-153, 1992.
- [0634] Zhang et al., *Gene*, 202:139-46, 1997
- [0635] Zhang et al., *Zea mays* B chromosome centromere repeat sequence *zea\_mays\_MBc216* pMBc216 unpublished
- [0636] Zukowsky et al., *Proc. Nat'l Acad. Sci. USA*, 80:1101-1105, 1983.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 13

<210> SEQ ID NO 1  
<211> LENGTH: 180  
<212> TYPE: DNA  
<213> ORGANISM: Brassica oleracea

<400> SEQUENCE: 1

agcttgattt ggatacataa agtggaggag aatcaccagg aagttgaata aatctcatag 60  
gagttggcat gaagaagtta tcccactttc aaatcagggtg attccagttt cccagtttg 120  
gaatagcaca gttcttctgt cgttccaatc aaaccaggat gaatcwtttt gtraraagct 180

<210> SEQ ID NO 2  
<211> LENGTH: 180  
<212> TYPE: DNA  
<213> ORGANISM: Brassica oleracea

<400> SEQUENCE: 2

agcttgattt tgatacataa agtagtggag aatcayyagg aagttgaata aatctcatag 60  
gagttaggat gaagaagtta tcmcactttc aaatcagggtg atcccarttt tctgtttg 120  
gaatatgaca acttmtttgt cattctaate aaaccaggaw gaatcgckat gtaaraagct 180

<210> SEQ ID NO 3  
<211> LENGTH: 180  
<212> TYPE: DNA  
<213> ORGANISM: Brassica oleracea

<400> SEQUENCE: 3

agcttgattt ggatacataa agtggaggag aatcaccagg aagttgaata aatctcatag 60  
gagttggsat gaagaagtta tcccactttc aaatcagggtg attccagttt cccagtttg 120  
gaatagcaca gttcttctgt cgttccaatc aaaccaggat gaatcacttt gtragaagct 180

<210> SEQ ID NO 4  
<211> LENGTH: 180  
<212> TYPE: DNA  
<213> ORGANISM: Brassica oleracea

<400> SEQUENCE: 4

agcttgattt tgatacataa agtartggag aatcayyagg aagttgaata aatctcatag 60  
gagttaggat gaagaagcta tcccactttc aaatcagggtg atcccarttt tctgtttg 120  
gaatak gaca rcttcttctgt cattctaate aaaccaggaw gaatcgckat gtaaraagct 180

<210> SEQ ID NO 5  
<211> LENGTH: 92  
<212> TYPE: DNA  
<213> ORGANISM: Glycine max

<400> SEQUENCE: 5

aaattcaaat ggtcataact tttmacwagg akgctccgatt caggcgcata atatctgag 60  
acgctcgaaa ttgaacaayg gaagctctcg ag 92

<210> SEQ ID NO 6  
<211> LENGTH: 91  
<212> TYPE: DNA

---

-continued

---

<213> ORGANISM: Glycine max

<400> SEQUENCE: 6

aaattcaaac gacaataact ttttactcgg atgtcycgatt gagtcccgta atatatcgag 60

acgctcgaaa ttgaatrytg aagctctgag c 91

<210> SEQ ID NO 7

<211> LENGTH: 92

<212> TYPE: DNA

<213> ORGANISM: Glycine max

<400> SEQUENCE: 7

aaattcaaat ggtcataact ttttactcgg akgtccgatt caggcgcata atatatcgag 60

acgctcgaaa ttgaacaayg gaagctctcg ag 92

<210> SEQ ID NO 8

<211> LENGTH: 91

<212> TYPE: DNA

<213> ORGANISM: Glycine max

<400> SEQUENCE: 8

aaattcaaac gacaataact ttttactcgg atgtcycgatt gagtcccgta atatatcgag 60

acgctcgaaa ttgaatrytg aagctctgag c 91

<210> SEQ ID NO 9

<211> LENGTH: 181

<212> TYPE: DNA

<213> ORGANISM: Lycopersicon esculentum

<400> SEQUENCE: 9

ccatcacggg ttttctgggc crtttggaaag gtcaaacgag ccccgagcgc agcatacgcc 60

tcattttgac gattttctgtg tgctattgca caccatTTTT tgggtgatcg ggattccgac 120

gtcaaaaatg ccaaatttgt tcgtggacgt cgtcaagac gttgtctatg catacggttg 180

g 181

<210> SEQ ID NO 10

<211> LENGTH: 181

<212> TYPE: DNA

<213> ORGANISM: Lycopersicon esculentum

<400> SEQUENCE: 10

ccatcacggg ttttctgggc crtttggaaag gtcaaacgag ccccgagcgc agcatacgcc 60

tcattttgac gattttctgtg tgctattgca caccatTTTT tgggtgatcg ggattccgac 120

gtcaaaaatg ccaaatttgt tcgtggacgt cgtcaagac gttgtctatg catacggttg 180

g 181

<210> SEQ ID NO 11

<211> LENGTH: 180

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<400> SEQUENCE: 11

ggccacacaa cccccatttt tgcgaaaaat agccatgaac gaccattttc aataatacyr 60

aaggctaaca cctacggatt tttraccaag aaatggctctc caccagaaat ccaagaatgt 120

-continued

---

gatctatggc aaggaacat atgtggggtg aggtgtayga gcctctggtc gaygatcaat 180

<210> SEQ ID NO 12  
 <211> LENGTH: 180  
 <212> TYPE: DNA  
 <213> ORGANISM: Zea mays

<400> SEQUENCE: 12

ggccacacaa cccccatttt tgtcgaaaat agccatgaay gaccattttc aataataaccg 60

aaggctaaca cctacggatt ttgaccaag aaatgggtctc caccagaaat ccaagaatgt 120

gatctatggc aaggaacat atgtggggtg aggtgtayga gcctctggtc gatgatcaat 180

<210> SEQ ID NO 13  
 <211> LENGTH: 158  
 <212> TYPE: DNA  
 <213> ORGANISM: Zea mays

<400> SEQUENCE: 13

ggttccgggtg gcaaaaactc gtagctttgt atgcaccmg acaccggtt tcggaatggg 60

tgacgtgyga caacagaaat tgcgmgaaac caccccaaac atgagttttg kacctaaagt 120

agtggattgg gcatgttcgt tgygaaaaac gaagaaat 158

---

1. A method for providing a service comprising the steps of:

- identifying a sequence associated with a centromer of a commercial plant;
- using a gene and the sequence to create a minichromosome; and
- introducing the minichromosome into a cell of the commercial plant to create a transgenic plant.

2-62. (canceled)

63. A method for generating revenue based on agricultural products comprising the steps of:

- developing a new crop based on a minichromosome including a sequence of a centromere; and
- receiving a fee for the new crop.

64. The method of claim 63 wherein the fee is received as a royalty payment.

65. The method of claim 63 wherein the fee is received through sales of seed generated by the new crop.

66. The method of claim 63 wherein the fee is received based on sales of the new crop.

67. The method of claim 63 wherein the fee is received through sales of a product made from the new crop.

68. The method of claim 63 wherein the fee is a one-time payment.

69. The method of claim 63 wherein the minichromosome includes at least one gene that adds a desirable characteristic to the new crop.

70. The method of claim 63 wherein the minichromosome includes at least two genes not present in an original crop containing the sequence.

71. The method of claim 63 wherein the fee is received, at least in part, as a cross-license of rights.

72. (canceled)

73. The method of claim 63 wherein the fee is received at least in part, as a service.

74. A method for performing a service comprising the steps of:

- developing a modified plant having a desired characteristic that does not naturally occur in a non-modified plant by inserting into a cell of the non-modified plant a minichromosome including
- a sequence of a centromere and a gene expressing the desired characteristic.

75-91. (canceled)

\* \* \* \* \*