



US 20090194317A1

(19) **United States**

(12) **Patent Application Publication**
Lindquist et al.

(10) **Pub. No.: US 2009/0194317 A1**

(43) **Pub. Date: Aug. 6, 2009**

(54) **ELECTRICAL CONDUCTORS AND DEVICES FROM PRION-LIKE PROTEINS**

Related U.S. Application Data

(75) Inventors: **Susan Lindquist**, Chestnut Hill, MA (US); **Thomas Scheibel**, Muenchen (DE)

(63) Continuation of application No. 11/089,551, filed on Mar. 24, 2005, now abandoned.

(60) Provisional application No. 60/559,286, filed on Mar. 31, 2004.

Correspondence Address:

MARSHALL, GERSTEIN & BORUN LLP
233 SOUTH WACKER DRIVE, 6300 SEARS TOWER
CHICAGO, IL 60606-6357 (US)

Publication Classification

(51) **Int. Cl.**
H01B 5/00 (2006.01)
C07K 1/02 (2006.01)
B05D 5/12 (2006.01)
(52) **U.S. Cl.** 174/126.1; 530/402; 427/118; 977/762


(73) Assignees: **THE WHITEHEAD INSTITUTE FOR BIOMEDICAL RESEARCH**, Cambridge, MA (US); **THE UNIVERSITY OF CHICAGO**, Chicago, IL (US)

(57) **ABSTRACT**

The present invention provides novel polypeptides comprising a prion-aggregation domain and a second domain; novel polynucleotides encoding such polypeptides; host cells transformed or transfected with such polynucleotides; novel fibrils with specific functionalities and unusually high chemical and thermal stability; and methods of making and using the foregoing.

(21) Appl. No.: 12/187,824

(22) Filed: Aug. 7, 2008



2184 TTACCTTTGTGCTGGAAGAAACGATTGCATCATTGATAAAAATTCGAAGGAAAAACTGCCAGCATGCCGCTATCGGAA
 728▶ Y L C A G R N D C I I D K I R R K N C P A C R Y R K
 2262 ATGCTTTCAGGCTGGAATGAACCTTGAAGCTCGAAAAACAAAGAAAAAATCAAAGGGATTTCAGCAAGCCACTGCAGG
 754▶ C L O A G M N L E A R K T K K K I K G I Q Q A T A G
 2341(GR526)
 2340 AGTCTCACAAAGACACTTCGGAAAATCCTAACAAAACAATAGTTCCTGCAGCATTACCACAGCTCACCCCTACCTTGGT
 780▶ V S Q D T S E N P N K T I V P A A L P Q L T P T L V
 2418 GTCACGTCTGGAGGTGATTGAACCCGAGGTGTTGTATGCAGGATATGATAGCTCTGTTCAGATTTCAGCATGGAGAAT
 806▶ S L L E V I E P E V L Y A G Y D S S V P D S A W R I
 2496 TATGACCACACTCAACATGTTAGGTGGGCGTCAAGTGATTGCAGCAGTGAATGGGCAAAGGCGATACTAGGCTTGAG
 832▶ M T T L N M L G G R O V I A A V K W A K A I L G L R
 2574 AAACCTACACCTCGATGACCAAAATGACCCCTGCTACAGTACTCATGGATGTTTCTCATGGCATTTCGCTTGGGTTGGAG
 858▶ N L H L D D Q M T L L Q Y S W M F L M A F A L G W R
 2652 ATCATACAGACAATCAAGCCGAAACCTGCTCTGCTTTGCTCCTGATCTGATTATTAATGAGCAGAGAATGCTCTTACC
 884▶ S Y R Q S S G N L L C F A P D L I N E Q R M S L P
 2730 CTGCATGTATGACCAATGTAAACACATGCTGTTTGTCTCCTCTGAATTACAAAGATTGCAGGTATCCTATGAAGAGTA
 910▶ C M Y D Q C K H M L F V S S E L O R L Q V S Y E E Y
 2808 TCTCTGTATGAAAACCTTACTGCTTCTCTCCTCAGTTCCTAAGGAAGGTCTGAAGAGCCAAGAGTTATTTGATGAGAT
 936▶ L C M K T L L L L L S S V P K E G L K S Q E L F D E I
 2886 TCGAATGACTTATATCAAAGAGCTAGGAAAAGCCATCGTCAAAGGGAAGGGAACCTCCAGTCAGAACTGGCAAAGGTT
 962▶ R M T Y I K E L G K A I V K R E G N S S O N W Q R F
 2964 TTACCAACTGACAAAGCTTCTGGACTCCATGCATGAGGTGGTTGAGAATCTCCTTACCTACTGCTTCCAGACATTTTT
 988▶ Y Q L T K L L D S M H E V V E N L L T Y C F Q T F L
 3042 GGATAAGACCATGAGTATGAAATCCAGAGATGTTAGCTGAAATCATCACTAATCAGATACCAAAATATTCAAATGG
 1014▶ D K T M S I E F P E M L A E I I T N Q I P K Y S N G
 3120 AAATATCAAAAAGCTTCTGTTTCATCAAAAATGA
 1040▶ N I K K L L F H Q K .

FIG. 1B

FIG. 2




FIG. 3A

1 TCGCGGTTT CGGTGATGAC GGTGAAAACC TCTGACACAT GCAGCTCCCG GAGACGGTCA CAGCTTGTCT GTAAGCGGAT GCCGGGAGCA GACAAGCCCG
04
101 TCAGGGCGCG TCAGCGGGTG TTGGCGGGTG TCGGGGCTGG CTTAACTATG CGGCATCAGA GCAGATGTA CTGAGAGTGC ACCATAACCAC AGCTTTTCAA
201 TTCAATTCAT CATTITTTTT TTATTCITTT TTTTGAITTC GGTTCITTTG AAATTTTTTT GATTCGGTAA TCTCCGAACA GAAGGANGAA CGAAGGAAGG

301 AGCACAGACT TAGATTGGTA TATATACGCA TATGTAGTGT TGAAGAAACA TGAATIGCC CAGTATICTT AACCCAAC TG CACAGAACAA AAACCTGCAG PstII (395)
401 GAAACGAAGA TAAATCATGT CGAAAGCTAC ATATAAGGAA CGTGCTGCTA CTCATCCTAG TCCTGTGCT GCCAAGCTAT TTAATATCAT GCACGAAAAG

1^D M S K A T Y K E R A A T H P S P V A A K L F N I M H E K
158^d G L G T A A L S N L I M C S F
501 CAAACAACT TGTTGCTTC ATTGGATGTT CGTACCACCA AGGAATTACT GGAGTTAGTT GAAGCATTAG GTCCCAAAT TTGTTACTA AAAACACATG

29^D Q T N L C A S L D V R T T K E L L E L V E A L G P K I C L L K T H
142^d C V F K H A E N S T R V V L S N S S N T S A N P G L I Q K S F V C T
EcoRV (503)
601 TGGATATCTT GACTGATTTT TCCATGGAGG GCACAGTTA^d GCCGCTAAAG GCATTATCCG CCAAGTACAA TTTTITACTC TTCGAAGACA GAAAATTGCG

62^D V D I L T D F S M E G T V K P L K A L S A K Y N F L L F E D R K F A
109^d S I K V S K E M S P V T L G S F A N D A L Y L K K S K S S L F N A
AplI (791)
701 TGACATTGGT AATACAGTCA AATTGCAGTA CTCGTCCGGT GTATACAGAA TAGCAGAATG GCCAGACATT ACGAATGCAC ACGGTGTGGT GGGCCACGGT

95^D D I G N T V K L Q Y S A G V Y R I A E W A D I T N A H G V V G P G
76^d S M P L V T L N C Y E A P T Y L I A S H A S M V F A C P T T P G P
AatII (851)
801 ATTGTTAGCG GTTTGAAGCA GCGGCAGAA GAAGTAACAA AGGAACCTAG AGGCTTTTG ATGTTAGCAG AATTGTCATG CAAGGCTCC CTATCTACTG

129^D I V S G L K Q A A E E V T K E P R G L L M L A E L S C K G S L S T
42^d I T L P K F C A A S S T V F S G L P R K I N A S N D H L P E R D V P
901 GAGAATATAC TAAGGTACT GTTGACATTG CGAAGAGCGA CAAGATTTT GTTATCGCT TTAATCTCA AAGAGACATG GGTGGAGAG ATGAAGGTTA

162^D G E Y T K G T V D I A K S D K D F V I G F I A O R D M G G R D E G Y
9^d S Y V L P V T S M
1001 CGATTGGTTG ATTATGACAC CCGTGTGGG TTTAGATGAC AAGGAGACG CATTGGGICA ACAGTATAGA ACCGTGGATG ATGTTGGTCTC TACAGGATCT

195^D D W L I M T P G V G L D D K G D A L G O Q Y R T V D D V V S T G S
1101 GACATTATTA TTGTTGGAAG AGGACTATTT GCAAGGGAA GGGATGCTAA GGTGAGGGT GAGCTTACA GAAAGCAGG CTGGGAAGCA TATTGAGAA

229^D D I I I V G R G L F A K G R D A K V E G E R Y R K A G W E A Y L R

5583 TTCAATGTTGGAAACACTATATAMGGGTTATAMAGTGGTTTAACTGTTTCCCTGTTGGTGGCTATATATCTCCAGTATACAAATGGTAAATATCATGATATCTTT
 5695 C TCGAGGGGGG GCGCGGTACC CAGCTTTTGT TCCCTTNGT GAGGGTAAAT TCGAGCTTG GCGTATCAT GGTCAATGCT GTTCTCTGTG TGAATTTGTT
 5796 ATCCCTCAC AATTCACAC AACTATAGAG CCGAGACAT AAGTGTAAA GCTGGGGTG CCTAATGAT GAGGTAATC AACTAATTTG CGTTCCGGCTC

PvuII

5896 ACTGCGCGCT TTCCAGTCCG GAAACTGTC GTCCACGTC CATTAATGA TGGCCACAG CCGGGGAGA GCGGGTTTC GATTTGGGGG CTCTTCGGCT
 5996 TCTCTCCTCA CTGACTCGT GCGCTCGTC GTTCGGCTG GCGAGCGGT ATCAGCTCAC TCANAGCCG TAAATAGGTT ATCCACAGAA TCAGGGGATA

PvuII

6096 AGCCAGAAA GAACATGTA GCAANAGCC AGCAANAGC CAGAACCGT AANAGCCG GTTTCTGCG GTTTTCCAT AGCTCCGCC CCCCAGCCA
 6196 CCATCACAAA ATCGACCT CAGTCCAG GTCCGAAAC CCGACAGAC TATPANGATA CCGGGGTTT CCCCCTGGA GCTCCCTGTT GCGCTCTCT
 6296 GTTCGACCC TCGCGCTTAC CCGATACCTG TCGCCCTTTC TCCCTTGGG AAGGCTGGC CTTCTCATA GCTCAACCTG TAGGTATTC AGTTGGGTT
 6396 AAGTGTTCG CTCANAGCTG GCTGTGTC AGAACCCC GGTTCAGCC CCTTATCCG TAACTATGTT CTTAGTCCA ACCCGGTAG
 6496 AACAGACTTA TCGCCACTGG CAGCAGCAC TGGTACAGG ATTAGCAGAG CGAGTATGT AAGGCGTCT ACAGATTTCT TGNAGTGTG GCCTAACTAC
 6596 GCCTACACTA GAAGGACAGT ATTGTGATC TCGCTCTGC TGAAGCCGT TACCTTGGG AANAGATGG GTAGCTCTG ATCCGGCANA CAAACCCCG
 6696 CTGGTAGCGG TGGTTTTTTT GTTTCNAGC ACCAGATTAC GCGCAGAAA AAGGATTC AAGAGTCC TTGTATCTTT TCTACGGGTT CTGAGGCTCA
 6796 GTGGAAAGAA AACTACGTT AAGGATTTT GGTATGAGA TTATCAAAA GGATCTTAC CTGATCCTT TTAATTTAAA ATGAAGTTT TAAATCAATC

6896 TAANGTAT ATGAGTAAAC TTGGTCTGAC AGTTAACAT GCTTATGAG TGGGCACT ATCTAGOGA TCTGTCTATT TGGTTCATCC ATAGTTGCT
 6996 GACTCCCGT CGTGTAGATA ACTAAGTAC GGGAGGCTT ACCATCTGC CCGAGTCTG CANTGATACC GCGAGAOCA CCGCTACCGG CTCAGATTT


2644 S G T Y I V V I R S P K G D P G L A A I I G R S G R E G A G S K
 7096 ATCAGAMTA AACGAGCCG CCGAGAGGC CAGAGGAGA AGTGTCTG CACTTATC CCGCTCATC CAGCTATA ATTTGTCGG GGAAGCTAGA

2314 D A I F W G A P L A S R L L P G A V K D A E M W D I L Q Q R S A L

FIG. 3D

7196 GTAAGTAGTT CGCCAGTTAA TAGTTTGGCG AACGTTGTG CCATTGCTAC AGGCATCGTG GTGTCACGCT CGTCGTTTGG TATGGCTTCA TTCAGCTCCG
1974 T L L E G T L L K R L T T A M A V P M T T D R E D N P I A E N L E P
PvuI (7368)
7296 GTTCCCAACG ATCAAGGCGA GTTACATGAT CCCCATGTT GTGCAAAAAA GCGGTTAGCT CCTTCGGTCC TCCGATCGTT GTCAGAAGTA AGTTGGCCGC
164 E W R D L R T V H D G M N H L F A T L E K P G G I T T L L L N A A
7396 AGTGTATCA CTCATGGTTA TGGCAGCACT GCATAATTCT CTTACTGTCA TGCCATCCGT AAGATGCTTT TCTGTGACTG GTGAGTACTC AACCAAGTCA
131 T N D S M T I A A S C L E R V T M G D T L H K E T V P S Y E V L D
7496 TTCTGAGAAT AGTGTATGCG GCGACCGAGT TGCTCTTGCC CGGCGTCAAT ACGGGATAAT ACCGCGCCAC ATAGCAGAAC TTTAAAAGTG CTCATCATTG
97 N O S Y H I R R G L Q E Q G A D I R S L V A G C L L V K F T S M M P
7596 GAAAACGTTT CTCGGGGCGA AAACCTCAA GGATCTTACC GCTGTTGAGA TCCAGTTCGA TGTAACCCAC TCGTGCACCC AACTGATCTT CAGCATCTTT
64 F R E E P R F S E L I K G S N L D L E I Y G V R A G L Q D E A D K
7696 TACTTTCACC AGCGTTTCTG GGTGAGCAAA AACAGGAAGG CAAAATGCCG CAAAAAGGG AATAAGGGCG ACACGGAAAT GTTGAATACT CATACTCTTC
31 V K V L T E P H A F V P L C F A A F F P I L A V R F H Q I S M
7796 CTTTTCAAT ATTATTGAAG CATTATCAG GGTTATTGTC TCATGAGCGG ATACATATTT GAATGTATTT AGAAAAATAA ACAAATAGGG GTTCCGCGCA
7896 CATTTCCTCCG AAAAGTGCCA CCTGACGTCT AAGAAACCAT TATTATCATG ACATTAACCT ATAAAAATAG GCGTATCAG AGGCCCTTTC ^{junction marker} GTC

FIG. 3E



Spontaneous conversion of Sup35 repeat mutants

FIG. 4




FIG. 5




FIG. 6




FIG. 7






FIG. 8



FIG. 9




FIG. 10

ELECTRICAL CONDUCTORS AND DEVICES FROM PRION-LIKE PROTEINS

[0001] This application claims priority benefit of U.S. Provisional Patent Application No. 60/559,286, filed Mar. 31, 2004. All priority applications are hereby incorporated by reference in their entirety.

ACKNOWLEDGMENT OF U.S. GOVERNMENT SUPPORT

[0002] This invention was made with U.S. Government support under Research Grant GM-25874 and GM-57840 awarded by the National Institutes of Health. The U.S. Government has certain rights in this invention.

FIELD OF THE INVENTION

[0003] The present invention relates generally to the fields of genetics and cellular and molecular biology, electronics, and nanotechnology. More particularly, the invention relates to amyloid or fibril-forming proteins and the genes that encode them, and especially to prion-like proteins and protein domains and the genes that encode them. The invention further relates to fibril-forming proteins that have been genetically or chemically modified to create fibrils that as electrical conductors, fuses, and electronic circuits.

DESCRIPTION OF RELATED ART

[0004] Nanometer-scale structures are of great interest as potential building blocks for future electronic devices. One significant challenge is the construction of nanowires to enable the electrical connection of such structures. Biomolecules may provide a solution to the difficulty of manufacturing wires at this scale because they naturally exist in the nanometer size range. Biomolecules that self-assemble have the potential to individually pattern into structures to aid the mass production of nanostructures.

[0005] The intrinsic properties of biomolecules are generally unsuitable for conducting electrical currents; therefore they are usually combined with an inorganic compound that acts as a conductor. This conductivity is achieved through a hierarchical assembly process where the first step is to form a regular scaffold by using biological molecules followed by a second step where the inorganic components are guided to aggregate selectively along the scaffold.

[0006] The first biomolecular templates used for microstructures were phospholipid tubules (Schnur, J. M., et al., *Thin Solid Films*, 152: 181-206 (1987)), and since then other self-assembling rod-like structures have been assessed for their strengths and weaknesses as nanostructural templates, including DNA, bacteriophages, and microtubules. These materials have many positive characteristics as nanostructure materials. DNA has good recognition capabilities, mechanical rigidity, and amenability to high-precision processing. Recent studies using DNA as a template for gold plating produced wires with ohmic conductivity [resistance, $R=86\Omega$ and a linear current-voltage (I-V) curve] (Hamack, O., et al., *Nanosci. Lett.*, 2: 919-923 (2002)); however, DNA is unstable under conditions (pH 10-12 and temperatures $>60^\circ\text{C}$.) necessary for industrial metallization. Bacteriophages are expected to have similar chemical and thermal constraints, and they do not readily polymerize to form continuous fibers.

[0007] Proteins are an attractive alternative material for the construction of nanostructures. Their physical size is appropriate and they are capable of many types of highly specific interactions; indeed, as many as 93,000 different protein-protein interactions have been predicted in yeast (Begley, T. J., et al., *Mol. Cancer Res.*, 1: 103-112 (2002); Uetz, P., et al., *Nature*, 403: 623-627 (2000); Marcotte, E., et al., *Nature*, 402: 83-86 (1999)). Moreover, proteins provide an extraordinary array of functionalities that could potentially be coupled to electronic circuitry in the building of nanoscale devices. Protein tubules have the advantage of a high degree of stiffness and greater stability than DNA. In addition they exhibit good adsorption to technical substrates like glass, silicon oxide, or gold. Various protein tubules such as microtubules and rhapsosomes (Fritzsche, W., et al., *Appl. Phys. Lett.*, 75: 2854-2856 (1999); Kirsch, R., et al., *Thin Solid Films*, 305: 248-253 (1997); Pazirandeh, M. & Campbell, J. R., *J. Gen. Microbiol.*, 139: 859-864 (1993)) have been assessed, but all have important limitations such as relatively high resistance once metallized (of the order of 200 k Ω) (Fritzsche, W., et al., supra), morphology that cannot withstand metallization under industrial conditions, or undesired aggregation once metallized (Kirsch, R., et al., supra). Therefore, there is a need to explore alternative biomaterials.

[0008] Prions (protein infectious particles) have been implicated in both human and animal spongiform encephalopathies, including Creutzfeldt-Jakob Disease, kuru, Gerstmann-Strassler-Scheinker Disease, and fatal familial insomnia in humans; the recently-publicized "mad cow disease" in bovines; "scrapie," which afflicts sheep and goats; transmissible mink encephalopathy; chronic wasting disease of mule, deer, and elk; and feline spongiform encephalopathy. See generally S. Prusiner et al., *Cell*, 93: 337-348 (1998); S. Prusiner, *Science*, 278:245-251 (1997); and A. Horwich and J. Weissman, *Cell*, 89: 499-510 (1997). A currently-accepted theory is that a prion protein (PrP) can exist in at least two conformational states: a normal, soluble cellular form (PrP^C) containing little β -sheet structure; and a "scrapie" form (PrP^{Sc}) characterized by significant β -sheet structure, insolubility, and resistance to proteases. Prion particles comprise multimers of the PrP^{Sc} form. Prion formation has been compared and contrasted to amyloid fibril formation that has been observed in other disease states, such as Alzheimer's disease. See J. Harper & P. Lansbury, *Annu. Rev. Biochem.*, 66: 385-407 (1997). More generally, the prion protein has been loosely classified (despite "some significant differences") as one of at least sixteen known human amyloidogenic proteins that, in an altered conformation, assemble into a fibril-like structure. See J. W. Kelly, *Curr. Opin. Struct. Biol.*, 6: 11-17 (1996), incorporated herein by reference.

[0009] There is growing patent and journal literature relating to scientists efforts to develop diagnostic, therapeutic, and prophylactic advances in the area of prion disease. For example, Fishleigh et al., U.S. Pat. No. 5,773,572 describes synthetic peptides that have at least one antigenic site of a prion protein, and suggest using such peptides to raise antibodies and to create vaccines. Prusiner et al., U.S. Pat. No. 5,750,361 describes prion protein peptides having at least one α -helical domain and forming a random coil conformation in aqueous medium, and suggests using such a peptide to assay for the scrapie form of prion protein (PrP^{Sc}).

[0010] Weiss et al., *J. Virology*, 69 (8): 4776-83 (1995) state that isolation of PrP^C from organisms has been a time-consuming and labor-intensive process. The authors purport to

describe the synthesis of Syrian golden hamster prion protein as a fusion with glutathione S-transferase (GST) to enhance solubility and stability of PrP^C, and the release of PrP^C from the fusion protein via thrombin cleavage. The authors report that only the cellular isoform PrP^C, and not the infectious PrP^{Sc} isoform, was produced. [See also Volkel et al., *Eur. J. Biochem*, 251:462-471 (1998); Meeker et al., *Proteins: Structure, Function, and Genetics*, 30: 381-387 (1998) (Describing system to overexpress a fusion between the small, minimally soluble serum amyloid A protein and the bacterial enzyme Staphylococcal nuclease; and Zahn et al., *FEBS Lett.*, 417 (3): 400-404 (1997) (reporting expression of human PrP proteins fused to a histidine tail to facilitate refolding).]

[0011] Prusiner et al., U.S. Pat. Nos. 5,792,901, 5,789,655, and 5,763,740 describe a transgenic mouse comprising a prion protein gene that includes codons from a PrP gene that is native to a different host organism, such as humans, and suggest uses of such mice for prion disease research. The '655 patent teaches to incorporate "a strong epitope tag" in the PrP nucleotide sequence to permit differentiation of PrP protein conformations using an antibody to the epitope. The patents describing these native, mutated, and chimeric PrP gene and protein sequences are incorporated herein by reference. Mouton et al., *Mol. Cell. Neurosci.*, 11 (3):127-133 (1998) report using a fusion between a putative nuclear localization signal of PrP and a green fluorescent protein to study targeting of the protein to the nuclear compartment.

[0012] Weissmann et al., U.S. Pat. No. 5,698,763, describes a transgenic mouse in which the PrP gene has been disrupted by homologous recombination, allegedly rendering the mouse non-susceptible to spongiform encephalopathies. Use of PrP anti-sense oligonucleotides to treat non-transgenic animals suffering from an incipient spongiform encephalopathy also is suggested.

[0013] Cashman et al., International Publication No. WO 97/45746, purports to describe prion protein binding proteins and uses thereof, e.g., to detect and treat prion-related diseases or to decontaminate samples known to contain or suspected of containing prion proteins. The authors also purport to describe a fusion protein having a PrP portion and an alkaline phosphatase portion, for use as an affinity reagent for labeling, detection, identification, or quantitation of PrP binding proteins or PrP^{Sc}'s in a biological sample, or for use to facilitate the affinity purification of PRP binding proteins.

[0014] In addition, there has been significant research in recent years concerning the biology of prion-like elements in yeast. [See, e.g., V. Kushnirov and M. Ter-Avanesyan, *Cell*, 94: 13-16 (1998); S. Lindquist, *Cell*, 89: 495-498 (1997); DePace et al., *Cell*, 93: 1241-1252 (1998); and R. Wickner, *Annu. Rev. Genet.*, 30:109-139 (1996) (all incorporated herein by reference).] Although the two yeast prion-like elements that have been extensively studied do not spread from cell to cell (except during mating or from mother-to-daughter cell) and do not kill the cells harboring them, as has been observed in the case of mammalian PrP prion diseases, certain heritable yeast phenotypes exist that display a very "prion-like" character. The phenotypes appear to arise as the result of the ability of a "normal" yeast protein that has acquired an abnormal conformation to influence other proteins of the same type to adopt the same conformation. Such phenotypes include the [PSI⁺] phenotype, which enhances the suppression of nonsense codons, and the [URE3] phenotype, which interferes with the nitrogen-mediated repression of certain catabolic enzymes. Both phenotypes exhibit cyto-

plasmic inheritance by daughter cells from a mother cell and are passed to a mating partner of a [PSI⁺] or [URE3] cell.

[0015] Yeast organisms present, in many respects, far easier systems than mammals in which to study genotype and phenotype relationships, and the study of the [PSI⁺] and [URE3] phenotypes in yeast has provided significant valuable information regarding prion biology. Studies have implicated the Sup35 subunit of the yeast translation termination factor and the Ure2 protein that antagonizes the action of a nitrogen-regulated transcription activator in the [PSI⁺] and [URE3] phenotypes, respectively. In both of these proteins, the above-stated "normal" biological functions reside in the carboxy-terminal domains, whereas the dispensable, amino-terminal domains have unusual compositions rich in asparagine and glutamine residues.

[0016] It is the amino-terminal domains of these proteins (e.g., no more than about residues 2-113 of Sup35 and about residues 1-65 of Ure2) that have been implicated in conferring the [PSI⁺] and [URE3] phenotypes in a prion-like manner. King et al., *Proc. Natl. Acad. Sci. USA*, 94:6618-6622 (1997), purportedly expressed the N-terminal 114 residues of SUP35 (with a cleavable polyhistidine tag for purification) and reported that this peptide spontaneously aggregates to form thin filaments showing a β -sheet-type circular dichroism in vitro. Deletion of the amino termini of Sup35 and Ure2 in yeast eliminates the [PSI⁺] and [URE3] phenotypes, respectively. In contrast, over-expression of these proteins, or of their amino-terminal fragments, can induce the [PSI⁺] or [URE3] phenotype de novo. Once cells have acquired the [PSI⁺] or [URE3] phenotype in this manner, they continue to pass the trait to their progeny, even after the plasmid containing the over-expressed element is lost. [See Derkatch et al., *Genetics*, 144:1375-1386 (1996).]

[0017] Interestingly, the Sup35 protein contains similarities to mammalian PrP proteins in that Sup35 is soluble in [psi⁻] strains but prone to aggregate into insoluble, protease-resistant aggregates in [PSI⁺] strains. In experiments using a fusion between the Sup35 amino terminus and green fluorescent protein (GFP, a protein that fluoresces green on exposure to blue light), it has been shown that the fusion protein is freely distributed in [psi⁻] cells but aggregated in [PSI⁺] cells. See, e.g., Glover et al., *Cell*, 89: 811-819 (1997); and Patino et al., *Science*, 273: 622-626 (1997). Chaperone proteins or "heat shock proteins," such as the protein Hsp104 in yeast, have been implicated in the conformational conversion of Sup35 protein that is associated with the [PSI⁺] phenotype [see, e.g., J. Glover and S. Lindquist, *Cell*, 94: 73-82 (1998); V. Kushnirov and M. Ter-Avanesyan, *Cell*, 94:13-16 (1998); Y. O. Chernoff et al., *Science*, 268: 880-883 (1995)], and may be implicated in the conformational conversion of PrP. See, e.g. E. Schirmer and S. Lindquist, *Proc. Natl. Acad. Sci. USA*, 94: 13932-13937 (1997); S. DeBurman et al., *Proc. Natl. Acad. Sci. USA*, 94:13938-13943 (1997).

[0018] As the foregoing discussion of literature indicates, there has been significant investigation into the biology of mammalian prions and prion-like yeast proteins for the purposes of developing a basic understanding of prion biology and developing effective measures for diagnosing, treating, and preventing mammalian prion diseases. Practical applications, including taking advantage of the structural characteristics and self-aggregating properties of prions and prion-like proteins, in addition to the immediate medical implications of

diagnosing, treating, and preventing spongiform encephalopathies and other amyloid diseases, is lacking.

SUMMARY OF THE INVENTION

[0019] The present invention relates to materials and methods involving prion-like fibers. For example, embodiments of the invention are directed to nanowires, fuses, circuits, and semiconductors constructed using modified prion-like elements as a scaffold, as well as methods of making and using them.

[0020] In one embodiment of the invention, an electrical conductor is provided comprising a fibril having a first location separated from a second location and an electrically conductive material disposed on the fibril between the first location and second location to conduct electricity along the fibril from the first location to the second location. The locations can be, but need not be, the ends of the fibril. In many practical applications, the first location may correspond with a contact between the electrical conductor and one element of an electrical circuit, and the second location may correspond to a contact with a second element of the circuit. In a preferred variation, the fibril used to make the electrical conductor comprises polypeptide subunits coalesced into an ordered aggregate, as described herein in detail.

[0021] Compared to other biological materials that have been contemplated for use in nanodevices, the fibrils described for use herein (e.g., for making electrical conductors) are characterized by chemical and thermal stability. In particular, the fibrils comprise polymers of polypeptide monomers which, as described below in detail, may exist in a soluble state or an aggregated fibrous state. For the purposes of this invention, a fibril that is characterized by chemical and thermal "stability" if it retains its fiber state for at least 60 minutes under conditions that may be encountered in industrial manufacturing processes and have a tendency to denature at least some proteins, nucleic acids, or other biological polymers. Exemplary conditions include elevated temperatures, extreme acidic or basic conditions, the presence of chemical denaturants, elevated salt conditions, and the presence of organic solvents. For example, fibrils for use in manufacturing an electrical conductor of the present invention preferably are chemically stable in the presence of:

[0022] denaturants such as urea (0-2M, more preferably 0-4M, more preferably 0-6M, more preferably 0-8 M) or guanidinium chloride (0-1M, more preferably 0-2 M);

[0023] salt solutions such as 0-1M or more preferably 0-2.5 M NaCl, KCl, sodium phosphate, or other halide salts;

[0024] industrial acids (e.g., aqueous solutions with pH between 4 and 7, or more preferably 3 and 7, more preferably 2 and 7, and more preferably 1-7 or 0.1-7);

[0025] basic solutions with pH in the range of 7-9, or more preferably 7-10 or 7-11 or 7-12 or 7-13;

[0026] organic solvents such as 100% ethanol;

[0027] extreme cold such as temperatures between 0-10° C., more preferably -10 to 0° C., -20 to 0° C., -30 to 0° C., -40 to 0° C., -50 to 0° C., -60 to 0° C., -70 to 0° C., or -80 to 0° C.;

[0028] heat such as temperatures between 50-60° C., and more preferably 50-70° C., 50-80° C., 50-90° C., 50-98° C., or 50-100° C.;

[0029] more generally, temperature ranges spanning both extreme cold and heat, e.g., thermal stability from -80° C. to 98° C. or any subranges thereof.

[0030] The techniques described herein can be used to make electrical conductors in a wide range of lengths and diameters. For example, electrical conductors may range in length from 0.05 to 10,000 μm in length, with every discrete length and range of lengths therebetween specifically contemplated, such as lengths of 0.06, 0.1, 0.2, 0.5, 0.8, 1, 10, 50, 100, 200 to 300 μm or more. Similarly, fibers may range in diameter from 1, 5, 9, 10, 20, 50, 75, 100, 150 to 200 nm, 300 nm, 400 nm, or 500 nm or more, with every diameter therebetween specifically contemplated as an embodiment of the invention. Diameter is influenced first by the diameter of the protein fibril used to make an electrical conductor, and second, by the amount and thickness of electrically conductive material disposed on its surface. In one embodiment, the aforementioned electrical conductor is provided wherein the electrical conductor is characterized by a length of 60 nm to 300 μm , and a diameter of 9 nm to 200 nm.

[0031] In another embodiment, the aforementioned electrical conductor is provided wherein at least one of the polypeptide subunits comprises a SCHAG amino acid sequence. Thus, the number of SCHAG amino acid sequences comprising an electrical conductor of the present invention can represent 0, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100% of the total polypeptide subunits in the electrical conductor. In a preferred embodiment, 90-100% of the polypeptide subunits comprise a SCHAG amino acid sequence.

[0032] In one embodiment of the invention, the aforementioned electrical conductor is provided wherein the SCHAG amino acid sequence includes at least one amino acid residue having a reactive amino acid side chain. It is possible that the SCHAG amino acid sequence, although containing at least one amino acid with a reactive amino acid side chain at the primary structure level, does not contain an amino acid with a reactive amino acid side chain that is surface exposed at the tertiary and/or quaternary structure level (e.g., when associated with fibrils). Accordingly, another embodiment of the invention provides the aforementioned electrical conductor wherein the SCHAG amino acid sequence includes at least one substitution of an amino acid residue having a reactive amino acid side chain.

[0033] Similarly, the number of amino acid substitutions may depend on the spatial relationship between the reactive amino acid side chains exposed to the environment and the length between the same or similar amino acid side chains of neighboring polypeptides in the fibril. Accordingly, a number of amino acid substitutions sufficient to reduce the gaps between amino acids with reactive side chains between neighboring polypeptides of the aforementioned electrical conductor is contemplated, thereby enabling a continuous connection along the length of the electrical conductor. It is also contemplated that the number of amino acid substitutions is inversely proportional to the amount of electrically conductive material required to provide the continuous connection along the length of the electrical conductor.

[0034] In a related embodiment, the aforementioned electrical conductor is provided wherein the reactive amino acid side chain is exposed to the environment of the fibril to permit attachment of the electrically conductive material thereto, and wherein the electrically conductive material is attached to the fibril at the reactive amino acid side chain. Similarly, another embodiment of the invention provides the aforementioned electrical conductor wherein the reactive amino acid side chain of the substituted amino acid is exposed to the environment of the fibril to permit attachment of the electri-

cally conductive material thereto, and wherein the electrically conductive material is attached to the fibril at the reactive amino acid side chain.

[0035] SCHAG amino acid sequences are rich in asparagine and glutamine residues. Thus, although many different amino acid sequences can comprise a SCHAG sequence, approximately 30% or more of the amino acid residues of SCHAG sequences may comprise asparagines and/or glutamine residues. Accordingly, in another embodiment of the invention, the aforementioned electrical conductor is provided wherein at least 30%, 35%, 40%, 45%, 50%, 60%, or more of the SCHAG amino acid sequence comprises asparagine or glutamine residues.

[0036] In another embodiment, the aforementioned electrical conductor is provided wherein the SCHAG amino acid sequence comprises an amino acid sequence at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97.5%, 98%, 99%, or 100% identical to a sequence selected from the group consisting of SEQ ID NOs: 2, 4, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 46, 47, and 50 and aggregation domain fragments thereof. Aggregation domain fragments are those fragments of the aforementioned sequences which contain enough of the original sequence to self-aggregate into fibers as described herein.

[0037] In yet another embodiment, the aforementioned electrical conductor is provided wherein the SCHAG amino acid sequence is selected from the group consisting of: a) an amino acid sequence that is at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97.5%, 98%, 99% or 100% identical to amino acids 2 to 113 of SEQ ID NO: 2; and b) an amino acid sequence that is at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97.5%, 98%, or 99% or 100% identical to amino acids 2 to 253 of SEQ ID NO: 2. In a related embodiment, the aforementioned electrical conductor is provided wherein the SCHAG amino acid sequence comprises at least one substitution of an amino acid residue having a reactive amino acid side chain and wherein the reactive amino acid side chain is exposed to the environment of the fibril to permit subsequent attachment of an electrically conductive material thereto.

[0038] As exemplified herein, specific amino acid sequences and amino acid substitutions are contemplated by the present invention. In one embodiment, the aforementioned electrical conductor is provided wherein the SCHAG amino acid sequence comprises the amino acid sequence of SEQ ID NO: 2, with the proviso that amino acid 184 of SEQ ID NO: 2 has been substituted for by an amino acid selected from the group consisting of cysteine, lysine, tyrosine, glutamate, aspartate, and arginine. In another embodiment, the aforementioned electrical conductor is provided wherein the SCHAG amino acid sequence comprises the amino acid sequence of SEQ ID NO: 2, with the proviso that amino acid 2 of SEQ ID NO: 2 has been substituted for by an amino acid selected from the group consisting of cysteine, lysine, tyrosine, glutamate, aspartate, and arginine.

[0039] Electrically conductive materials contemplated by the present invention include, but are not limited to, materials that comprise metal atoms and semiconductor materials. Thus, in one embodiment of the invention, the aforementioned electrical conductor is provided wherein the electrically conductive material comprises a material selected from the group consisting of a metal atom or a semiconductor material. Exemplary materials that comprise metal atoms are pure metals and metal alloys, inorganic compounds that contain metals, and organometallic compounds and complexes

comprised of one or more metal atoms attached to or complexed with an organic compound that can form a covalent bond with a polypeptide. Any conducting metal atom is suitable for practicing the invention, including but not limited to gold, silver, nickel, copper, platinum, aluminum, gallium, palladium, iridium, rhodium, tungsten, titanium, zinc, tin, alloys comprising the same, and combinations thereof. Additional metal atoms are also contemplated. The present invention further provides an electrical conductor wherein the semiconductor material is selected from the group consisting of GaAs, ZnS, CdS, InP and Si.

[0040] In one embodiment of the invention, the aforementioned electrical conductor is provided wherein the fibril is gold-toned. It is contemplated by the present invention that an electrical conductor described herein may possess a range of resistances from close to 0 ohms to 5000 ohms and every value in between. For example, resistances may range from 1, 5, 10, 20, 50, 75, 100, 150, 200, 250, 500, or 1000 Ω . In still another embodiment, the aforementioned electrical conductor is provided wherein the fibril is characterized by a resistance range of 0-100 Ω and linear I-V curves at useful power levels. Further, an electrical conductor is provided wherein the fibril is characterized by a resistance range of 0-100 Ω and linear I-V curves between 0 to 0.3×10^{-6} A and between 0-30 $\times 10^{-6}$ V.

[0041] A related aspect of the present invention is a method of making electrical conductors described herein, and methods of making electrical circuits, fuses, or devices comprising the electrical conductors.

[0042] For example, in one embodiment, a method of making an electrical conductor is provided comprising steps of: (a) making a fibril with first and second separated locations; and (b) disposing on the fibril an electrically conductive material in an amount effective to conduct electricity along the fibril from the first location to the second location.

[0043] Procedures for making the fibril (step (a)) are described below in detail. For example, such procedures comprise providing a solution or suspension of polypeptides that have the ability to coalesce into ordered aggregates, and incubating the solution or suspension under conditions to form fibrils from the polypeptides. A number of physical and chemical variations of such procedures are contemplated. In one embodiment, the method comprises rotating the solution or suspension to increase turbulence and surface area, thereby promoting fibril formation. In a preferred variation, the fiber formation further comprises contacting the fibrils with additional soluble or suspended polypeptide under conditions to extend the length of the fibrils.

[0044] The step (b) of disposing electrically conductive material can be performed in any manner by which an electrical conductor such as a metal can be disposed onto a fibril, such as chemical attachment, plating techniques, vapor deposition, combinations thereof, and the like. In one embodiment, step (b) comprises disposing a substrate on the fibril, and disposing a first electrically conductive material on the substrate. The substrate serves as a linker between the fibril and the first electrically conductive material, although the substrate can itself have electrical conducting properties. Thus, in one variation, the disposing the substrate comprises attaching a compound comprising a metal atom to a reactive amino acid side chain of a polypeptide in the fibril. For instance, the substrate optionally comprises gold particles with surface-accessible cross-linking groups. For example, a substrate exemplified herein is Nanogold, an organic, gold-

atom containing compound which contains gold atoms and can contribute to electrical conducting properties, and which was attached to exposed cysteine residues of a prion fibril. The Nanogold served as sites for subsequent attachment of silver and/or gold attachment. In a related embodiment, a second electrically conductive material is disposed on the first electrically conductive material.

[0045] As described herein, various electrically conductive materials are contemplated for use with the electrical conductors of the present invention. In one embodiment, the aforementioned method is provided wherein the disposing the first electrically conductive material comprises attaching a compound comprising a metal atom to the substrate. Further the aforementioned method is provided wherein the first electrically conductive material comprises silver ions. In yet another embodiment, the aforementioned method is provided wherein the disposing the second electrically conductive material comprises attaching a compound comprising a metal atom to the first electrically conductive material. In still another embodiment, the aforementioned method is provided wherein the second electrically conductive material comprises gold ions.

[0046] In a related embodiment, the aforementioned method is provided wherein the substrate comprises gold particles with surface-accessible cross-linking groups, the first electrically conductive material comprises silver ions, and the second electrically conductive material comprises gold ions. In a related embodiment, the aforementioned method is provided wherein the fibril is characterized by a resistance in the range of 0-100Ω and a linear current-voltage (I-V) curve.

[0047] In still another aspect, the invention includes all variety of electrical devices that can be synthesized with an electrical conductor of the invention. Such devices include everything from nanoscale wires, wires attached to substrates, fuses, circuits, and the like to larger and more complicated devices such as microchips, computers, consumer electronics, medical devices, laboratory tools, and the like that comprise electrical conductors, fuses, or circuits of the invention.

[0048] For example, in one embodiment, a fuse is provided comprising an electrical conductor, a first electrode attached to the first position, and a second electrode attached to the second position, wherein the electrical conductor electrically connects the first electrode to the second electrode. In a preferred variation of the fuse, the electrical conductor is constructed to fail to conduct electricity when exposed to an electrical current above a first amount, which can be described as the failure amount or overload amount of power. By "first amount" is simply meant an amount of electrical power (current×voltage) above which a fuse is designed to fail. In one variation, the electrical conductor destructs when exposed to an electric current above the first amount, thereby eliminating electrical conductivity across the fuse.

[0049] In another embodiment of the present invention, an electrical circuit is provided comprising a source of electricity, one or more circuit elements, and electrical conductors disposed between the source of electricity and the one or more circuit elements, wherein at least one of the electrical conductors is an electrical conductor of the invention. For example, the electrical conductor comprises a fibril and an electrically conductive material disposed on the fibril to conduct electricity along the fibril from a first position on the fibril such as the source of electricity to a second position on

the fibril, such as one of the circuit elements. The electrical conductor also may be disposed between two circuit elements. Exemplary circuit elements includes any circuit component selected from the group consisting of a capacitor, an inductor, a resistor, an integrated circuit, an oscillator, a transistor, a diode, a switch, and a fuse. The one or more circuit elements may be passive circuit elements, active circuit elements, or combinations thereof.

[0050] The present invention is also directed to employing unique features of prion biology in a practical context beyond fundamental prion research and applied research directed to the development of diagnostic, therapeutic, and prophylactic treatments of mammalian prion diseases (although aspects of the invention have utility in such contexts also). Likewise, the present invention also relates to the construction of novel prion-like elements that can change the phenotype of a cell in a beneficial way.

[0051] In one aspect, the invention provides a polynucleotide comprising a nucleotide sequence that encodes a chimeric polypeptide, the polynucleotide comprising: a nucleotide sequence encoding at least one SCHAG amino acid sequence fused in frame with a nucleotide sequence encoding at least one polypeptide of interest other than a marker protein, or a glutathione S-transferase (GST) protein, or a staphylococcal nuclease protein. In a preferred embodiment, the polynucleotide has been purified and isolated. In another preferred embodiment, the polynucleotide is stably transformed or transfected into a living cell.

[0052] By "chimeric polypeptide" is meant a polypeptide comprising at least two distinct polypeptide segments (domains) that do not naturally occur together as a single protein. In preferred embodiments, each domain contributes a distinct and useful property to the polypeptide. Polynucleotides that encode chimeric polypeptides can be constructed using conventional recombinant DNA technology to synthesize, amplify, and/or isolate polynucleotides encoding the at least two distinct segments, and to ligate them together. See, e.g., Sambrook et al., *Molecular Cloning—A Laboratory Manual*, Second Ed., Cold Spring Harbor Press (1989); and Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1998); both incorporated herein by reference.

[0053] The chimeric polypeptide comprises a SCHAG amino acid sequence as one of its polypeptide segments. By "SCHAG amino acid sequence" is meant any amino acid sequence which, when included as part or all of the amino acid sequence of a protein, can cause the protein to coalesce with like proteins into higher ordered aggregates commonly referred to in scientific literature by terms such as "amyloid," "amyloid fibers," "amyloid fibrils," "fibrils," or "prions." In this regard, the term SCHAG is an acronym for Self-Coalesces into Higher-ordered Aggregates. By "higher ordered" is meant an aggregate of at least 25 polypeptide subunits, and is meant to exclude the many proteins that are known to comprise polypeptide dimers, tetramers, or other small numbers of polypeptide subunits in an active complex. The term "higher-ordered aggregate" also is meant to exclude random agglomerations of denatured proteins that can form in non-physiological conditions. [From the term "self-coalesces," it will be understood that a SCHAG amino acid sequence may be expected to coalesce with identical polypeptides and also with polypeptides having high similarity (e.g., less than 10% sequence divergence) but less than complete identity in the SCHAG sequence.] It will be understood that many proteins that will self-coalesce into higher-ordered aggregates can

exist in at least two conformational states, only one of which is typically found in the ordered aggregates or fibrils. The term “self-coalesces” refers to the property of the polypeptide to form ordered aggregates with polypeptides having an identical amino acid sequence under appropriate conditions as taught herein, and is not intended to imply that the coalescing will naturally occur under every concentration or every set of conditions. In fact, data exists suggesting that trans-acting factors, such as chaperone proteins, may be involved in the protein’s conformational switching, *in vivo*). Aggregates formed by SCHAG polypeptides typically are rich in β -sheet structure, as demonstrated by circular dichroism; bind Congo red dye and give a characteristic spectral shift in polarized light; and are insoluble in water or in solutions mimicking the physiological salt concentrations of the native cells in which the aggregates originate. In preferred embodiments the SCHAG polypeptides self-coalesce to form amyloid fibrils that typically are 5-20 nm in width and display a “cross- β ” structure, in which the individual β strands of the component proteins are oriented perpendicular to the axis of the fibril. The SCHAG amino acid sequence may be said to constitute an “amyloidogenic domain” or “fibril-aggregation domain” of a protein because a SCHAG amino sequence confers this self-coalescing property to proteins which include it.

[0054] Exemplary SCHAG amino acid sequences include sequences of any naturally occurring protein that has the ability to aggregate into amyloid-type ordered aggregates under physiological conditions, such as inside of a cell. In one preferred embodiment, the SCHAG amino acid sequence includes the sequences of only that portion of the protein responsible for the aggregation behavior. Many such sequences have been identified in humans and other animals, including amyloid β protein (residues 1-40, 1-41, 1-42, or 1-43), associated with Alzheimer’s disease; immunoglobulin light chain fragments, associated with primary systemic amyloidosis; serum amyloid A fragments, associated with secondary systemic amyloidosis; transthyretin and transthyretin fragments, associated with senile systemic amyloidosis and familial amyloid polyneuropathy I; cystatin C fragments, associated with hereditary cerebral amyloid angiopathy; β_2 -microglobulin, associated with hemodialysis-related amyloidosis; apolipoprotein A-1 fragments, associated with familial amyloid polyneuropathy III; a 71 amino acid fragment of gelsolin, associated with Finnish hereditary systemic amyloidosis; islet amyloid polypeptide fragments, associated with Type II diabetes; calcitonin fragments, associated with medullary carcinoma of the thyroid; prion protein and fragments thereof, associated with spongiform encephalopathies; atrial natriuretic factor, associated with atrial amyloidosis; lysozyme and lysozyme fragments, associated with hereditary non-neuropathic systemic amyloidosis; insulin, associated with injection-localized amyloidosis; and fibrinogen fragments, associated with hereditary renal amyloidosis. See J. W. Kelly, *Curr. Op. Struct. Biol.*, 6: 11-17 (1996), incorporated herein by reference. In addition, several other SCHAG amino acid sequences of yeast and fungal origin are described in detail below. Also, the Examples below set forth in detail how to use the SCHAG sequences specifically identified herein or elsewhere in the literature to screen databases or genomes for additional naturally occurring SCHAG amino acid sequences. The Examples also provide assays to screen candidate SCHAG sequences for prion-like properties. In addition, the Examples provide assays to rapidly screen random DNA fragments to determine whether they encode a

SCHAG amino acid sequence. Such screening assays are themselves considered aspects of the invention.

[0055] In addition, SCHAG amino acid sequences include those sequences derived from naturally occurring SCHAG amino acid sequences by addition, deletion, or substitution of one or more amino acids from the naturally occurring SCHAG amino acid sequences. Detailed guidelines for modifying SCHAG amino acid sequences to produce synthetic SCHAG amino acid sequences are described below. Modifications that introduce conservative substitutions are specifically contemplated for creating SCHAG amino acid sequences that are equivalent to naturally occurring sequences. By “conservative amino acid substitution” is meant substitution of an amino acid with an amino acid having a side chain of a similar chemical character. Similar amino acids for making conservative substitutions include those having an acidic side chain (glutamic acid, aspartic acid); a basic side chain (arginine, lysine, histidine); a polar amide side chain (glutamine, asparagine); a hydrophobic, aliphatic side chain (leucine, isoleucine, valine, alanine, glycine); an aromatic side chain (phenylalanine, tryptophan, tyrosine); a small side chain (glycine, alanine, serine, threonine, methionine); or an aliphatic hydroxyl side chain (serine, threonine). Alternatively, similar amino acids for making conservative substitutions can be grouped into three categories based on the identity of the side chains. The first group includes glutamic acid, aspartic acid, arginine, lysine, histidine, which all have charged side chains; the second group includes glycine, serine, threonine, cysteine, tyrosine, glutamine, asparagine; and the third group includes leucine, isoleucine, valine, alanine, proline, phenylalanine, tryptophan, methionine, as described in Zubay, G., *Biochemistry*, third edition, Wm.C. Brown Publishers (1993).

[0056] Also contemplated are modifications to naturally occurring SCHAG amino acid sequences that result in addition or substitution of polar residues (especially glutamine and asparagine, but also serine and tyrosine) into the amino acid sequence. Certain naturally occurring SCHAG amino acid sequences are characterized by short, sometimes imperfect repeat sequences of, e.g., 5-12 residues. Modifications that result in substantial duplication of such repetitive oligomers are specifically contemplated for creating SCHAG amino acid sequences, too.

[0057] In another variation of the invention, the SCHAG amino acid sequence is encoded by a polynucleotide that hybridizes to any of the nucleotide sequences of the invention; or the non-coding strands complementary to these sequences, under the following exemplary moderately stringent hybridization conditions:

[0058] (a) hybridization for 16 hours at 42° C. in an aqueous hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% Dextran sulphate; and

[0059] (b) washing 2 times for 30 minutes at 60° C. in an aqueous wash solution comprising 0.1% SSC, 1% SDS. Alternatively, highly stringent conditions include washes at 68° C.

[0060] Also provided are purified and isolated polynucleotide comprising a nucleotide sequence that encodes at least one SCHAG amino acid sequence, wherein the SCHAG-encoding portion of the polynucleotide is at least about 99%, at least about 98%, at least about 95%, at least about 90%, at least about 85%, at least about 80%, at least about 75%, or at least about 70% identical over its full length to one of the nucleotide sequences of the invention. Methods of screening

for natural or artificial sequences for SCHAG properties are also described elsewhere herein.

[0061] A preferred category of SCHAG amino acid sequences are prion aggregation domains from prion proteins. The term “prion-aggregation domain” is intended to define a subset of SCHAG amino acid sequences that can exist in at least two conformational states, only one of which is typically found in the aggregated state. In one conformational state, proteins comprising the prion-aggregation domain or fused to the prion-aggregation domain perform their normal function in a cell, and in another conformational state, the native proteins form aggregates (prions) that phenotypically alter the cell, perhaps by sequestering the protein away from its normal site of subcellular activity, or by disrupting the conformation of an active domain of the protein, or by changing its activity state, or by acquiring a new activity upon aggregation, or perhaps merely by virtue of a detrimental effect on the cell of the aggregate itself. A hallmark feature of prion-aggregation domains is that the phenotypic alteration that is associated with prion formation is heritable and/or transmissible: prions are passed from mother to daughter cell or to mating partners in organisms such as in the case of yeast Sup35, and Ure2 prions, perpetuating the [PSI⁺] or [URE3] prion phenotypes, or the prions are transmitted in an infectious manner in organisms such as in the case of PrP prions in mammals, leading to transmissible spongiform encephalopathies. This defining characteristic of prions is attributable, at least in part, to the fact that the aggregated prion protein is able to promote the rearrangement of unaggregated protein into the aggregated conformation (although chaperone-type proteins or other trans-acting factors in the cell may also assist with this conformational change). It is likewise a feature of prion-aggregation domains that over-production of proteins comprising these domains increases the frequency with which the prion conformation and phenotype spontaneously arises in cells.

[0062] Prion aggregation amino acid sequences comprising amino terminal sequences derived from yeast or fungal Sup35 proteins, Ure2 proteins, or the carboxy terminal sequences derived from yeast Rnq1 proteins are among those that are highly preferred. Referring to the *S. cerevisiae* Sup35 amino acid sequence set forth in SEQ ID NO: 2, experiments have shown that no more than amino acids 2-113 (the N domain) of that sequence are required to confer some prion aggregation properties to a protein, although inclusion of the charged “M” (middle) region immediately downstream of these residues, e.g., thru residue 253, is preferred in some embodiments. The N domain alone is very amyloidogenic and immediately aggregates into fibers, even in the presence of 2 M urea, a phenomenon that is desirable in embodiments of the invention where formation of stable fibrils of chimeric polypeptides is preferred. When the N domain is fused to the highly charged M domain, fiber formation proceeds in a slower, more orderly way. The M domain is postulated to shift the equilibrium to permit greater “switchability” between aggregated and soluble forms, and is preferably included where phenotypic switching is desirable. Referring to the *S. cerevisiae* Ure2 amino acid sequence set forth in SEQ ID NO: 4, experiments have shown that no more than amino acids 2-65 of that sequence are required to confer prion aggregation activity to a protein. Referring to the *S. cerevisiae* Rnq1 amino acid sequence set forth in SEQ ID NO: 50, experiments have shown that no more than amino acids 153-405 of that sequence are required to confer prion aggregation activity to

a protein. Moreover, sequences differing from the native sequences by the addition, deletion, or substitution of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more amino acids, especially the addition or substitution of additional glutamine or asparagine residues, but which retain the properties of prion-aggregation domains as described in the preceding paragraph, are contemplated. Also, orthologs (corresponding proteins or prion aggregation domains thereof from different species) comprise an additional genus of preferred sequences (Kushinov et al., *Yeast* 6:461-472 (1990); Chernoff et al., *Mol Microbiol* 35:865-876 (2000); Santoso et al., *Cell* 100:277-288 (2000); and Kushinov et al., *EMBO J* 19:324-31 (2000)). By way of example, Sup35 amino acid sequences from *Pichia pinus* and *Candida albicans* are set forth in Genbank Accession Nos. X56910 (SEQ ID NO: 46) and AF 020554 (SEQ ID NO: 47), respectively. Polypeptides of the invention include polypeptides that are encoded by polynucleotides that hybridize under stringent, preferably highly stringent conditions, to the polynucleotide sequences of the invention, or the non-coding strand thereof. Polypeptides of the invention also include polypeptides that are at least about 99%, at least about 98%, at least about 95%, at least about 90%, at least about 85%, at least about 80%, at least about 75%, or at least about 70% identical to one of SCHAG amino acid sequences of the invention.

[0063] As set forth above, in some aspects of the invention, the nucleotide sequence encoding the SCHAG amino acid sequence of the polypeptide is fused in frame with a nucleotide sequence encoding at least one polypeptide of interest. By “in frame” is meant that when the nucleotide is transformed into a host cell, the cell can transcribe and translate the nucleotide sequence into a single polypeptide comprising both the SCHAG amino acid sequence and the at least one polypeptide of interest. It is contemplated that the nucleotide sequences can be joined directly; or that the nucleotide sequences can be separated by additional codons. Such additional codons may encode an endopeptidase recognition sequence or a chemical recognition sequence or the like, to permit enzymatic or chemical cleavage of the SCHAG amino acid sequence from the polypeptide of interest, to permit isolation of the polypeptide of interest. Preferred recognition sequences are sequences that are not found in the polypeptide of interest, so that the polypeptide of interest is not internally cleaved during such isolation procedures. It will be understood that modification of the polypeptide of interest to eliminate internal recognition sequences may be desirable to facilitate subsequent cleavage from the SCHAG amino acid sequence. Suitable enzymatic cleavage sites include: the amino acid sequences -(Asp)_n-Lys-, wherein n signifies 2, 3 or 4, recognized by the protease enterokinase; -Ile-Glu-Gly-Arg-, recognized by coagulation factor X_a; an arginine residue or a lysine residue cleaved by trypsin; a lysine residue cleaved by lysyl endopeptidase; a glutamine residue cleaved by V8 protease, and a glu-asn-leu-tyr-phe-gln-gly site recognized by the tobacco etch virus (TEV) protease. Suitable chemical cleavage sites include tryptophan residues cleaved by 3-bromo-3-methyl-2-(2-nitrophenylmercapto)-3H-indole; cysteine residues cleaved by 2-nitroso-5-thiocyanobenzoic acid; the dipeptides -Asp-Pro- or -Asn-Gly- which can be cleaved by acid and hydroxylamine, respectively; and a methionine residue which is specifically cleaved by cyanogen bromide (CNBr). In another variation, the additional codons

comprise self-splicing intein sequences that can be activated, e.g., by adjustments to pH. See Chong et al., *Gene*, 192:27-281 (1997).

[0064] Additional codons also may be included between the sequence encoding the prion aggregation amino acid sequence and the sequence encoding the protein of interest to provide a linker amino acid sequence that serves to spatially separate the SCHAG amino acid sequence from the polypeptide of interest. Such linkers may facilitate the proper folding of the polypeptide of interest, to assure that it retains a desired biological activity even when the protein as a whole has formed aggregates with other proteins containing the SCHAG amino acid sequence. Also, additional codons may be included simply as a result of cloning techniques, such as ligations and restriction endonuclease digestions, and strategic introduction of restriction endonuclease recognition sequences into the polynucleotide.

[0065] In still another variation, the additional codons comprise a hydrophilic domain, such as the highly-charged M region of yeast Sup35 protein. While the N domain of Sup35 has proven sufficient in some cases to effect prion-like behavior, suggesting that the M region is not absolutely required in all cases, it is contemplated that the M region or a different peptide that includes hydrophilic amino acid side chains will in some cases be helpful for modulating prion-like character of chimeric peptides of the invention. Without intending to be limited to a particular theory, the highly charged M domain is thought to act as a "solubilization" domain involved in modulating the equilibrium between the soluble and the aggregate forms of Sup35, and these properties may be advantageously adapted for other SCHAG sequences.

[0066] By "polypeptide of interest" is meant any polypeptide that is of commercial or practical interest and that comprises an amino acid sequence encodable by the codons of the universal genetic code. Exemplary polypeptides of interest include: enzymes that may have utility in chemical, food-processing (e.g., amylases), or other commercial applications; enzymes having utility in biotechnology applications, including DNA and RNA polymerases, endonucleases, exonucleases, peptidases, and other DNA and protein modifying enzymes; polypeptides that are capable of specifically binding to compositions of interest, such as polypeptides that act as intracellular or cell surface receptors for other polypeptides, for steroids, for carbohydrates, or for other biological molecules; polypeptides that comprise at least one antigen binding domain of an antibody, which are useful for isolating that antibody's antigen; polypeptides that comprise the ligand binding domain of a ligand binding protein (e.g., the ligand binding domain of a cell surface receptor); metal binding proteins (e.g., ferritin (apoferritin), metallothioneins, and other metalloproteins), which are useful for isolating/purifying metals from a solution containing them for metal recovery or for remediation of the solution; light-harvesting proteins (e.g., proteins used in photosynthesis that bind pigments); proteins that can spectrally alter light (e.g., proteins that absorb light at one wavelength and emit light at another wavelength); regulatory proteins, such as transcription factors and translation factors; and polypeptides of therapeutic value, such as chemokines, cytokines, interleukins, growth factors, interferons, antibiotics, immunopotentiators and immunosuppressors, and angiogenic or anti-angiogenic peptides.

[0067] However, specifically excluded from the scope of the invention are chimeric polynucleotides that have hereto-

fore been described in the literature. For example, excluded from the scope of the invention are polynucleotides encoding a fusion consisting essentially of a SCHAG domain of a characterized protein fused in-frame to only: (1) a marker protein such as a fluorescing protein (e.g., green fluorescent protein or firefly luciferase), an antibiotic resistance-confering protein, a protein involved in a nutrient metabolic pathway that has been used in the literature for selective growth on incomplete growth media, or a protein (e.g., β -galactosidase, an alkaline phosphatase, or a horseradish peroxidase) involved in a metabolic or enzymatic pathway of a chromogenic or luminescent substrate that results in the production of a detectable chromophore or light signal that has been used in the literature for identification, selection, or quantitation; or (2) a protein (e.g., glutathione S-transferase or Staphylococcal nuclease) that has been used in the literature as a fusion partner for the express purpose of facilitating expression or purification of other proteins. Notwithstanding this exclusion of certain products from the invention, the inventors contemplate novel uses of such specifically excluded products as aspects of the present invention. Moreover, polynucleotides that include a SCHAG sequence, and sequence encoding a polypeptide of interest, and a sequence encoding a marker protein such as green fluorescent protein are considered within the scope of the invention. Also, notwithstanding the above exclusion, polynucleotides that encode polypeptides whose SCHAG properties are described herein for the first time, fused to a marker protein, are considered within the scope of the invention. Also, purified fusion polypeptides that have been described in the literature and examined only in vivo, but never purified, are intended as aspects of the invention. For example, isolated fibers comprising polypeptides encoding a fusion protein consisting of essentially one or more SCHAG sequences fused to a marker protein, e.g., GFP are contemplated. Several such examples are provided in Example 5.

[0068] The encoding sequences of the polynucleotide may be in either order, i.e., the SCHAG amino acid encoding sequence may be upstream (5') or downstream (3') of the sequence, such that the SCHAG amino acid sequence of the resultant protein is disposed at an amino-terminal or carboxyl-terminal position relative to the protein of interest. In the case of SCHAG amino acid sequences identified or derived from sequences in nature, the encoding sequences preferably are ordered in a manner mimicking the order of the polypeptide from which the SCHAG amino acid sequence was derived. For example, the yeast Sup35 protein has an amino terminal SCHAG domain and a carboxy-terminal domain containing Sup35 translation termination activity. Thus, in embodiments of the invention where the SCHAG amino acid encoding sequence is derived from a Sup35 protein, this sequence preferably is disposed upstream (5') of the sequence encoding the at least one polypeptide of interest. In embodiments wherein the fibril-aggregation amino acid encoding sequence is derived from the sequence set forth in Genbank Accession No. p25367 (SEQ ID NO: 29) (where the prion-like domain is C-terminal), this sequence is preferably disposed downstream (3') of the sequence encoding the at least one polypeptide of interest. In an embodiment comprising sequences encoding two or more polypeptides of interest, the SCHAG encoding sequence may be disposed between the two polypeptides of interest.

[0069] To the extent that such sequences are not already inherent in the above-described polynucleotides, it will be

understood that such polynucleotides preferably further comprise a translation initiation codon fused in frame and upstream (5') of the encoding sequences, and a translation stop codon fused in frame and downstream (3') of the encoding sequences. Also, it may be desirable in some embodiments to direct a host cell to secrete the chimeric polypeptide. Thus, it is contemplated that the polynucleotide may further comprise a nucleotide sequence encoding a translation initiation codon and a secretory signal peptide fused in frame and upstream of the encoding sequences.

[0070] In preferred embodiments, the polynucleotide of the invention further comprises additional sequences to facilitate and/or control expression in selected host cells. For example, the polynucleotide includes a promoter and/or an enhancer sequence operatively connected upstream (5') of the encoding sequences, to promoter expression of the encoding sequences in the selected host cell; and/or a polyadenylation signal sequence operatively connected downstream (3') of the encoding sequences. Since concentration is a factor that may influence the aggregation state of encoded chimeric polypeptides, regulatable (e.g., inducible and repressible) promoters are highly preferred.

[0071] To facilitate identification of cells that have been successfully transformed/transfected with the polynucleotide of the invention, the polynucleotide may further include a sequence encoding a selectable marker protein. The selectable marker may be a completely distinct open reading frame on the polynucleotide, such as an open reading frame encoding an antibiotic resistance protein or a protein that facilitates survival in a selective nutrient medium. The selectable marker also may itself be part of the chimeric polypeptide of the invention. In one embodiment, a visual marker such as a fluorescent protein (e.g., green fluorescent protein) is used that is distributed in the cell in a different manner when the protein is in the prion form than when the protein is in the non-prion form. In either case, cells comprising the selectable marker can be sorted, e.g., using techniques such as fluorescence activated cell sorting. Thus, this marker, in addition to permitting selection of transformed or transfected cells, also permits identification of the conformational state of the chimeric polypeptide. In another embodiment, the marker has two components: 1) a function that is changed when the protein is in a prion form and 2) a visual or selectable marker for that function. An example is the glucocorticoid receptor, GR and a reporter gene. GR is a transcription factor that binds to a specific DNA sequence to activate transcription. When this DNA sequence is fused to the coding sequence for an easily detected protein such as β -galactosidase or luciferase GR function can be easily assayed by the induction of the β -galactosidase or luciferase proteins.

[0072] Optionally, the polynucleotide of the invention further includes an epitope tag fused in frame with the encoding sequences, which tag is useful to facilitate detection in vivo or in vitro and to facilitate purification of the chimeric polypeptide or of the protein of interest after it has been cleaved from the SCHAG amino acid sequence of the chimeric polypeptide. (An epitope tag alone is not considered to constitute a polypeptide of interest). A variety of natural or artificial heterologous epitopes are known in the art, including artificial epitopes such as FLAG, Strep, or poly-histidine peptides. FLAG peptides include the sequence Asp-Tyr-Lys-Asp-Asp-Asp-Lys (SEQ ID NO: 5) or Asp-Tyr-Lys-Asp-Glu-Asp-Asp-Lys (SEQ ID NO: 6). [See generally Brewer, *Bioprocess. Technol.*, 2: 239-266 (1991); Kunz, *J. Biol. Chem.*, 267: 9101-

9106 (1992); Brizzard et al., *Biotechniques* 16: 730-735 (1994); Schafer, *Biochem. Biophys. Res. Commun.*, 207: 708-714 (1995).] The Strep epitope has the sequence Ala-Trp-Arg-His-Pro-Gln-Phe-Gly-Gly (SEQ ID NO: 7). [See Schmidt, *J. Chromatography*, 676: 337-345 (1994).] Another commonly used artificial epitope is a poly-His sequence having six consecutive histidine residues. Commonly used naturally-occurring epitopes include the influenza virus hemagglutinin sequence Tyr-Pro-Tyr-Asp-Val-Pro-Asp-Tyr-Ala-Ile-Glu-Gly-Arg (SEQ ID NO: 8) and truncations thereof, which is recognized by the monoclonal antibody 12CA5 [Murray et al., *Anal. Biochem.*, 229: 170-179 (1995)] and the sequence (Glu-Gln-Lys-Leu-Leu-Ser-Glu-Glu-Asp-Leu-Asn) (SEQ ID NO: 9) from human c-myc, which is recognized by the monoclonal antibody 9E10 (Manstein et al., *Gene*, 162: 129-134 (1995)).

[0073] In another embodiment, the polynucleotide includes 5' and 3' flanking regions that have substantial sequence homology with a region of an organism's genome. Such sequences facilitate introduction of the chimeric gene into the organism's genome by homologous recombination techniques.

[0074] In yet another aspect, the invention provides a polynucleotide comprising a nucleotide sequence that encodes a chimeric polypeptide, the chimeric polypeptide comprising an amyloidogenic domain that causes the polypeptide to aggregate with polypeptides sharing an identical or nearly identical domain into ordered aggregates such as fibrils, fused to a domain comprising a polypeptide of interest; wherein the amyloidogenic domain comprises an amyloidogenic amino acid sequence of a naturally occurring protein and further includes a duplication of at least a portion of the naturally occurring amyloidogenic amino acid sequence, the duplication increasing the amyloidogenic affinity of the chimeric polypeptide relative to an identical chimeric polypeptide lacking the duplication. By way of example, if the naturally occurring protein comprises a Sup35 protein of *Saccharomyces cerevisiae* that is characterized by the partial amino acid sequence PQGGYQQYN (SEQ ID NO: 10), which sequence exists as multiple imperfect repeats, the duplication preferably includes the amino acid sequence PQGGYQQYN and/or an imperfect repeat thereof, such as a repeat wherein one or two residues has been added, deleted, or substituted. An exemplary sequence containing the NM regions of yeast Sup35, with two additional repeat segments, is set forth in SEQ ID NOs: 16 and 17.

[0075] In a related aspect, the invention provides a polynucleotide comprising a nucleotide sequence that encodes a chimeric polypeptide, the chimeric polypeptide comprising an amyloidogenic domain that causes the polypeptide to aggregate with identical polypeptides into fibrils, fused to a domain comprising a polypeptide of interest; wherein the amyloidogenic domain comprises amyloidogenic amino acid sequences of at least two naturally occurring amyloidogenic proteins.

[0076] In yet another related aspect, the invention provides a polynucleotide comprising a nucleotide sequence of the formula FPBT or FBPT, wherein: B comprises a nucleotide sequence encoding a polypeptide that is encoded by a portion of the genome of the cell; F and T comprise, respectively, 5' and 3' flanking sequences adjacent to the sequence encoding B in the genome of the cell; and P comprises a nucleotide sequence encoding a prion-aggregating amino acid sequence, wherein P is fused in frame to B. Using such polynucleotides

and conventional homologous recombination techniques [see, e.g., Ausbel et al. (1998), Volume 3, supra], one can perform homologous recombination in a living cell to convert a protein-encoding gene of the cell to a prion gene of the cell, as described in greater detail below. Alternatively, strains can be constructed wherein the endogenous protein-encoding gene is deleted and a prion version of the gene is added back into the cell, either on a plasmid or by integration into the host genome.

[0077] The homologous recombination technique is itself intended as an aspect of the invention. For example, the invention provides a method of modifying a living cell to create an inducible and stable phenotypic alteration in the cell, comprising the steps of: transforming a living cell with the polynucleotide described in the preceding paragraph; culturing the cell under conditions that permit homologous recombination between the polynucleotide and the genome of the cell; and selecting a cell in which the polynucleotide has homologously recombined with the genome to create a genomic sequence comprising the formula PB or BP.

[0078] More generally, the invention provides a method of modifying a living cell to create an inducible and stable phenotypic alteration in the cell, such as a method comprising steps of: identifying a target polynucleotide sequence in the genome of the cell that encodes a polypeptide of interest; and transforming the cell to substitute for or modify the target sequence, wherein the substitution or modification produces a cell comprising a polynucleotide that encodes a chimeric polypeptide, wherein the chimeric polypeptide comprises a SCHAG amino acid sequence fused in frame with the polypeptide of interest. Such modifications can be performed in several ways, such as (1) homologous recombination as described in the preceding paragraphs; (2) knockout or inactivation of the target sequence followed by introduction of an exogenous chimeric sequence encoding the desired chimeric polypeptide; or (3) targeted introduction of a SCHAG-encoding polynucleotide sequence upstream and in-frame with the target sequence encoding the polypeptide of interest; (4) subsequent cloning or sexual reproduction of such cells; and/or other techniques developed by those in the art.

[0079] The foregoing aspects of the invention relate largely to polynucleotides. Also intended as part of the invention are vectors comprising the polynucleotides, and host cells comprising either the polynucleotides or comprising the vectors. Vectors are useful for amplifying the polynucleotides in host cells. Preferred vectors include expression vectors, which contain appropriate control sequences to permit expression of the encoded chimeric protein in a host cell that has been transformed or transfected with the vectors. Both prokaryotic and eukaryotic host cells are contemplated as aspects of the invention. The host cell may be from the same kingdom (prokaryotic, animal, plant, fungi, protista, etc.) as the organism from which the SCHAG amino acid sequence of the polynucleotide was derived, or from a different kingdom. In a preferred embodiment, the host cell is from the same species as the organism from which the SCHAG amino acid sequence of the polynucleotide was derived.

[0080] In yet another embodiment, the invention includes a host cell transformed or transfected with at least two polynucleotides encoding chimeric polypeptides according to the invention, wherein the at least two polynucleotides comprise compatible SCHAG amino acid sequences and distinct polypeptides of interest. Such host cells are capable of producing two chimeric polypeptides of the invention, which can

be induced *in vitro* or *in vivo* to aggregate with each other into higher ordered aggregates. As explained in greater detail below, such aggregates can be advantageously employed in multi-step chemical reactions when the two or more polypeptides of interest each participate in a step of the reaction. Experiments using fluorescence resonance energy transfer (FRET) have demonstrated the efficacy of heterogeneous polypeptide aggregation into co-polymers.

[0081] In addition, the chimeric polypeptides encoded by any of the foregoing polynucleotides are intended as an aspect of the invention. Purified polypeptides are preferred, and are obtained using conventional polypeptide purification techniques. For example, the invention provides a chimeric polypeptide comprising: at least one SCHAG amino acid sequence and at least one polypeptide of interest other than a marker protein, a glutathione S-transferase (GST) protein, or a Staphylococcal nuclear protein. As described above, the SCHAG amino acid sequence may be directly linked (via a peptide bond) to the polypeptide of interest, or may be indirectly linked by virtue of the inclusion of an intermediate spacer region, a solubility domain, an epitope to facilitate recognition and purification, and so on.

[0082] As explained herein in detail, polypeptides of the invention are capable of existing in a conformation in which the polypeptide coalesces with similar polypeptides into ordered aggregates that may be referred to as “amyloid,” “fibrils,” “prions,” or “prion-like aggregates.” Such ordered aggregates of polypeptides of the invention are intended as an additional aspect of the invention. Such ordered aggregates tend to be insoluble in water or under physiological conditions mimicking a host cell, and consequently can be purified and isolated using standard procedures, including but not limited to centrifugation or filtration. In a preferred embodiment, the SCHAG amino acid sequence is an amino acid sequence that will self-coalesce into ordered “cross- β ” fibril structures that are filamentous in character, in which individual β -sheet strands of component chimeric proteins are oriented perpendicular to the axis of the fibril. In a highly preferred embodiment, the polypeptide of interest is disposed radiating away from the fibril core of SCHAG peptide sequences, and retains one or more characteristic biological activities (e.g., binding activities for polypeptides of interest that have specific binding partners; enzymatic activity for polypeptides of interest that are enzymes).

[0083] In still another embodiment, the invention provides a composition comprising an ordered aggregate of at least two chimeric polypeptides of the invention, wherein the at least two chimeric polypeptides have compatible SCHAG amino acid sequences and distinct polypeptides of interest. By “compatible” SCHAG amino acid sequences is meant SCHAG amino acid sequences that are either identical or sufficiently similar to permit co-aggregation with each other into higher ordered aggregates. In a preferred embodiment, the two or more polypeptides of interest retain their native biological activity (e.g., binding activity; enzymatic activity) in the ordered aggregate. Such aggregates can be advantageously employed in multi-step chemical reactions, as described in detail below.

[0084] The invention further includes methods of making and using polynucleotides and polypeptides of the invention.

[0085] For example, the invention provides a method comprising the steps of: transforming or transfecting a cell with a polynucleotide of the invention; and growing the cell under conditions which result in expression of the chimeric

polypeptide that is encoded by the polynucleotide in the cell. In a preferred embodiment, the method further includes the step of isolating the chimeric polypeptide from the cell or from growth medium of the cell. In one variation, the method further comprises the step of detaching the SCHAG amino acid sequence of the protein from the polypeptide of interest. As described above in detail, the detachment may be effected with any appropriate means, including chemicals, proteolytic enzymes, self-splicing inteins, or the like. Optionally, the method further includes the step of isolating the protein of interest from the SCHAG amino acid sequence.

[0086] In a related embodiment, the invention provides a method of making a protein of interest, comprising the steps of: transforming or transfecting a cell with a polynucleotide, the polynucleotide comprising a nucleotide sequence that encodes a chimeric polypeptide, the chimeric polypeptide comprising an amyloidogenic domain that causes the polypeptide to aggregate with identical polypeptides into higher-ordered aggregates such as fibrils, fused to domain comprising a polypeptide of interest; growing the cell under conditions which result in expression of the chimeric polypeptide in the cell and aggregation of the chimeric polypeptide into fibrils; and isolating the chimeric polypeptide from the cell or from growth medium of the cell. In a preferred embodiment, the isolating step comprises the step of separating the fibrils from soluble proteins of the cell. In a highly preferred embodiment, the method further comprises the steps of proteolytically detaching the amyloidogenic domain of the chimeric protein from the polypeptide of interest; and isolating the polypeptide of interest. Preferably the detached polypeptide of interest maintains one or more of its biological functions, e.g. enzymatic activity, the ability to bind to its ligand, the ability to induce the production of antibodies in a suitable host system, etc.

[0087] In yet another aspect, the invention provides a method of modifying a living cell to create an inducible and stable phenotypic alteration in the cell. For example, such a method comprising the step of transforming or transfecting a living cell with a polynucleotide according to the invention, wherein the polynucleotide includes a promoter sequence to promote expression of the encoded chimeric polypeptide in the cell, the promoter being inducible to promote increased expression of the chimeric polypeptide to a level that induces aggregation of the chimeric polypeptide into higher-ordered aggregates such as fibrils. In one preferred embodiment, the method further comprises the step of growing the cell under conditions which induce the promoter, thereby causing increased expression of the polypeptide and inducing aggregation of the chimeric polypeptide into aggregates or fibrils in the cell. In a highly preferred embodiment, the host cell lacks any native protein that contains the same SCHAG amino acid sequence that might co-aggregate with the chimeric polypeptide. For example, the SCHAG amino acid sequence comprises an amino terminal domain of a Sup35 protein, and the host cell is a yeast cell that comprises a mutant Sup35 gene that expresses a Sup35 protein lacking an amino terminal domain capable of prion aggregation. In such host cells, the chimeric polypeptide can be expressed at a high level and induced to aggregate without concomitant precipitation of the host cell's Sup35 protein into the aggregates, which could be detrimental to host cell viability.

[0088] In yet another aspect, the invention provides methods for reverting the phenotype obtained according to the method described in the preceding paragraph. One such

method comprises the step of overexpressing a chaperone protein in the cell to convert the polypeptide from a fibril-forming conformation into a soluble conformation. In a preferred embodiment, the chaperone protein comprises the Hsp104 protein of yeast, or a related Hsp100-type protein from another species. Examples include the ClpB protein of *E. coli* and the At101 protein of *Arabidopsis*. [See generally Schirmer et al., *Trends in Biochemistry*, 21: 289-296 (1996), incorporated herein by reference.] The over-expression is achieved, e.g., by placing the gene encoding the chaperone protein under the control of an inducible promoter and inducing the promoter.

[0089] Another such method for reverting the phenotype comprises the step of contacting the cell with a chemical denaturant at a concentration effective to convert the polypeptide from a fibril-forming conformation to a soluble conformation. Exemplary denaturants include guanidine HCl (preferably about 0.1 to 100 mM, more preferably 1-10 mM) and urea. In another variation, the cell is subjected to heat or osmotic shock for a period of time effective to convert the polypeptide's conformation. Both over-expression of Hsp104 and growth on guanidine-HCl containing medium have proven effective for inducing phenotypic reversion of chimeric NM-GR prion constructs described in the Examples herein.

[0090] In yet another aspect, the invention provides materials and methods for identifying novel SCHAG amino acid sequences. One such method comprises the steps of joining a candidate nucleotide sequence "X" to a nucleotide sequence encoding the carboxyl terminal domain of a Sup35 protein (CSup35), especially a yeast Sup35 protein, to create a chimeric polynucleotide of the formula 5'-XCSup35-3' or 5'-CSup35X-3'; transforming or transfecting a host cell with the chimeric polynucleotide; growing the host cell under conditions in which the host cell loses its native Sup35 gene, such that the chimeric polynucleotide becomes the only polynucleotide encoding CSup35; growing the resultant host cell under conditions selective for a nonsense suppressive phenotype; and selecting a host cell displaying the nonsense suppressive phenotype, wherein growth in the selective conditions is correlated with the candidate nucleotide sequence X encoding a SCHAG amino acid sequence. Additional methods steps and alternative methods are described in detail below in the Examples. In one variation, the CSup35 is substituted by a different protein domain for which selection on the basis of inactivation is possible.

[0091] Many of the foregoing aspects of the invention relate, at least in part, to embodiments that involve chimeric polynucleotides and polypeptides, wherein properties of SCHAG amino acid sequences are advantageously employed through attaching them to other sequences using recombinant molecular biological techniques. In another variation of the invention, the advantageous properties of SCHAG amino acid sequences are exploited by making SCHAG sequences with sites that are modifiable using organic chemistry or enzymatic techniques.

[0092] For example, in one embodiment, the invention provides a method of making a reactable SCHAG amino acid sequence comprising the steps of identifying a SCHAG amino acid sequence, wherein polypeptides comprising the SCHAG amino acid sequence are capable of forming ordered aggregates; analyzing the SCHAG amino acid sequence to identify at least one amino acid residue in the sequence having a side chain exposed to the environment in an ordered

aggregate of polypeptides that comprise the SCHAG amino acid sequence; and modifying the SCHAG amino acid sequence by substituting an amino acid containing a reactive side chain for the amino acid identified as having a side chain exposed to the environment in an ordered aggregate of polypeptides that comprise the SCHAG amino acid sequence. By "reactive" side chain is meant an amino acid with a charged or polar side chain that can be used as a target for chemical modification using conventional organic chemistry procedures, preferably procedures that can be performed in an environment that will not permanently denature the protein. In preferred embodiments, the amino acid containing a reactive side chain is cysteine, lysine, tyrosine, glutamate, aspartate, and arginine. The identifying step entails any selection of a SCHAG amino acid sequence. For example, the identifying can simply entail selecting one of the SCHAG amino acid sequences described in detail herein; or can entail screening of genomes, proteins, or phenotypes of organisms to identify SCHAG sequences (e.g., using methodologies described herein); or can entail de novo design of SCHAG sequences based on the properties described herein.

[0093] Proteins comprising the SCHAG sequence are capable of coalescing into higher-ordered aggregates. The polypeptides of such aggregates have amino acids that are disposed internally (in close proximity only to other amino acids in the aggregate), and other amino acids whose side chains are exposed to the environment of the aggregate such that they contact molecules in the environment. In the method, the analyzing step entails a prediction or a determination of at least one amino acid within the SCHAG sequence that is exposed to the environment of an aggregate of the proteins, meaning that it is an amino acid that will likely contact chemical reagents that mixed with the aggregates.

[0094] Amino acids in a SCHAG amino acid sequence having side chains exposed to the environment in ordered aggregates of polypeptides comprising the SCHAG amino acid sequence can be identified experimentally, for example, by structural analysis of mutants constructed using site-directed mutagenesis, e.g., high throughput cysteine scanning mutagenesis, as described in detail below in the Examples. Alternatively, specific amino acids in a SCHAG amino acid sequence can be predicted to have side chains that are exposed to the environment in ordered aggregates of polypeptides comprising the SCHAG amino acid sequence based on structural studies or computer modeling of the SCHAG amino acid sequence. The step of modifying the amino acid sequence entails changing the identity of an amino acid within the sequence. For the purposes of such a method, the act of inserting a reactive amino acid within the amino acid sequence, at a position essentially adjacent to the position of the identified amino acid, is considered the equivalent of substituting that amino acid for the identified amino acid. In other words, for the purposes of making a reactable SCHAG amino acid sequence, the term "substituting" should be understood to include inserting an amino acid within the amino acid sequence, at a position essentially adjacent to the position of the identified amino acid.

[0095] It is contemplated that some naturally-occurring SCHAG amino acid sequences will fortuitously include one or more reactive amino acids whose side chains are exposed to the environment in polypeptide aggregates. Use of such naturally occurring SCHAG reactive amino acids is contemplated as an additional aspect of the invention. Moreover, modification of naturally occurring SCHAG amino acid

sequences that contain an undesirable number of reactive amino acids to eliminate one or more reactive amino acids is contemplated.

[0096] In a preferred embodiment, the method further comprises a step of making a polypeptide comprising the reactable SCHAG amino acid sequence. Substitution of such amino acids with amino acid residues containing reactive side chains can be carried out in the laboratory by, e.g., site-directed mutagenesis of a SCHAG-encoding polynucleotide or by peptide synthesis of the SCHAG amino acid sequence. In another preferred embodiment, the invention additionally comprises the step of making a polymer comprising an ordered aggregate of polypeptide monomers wherein at least one of the polypeptide monomers comprises a reactable SCHAG amino acid sequence. For example, polypeptide monomers comprising the reactable SCHAG amino acid sequence are seeded with an aggregate or otherwise subjected to an environment favorable to the formation of an ordered aggregate or "polymer" of the polypeptide monomers. In yet another preferred embodiment, the invention further comprises the step of contacting the reactive side chains with a chemical agent to attach a substituent to the reactive side chains. The substituent itself may be a linker molecule to facilitate attachment of one or more additional molecules. The substituent may be attached using a chemical agent. Attachment of a substituent depends on the nature of the substituent, as well as the identity of the reactive side chain, and can be accomplished by conventional organic chemistry procedures. Exemplary procedures for modifying the sulfhydryl group of a cysteine residue that has been introduced into a SCHAG amino acid sequence are described in greater detail below in the Examples. In preferred embodiments, the substituent is an enzyme, a metal atom, an affinity binding molecule having a specific affinity binding partner, a carbohydrate, a fluorescent dye, a chromatic dye, an antibody, a growth factor, a hormone, a cell adhesion molecule, a toxin, a detoxicant, a catalyst, or a light-harvesting or light altering substituent. In a preferred embodiment, the reactive amino acid that has been introduced into the SCHAG sequence will be substantially absent from the rest or the SCHAG amino acid sequence, or at least substantially absent from those portions of the sequence that are exposed to the environment in ordered aggregates of the polypeptide. This absence may be a natural feature, or may be the result of an additional modification step to substitute or delete other occurrences of the amino acid. Designing the reactable SCHAG amino acid sequence in this manner permits controlled chemical modification at the reactive sites that have been designed into the sequence, without modification of other residues.

[0097] In yet another embodiment of the invention, the invention further comprises the steps of contacting the polypeptides comprising the reactive side chains with a chemical agent to attach a substituent to the reactive side chains, thereby providing modified polypeptides, and making a polymer comprising an ordered aggregate of polypeptide monomers, wherein at least some of the polypeptide monomers comprise the modified polypeptides. Exemplary procedures for making a polymer comprising an ordered aggregate of modified polypeptide monomers are described in greater detail below in the Examples.

[0098] In yet another embodiment, the invention provides a method of making a reactable SCHAG amino acid sequence, wherein the SCHAG amino acid sequence is modified to contain exactly one, two, three, four, or some other speci-

cally desired number of the reactive amino acids. An exemplary method comprises the steps of (a) identifying a SCHAG amino acid sequence, wherein polypeptides comprising the SCHAG amino acid sequence are capable of forming ordered aggregates; (b) analyzing the SCHAG amino acid sequence to identify at least one amino acid residue in the sequence having a side chain exposed to the environment in an ordered aggregate of polypeptides that comprise the SCHAG amino acid sequence; (c) modifying the SCHAG amino acid sequence by substituting an amino acid containing a reactive side chain for the amino acid identified as having a side chain exposed to the environment in an ordered aggregate of polypeptides that comprise the SCHAG amino acid sequence; (d) analyzing the SCHAG amino acid sequence to identify at least a second amino acid residue in the sequence having an amino acid side chain that is exposed to the environment in an ordered aggregate of polypeptides that comprise the SCHAG amino acid sequence; and (e) modifying the SCHAG amino acid sequence by substituting an amino acid containing a reactive side chain for at least one amino acid identified according to step (d), wherein the amino acid substituted in steps (c) and (d) differ, thereby making a reactable SCHAG amino acid sequence with at least two selectively reactable sites. This method can be further elaborated to create SCHAG amino acids sequences with more than two selectively reactable sites. By introducing two or more different reactive amino acids, a SCHAG sequence is created with two or more sites that can be separately reacted/modified. It will be appreciated that the method also can be performed to introduce the same reactive amino acid for each identified amino acid, to create two or more identical reactive sites in the SCHAG sequence.

[0099] In another embodiment of the invention, the invention provides polypeptides comprising a SCHAG amino acid sequence that has been modified by substituting at least one amino acid that is exposed to the environment in an ordered aggregate of the polypeptides with an amino acid containing a reactive side chain, as well as polynucleotides that encode the polypeptides. In a further embodiment, a substituent is attached to the reactive amino acid of the modified polypeptide of the invention or reactable SCHAG sequence. In a highly preferred embodiment, the SCHAG amino acid sequence is modified to contain exactly one, two, three, four, or some other specifically desired number of the reactive amino acids, thereby providing a SCHAG amino acid sequence which is modifiable at controlled, stoichiometric levels and positions. To achieve this goal, modifications to remove undesirable, native reactive amino acids from a naturally occurring SCHAG sequence are contemplated. Polypeptides comprising a naturally occurring SCHAG amino acid sequence characterized by one or more reactive amino acids, that have been modified by substituting or eliminating a natural reactive amino acid, are considered a further aspect of the invention, as are polynucleotides that encode the polypeptides.

[0100] In still another variation, the invention provides various living cells with two or more customized, reversible phenotypes. For example, the invention provides a living cell that comprises: (a) a first polynucleotide comprising a nucleotide sequence encoding a polypeptide that comprises a prion aggregation domain and a domain having transcription or translation modulating activity, wherein the living cell is capable of existing in a first stable phenotypic state characterized by the polypeptide existing in an unaggregated state

and exerting a transcription or translation modulating activity and a second phenotypic state characterized by the polypeptide existing in an aggregated state and exerting altered transcription or translation modulating activity; and (b) an exogenous polynucleotide comprising a nucleotide sequence that encodes a polypeptide of interest, with the proviso that the sequence encoding the polypeptide of interest includes a regulatory sequence causing differential expression of the polypeptide in the first phenotypic state compared to the second phenotypic state. Exemplary prion aggregation domains are described with respect to Sup35, Rnq1, and Ure2. The first polynucleotide may itself be an endogenous (native) polynucleotide of the cell, such as the native yeast Sup35 sequence in a yeast cell, which comprises a prion aggregation domain fused to a translation termination factor sequence. Alternatively, the first polynucleotide may be introduced into the cell (or a parent cell) using genetic engineering techniques. The term "exogenous polynucleotide" is meant to encompass any polynucleotide sequence that differs from a naturally occurring sequence in the cell as a result of human genetic manipulation. For example, an exogenous sequence may constitute an expression construct that has been introduced into a cell, such as a construct that contains a promoter, a foreign polypeptide-encoding sequence, a stop codon, and a polyadenylation signal sequence. Alternatively, an exogenous sequence may constitute an endogenous polypeptide-encoding sequence that has been modified only by the introduction of a promoter, an enhancer, or other regulatory sequence that is not naturally associated with the polypeptide-encoding sequence. Introduction of a regulatory sequence that is influenced by the aggregation state of the polypeptide encoded by the first polynucleotide is specifically contemplated. In one preferred variation, the cell further comprises a nucleotide sequence that encodes a polypeptide that modulates the expression level or conformational state of the polypeptide that comprises the prion aggregation domain. Such a polynucleotide facilitates manipulation of the cell to switch phenotypes. Polynucleotides encoding chaperone proteins that influence prion protein folding represent one example of this latter category of polynucleotide. In one specific variation, the invention provides a living cell according to claim 97, wherein the first polynucleotide comprises a nucleotide sequence encoding a polypeptide that comprises a prion aggregation domain fused in-frame to a nucleotide sequence encoding a translation termination factor polypeptide; and wherein the regulatory sequence comprises a stop codon that interrupts translation of the polypeptide of interest.

[0101] In another variation, the invention provides a living cell comprising: (a) a polynucleotide comprising a nucleotide sequence encoding a polypeptide that comprises a prion aggregation domain fused in-frame to a nucleotide sequence encoding a translation termination factor polypeptide; and (b) an exogenous polynucleotide comprising a nucleotide sequence that encodes a polypeptide of interest, with the proviso that the sequence encoding the polypeptide of interest includes at least one stop codon that interrupts translation of the polypeptide of interest; wherein the living cell is capable of existing in a first stable phenotypic state characterized by translational fidelity and substantial absence of synthesis of the polypeptide of interest and a second phenotypic state characterized by aggregation of the translation termination factor, reduced translational fidelity, and expression of the polypeptide of interest.

[0102] The invention also provides polymers or fibers of ordered aggregates comprising polypeptide subunits wherein at least one of the polypeptide subunits comprises a reactable SCHAG amino acid sequence. By the term “fibril” or “fiber” is meant a filamentous structure composed of higher ordered aggregates. By “polymer” is meant a highly ordered aggregate that may or may not be filamentous. In another embodiment, the polymer or fiber is modified or substituted by attaching a substituent to the reactable SCHAG amino acid sequence of the polypeptide subunits. Also contemplated are polymers or fibers that comprise more than one type of substituent by attachment of different substituents to the reactable SCHAG amino acid sequence of the polypeptide subunits of the polymer or fiber. Attachment of the substituents to the reactive side chains contained in the reactable SCHAG amino acid sequence can occur either before or after coalescing of the polypeptides comprising the reactable SCHAG amino acid sequences into polymers comprising ordered aggregates of the polypeptides. Modification by attachment of specific substituents to such polymers or fibers can confer distinct functions to these molecules. Thus, polymers or fibers, wherein one or more discrete regions of the polymer or fiber are modified to enable a distinct function are contemplated. In another variation, different regions of a polymer or fiber are differentially modified to confer different functions. Also contemplated are polymers or fibers containing patterns of attachments, and consequently patterns of functionalities. The invention also provides polymers comprising fibers wherein at least one fiber has a distinct function different from that of another fiber in the polymer. Fibers comprising polypeptides subunits that are capable of emitting light or altering the wavelength of the light emitted in response to binding of a ligand to the fiber can be used as highly sensitive biosensors. Polymers comprising fibers wherein some of the fibers comprise polypeptide subunits capable of absorbing light of one wavelength and emitting light of second wavelength, and other fibers comprising polypeptide subunits capable of absorbing the light emitted by the first set of fibers and emitting light of a different wavelength are also contemplated.

[0103] In one preferred embodiment, the polymer or fiber is long and thin and contains no or few branches, except at positions defined by deliberate introduction of sites for interaction between the polypeptide subunits. Polymers or fibers in which the polypeptide subunits have been modified to enable directed interactions between the polypeptide subunits within a single polymer or fiber, or between two discrete polymers or fibers are contemplated. Polymers of fibers that have been modified to enable interactions to occur between separate polymers or fibers can be used to create a meshwork of polymers or fibers. In one variation, the meshwork can be generated reversibly by using interactions dependent on sulfhydryl groups present on the polypeptide subunits of the polymer or fiber. Such meshworks can be useful, for example, for filtration purposes. In another preferred embodiment, a fibril, ordered aggregate, polymer or fiber is attached to a solid support. For example, binding of a polymer or fiber to a solid support can be mediated by biotin-avidin interactions, wherein the biotin is attached to the polymers or fibers and avidin is bound to the solid support or vice versa.

[0104] In a related embodiment, the invention provides a method of making a polymer or fiber with a predetermined quantity of reactive sites for chemically modifying the polymer or fiber, comprising the steps of providing a first polypep-

ptide comprising a first SCHAG amino acid sequence that is capable of forming ordered aggregates with polypeptides identical to the first polypeptide; providing a second polypeptide comprising a second SCHAG amino acid sequence that is capable of forming ordered aggregates with polypeptides identical to the first polypeptide or the second polypeptide, wherein the second SCHAG amino acid sequence includes at least one amino acid residue having a reactive amino acid side chain that is exposed to the environment and serves as a reactive site in ordered aggregates of the second polypeptide and; mixing the first and second polypeptides under conditions favorable to aggregation of the polypeptides into ordered aggregates, wherein the polypeptides are mixed in quantities or ratios selected to provide a predetermined quantity of second polypeptide reactive sites. In a preferred embodiment, the invention further comprises the step of reacting the reactive side chains to attach a substituent to the reactive amino acid side chains of the polymer or fiber. Alternatively, the step of reacting the reactive side chains to attach a substituent to the reactive amino acid side chains is performed prior to mixing of the polypeptides comprising reactable SCHAG amino acid sequences to form ordered aggregates. In yet another embodiment, the invention provides a method of making a polymer or fiber comprising a first polypeptide comprising a first SCHAG amino acid sequence and a second polypeptide comprising a second SCHAG amino acid sequence, wherein both the first and second SCHAG amino acid sequence includes at least one amino acid residue having a reactive amino acid side chain that is exposed to the environment and serves as a reactive site, and wherein the reactive amino acid side chains of the first and second SCHAG amino acid sequences that are exposed to the environment in ordered aggregates are not identical, thereby permitting selective reaction of the reactive amino acid side chain of the first SCHAG amino acid sequence without reacting the reactive amino acid side chain of the second SCHAG amino acid sequence.

[0105] In another embodiment, the invention provides a method of making a polymer comprising two or more regions with distinct function comprising the steps of (a) providing a first polypeptide comprising a SCHAG amino acid sequence and a first functional domain and a second polypeptide comprising a SCHAG amino acid domain and a second functional domain that differs from the first functional domain, wherein the SCHAG amino acid sequences of the polypeptides are capable of forming ordered aggregates with polypeptides identical to the first or second polypeptide; (b) aggregating the first polypeptide by subjecting a composition comprising the first polypeptide to conditions favorable to aggregation of the first polypeptide into ordered aggregates, thereby forming a polymer comprising a region containing polypeptides that include the first functional domain; and (c) mixing a composition comprising the second polypeptide with the polymer formed according to step (b), under conditions favorable to aggregation of the second polypeptide with the polymer of step (b), thereby forming a polymer comprising the first region containing polypeptides that include the first functional domain and a second region containing polypeptides that include the second functional domain.

[0106] In one preferred embodiment, the SCHAG amino acid sequences of the first and second polypeptides are identical. In another preferred embodiment, at least one of the first and second functional domains comprises an amino acid that comprises a reactive amino acid side chain. In yet another

preferred embodiment, at least one of the first and second functional domains comprises an amino acid sequence of a polypeptide of interest. In another variation, the method further comprises the step of mixing a composition comprising the first polypeptide with the polymer formed according to step (c), under conditions favorable to aggregation of the first polypeptide with the polymer of step (c), thereby forming a polymer comprising the first region containing polypeptides that include the first functional domain, the second region containing polypeptides that include the second functional domain, and a third region containing polypeptides that include the first functional domain. Alternatively, the invention provides a method of making a polymer comprising two or more regions with distinct function wherein the method further comprises the steps of providing a third polypeptide that comprises a SCHAG amino acid sequence and a third functional domain that differs from the first and second functional domains, wherein the SCHAG amino acid sequence of the third polypeptide is capable of forming ordered aggregates with polypeptides identical to the first polypeptide or the second polypeptide; and mixing a composition comprising the third polypeptide with the polymer formed according to step (c), under conditions favorable to aggregation of the third polypeptide with the polymer of step (c), thereby forming a polymer comprising the first region containing polypeptides that include the first functional domain, the second region containing polypeptides that include the second functional domain, and a third region containing polypeptides that include the third functional domain.

[0107] Additional features and variations of the invention will be apparent to those skilled in the art from the entirety of this application, including the drawing and detailed description, and all such features are intended as aspects of the invention. Likewise, features of the invention described herein can be re-combined into additional embodiments that also are intended as aspects of the invention, irrespective of whether the combination of features is specifically mentioned above as an aspect or embodiment of the invention. Also, only such limitations which are described herein as critical to the invention should be viewed as such; variations of the invention lacking limitations which have not been described herein as critical are intended as aspects of the invention.

[0108] In addition to the foregoing, the invention includes, as an additional aspect, all embodiments of the invention narrower in scope in any way than the variations specifically mentioned above. Although the applicant(s) invented the full scope of the claims appended hereto, the claims appended hereto are not intended to encompass within their scope the prior art work of others. Therefore, in the event that statutory prior art within the scope of a claim is brought to the attention of the applicants by a Patent Office or other entity or individual, the applicant(s) reserve the right to exercise amendment rights under applicable patent laws to redefine the subject matter of such a claim to specifically exclude such statutory prior art or obvious variations of statutory prior art from the scope of such a claim. Variations of the invention defined by such amended claims also are intended as aspects of the invention.

BRIEF DESCRIPTION OF THE DRAWING

[0109] FIG. 1 depicts the DNA and deduced amino acid sequences (SEQ ID NOS: 50-51) of an NMSup35-GR chimeric gene described in Example 1.

[0110] FIG. 2 depicts a map of an integration plasmid described in Example 2 which contains a chimeric gene comprising the amino-terminal domain of yeast Ure2 protein, a hemagglutinin tag sequence, and the carboxyl-terminal domain of yeast Sup35 protein.

[0111] FIG. 3 depicts the nucleotide sequence (SEQ ID NO: 49) of the plasmid of FIG. 2. As shown in FIG. 2, the NURE2-CSUP35 chimeric gene is encoded on the strand complementary to the strand whose sequence is depicted in FIG. 3.

[0112] FIG. 4 schematically depicts that the structure of wild-type (WT) yeast Sup35 protein (Top), which contains an amino-terminal region characterized by five imperfect short repeats, a highly charged middle (M) region, and a carboxyl-terminal region involved in translation termination during protein synthesis; a Sup35 mutant designated RA2-5, characterized by deletion of four of the repeat sequences in the N region; and a Sup35 mutant designated R2E2 (bottom), into which two additional copies of the second repeat segment have been engineered into the N region. Also depicted is the frequency with which yeast strains carrying these various Sup35 constructs were observed to spontaneously convert from a [psi⁻] to a [PSI⁺] phenotype.

[0113] FIG. 5 depicts gold and silver enhancement of NM fibers. Long NM^{K184C} fibrils were assembled by seeding soluble NM^{K184C} with short NM^{K184C} fibrils. Monomaleimido Nanogold was covalently cross-linked (2) and the 1.4-nm Nanogold particles were subjected to gold toning (3-4). Fibrils are labeled as 1; nanogold particles are labeled as 2; silver particles are labeled as 3; and gold particles are labeled as 4.

[0114] FIG. 6 depicts gold toning is specific to labeled fibers. The resulting gold-toned fibers show a significant increase in height from 9-11 nm (bare fibers, labeled as 1) to 80-200 nm (labeled fibers, labeled as 2), imaged by AFM.

[0115] FIG. 7 depicts gold nanowires that did not bridge the gap when randomly deposited on patterned electrodes and imaged by TEM.

[0116] FIG. 8 shows depicts gold nanowires bridging the gap between two electrodes.

[0117] FIG. 9 depicts vaporization of some conducting nanowires after increasing the voltage. Conductive nanowires are labeled as 1, while vaporized nanowires are labeled as 2.

[0118] FIG. 10 schematically depicts an electrical circuit. A power source (i.e., electrical source) is labeled as 1; electrical conductors are labeled as 2; and circuit elements are labeled as 3.

DETAILED DESCRIPTION OF THE INVENTION

[0119] The invention described herein is related to the invention described in U.S. patent application Ser. No. 09/591,632, filed Jun. 9, 2000, which claims priority benefit of U.S. Provisional Application No. 60/138,833, filed Jun. 9, 1999. Both of these applications are incorporated herein by reference.

[0120] The present invention expands the study of prion biology beyond the contexts where it has heretofore focused, namely fundamental research directed to developing a greater understanding of prion biology and medical research directed to developing diagnostic and therapeutic materials and methods for prion-associated disease states, and provides diverse and practical applications that advantageously employ certain unique properties of prions, including one or more of the following:

[0121] (1) prion genes and proteins afford the possibility of two stable, heritable phenotypes and the ability to effect at least one switch between such phenotypes;

[0122] (2) prions provide the ability to sequester a protein or protein-binding molecule into an ordered aggregate;

[0123] (3) prion protein aggregates are easily isolated from cells containing them; with at least some prions, the ordered aggregate is fibrillar in structure, stable and unreactive, a collection of properties that is exploited in certain embodiments of the invention;

[0124] (4) a protein of interest that is fused to a prion protein can potentially retain its normal biological activity even when the fusion has formed an ordered prion aggregate;

[0125] (5) a protein of interest that is fused to a prion protein can switch from an active to an inactive state, and this change is reversible;

[0126] (6) prion protein aggregates form fibrils with unusually high chemical and thermal stability for biological material; and.

[0127] (7) prion protein aggregates form fibrils that can be modified to incorporate specific functionalities, thereby combining the advantages of biomolecules with, for example, electronic circuitry.

[0128] Prion proteins have been observed to exist in at least two stable conformations in cells that synthesize them. For example, the PrP protein in mammals has been observed in a soluble PrP^C conformation in “normal” cells and in an aggregated, insoluble PrP^{Sc} conformation in animals afflicted with transmissible spongiform encephalopathies. Similarly, the Sup35 protein in yeast has been observed in a “normal” non-aggregated conformation in which it forms a component of a translation termination factor, and also aggregated into fibril structures in [PSI⁺] yeast cells (characterized by suppression of normal translation termination activity). To the extent that scientific literature has ascribed any practical importance to these observations, the importance has focused on identifying materials and methods to modulate conformational switching, which might lead to treatments for prion-mediated diseases; or to detect the infectious PrP^{Sc} form to protect the food supply; or to diagnose infection and prevent its spread. At least in the case of the yeast Sup35 prion, the [PSI⁺] phenotype can be eliminated by effecting an over-expression or under-expression of the heat shock protein Hsp104, and can be induced by effecting an over-expression of Sup35 or the Sup35 amino-terminal prion-aggregation domain.

[0129] The practical applications that arise from the ability to alter the phenotype of a cells or an entire organism by transforming/transfecting cells with a polynucleotide that encodes a non-native protein (and/or that integrates into the cell's genome to cause production of a non-native protein) are legion and underlie a major portion of the entire biotechnology industry. Such applications include medical/therapeutic applications (e.g., gene therapy to treat genetic disorders such as hemophilia; gene therapy to treat pathological conditions such as ischemia, inborn errors of metabolism, restenosis, or cancer); pharmacological applications (e.g., recombinant production of therapeutic polypeptides such as erythropoietin, human growth hormone, angiogenic and anti-angiogenic peptides, or cytokines for therapeutic administration); industrial applications (e.g., genetic engineering of microorganisms for bioremediation or frost prevention; or recombinant production of catalytic enzymes, vitamins, proteins, or other organic molecules for use in chemical and food pro-

cessing); and agricultural applications (e.g., genetic engineering of plants and livestock to promote disease resistance, faster growth, better nutritional value, environmental durability, and other desirable properties); just to name a few. In such biotechnology applications, a cell typically is transformed/transfected with a single novel gene to introduce a single phenotypic alteration that persists as long as the gene is present. Means of controlling the new phenotype conventionally involve eliminating the new gene, or possibly placing the gene under the control of inducible or repressible promoter to control the level of gene expression. The present invention provides the realization that prion genes and proteins afford an additional, alternative means of biological control, because the introduction of a prion sequence into a protein introduces the possibility of two stable, heritable phenotypes and the ability to effect at least one switch between such phenotypes. Specifically, one can phenotypically alter a cell to produce a protein of interest by transforming/transfecting a cell with a gene encoding a prion-aggregation domain fused to a protein of interest. To reduce or eliminate the activity of this protein, one induces the protein to undergo a conformational alteration and adopt a prion-like aggregating phenotype, thereby sequestering the protein. To re-introduce the original recombinant phenotype, one induces the protein to undergo a conformational alteration and adopt the soluble phenotype.

[0130] By way of example, the phenotypic alteration potential of prion-like proteins can be harnessed to permit a species (plant, animal, microorganisms, fungi, etc.) to survive in a wider range of environmental conditions and/or quickly adapt to environmental changes. Species that thrive in one environment often have difficulty in another. For example, some photosynthetic organisms grow well under bright light because they produce pigments that protect the organism from potentially toxic effects of bright light, whereas others grow well under low light conditions because of other light-gathering pigment systems that efficiently harvest all available light. By placing the regulators for such systems under a prion control mechanism, prion conformational switching is advantageously harnessed for increased environmental adaptability.

[0131] A preferred prion system for harnessing environmental adaptation is a prion system such as the Sup35 or Ure2 yeast prions that undergo natural switching. In these systems, the yeast prion state and phenotype arises naturally (in a non-prion population) at a frequency of about one per million cells, and is lost at a similar frequency in a prion population. Thus, in any yeast culture of reasonable size, both phenotypes will be present. If the prion state imparts a growth advantage under some conditions and the non-prion state imparts a growth advantage under other conditions, the culture as a whole will survive and thrive under either set of conditions. Although one phenotype may be disfavored and selected against, it will nonetheless be present (due to natural switching behavior of the prion) and ready to “take over” the culture if conditions change to favor it. In this regard, also contemplated as an aspect of the invention is a cell culture comprising cells transformed or transfected with a polynucleotide according to the invention, wherein the cells express the chimeric polypeptide encoded by the polynucleotide, and wherein the cell culture includes cells wherein the chimeric polypeptide is present in an aggregated state and cells free of aggregated chimeric polypeptide.

[0132] The prion-mediated flexibility described in the preceding paragraph possesses a crucial advantage over traditional “switches” because it does not depend upon fortuitous genetic mutations and reversions. Each phenotype arises from the same genotype and each is available within the population, even under selective conditions. Thus, in a cultured photosynthetic organism as described above, transformation with one or more genes encoding an aggregating domain fused to pigment or protective proteins will provide an increased adaptability to varying light conditions.

[0133] This “natural switching” quality of prions has applicability to a wide variety of variable growth conditions that might be encountered by cultured cells or organisms, including varied levels of salinity, metals, carbon sources, and toxic metabolic byproducts. Adaptability to such environments is often mediated by one or a few proteins, such as metal-binding proteins and enzymes involved in the synthesis or breakdown of particular organic compounds. The advantages of prion natural switching are considered particularly well suited for fields of bioremediation, where multiple environmental conditions are expected to be encountered, and fermentation processes where nutrients are consumed and fermentation by products are created, changing an environment over time.

[0134] By way of another example, pigment genes for flowers, textile fibers (e.g., cotton), or animal fibers (e.g., wool) are placed under the control of prion-like aggregating elements. A plurality of colors and/or color patterns is achieved in a single plant by altering growing conditions to induce or cure the prion regulated pigment, or by subjecting portions of the plant to chemical agents that modulate conformation of the prion protein.

[0135] The present invention also provides practical applications stemming from the realization that prions provide the ability to sequester a protein of interest or the protein’s binding partner into an ordered aggregate. This property is demonstrated herein by way of example involving the prion aggregation domain of the yeast Sup35 gene fused to a glucocorticoid receptor. When cells expressing this fusion are in a non-prion phenotype (i.e., the fusion protein is soluble), the cells are susceptible to hormonal induction through the glucocorticoid receptor, and one can induce the expression of a second gene that is operably fused to a glucocorticoid response element. However, when cells expressing the fusion are in a prion phenotype (i.e., the fusion protein is forming aggregates), the susceptibility to hormonal induction is reduced, because the glucocorticoid receptor that is sequestered into cytoplasmic aggregates is unable to effect its normal activity in the cell’s nucleus.

[0136] This ability to a sequester protein or protein-binding partner has direct application in the recombinant production of biological molecules, especially where recombinant production is difficult using conventional techniques, e.g., because the molecule of interest appears to exert a toxic or growth-altering effect on the recombinant host cell. Such effects can be reduced, and production of the polypeptide of interest enhanced, by expressing the polypeptide of interest as fusion with a prion aggregation domain in a host cell that has, or is induced to have, a prion aggregation phenotype. In such host cells, the recombinant fusion protein forms ordered aggregates through its prion aggregation domain, thereby sequestering the protein of interest as part of the aggregate, and reducing its adverse effects on other cellular components or reactions. (If the molecule of interest is the binding partner

of the non-prion domain of the fusion protein, the binding partner also will be sequestered by the aggregate, provided that the binding activity of this domain is retained in the aggregate).

[0137] The present inventors also provide practical applications stemming from the fact that prion aggregates can be readily isolated from cells containing them. Because prions form insoluble aggregates in appropriate host cells, it is relatively easy to separate aggregated prion protein from most other proteinaceous and non-proteinaceous matter of a host cell, which is comparatively more soluble, using centrifugation techniques. When the prion protein is fused to a protein of interest, the protein of interest can likewise be separated from most other host cell impurities by centrifugation techniques. Thus, the present invention provides materials and methods useful for the purification of virtually any recombinant protein of interest. If a recognition sequence for chemical or enzymatic cleavage is included between the prion aggregation domain and the protein of interest, the protein of interest can be cleaved and separated from the insoluble prion aggregate in a second purification step. Such protein production techniques are considered an aspect of the invention. For example, the invention provides a method comprising the steps of: expressing a chimeric gene in a host cell, the chimeric gene comprising a nucleotide sequence encoding a SCHAG amino acid sequence fused in frame to a nucleotide sequence encoding a protein of interest; subjecting the host cell, or a lysate thereof, or a growth medium thereof to conditions wherein the chimeric protein encoded by the chimeric gene aggregates; and isolating the aggregates. In one variation, the method further includes the step of cleaving the protein of interest from the SCHAG amino acid sequence and isolating the protein of interest.

[0138] Moreover, the improved purification techniques are not limited to proteins fused to a prion domain. For example, a host cell expressing a prion aggregation domain fused to a protein of interest can be used in a like manner to purify a binding partner of the protein of interest. For example, if the protein of interest is a growth factor receptor, it can be used to sequester the growth factor itself by virtue of the receptor’s affinity for the growth factor. In this way, the growth factor can be similarly purified, even though it is not itself expressed as a prion fusion protein. If the protein of interest comprises an antigen binding domain of an antibody, then the same techniques can be used to sequester and purify virtually any antigen (protein or non-protein) that is produced by the host cell or introduced into the host cell’s environment. In this regard, it is well-known in the literature that relatively short variable (V) regions within antibodies are largely responsible for highly specific antigen-antibody immunoreactivity, and such antigen-binding regions occur within particular regions of an antibody’s primary structure and are susceptible to isolation and cloning. (See, e.g., Morrison and Oi, *Adv. Immunol.*, 44:65-92 (1989). For example, the variable domains of antibodies may be cloned from the genomic DNA of a B-cell hybridoma or from cDNA generated from mRNA isolated from a hybridoma of interest. Likewise, it is known in the art how to isolate only those portions of the variable region gene fragments that encode antigen-binding complementarity determining regions (“CDR”) of an antibody, and clone them into a different polypeptide backbone. [See, e.g., Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeyen et al., *Science*, 239:1534-36 (1988); and Tempest et al., *Bio/Technology*, 9:266-71

(1991).] A polypeptide comprising an antigen binding domain of an antibody of interest might comprise only one or more CDR regions from an antibody, or one or more V regions from an antibody, or might comprise entire V region fragments linked to constant domains from the same or a different antibody, or might comprise V regions that have been cloned into a larger, non-antibody polypeptide in a way that preserves their antigen binding characteristics, or might comprise antibody fragments containing V regions, and so on. Also, it is known in the art to select and isolate polypeptides comprising antigen binding domains of antibodies using techniques such as phage display that obviate the need to immunize animals and work with native antibodies at all.

[0139] The present invention also provides practical applications stemming from the fact that at least some proteins of interest will retain their normal biological activity when expressed as a fusion with a prion aggregation domain, even when the fusion protein forms prion-like aggregates. This feature of the invention is demonstrated by way of example below using the *S. cerevisiae* Sup35 prion aggregation domain fused to a green fluorescent protein (GFP). Even in [PSI⁺] cells or in other cells where aggregation of the fusion protein into fibrils has occurred, the GFP fluoresces green under blue light, indicating that the GFP portion of the fusion has retained a biologically active conformation.

[0140] When the example is repeated substituting a protein of interest for the GFP marker protein, ordered aggregates comprising a biologically active protein of interest are produced. In a preferred embodiment, the protein of interest is a protein that is capable of binding a composition of interest. For example, the protein of interest comprises an antigen binding domain of an antibody that specifically binds an antigen of interest; or it comprises a ligand binding domain of a receptor that binds a ligand of interest. Fibrils comprising such fusion proteins can be used as affinity matrices for purifying the composition of interest. Thus, aggregates of a chimeric protein comprising a SCHAG amino acid sequence fused to an amino acid sequence encoding a binding domain of a protein having a specific binding partner are intended as an aspect of the invention.

[0141] In another preferred embodiment, the polypeptide of interest is an enzyme, especially an enzyme considered to be of catalytic value in a chemical process. Fibrils comprising such fusion proteins can be used as a catalytic matrix for carrying out the chemical process. Thus, aggregates of a chimeric protein comprising a SCHAG amino acid sequence fused to an enzyme are intended as an aspect of the invention.

[0142] In another preferred embodiment, ordered aggregates are created comprising two or more enzymes, such as a first enzyme that catalyzes one step of a chemical process and a second enzyme that catalyzes a downstream step involving a “metabolic” product from the first enzymatic reaction. Such aggregates will generally increase the speed and/or efficiency of the chemical process due to the proximity of the first reaction products and the second catalyst enzyme. Aggregates comprising two or more proteins of interest can be produced in multiple ways, each of which is itself considered an aspect of the invention.

[0143] It may be advantageous to attach fibers to a solid support such as a bead (e.g., a Sepharose bead) or a surface to create a “chip” containing loci with biological or chemical function.

[0144] In one variation, each chimeric protein comprising an aggregation domain and a protein of interest is produced in

a separate and distinct host cell system and recovered (purified and isolated). The proteins are either recovered in soluble form or are solubilized. (Complete purification is desirable but not essential for subsequent aggregation/polymerization). Thereafter, a desired mixture of the two or more proteins is created and induced into polymerization, e.g., by “seeding” with a protein aggregate, by concentrating the mixture to increase molarity of the proteins, or by altering salinity, acidity, or other factors. The desired mixture may be 1:1 or may be at a ratio weighted in favor of one chimeric protein (e.g., weighted in favor of an enzyme that catalyzes a slower step in a chemical process). The different chimeric proteins co-polymerize with the seed and with each other because they comprise compatible aggregation (SCHAG) domains, and most preferably identical aggregation domains. In certain embodiments it may be desirable to include in the pre-aggregation mixture a polypeptide comprising the SCHAG domain only, without an attached enzyme, for the purpose of increasing the average space between individual enzyme molecules in the aggregate that is formed. The additional space may be desirable, for example, if the enzyme’s substrate is a large molecule.

[0145] In another variation, the two distinct host cell systems are co-cultured, and the chimeric transgenes include signal peptides to induce the cells to secrete the chimeric proteins into the common culture medium. The proteins can be co-purified from the medium or induced to aggregate without prior purification.

[0146] In still another variation, the transgenes for two or more recombinant chimeric polypeptides are co-transfected into the same host cell, either on a single polynucleotide construct or multiple constructs. Such a host cell produces both recombinant polypeptides, which can be induced to polymerize in vivo in a prion phenotype host, or can be recovered in soluble form and induced to polymerize in vitro. The present invention also exploits the fact that at least certain prion proteins form aggregates that are fiber-like in shape; strong; and resistant to destruction by heat and many chemical environments. This collection of properties has tremendous industrial application that heretofore has not been exploited. Thus, in one embodiment, the invention provides polypeptides comprising SCHAG amino acid sequences which have been modified to comprise a discrete number of reactive sites at discrete locations. The polypeptides can be recombinantly produced and purified and aggregated into robust fibers resistant to destruction. The reactive sites permit modification of the polypeptides (or the fibers comprising the polypeptides) by attachment of virtually any chemical entity, such as pigments, light-gathering and light-emitting molecules for use as sensors, indicators, or energy harnessing and transduction; enzymes; metal atoms; organic and inorganic catalysts; and molecules possessing a selective binding affinity for other molecules. Electrical fields may be applied to fibers that are labeled with metal atoms, so that the fibers can be oriented in a specific direction. Because the fiber monomers are protein, conventional genetic engineering techniques can be used to introduce any number of desired reactive sites at precise locations, and the precise location of the reactive sites can be studied using conventional protein computer modeling as well as experimental techniques. Proteins and fibers of this type enjoy the utilities of the chimeric proteins described above (e.g., as chemical purification matrices, chemical reaction matrices, etc.) and additional utility due to the ability to bind a potentially infinite variety of

non-protein molecules of interest to the reactive sites. The fibers can be grown or attached to solid supports to create devices comprising the fibers.

[0147] In another preferred embodiment, the polypeptides of the present invention are used for the construction of nanostructures. For example, the N-terminal and middle region (NM) of yeast *Saccharomyces cerevisiae* Sup35p (i.e., NM) forms self-assembling β -sheet-rich amyloid fibers that are suitably sized and shaped for nanocircuitry with diameters of 9-11 nm (Glover, J. R., et al., *Cell*, 89: 811-819 (1997)). The highly flexible structure of soluble NM rapidly converts to form amyloid fibers when it associates with preformed fibers that act as seeds for fiber formation (Serio, T. R., et al., *Science*, 289: 1317-1321 (2000); Scheibel, T. & Lindquist, S. L., *Nat. Struct. Biol.*, 8:958-962 (2001); DePace, A. H. & Weissman, J. S., *Nat. Struct. Biol.*, 9, 389-396 (2002)). The fibers grow by extension from either end (Scheibel, T., et al., *Curr. Biol.*, 11: 366-369 (2001)), and this bidirectional formation is useful for forming varied fiber patterns: a valuable property for the production of circuitry.

[0148] NM has several advantageous properties for manufacturing. NM fibers have a higher than average chemical stability as demonstrated by its resistance to proteases and protein denaturants (Serio, T. R., et al., supra). Indeed, PrP, the mammalian prion counterpart of Sup35p, is infamous for its extraordinary resistance to destruction. (However, neither Sup35p nor NM are infectious to humans and therefore can be handled safely). The stability of NM suggests that it can withstand diverse metallization procedures necessary for creating electric circuits in industrial settings. In addition, NM fibers do not form aggregates as readily as other amyloids. Furthermore, under some circumstances such as different surface treatments, methods of fiber deposition, and solutions in which they are suspended, NM fibers tend not to aggregate with each other. The solubility of NM in physiological buffers greatly facilitates handling before and during fiber formation (Scheibel, T., et al., *Curr. Biol.*, 11: 366-369 (2001)).

[0149] Moreover, among the various DNA and protein fibers that have been described, NM fibers are unusual in that they are highly resistant to extended periods at high temperatures, exposure to high and low salt, strong denaturants, strong alkalis and acids, and 100% ethanol. These properties will allow them to withstand the harsh conditions in industrial processes. Depending on the conditions, NM fibers can nucleate spontaneously or self-assemble from preformed nuclei (Scheibel, T. & Lindquist, S. L., *Nat. Struct. Biol.*, 8:958-962 (2001)), an advantageous property for the practical assembly of circuits on a large scale. Further, the ability to manipulate the fiber length as described herein increases flexibility in designing nanostructures.

[0150] Bidirectional growth from NM seeded fibers can be used to incorporate NM derivatives with different modifications, interspersing them along individual fibers, e.g., with and without exposed cysteines. As different substrates can be prepared to bind to cysteine and to native lysine, these alternative binding sites provide flexibility and diversity in the patterning and mixing of substrates covalently bound to the fiber. Genetic engineering can be used to fuse a wide array of protein domains to the C-terminus of NM during its initial in vivo synthesis in such a way that the domains are tethered laterally, external to the surface of assembled fibers. Thus they remain functional even when NM is in its fibrous form.

[0151] Because many enzymes can function when attached to protein fibers, it is possible to incorporate more complex

reaction centers into NM nanocircuitry, thereby creating electronic circuits that can take advantage of biological capacities. Mechanisms such as the vaporization of NM fibers with high voltages could act as a fuse or a switch to permanently activate or inactivate specific reaction centers within the circuitry.

[0152] Fibril-based electrical conductors of the invention can be used as components in any product, device, or method of manufacture requiring electrical conductors. Due to their small size, electrical conductors of the invention are especially useful for small-scale devices such as microcircuits in nanodevices. Referring to FIG. 10, an exemplary circuit comprises a power source **1**, one or more circuit elements **3**, and electrical conductors (e.g., wires) disposed between the power source and the circuit elements **2** (and optionally between circuit elements). For example, a first location of the electrical conductor is attached to or contacts the power source and a second location of the electrical conductor is attached to or contacts a circuit element in a manner whereby the electrical conductor can conduct electricity between the power source and the circuit element (or between circuit elements). Circuit elements can be active or passive and can be any component that could be included in a circuit, such as a capacitor, an inductor, a resistor, an integrated circuit, an oscillator, a transistor, a diode, a switch, or a fuse.

[0153] There is a great opportunity to expand further the potential interconnections in these circuits by exploiting the natural diversity and strength of protein-protein interactions (Begley, T. J., et al., *Mol. Cancer Res.*, 1: 103-112 (2002); Uetz, P., et al., *Nature*, 403: 623-627 (2000); Marcotte, E., et al., *Nature*, 402: 83-86 (1999)). Protein-protein interactions can be extremely specific and strong, as can the interactions of protein-ligand-protein. Such protein properties can be used as a mechanism to bring premetallized wires into juxtaposition in response to changes in physical conditions, the presence of ligands, and the appearance of partner proteins, etc. These connections are readily reversible (Schreiber, S. L. & Crabtree, G. R. *Harvey Lect.*, 91: 99-114 (1995-1996); Spencer, D. M., et al., *Science*, 262: 1019-1024 (1993)).

[0154] Complex circuit schematics can be generated with NM fibers, initiated by patterned surface modifications (independently or in combination) such as lithography, growth in flows or magnetic field gradients, alignment by electrical fields, active patterning with optical tweezers, dielectrophoresis and 3D patterning using hydrogels or microfluidic channels (Korda, P., et al., *Rev. Sci. Instrum.* 73: 1956-1957 (2002); Kane, R. S., et al., *Biomaterials* 20: 2363-2376 (1999); Inouye, H., et al., *Biophys. J.* 64: 502-519 (1993); Luther, P. W., et al., *Nature* 303: 61-64 (1983); Kubista, M., et al., *J. Biomol. Struct. Dyn.* 8: 37-54 (1990); Hermanson, K. D., et al., *Science* 294: 1082-1086 (2001)). The feasibility of such maneuvers is demonstrated by the natural tendency of NM fibers to align with each other rather than to form dense intractable clumps characteristic of other protein amyloids and the conditions that produce such alignments can be optimized. Attachment of NM to patterned surfaces can be mediated via covalent bonds to native lysine residues, genetically engineered cysteine residues, or other novel residues or modifications.

[0155] The present invention provides a mechanism for generating robust nanowires that meet the needs of industrial processes with the potential to couple powerful combinations of biological processes and functionalities with electronic circuitry. In particular, these nanowires may be electrical

conductors which may include any type of electrically conductive materials such as metal, like gold, silver, copper, etc., or semi-conductive materials such as known semi-conductors suited to conduct electricity either along the length of the nanowire, radially with respect to the nanowire, or a combination of both.

[0156] These and other aspects of the invention will be better understood by reference to the following examples. The examples are not intended to limit the scope of the invention, and variations will be apparent to the reader from the entirety of this document.

Example 1

Construction and Assaying of a Chimeric, Prion-Like Gene and Protein with Yeast Sup35 Protein

[0157] The following experiments were performed to demonstrate that a prion-determining domain of a prion-like protein can be fused to a polypeptide from a wholly different protein to construct a novel, chimeric gene and protein having prion-like properties. The relevance of these experiments to the present invention also is explained.

A. Construction of a NMSup35-GR Chimeric Gene

[0158] The yeast (*Saccharomyces cerevisiae*) Sup35 protein (SEQ ID NO: 2, 685 amino acids, Genbank Accession No. M21129) possesses the prion-like capacity to undergo a self-perpetuating conformational alteration that changes the functional state of Sup35 in a manner that creates a heritable change in phenotype. Experiments have demonstrated that it is the amino-terminal (N region, amino acids 1-123 of SEQ ID NO: 2) or the amino-terminal plus middle (M, amino acids 124-253 of SEQ ID NO: 2) regions of Sup35 that are responsible for this prion-like capacity. See Glover et al., *Cell*, 89: 811-819 (1997); see also King et al., *Proc. Natl. Acad. Sci. USA*, 94:6618-6622 (1997) (N-terminal polypeptide fragment consisting of residues 2-114 of Sup35 spontaneously aggregates to form thin filaments in vitro). The M domain is highly charged and therefore acts to maintain the protein in solution. This property causes the aggregation process to proceed more slowly, providing beneficial control to the system.

[0159] A chimeric polynucleotide FIG. 1 and (SEQ ID NO: 50) was constructed comprising a nucleotide sequence encoding the N and M domains of Sup35 (FIG. 1 and SEQ ID NO: 50, bases 1 to 759) fused in-frame to a nucleotide sequence (derived from a cDNA) encoding the rat glucocorticoid receptor (GR) (Genbank Accession No. M14053, FIG. 1 and SEQ ID NO: 50, bases 766-3150), a hormone-responsive transcription factor, followed by a stop codon. This construct was inserted into the pRS316CG (ATCC Accession No. 77145, Genbank No. U03442) and pG1 (Guthrie & Sink, "Guide to Yeast Genetics and Molecular Biology" in *Methods of Enzymology*, Vol. 194, pp. 389-398 (1981)) plasmids under the control of either the CUP1 promoter (plasmid pCUP1-NMGR, inducible by adding copper to the growth medium) or the constitutive GPD promoter (plasmid pGDP-NMGR). The nucleotide sequences of CUP1 and GDP (Genbank Accession No. M13807) promoters are set forth in SEQ ID NOs: 11 and 48, respectively. The GR coding sequence without NM, in the same promoter and vector constructs (plasmids pCUP1-GR and pGDP-GR), served as a control. GR activity in transformed yeast was monitored with two reporter constructs containing a glucocorticoid response promoter

element (GRE) [Schena & Yamamoto, *Science*, 241:965-967 (1988)] fused to either a β -galactosidase (Swiss-Prot. Accession No. P00722) or to a firefly luciferase (Genbank Accession No. M15077) coding sequence. When GR is activated by hormone, e.g., deoxycorticosterone (DOC), it normally binds to the GRE and promotes transcription of the reporter enzyme in either mammals or yeast. See M. Schena and K. Yamamoto, *Science* 241:965-967 (1988).

B. Construction of a NMSUP35-GFP Chimeric Gene

[0160] A chimeric gene comprising the NM region of Sup35 fused to a green fluorescent protein (GFP) sequence and under the control of the CUP1 promoter was constructed essentially as described in Patino et al., *Science*, 273: 622-626 (1996) (construct NPD-GFP), incorporated by reference herein. (The use of GFPs as reporter molecules is reviewed in Kain et al., *Biotechniques*, 19:650-655 (1995); and Cubitt et al., *Trends Biochem. Sci.*, 20:448-455 (1995), incorporated by reference herein). The resulting construct encodes the NH₂-terminal 253 residues of Sup35 (SEQ ID NO: 2) fused in-frame to GFP. The NM-Sup35-GFP encoding sequence was amplified by PCR and cloned into plasmid pCLUC [D. Thiele, *Mol. Cell. Biol.*, 8: 745 (1988)], which contains the CUP1 promoter for copper-inducible expression. A similar construct was created substituting the constitutive GDP promoter for the CUP1 promoter. An identical GFP construct lacking the NM fusion also was created.

C. Transformation and Phenotypic Analysis of [psi⁻] and [PSI⁺] Yeast

[0161] 1. Constructs Regulated by the CUP1 Promoter

[0162] The GR and NM-GR constructs regulated by the CUP1 promoter on a low copy plasmid (ura selection) were transformed into [psi⁻] and [PSI⁺] yeast cells (strain 74D) along with a 2 μ (high copy number) plasmid containing a GR-regulated β -galactosidase reporter gene with leucine selection. Transformants were selected by sc.-leu-ura and used to inoculate sc.-leu-ura medium. Cultures were grown overnight at 30° C., and induced by adding copper sulfate to the medium to a final 0-250 μ M copper concentration.

[0163] After 4 to 24 hours of induction, both proteins were expressed at a similar level in [psi⁻] cells, and both the GR and NM-GR transformed [psi⁻] cells produced similar levels of reporter enzyme activity in response to hormone (DOC added to a final concentration of 10 μ M at the time of copper sulfate induction). Virtually no reporter enzyme activity was detected without hormone. The fact that both GR and NM-GR constructs resulted in similar levels of activity indicates that the NM fusion does not intrinsically alter the ability of GR to function in hormone-activated transcription, demonstrating the utility of the NM domain as a fusion protein tag.

[0164] In contrast, when the same constructs were transformed into yeast cells that contain the heritable, conformationally-altered form of Sup35 [PSI⁺], GR activity was reduced in cells expressing the NM-GR fusion construct, compared to cells expressing GR. Thus, pre-existing prions (which comprise self-coalescing aggregates of NM-containing Sup35 protein) can interact with NM-GR. Similar results were obtained with NM-Green Fluorescent Protein (GFP) constructs: NM-GFP interacted with pre-existing [PSI⁺] elements, but GFP alone did not.

[0165] An important difference existed between the NM-GR and NM-GFP studies in the [PSI⁺] cells, however. Unlike the NM-GR fusion, the NM-GFP fusion retained similar GFP activity with the [PSI⁺] prion, i.e., the NM-GFP fusion still

glowed green. This difference in activity is explained by the facts that, for biological activity, GR needs to be in the nucleus, bind to DNA, and interact in specific ways with other elements of the transcription machinery. When NM-GR is sequestered in [PSI⁺] cells by interacting (aggregating) with the Sup35 prion filaments, the GR function is diminished.

[0166] 2. Constructs Regulated by the Constitutive GPD Promoter on a High Copy Plasmid.

[0167] A set of experiments demonstrated that plasmids that cause expression of NM at a high level can be successfully transformed into [psi⁻] yeast cells, but not into [PSI⁺] cells. Apparently, over-expressed NM causes excessive prion-like aggregation of endogenous Sup35 in cells that are already [PSI⁺], eliminating so much translation termination factor function that the yeast cells cannot survive.

[0168] When a high copy plasmid vector comprising the NM-GR open reading frame under the control of the constitutive GPD promoter was used to transform [psi⁻] or [PSI⁺] yeast, no [PSI⁺] transformants were obtained, whereas [psi⁻] transformants were readily obtained. The control GR construct in the same vector and under control of the same promoter transformed equally well into both [PSI⁺] and [psi⁻] cells.

[0169] When amino acids 22-69 in the N domain of Sup35 are deleted, the resultant protein fails to form ordered aggregates, and yeast comprising this Sup35 variant fail to adopt a [PSI⁺] phenotype. When these same amino acids were deleted from the high copy number NM-GR plasmid, the inability to transform [PSI⁺] cells was eliminated: transformants were obtained as readily in [PSI⁺] as [psi⁻] cells.

[0170] Both NM-GR and GR [psi⁻] transformants were used to inoculate sc.-leu-trp medium, and the cultures were grown at 30° C. overnight, diluted into fresh medium to achieve a cell density of 2-4×10⁶ cells/ml, induced with DOC (10 μM final concentration), and grown for an additional period varying from 1 hour to overnight. Analysis of marker gene activity in the transformed [psi⁻] cells demonstrated that hormone responsive transcription was lower in NM-GR transformants than in GR transformants. Western blotting using an anti-GR monoclonal antibody (Affinity Bioreagents Inc., MA1-510) was used to examine the levels of NMGR and GR expression in these cells. Although cells carrying the NM-GR fusion had lower levels of GR activity, the NM-GR protein was actually expressed at a much higher level than the GR protein without the NM domain. Thus, the reduced levels of hormone-activated transcriptional activity were not due to an effect of NM on the accumulation of the transcription factor, but to an alteration in GR activity in the NM-GR-expressing cells. This reduced activity suggested that NM-GR is capable of undergoing a de novo, prion-like alteration in function when it is expressed at a sufficiently high level.

[0171] To confirm that NM-GR was forming prions de novo in the transformed [psi⁻] cells into which it had been introduced, such cells were induced with copper to express NM-GR and then were plated onto copper-free media lacking adenine, and therefore selective for the [PSI⁺] element/phenotype. See Chernoff et al., *Science*, 268: 880 (1995), and Cox et al., *Yeast*, 4 (3): 159-178 (1988). A substantial fraction of the cells were able to grow on medium selective for [PSI⁺], suggesting that the highly expressed NM-GR was responsible for the formation of new prions putatively containing both NM-GR and Sup35 protein. Moreover, the number of colonies obtained varied with the level of copper induction prior to plating. This change in the growth properties of the cells was

observed to be heritable and was maintained even under conditions where the NM-GR plasmid construct was lost by the host cells, indicating that NM-GR had induced the formation of a new Sup35-containing prion.

D. Analysis of NMGR-Induced Phenotype in Cells Carrying a Deletion of the NM Region of Sup35.

[0172] To further confirm that NM-GR was truly functioning as an independent, novel prion, experiments were conducted to determine whether an NM-GR prion was formed independently of both the yeast [PSI⁺] element and the endogenous Sup35 protein. Specifically, the GPD-regulated GR and NM-GR constructs were co-transformed with plasmid p5275 (containing GRE linked to a firefly luciferase reporter gene) into a yeast strain (ΔNMSUP35) carrying a deletion of the NM region of the SUP35 gene. Three independent transformants of each construct (GR or NM-GR) were examined. Colonies were picked and grown overnight in SC selective media (-trp, -ura) at 30° C. Thereafter, deoxycorticosterone (DOC) was added to the growth medium to a final concentration of 10 μM. Luciferase activity was assayed in intact cells after 25 hours of DOC induction.

[0173] All three transformants expressing the NM-GR protein showed lower levels of GR activity (specific activities of about 4, 5, 4) than the three transformants expressing GR without the NM fusion (specific activities of about 23, 28, and 39). The differences in GR activity was observed after 1 hour of hormone induction and appeared to increase after 5.5 or after 25 hours of induction.

[0174] Western blotting was conducted to determine whether the differences in activity were the result of differences in protein concentration. Ethanol lysates were prepared from 3 ml yeast cultures expressing GR or NMGR twenty-five hours after the addition of DOC. About 50 μg total protein was analyzed by SDS/PAGE and immunoblot. The protein gel was transferred onto PVDF membranes and probed with a monoclonal antibody against GR (Bu-GR2, Affinity Bioreagents, Golden Colo.). The same membrane was later stained with Coomassie blue to semiquantitatively evaluate total protein. The Western studies again showed that the levels of NM-GR were higher than the levels of GR alone.

E. Effect of Guanidine Hydrochloride and Hsp104 on NM-GR Prions.

[0175] When the yeast having [URE3] or [PSI⁺] phenotypes are passaged on medium containing low concentrations of guanidine hydrochloride (GdHCl), their prion determinants change ("cure") at a high frequency from the aggregated, inactive prion state into the active, unaggregated state, and such changes are heritable. These phenotypes also can be cured by over-expression of the chaperone Hsp104.

[0176] Another series of experiments were conducted to assay for such curative behavior in yeast harboring an NM-GR construct. The natural GR protein contains a ligand-binding domain and hormone must be added to the medium to determine whether or not the protein is active. For this series of experiments, the hormone-binding domain was removed from the NM-GR construct, creating an NM-GR fusion that was constitutively active.

[0177] Yeast expressing the NM-GR chimeric construct and a glucocorticoid response element fused to a β-galactosidase marker exhibited different levels of prion-like behavior, manifested by different colony colors. In addition to white

colonies (indicative of a prion-like state lacking β -gal induction) and blue colonies (indicative of soluble NM-GR and high levels of β -gal induction), medium blue and pale blue colonies also were observed. (Western blotting indicated that differently colored colonies contained comparable amounts of GR protein). These differently colored colonies were replica-plated onto plates containing 5 mM GdHCl and then subsequently replica-plated again onto X-Gal indicator plates. In control cells expressing vector alone (no NM-GR insert), white colonies remained white. However, all of the NM-GR-expressing colonies produced blue colonies. The efficiency of curing varied with the NM-GR strain: medium blue colonies produced almost entirely blue colonies, whereas pale blue colonies produced a mixture of blue and white colonies.

[0178] To determine if the heritable loss of NM-GR activity is susceptible to Hsp104 curing, white colonies of cells expressing NM-GR were transformed with a GDP-HSP104 over-expression plasmid and streaked onto X-Gal indicator plates. Control cells transformed with empty vector remained white. In contrast, white cells transformed with the Hsp104 over-expression construct changed to blue. The blue cells remained blue upon-restreaking, indicating that transient over-expression of Hsp104 was sufficient to cure cells of the heritable reduction of NM-GR activity.

[0179] When the same NM-GR constructs were used to transform yeast containing a deletion mutation of Hsp104, white colonies were never produced. This finding is consistent with the observation that Hsp104 mutations are incompatible with the maintenance of the [PSI⁺] phenotype.

[0180] Together, the foregoing data indicate that the difference in GR activity observed when NM-GR is expressed at a high constitutive level is due to a heritable alteration in GR function, rather than to an alteration in GR expression.

[0181] Collectively, the foregoing experiments demonstrate that the amino-terminal domain of a prion-like yeast gene, Sup35, can be fused to a polypeptide from a wholly different protein to construct a novel, chimeric gene and protein having prion-like properties. Significantly, these results are believed to be the first demonstration that a SCHAG protein domain can be fused to a non-native protein domain to form a chimera, expressed in a host cell that fails to express the native SCHAG protein, and still behave in a prion-like manner. (Specifically, these results demonstrate that the NM domains of SUP35 will behave like a prion even when the C-terminal domain of the protein is not the native Sup35 C-terminus, and even when the host cell does not express an endogenous Sup35 protein containing an NM region). The experiments also define exemplary assays for screening other putative prion-like peptides for their ability to confer a prion-like phenotype. (It will be apparent that the use of markers other than GFP, GR, luciferase, or β -galactosidase would work in such assays. The GFP marker is useful insofar as it provides an effective marker for localizing a fusion protein *in vivo*. The GR marker is additionally useful insofar as GR activity depends on GR localization in the nucleus, DNA binding, and interaction with transcription machinery; whereas GFP is active in the cytoplasm). Exemplary prion-like peptides for screening in this manner are peptides identified according to assays described below in Example 5; mammalian PrP peptides responsible for prion-forming activity; and other known fibril-forming peptide sequences, such as human amyloid β (142) peptide.

[0182] In addition, the experiments demonstrate an improved procedure for recombinant production of certain proteins that might otherwise be difficult to recombinantly produce, e.g., due to the protein's detrimental effect on the growth or phenotype of the host cell. For example, DNA binding and DNA modifying enzymes that might locate to a cell's nucleus and detrimentally effect a host cell may be expressed as a fusion with a SCHAG amino acid sequence from a prion-like protein. In host cells wherein the aggregate-forming phenotype is present, the recombinant protein is "sequestered" into higher order aggregates. By virtue of this sequestration, the biological activity of the resultant protein in the nucleus is reduced. The fusion protein is purified from the insoluble fraction of host cell lysates, and can be cleaved from the fibril core if an appropriate endopeptidase recognition sequence has been included in the fusion construct between the SCHAG amino acid sequence and the sequence of the protein of interest. (An appropriate endopeptidase recognition sequence is any recognition sequence that is not present in the protein of interest, such that the endopeptidase will cleave the protein of interest from the fibril structure without also cleaving within the protein of interest).

Example 2

Construction and Assaying of a Chimeric, Prion-Like Gene and Protein with Yeast Ure2 Protein

[0183] The following experiments were performed to demonstrate that the prion-determining domain of yeast Ure2 protein also can be fused to a polypeptide other than the Ure2 functional domain to construct a novel, chimeric gene and protein having some prion-like properties. Two prion-like elements are known in yeast: [PSI⁺] and [URE3]. The underlying proteins, Sup35 and Ure2, each contain an amino-terminal domain (the N domain) that is not essential for normal function but is crucial for prion formation. The N domains of both Sup35 and Ure2 are unusually rich in the polar amino acids asparagine and glutamine.

[0184] A. Construction of a NUre2-CSup35 Chimeric Gene

[0185] A chimeric polynucleotide (FIG. 3, SEQ ID NO: 49) was constructed comprising a nucleotide sequence encoding the N domain of yeast (*Saccharomyces cerevisiae*) Ure2 protein (Genbank Accession No. M35268, SEQ ID NO: 3, bases 182 to 376, encoding amino acids 1 to 65 (SEQ ID NO: 4) of Ure2 (NUre2)), fused in-frame to a nucleotide sequence encoding a hemagglutinin tag (SEQ ID NO: 13, TAC CCA TAC GAC GTC CCA GAC TAC GCT), fused in-frame to a nucleotide sequence encoding the C domain of yeast Sup35 (CSup35) protein that is responsible for translation-regulation activity of Sup35 (Genbank Accession No. M21129, SEQ ID NO: 1, bases 1498-2793, encoding amino acids 254 to 685 of Sup35 (SEQ ID NO: 2)). At the 5' and 3' ends of this construct were 5' and 3' flanking regions, respectively, of the yeast Sup35 genomic DNA. This construct was inserted into the pRS306 plasmid (available from the ATCC, Manassas, Va., USA, Accession No. 77141; see also Genbank Accession No. U03438) as shown in FIGS. 2 and 3, and used to transform yeast as described below.

[0186] B. Transformation and Phenotypic Analysis of Yeast

[0187] To replace the Sup35 gene with the NUre2-CSup35 chimeric gene, the first step was to integrate the gene fragment into the yeast genome. Freshly grown cells from overnight culture were collected and resuspended in 0.5 ml LiAc-

PEG-TE solution (40% PEG4000, 100 mM Tris-HCL, pH7.5, 1 mM EDTA) in a 1.5 ml tube. 100 µg/10 µl carrier DNA (salmon testis DNA, boiled 10 minutes and chilled immediately on ice) and 1 µg/2 µl of transforming plasmid DNA were added and mixed. This transformation mixture was incubated overnight at room temperature and then heat shocked at 42°C for 15 minutes. 100 µl of transformation mixture were then spread onto a uracil dropout plate. After transformation, selection for Ura+ results in an integration event, such that native and chimeric genes bracket the URA3-containing plasmid sequence. Transformants were picked and cells having the integrated chimeric gene were confirmed by genomic PCR and Western blot.

[0188] The second step of the replacement involved the excision or “popping out” of the wildtype Sup35 gene through homologous recombination between the native Sup35 and the chimeric sequence. Popout of the plasmid was monitored by screening for colonies that are ura- and therefore resistant to the drug 5-fluoroorotic acid (5-FOA). Cells with NUre2-CSup35 integrated were thus plated onto 5-FOA medium to select for those that have the plasmid sequence containing one copy of the Sup35 gene popped out. Clones in which the native Sup35 gene had been replaced with the chimeric gene were then screened by means of colony PCR and further confirmed by Western blot.

[0189] To screen for yeast strains that have gene integration and replacement, a Ure2 coding sequence N-terminal primer and a Sup35 coding sequence primer were used for PCR reactions. The NUre2-CSup35 DNA fragment can only be amplified from genomic DNA of cells containing the chimeric gene. To confirm that only the fusion protein of NUre2-CSup35 was expressed in those cells that have the gene replacement, yeast cells were lysed and the cell lysates were run on SDS-polyacrylamide gel and proteins were transferred to PVDF immunoblot. Since there is a hemagglutinin (HA) tag inserted between NUre2 and CSup35, Western blots were then probed with anti-HA antibody from Boehringer Mannheim. To confirm that NUre2-CSup35 is the only copy of Sup35 gene in yeast genome, Western blots were also probed with an antibody against the middle region of Sup35 protein. Loss of antibody signal verified that the NM region of Sup35 gene had been replaced with the N-terminus of Ure2. Thus, the transformed cells were characterized by a deleted native Sup35 gene that had been replaced by the NUre2-CSup35 chimeric gene.

[0190] Transformed colonies carrying the chimeric NUre2-CSup35 gene of interest were grown on rich medium (YPD) at 30° C. The resultant colonies were streaked onto [PSI⁺] selective medium (SD-ADE) and incubated at 30° C. to determine whether some or all contained a [PSI⁺] phenotype. Two different types of colonies were observed. Some showed normal translational termination characteristic of a [psi⁻] phenotype. Others showed the suppressor phenotype characteristic of [PSI⁺] cells. Both phenotypes were very stable and were inherited from generation to generation of the transformed yeast cells.

[0191] To determine whether the observed difference in translational fidelity was due to a heritable change in protein conformation, cells were lysed and the lysates subjected to centrifugation at 12,000 or 100,000×g for 10 minutes. Supernatants and precipitate fractions were screened for the fusion protein using an anti-HA antibody (HA 11, Covance Research Products Inc.). The cells that showed reduced translational fidelity also showed aggregation of the NUre2-

CSup35 fusion protein, whereas the fusion protein did not appear aggregated in cells having normal translation termination characteristics.

[0192] The foregoing experiments demonstrate that the amino-terminal domain of another prion-like yeast gene, Ure2, can be fused to a polypeptide derived from a wholly different protein to construct a novel, chimeric gene and protein having prion-like properties. These results represent the first such demonstration of this kind. [Compare Maisson & Wickner, *Science*, 270: 93 (1995) (Ure2₁₋₆₅/β-gal fusion did not change the activity of the β-galactosidase enzyme) and Paushkin et al., *EMBO J.*, 15 (12): 3127-3134 (1996) (GST-NSup35 chimeric construct did not allow native Sup35 to adopt an altered state.)]

[0193] Several factors are suggested for achieving prion-like behavior with chimeric genes that comprise SCHAG sequences. First, it is preferable to include the SCHAG sequence at a location in the chimeric gene (e.g., amino-terminus or carboxy-terminus) that corresponds to the location at which it is found in its native gene. For example, if NSup35 is selected as the SCHAG sequence, then the chimeric gene preferably is constructed with NSup35 at the amino-terminus, preceding the sequence encoding the polypeptide of interest. Second, it is preferable to include a spacer region of, e.g., at least 5, 10, 20, 30, 40, or 50 amino acids, and preferably at least 60, 70, 80, 90, 100, 120, 130, 140, or 150 amino acids, to separate the SCHAG domain from other domains and reduce the likelihood of steric hinderance caused by other domains. The length of spacer apparently can be quite large because a chimeric construct comprising whole Sup35 fused to Green Fluorescence Protein appears to act as a prion in preliminary experiments. Third, it is preferable if the protein of interest is a protein that does not itself naturally form multimers, because multimer formation of the protein of interest is apt to cause steric interference with the ordered aggregation of the SCHAG domain. (Maisson & Wickner's research involved β-galactosidase, which forms a tetrameric functional unit.) The experiments also demonstrate an alternative assay system (i.e., CSup35 fusions) to the GFP and GR assay systems described in the preceding example to screen peptide sequences for their ability to confer prion-like phenotypic properties.

[0194] Also contemplated are fusion proteins comprising the M domain of Sup35, or portions of fragments thereof, fused to a different protein to generate a novel protein with prion-like activities. Likewise, fusion proteins displaying prion-like properties, comprising portions or fragments of the N domain, or comprising portions or fragments of the N and of the M domain are also contemplated.

Example 3

Modulation of Propensity of Protein to Form Prion-Like Aggregates

[0195] The following experiments demonstrate that the propensity of novel chimeric proteins to aggregate into prion-like fibrils can be modulated by varying the number of oligopeptide repeats in the SCHAG portion of the chimeric protein. An increased propensity to form such fibrils is useful in instances where the fibrils themselves comprise a desirable end product to be harvested from cells, e.g., via lysis and centrifugation; and in instances where fibril formation in vivo is desired to phenotypically alter a cell, e.g., by sequestering

a biologically active molecule in the cell away from the molecule's normal subcellular region of biological activity.

[0196] The yeast Sup35 protein contains an oligopeptide repeat sequence (PQGGYQQYN, SEQ ID NO: 2, residues 75 to 83; with imperfect repeats at residues 41 to 50; 56 to 64; 65 to 74; and 84 to 93). The following experiments demonstrated that an expansion of this oligopeptide repeat in the NM region of Sup35 increases the rate of appearance of new, heritable, [PSI⁺]-like elements, whereas decreasing the number of repeats lessened the rate of appearance of such elements.

[0197] Three expression vectors were created for the experiment containing a chimeric gene comprising a CUP1 promoter sequence (SEQ ID NO: 11) operably linked to a sequence encoding a Sup35 NM region, fused in-frame with a "superglow" GFP encoding sequence (SEQ ID NO: 39). In the first construct (R_A2-5), the Sup35 NM region had been modified by deleting four of the five oligopeptide repeats found in the native N region (SEQ ID NOs: 14 & 15). In the second construct (R2E2), the Sup35 NM region had been modified by twice expanding the second oligopeptide repeat found in the native N region, creating a total of seven oligopeptide repeats (SEQ ID NOs: 16 & 17). In the third construct, the native Sup35 NM region was employed (SEQ ID NO: 1, nucleotides 739 to 1506, encoding residues 1 to 256 of SEQ ID NO: 2). The CUP1 promoter permitted control of the expression of the chimeric proteins by manipulation of copper ion concentration in the growth medium. [See Thiele, D. J., *Mol. Cell. Biol.*, 8: 2745-2752 (1988).] The attachment of GFP to NM permitted visualization of the mutant proteins in living cells.

[0198] Each of the three above-described NM-GFP constructs were introduced via homologous recombination at the site of the wild-type Sup35 gene into [psi⁻] yeast cells carrying a nonsense mutation in the ADE1 gene (strain 74-D694 [psi⁻]), and monitored for the frequency at which cells converted to a [PSI⁺] phenotype. Cell cultures in the log phase of growth at 30° C. were induced to express the GFP-fusion proteins by adding CuSO₄ to the cultures cells to a final concentration of 50 μM. For analysis via fluorescence microscopy, cells were fixed with 1% formaldehyde after four hours and twenty hours of culture. For analysis of [PSI⁺] induction, cells over-expressing the GFP fusion proteins were serially diluted and spotted onto YPD and SD-ADE media after four hours and twenty hours. Conversion was measured by the ability of cells to grow on medium without adenine (SD-ADE). The [PSI⁺] phenotype causes readthrough of nonsense mutations, producing sufficient protein to suppress the ADE1 mutation and allow growth without adenine.

[0199] Cells were induced with copper for 4 hours to promote expression of the chimeric gene and serially diluted, and then aliquots of each dilution were plated on SD-ADE, conditions that allowed loss of the plasmid. To demonstrate that the initial cultures contained similar numbers of cells, serial dilutions from each culture also were plated on rich medium (YPD) which allowed the growth of all cells in the culture. After incubating the plates for 48 hours at 30° C., colonies on each plate were counted.

[0200] Cells expressing the oligopeptide repeat expansion mutation converted to [PSI⁺] at a much higher frequency than cells expressing the native Sup35NM-GFP, which in turn converted to [PSI⁺] at a higher frequency than cells expressing the oligopeptide repeat deletion mutation. The observed conversion results were specifically attributable to the pro-

duction of the chimeric proteins, because the conversion to [PSI⁺] did not occur in cells that were not induced with copper (control).

[0201] In a related experiment, the repeat expansion and repeat deletion mutations were introduced into a full-length Sup35 protein-encoding sequence to create constructs encoding the NM(R2E2) and NM(RΔ2-5) fused to the CSup35 domain. These constructs were introduced into the genome of [psi⁻] yeast strain 74-D694 with the wild-type Sup35 promoter, in each case replacing the native Sup35 gene. Transformants were selected on uracil-deficient medium and confirmed by genomic PCR. Recombinant excision events were selected on medium containing 5-fluoroorotic acid. [See Ausubel et al., *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, New York (1991).] Strains in which wild-type Sup35 was replaced with the R2E2-CSup35 and RΔ2-5CSup35 variants were screened by PCR and confirmed by Western blotting. The cells were cultured on ypd or synthetic complete media at 25° C. for 24 hours, serially diluted, and plated on SD-ADE media to screen for [PSI⁺] conversions. As shown in FIG. 4, the spontaneous rate of appearance of [PSI⁺] colonies was increased about 5000-fold in cells carrying the repeat expansion (R2E2) compared to wild-type cells. The wild-type cells produced colonies on the selective medium at a frequency of about 1 per million cells plated. The RΔ2-5 cells produced such colonies at even lower frequency, and it appears that none of these were attributable to development of a [PSI⁺] phenotype, since they could not be cured by growth on medium containing 5 mM guanidine HCl. In contrast, growth of the wild-type and the R2E2 colonies on the selective medium could indeed be cured by the guanidine HCl treatment.

[0202] In additional experiments, the effects of the Sup35 repeat variants were examined when they were used to replace the wild-type Sup35 gene in [PSI⁺] cells. Cells with the R2E2 replacement remained [PSI⁺], whereas all cells carrying the RΔ2-5 replacement became [psi⁻]. Thus, maintenance of the [PSI⁺] phenotype requires a Sup35 gene having more than one of the oligopeptide repeats.

[0203] Still another series of tests examined the effects of the repeat variants on the structural transition of NM in vitro. When purified recombinant NM is denatured and diluted into aqueous buffers, it slowly changes from a random coil into a β-sheet rich structure and forms fibers that bind Congo red with the spectral shift characteristic of amyloid proteins. When deposited at high concentrations, the Congo red-stained fibers also show apple-green birefringence. To determine if the repeat variants alter the intrinsic capacity of the protein to fold in this form, the wild-type and two repeat variants were purified in fully denatured states and then diluted into a non-denaturing buffer. Structural changes were monitored by the binding of Congo red [Klunk et al., *J. Histochem. Cytochem.*, 37: 1293-1297 (1989)] and confirmed by circular dichroism and electron microscopy analysis. In these experiments, the R2E2 variant converted to a β-sheet rich structure about twice as quickly as the wild-type NM polypeptide, which in turn converted significantly faster than the RΔ2-5 variant. These differences were reproducibly obtained in both rotated and unrotated reactions, although the transition was slower in the unrotated reactions. This data indicates that alterations in the number of repeat units alters the propensity of Sup35 NM polypeptides to progress from an unfolded state into a β-sheet rich, higher-ordered structure.

[0204] The foregoing experiments demonstrate that the propensity of novel chimeric proteins to aggregate into prion-like fibrils can be modulated by alteration of the SCHAG amino acid sequence of the chimera. Modulation of any SCHAG amino acid sequence in this manner is specifically contemplated as an aspect of the invention, as are the resulting gene and protein products. In addition to alteration by adding or deleting oligopeptide repeat regions, alterations by adding or deleting larger regions is specifically contemplated as an aspect of the invention. By way of example, the entire N terminal region of Sup35 or Ure2 could be duplicated to increase the propensity of transformed cells to produce aggregated chimeric sequences.

Example 4

Demonstration that a Prion can be Moved from One Organism to Another

[0205] The following experiments demonstrate that a prion protein from one organism will continue to behave in a prion-like manner when recombinantly expressed in another organism, and can even do so when expressed in a different cellular compartment than that in which the protein is produced in its native host.

[0206] Polynucleotides encoding mouse (SEQ ID Nos: 18 and 19) and Syrian Hamster (SEQ ID Nos: 20 and 21) PrP proteins were expressed in yeast cells under the control of the constitutive GPD promoter. The protein was produced in the yeast cytosol, without signal sequences that would normally guide it to the endoplasmic reticulum, and without the tail that is normally clipped off during maturation of these proteins in their native hosts. In other words, the PrP protein product in yeast was similar to the final mature product in mammalian neurons, except that it did not contain the sugar modification and GPI anchor. There has been considerable data suggesting that these sugar and GPI anchor characteristics are not required for prion formation.

[0207] The normal cellular form of PrP (PrP^C) is detergent soluble, but the conformationally changed-protein that is characteristic of neurodegenerative prion disease states (PrP^{Sc}) is insoluble in detergent such as 10% Triton. When PrP protein is expressed in yeast, it was insoluble in non-ionic detergents, suggesting that a PrP form was present.

[0208] PrP-transfected yeast cells were lysed in the presence of 10% Sarkosyl and centrifuged at 16,000×g over a 5% sucrose cushion for 30 minutes. Proteins in both the supernatant and pellet fractions were analyzed on SDS polyacrylamide gels. Coomassie blue staining revealed that most proteins were soluble under these conditions and were present in the supernatant fraction. When identical gels were blotted to membranes and reacted with antibodies against mammalian PrP, most of the PrP protein was found in the pellet fraction, further suggesting that a PrP^{Sc} form was present in the yeast.

[0209] Protease studies provide further evidence that the yeast PrP was adopting a PrP^{Sc} conformation. When PrP protein is expressed in yeast it displays the same highly specific pattern of protease digestion as does the disease form of the protein in mammals. The normal cellular form of PrP is very sensitive to protease digestion. In the disease form, the protein is resistant to protease digestion. This resistance is not observed across the entire protein, but rather, the N-terminal region from amino acids 23 to 90 is digested, while the remainder of the protein is resistant. As expected, when PrP was expressed in the yeast cytosol it was not glycosylated,

and it migrated on an SDS gel as a protein of ~27 kD. After protease digestion, a resistant fragment of ~19-20 kD was detected, corresponding exactly to the size expected if the protein were being cleaved at the same site as the PrP^{Sc} form of the protein that can be recovered from diseased mammalian brains.

[0210] The foregoing data indicates that, when mammalian PrP is expressed in yeast, a species from an entirely different taxonomic kingdom, it behaves unlike common yeast proteins, and very much like the disease form of PrP in mammals.

[0211] Besides the diseased form, a small portion of PrP protein expressed in yeast cytosol also behaves like the normal cellular form of PrP. Even after centrifugation at 180,000 g for 90 minutes, there is still some PrP protein detectable in the supernatant fraction. This part of PrP expressed in yeast, like normal cellular PrP, was soluble in non-ionic detergent, suggesting this small portion of PrP is present in the PrP^C conformation.

Example 5

Assays to Identify Novel Prion-Like Amyloidogenic Sequences

[0212] The following experiments demonstrate how to identify novel prion-like amyloidogenic sequences and confirm their ability to form prions in vivo. The experiments involve (A) identifying sequences suspected of having prion forming capability; and (B) screening the sequences to confirm prion forming ability.

[0213] A. Identifying Sequences Suspected of Having Prion Forming Capability

[0214] Known prion or prion-like amino acid sequences, or polynucleotides encoding such sequences, are used to probe sequence databases or genomic libraries for similar sequences. For example, in one embodiment, a prion or prion-like amino acid sequence (e.g., a mammalian PrP sequence; the N or NM regions from a yeast Sup35 sequence; or the N region from a yeast Ure2 sequence) is used to screen a protein database (e.g., Genbank or NCBI) using a standard search algorithm (e.g., BLAST 1.4.9.MP or more recent releases such as BLAST 2.0, and a default search matrix such as BLOSUM62 having a Gap existence cost of 11, a per-residue gap cost of 1, and a Lambda ratio of 0.85. See generally Altschul et al., *Nucleic Acids Res.*, 25 (17): 3389-3402 (1997)). As an exemplary cutoff, database hits are selected having P(N) less than 4×10^{-6} , where P(N) represents the smallest sum probability of an accidental similarity. For database searching, polypeptide sequences are preferred, but it will be apparent that polynucleotides encoding the amino acid sequences also could be used to probe nucleotide sequence databases.

[0215] In an alternative embodiment, one or more polynucleotides encoding a prion or prion-like sequence is amplified and labeled and used as a hybridization probe to probe a polynucleotide library (e.g., a genomic library, or more preferably a cDNA library) or a Northern blot of purified RNA for sequences having sufficient similarity to hybridize to the probe. The hybridizing sequences are cloned and sequenced to determine if they encode a candidate amino acid sequence. Hybridization at temperatures below the melting point (T_m) of the probe/conjugate complex will allow pairing to non-identical, but highly homologous sequences. For example, a hybridization at 60° C. of a probe that has a T_m of 70° C. will permit ~10% mismatch. Washing at room temperature will

allow the annealed probes to remain bound to target DNA sequences. Hybridization at temperatures (e.g., just below the predicted T_m of the probe/conjugate complex) will prevent mismatched DNA targets from being bound by the DNA probe. Washes at high temperature will further prevent imperfect probe/sequence binding. Exemplary hybridization conditions are as follows: hybridization overnight at 50° C. in APH solution [5×SSC (where 1×SSC is 150 mM NaCl, 15 mM sodium citrate, pH 7), 5×Denhardt's solution, 1% sodium dodecyl sulfate (SDS), 100 µg/ml single stranded DNA (salmon sperm DNA)] with 10 ng/ml probe, and washing twice at room temperature for ten minutes with a wash solution comprising 2×SSC and 0.1% SDS. Exemplary stringent hybridization conditions, useful for identifying interspecies prion counterpart sequences and intraspecies allelic variants, are as follows: hybridization overnight at 68° C. in APH solution with 10 ng/ml probe; washing once at room temperature for ten minutes in a wash solution comprising 2×SSC and 0.1% SDS; and washing twice for 15 minutes at 68° C. with a wash solution comprising 0.1×SSC and 0.1% SDS.

[0216] In another alternative embodiment, known prion sequences or other SCHAG amino acid sequences are modified, e.g., by addition, deletion, or substitution of individual amino acids; or by repeating or deleting motifs known or suspected of influencing fibril-forming propensity. To form novel prion sequences, modifications to increase the number of polar residues (glutamine, asparagine, serine, tyrosine) are specifically contemplated, with modifications that increase glutamine and asparagine content being highly preferred. [See Depace et al., *Cell*, 93:1241-1252 (1998), incorporated herein by reference.] In a preferred embodiment, the alterations are effected by site directed mutagenesis or de novo synthesis of encoding polynucleotides, followed by expression of the encoding polynucleotides.

[0217] In yet another alternative embodiment, antibodies are generated against the prion forming domain of a prion or prion-like protein, using standard techniques. See, e.g., Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1988). The antibodies are used to probe a Western blot of proteins for interspecies counterparts of the protein, or other proteins that possess highly conserved prion epitopes. Candidate proteins are purified and partially sequenced. The amino acid sequence information is used to generate probes for obtaining an encoding DNA or cDNA from a genomic or cDNA library using standard techniques.

[0218] Sequences identified by the foregoing techniques can be further evaluated for certain features that appear to be conserved in prion-like proteins, such as a region of 50 to 150 amino acids near the protein's amino-terminus or carboxyl-terminus that is rich in glycine, glutamine, and asparagine, and possibly the polar residues serine and tyrosine, which region may contain several oligopeptide repeats and have a predicted high degree of flexibility (based on primary structure). In the case of Sup35, a highly charged domain separates the flexible N-terminal region having these properties from the functional C-terminal domain. Sequences possessing one or more of these features are ranked as preferred prion candidates for screening according to techniques described in the following section.

[0219] By way of example, the Genbank protein database (accessible via the worldwide web at www.ncbi.nlm.nih.gov) was screened using the Basic Local Alignment Search Tool (BLAST) program (version 1.4.9) using the standard (de-

fault) matrix and stringency parameters (BLOSUM62). The prion forming domains of Ure2 (Genbank Acc. No. M35268, SEQ ID NO: 4, amino acids 1-65) and Sup35 (Genbank Acc. No. M21129, SEQ ID NO: 2, amino acids 1-114) from *S. cerevisiae* were used as BLAST query sequences. Open reading frames (ORFs) from *S. cerevisiae* with high similarity scores [P(N) less than 4×10^{-6}] resulting from the initial search included the following Genbank database entries:

[0220] (1) residues 53-97 from Accession No. Z73582 (SEQ ID NO: 22), an uncharacterized open reading from *S. cerevisiae*;

[0221] (2) residues 1030-1071 from PID No. e236901, in Accession No. Z71255 (SEQ ID NO: 23), an uncharacterized open reading from *S. cerevisiae*;

[0222] (3) residues 4-58 from locus ybm6, Accession No. P38216 (SEQ ID NO: 24), an uncharacterized open reading from *S. cerevisiae*;

[0223] (4) residues 251-380 from locus hrp1, Accession No. U35737 (SEQ ID NO: 25), an RNA binding and transport protein having homology to hnRNP1 in humans.

[0224] (5) residues 28-126 from locus npl3, Accession No. U33077 (SEQ ID NO: 26), an RNA binding and transport protein that functions genetically in the same pathway as Hrp1;

[0225] (6) residues 97-286 from locus mcm1, Accession No. X14187 (SEQ ID NO: 27), a DNA binding protein active in cell cycle regulation and mating-type specificity;

[0226] (7) residues 205-414 from locus nsr1, Accession No. P27476 (SEQ ID NO: 28), a protein that binds nuclear localization sequences and is active in mRNA processing;

[0227] (8) residues 153-405 from Accession No. P25367 (SEQ ID NO: 29), an uncharacterized open reading frame;

[0228] (9) residues 806-906 from Accession No. P40467 (SEQ ID NO: 30), an uncharacterized open reading frame;

[0229] (10) residues 605-677 from Accession No. S54522 (SEQ ID NO: 31), an uncharacterized open reading frame;

[0230] (11) residues 100-300 from locus yk76, Accession No. P36168 (SEQ ID NO: 32), an uncharacterized open reading frame;

[0231] (12) residues 1 to 250 from locus fps1, Accession No. S16712 (SEQ ID NO: 33), a membrane channel protein that controls passive efflux of glycerol;

[0232] (13) residues 334-388 from Accession No. p40002 (SEQ ID NO: 34), an uncharacterized open reading frame;

[0233] (14) residues 325-375 from locus mad1, Accession No. P40957 (SEQ ID NO: 35), an uncharacterized open reading frame; and

[0234] (15) residues 215-284 from locus kar1, Accession No. M15683 (SEQ ID NO: 36), an uncharacterized open reading frame.

[0235] The nuclear polyadenylated RNA-binding protein hrp1 (Genbank Accession No. U35737) is an especially promising prion candidate. It is the clear yeast homologue of a nematode protein previously cloned by cross-hybridization with the human PrP gene; it scored highly (p value 3.9×10^{-5}) in a Genbank BLAST search for sequences having homology to the N-terminal domain of Sup35; and it contains a stretch of 130 amino acids at its C-terminus that is glycine- and asparagine-rich and contains repeat sequences similar to the oligomeric repeats in the N-terminal domain of Sup35; and is predicted by secondary structure programs to consist entirely of turns.

[0236] The sequence corresponding to residues 153-405 of SEQ ID NO: 29 comprises another promising prion candi-

date. This region is rich in glutamine and asparagine, and is part of a protein that is normally found in aggregates in yeast although it is not aggregated in some strains. When expressed as a fusion protein with green fluorescent protein, this sequence causes the GFP to aggregate. This aggregation is completely dependent upon Hsp104, much the same as Sup35 aggregation. When residues 153-405 of SEQ ID NO: 29 are substituted for the NM region of SUP35 and transformed into [psi-] yeast, the yeast exhibit a suppression phenotype analogous to [PSI⁺].

[0237] B. Screening Sequences to Confirm Prion-Forming Capability.

[0238] Sequences identified according to methods set forth in Section A are screened to determine if the sequences represent/encode proteins having the ability to aggregate in a prion-like manner.

[0239] 1. Aggregation Assay Using Fusion Proteins

[0240] In a preferred screening technique, a polynucleotide encoding the ORF of interest is amplified from DNA or RNA from a host cell using polymerase chain reaction, or is synthesized using the well-known universal genetic code and using an automated synthesizer, or is isolated from the host cell of origin. The polynucleotide is ligated in-frame with a polynucleotide encoding a marker sequence, such as green fluorescent protein or firefly luciferase, to create a chimeric gene. In a preferred embodiment, the polynucleotide is ligated in frame with a polynucleotide encoding a fusion protein such as a Bleomycin/luciferase fusion, which would permit both selection for drug-resistance and quantification of soluble and insoluble proteins by enzymatic assay. See, e.g., Elgersma et al., *Genetics*, 135: 731-740 (1993).

[0241] The chimeric gene is then inserted into an expression vector, preferably a high-copy vector and/or a vector with a constitutive or inducible promoter to permit high expression of the ORF-marker fusion protein in a suitable host, e.g., yeast. The expression construct is transformed or transfected into the host, and transformants are grown under conditions that promote expression of the fusion protein. Depending on the marker, the cells may be analyzed for marker protein activity, wherein absence of marker protein activity despite the presence of the marker protein is correlated with a likelihood that the ORF has aggregated, causing loss of the marker activity. Alternatively, host cells or host cell lysates are analyzed to determine if the fusion protein in some or all of the cells has aggregated into aggregates such as fibril-like structures characteristic of prions. The analysis is conducted using one or more standard techniques, including microscopic examination for fibril-like structures or for coalescence of marker protein activity; analysis for sensitivity or resistance to protease K; spectropolarimetric analysis for circular dichroism that is characteristic of amyloid proteins; and/or Congo Red dye binding.

[0242] A number of the candidates identified above were screened in this manner using a GFP fusion construct. To create the vector that was employed in these analyses, a copper inducible Cup1 promoter was amplified from a genomic library by standard polymerase chain reaction (PCR) methods using the primers 5'-GG GAATTCCAATTACCGACATTTGGGCGC-3' (SEQ ID NO: 37) and 5'-GG GGAATCCTGTATTGATTGATTGATTGTAC-3' (SEQ ID NO: 38), digested with the restriction enzymes EcoRI and BamHI, and ligated into the pRS316 vector that had digested with EcoRI and BamHI. The annealed vector, designated

pRS316Cup1, was transformed into *E. Coli* strain AG-1, and transformants were selected using the ampicillin resistance marker of the vector. Correctly transformed bacteria were grown overnight to provide DNA for further vector construction.

[0243] Next, a sequence encoding superbright GFP (SEQ ID NOs: 39, 40) was inserted into the pRS316Cup1 vector. Superbright GFP was amplified from pPSGFP using the primers 5'-GACCGCGGATGGCTAGCAAAGGAGAAG-3' (SEQ ID NO: 41) and 5'-CCTGAGCTCTCATTTGTATAGTTCATCC-3' (SEQ ID NO: 42). The resultant PCR products were digested with SacI and SacII and inserted into PRS316Cup1 that also had been digested with SacI and SacII. This created a pRS316Cup1 GFP plasmid into which a polynucleotide encoding a candidate open reading frame could be inserted for expression studies. In particular, it was contemplated that candidate open reading frames be amplified by PCR from genomic DNA or cDNA using primers engineered to contain BamHI and SacII restriction sites, to permit rapid cloning into the BamHI and SacII sites of the derived PRS316Cup1 GFP vector. For example, in the case of open reading frame (ORF) P25367 the following primers were used: 5'-GGA GGATCCATGGATACGGATAAGTTAATCTCAG-3' (SEQ ID NO: 43, BamHI site underlined) and 5'-GGA CCGCGGTAGCGGTTCTGTTGAGAAAAGTTGCC-3' (SEQ ID NO: 44, SacII site underlined). PCR products were digested with BamHI and SacII and inserted into the derived plasmid. This created a plasmid that can inducibly express a fusion of an open reading frame of interest fused to GFP. The sequence of pRS316-Cup1-p25367-GFP is set forth in SEQ ID NO: 45.

[0244] 2. In Vitro Aggregation Assay Using Chaperone Protein

[0245] A polynucleotide encoding the ORF of interest is synthesized using the well-known universal genetic code and using an automated synthesizer, or is isolated from the host cell of origin, or is amplified using polymerase chain reaction from DNA or RNA from such a host cell. In a preferred embodiment, the polynucleotide further includes a sequence encoding a tag sequence, such as a polyhistidine tag, HA tag, or FLAG tag, to facilitate purification of the recombinant protein. The polynucleotide is inserted into an expression vector and expressed in a host cell compatible with the selected vector, and the resultant recombinant protein is purified.

[0246] Serial dilutions of the recombinant polypeptide (e.g. 100 mM, 10 mM, 1 mM, 0.1 mM, 0.01 mM final concentration) are mixed with 1 µg of a chaperone protein such as yeast Hsp104 protein [See Schirmer and Lindquist, *Meth. Enzymol.*, 290: 430-444 (1998)] in a low salt buffer (e.g., 10 mM MES, pH 6.5, 10 mM MgSO₄) containing 5 mM ATP in a 25 µl reaction volume. As controls, reactions are performed in parallel using buffer alone or using Sup35 protein. Reactions are incubated at 37° C. for eight minutes, and the ATPase activity of the chaperone protein is measured by determining released phosphate, e.g., using Malachite Green [Lanzetta et al., *Analyt. Biochem.*, 100: 95-97 (1979)]. In this assay, several fibril-aggregation proteins, including yeast Sup35, the yeast Sup35 N terminal domain, mammalian PrP protein, and β-amyloid (1-40) and (1-42) forms, were found to inhibit the ATPase activity of Hsp104; whereas control proteins (aldolase, BSA, apoferritin, and IgM) did not.

[0247] 3. Assay Results

[0248] To determine if the proteins represented by the ORF's identified above in part A were aggregation prone, a hallmark of prions, polynucleotides encoding the specified residues of interest within the ORF's were amplified from *S. cerevisiae* genomic DNA via PCR and ligated in-frame to a sequence encoding superbright, as described above in section B.1.

[0249] These plasmids were transformed into the yeast strain 74D (a, his, met, leu, ura, ade). Transformant colonies were selected (ura+) and inoculated into liquid SD ura and grown to early log phase. Copper sulfate was added to the cultures (final concentration 50 μ M copper) to induce protein expression. Cells were fixed after four hours of induction and intracellular GFP expression was visualized.

[0250] Examination of GFP fluorescence revealed that the sGFP tag had coalesced in transformants expressing six of the ORF's. This coalescence was similar to that observed with Sup35-GFP fusions in [PSI⁺] yeast and was considered to be indicative of an ORF having prion-like aggregate-forming ability. Two of the positive sequences represent uncharacterized open reading frames: Z73582 and ybm6. Four are known proteins: mcm1, fps1, p25367 and hrp1 as described above in section B.1. Aggregation of the MCM1-GFP fusion was relatively rare, and was not influenced by Hsp104 dosage in the cells. Of particular interest was the hrp1 construct, which aggregated into multiple cytoplasmic points in the transformed *S. cerevisiae*, and also in transformed *C. elegans*. Deletion of the Hsp104 gene was shown to eliminate the aggregation pattern of hrp1. Also of special interest was the aggregation pattern of the P25367 construct, because this aggregation was completely eliminated by overexpression of Hsp104.

[0251] The foregoing experiments demonstrate that searches with prion forming sequences will identify additional sequences with prion-like properties, which sequences can be used according to various aspects of the invention that are specifically exemplified herein with respect to Sup35 or URE2 sequences.

[0252] The ability of newly identified aggregating proteins to exist in both an aggregating and non-aggregating conformational state can be further examined, if desired, by studying aggregation phenomena in host cells expressing varying levels of the protein (a result achieved using an inducible promoter, for example), and in host cells having normal and over- or under-expressed chaperone protein levels. (The ability of Sup35 in yeast to enter a [PSI⁺] conformation depends on an appropriate intermediate level of the chaperone protein Hsp104; elimination of Hsp104 or over-expression of Hsp104 causes loss of [PSI⁺] and prevents de novo appearance of [PSI⁺]. See Chernoff et al., *Science*, 268: 880 (1995) and Patino et al., *Science*, 273: 622-626 (1996). Growth on a mildly denaturing media, as described elsewhere herein, provides another alternative assay.

[0253] The foregoing assays, chimeric constructs, and candidate SCHAG amino acid sequences are all intended as aspects of the invention.

Example 6

Identification of Rnq1 as an Epigenetic Modifier of Protein Function in Yeast

[0254] The following experiments demonstrate that putative prions can be identified by searching for three attributes

of the known yeast prion proteins: unusual amino-acid composition with a high concentration of the polar amino-acid residues glutamine and asparagine, constant expression levels through log and stationary phase growth, and a capacity to switch between distinct stable physical states (in this case, insoluble and soluble forms). One of the candidates isolated in this search, Rnq1, has both in vitro and in vivo characteristics of a prion. Rnq1, exists in distinct, heritable physical states, soluble and insoluble. The insoluble state is dominant and transmitted between cells through the cytoplasm. When the prion-like region of Rnq1 was substituted for the prion domain of Sup35, the protein determinant of the prion [PSI⁺], the phenotypic and epigenetic behavior of [PSI⁺] was fully recapitulated. These findings identify Rnq1 as a prion, demonstrate that prion domains are modular and transferable, and establish a paradigm for identifying and characterizing novel prions.

[0255] A. Identification of Prion Candidates

[0256] The characteristics of Sup35 and Ure2 suggested several criteria for identifying new prion candidates. Previous experiments have demonstrated that particular regions (residues 1-65 for Ure2 (Genbank Acc. No. M35268, SEQ ID NO: 4) and residues 1-123 for Sup35 (Genbank Acc. No. M21129, SEQ ID NO: 2)) are critical for prion formation by these proteins. Over-expression of these regions is sufficient to induce the prion phenotype de novo. Deletion of these regions has no effect upon the normal cellular function of the proteins but prevents them from entering the prion state. These critical prion-determining domains have an unusually high concentration of the polar residues glutamine and asparagine and are predicted to have very little secondary structure. The domains are located at the ends of proteins that have an otherwise ordinary amino acid composition. We hypothesized that by searching for open reading frames with these characteristics we might find new prion proteins.

[0257] A BLAST search (1.4.9 MP version) of the NCBI database of non-redundant coding sequences was performed using the prion-determining domains of Ure2 and Sup35 (residues 1-65 of SEQ ID NO: 4 and residues 1-123 of SEQ ID NO: 2, respectively) as the query sequence with the following parameters: V=100, B=50, H=0, S=90, and P=4. This search revealed approximately twenty open reading frames that had prion-like domains appended to polypeptides with an otherwise normal amino acid composition. To restrict the number of likely candidates, we took advantage of recent global descriptions of mRNA expression patterns. In examining this data we noted that Sup35 and Ure2 are expressed at nearly constant levels as cells transit from the log to the stationary phase of growth. Large fluctuations in expression would be inconsistent with the stability of both their heritable prion and non-prion states. The open reading frames from the BLAST search whose expression varies by less than two-fold in the log phase transition were selected for further analysis. They were fused to the coding sequence of green fluorescent protein (GFP) using PCR and expressed in the yeast strain 74D-694 (ade1-14, trp1-289, his3-200, ura3-52, leu2-3, lys2). Three of the proteins, RNQ1 (Genbank Acc. No. NP009902, SEQ ID NO: 50), YBR016w (Genbank Acc. No. NP009572, SEQ ID NO: 51), and HRP1 (Genbank Acc. No. NP014518, SEQ ID NO: 52), showed coalescence of GFP, as previously described for Sup35.

[0258] B. Rnq1 Exists in Distinct States Controllable by Hsp104

[0259] We next asked if expression of the fusion protein in a strain that lacked the chaperone Hsp104 eliminated the coalescence of GFP, as it does for Sup35-GFP fusions. This is not a necessary criterion for prion proteins (an interaction with Hsp104 has not been demonstrated for [URE3]) but interaction with the chaperone provides a useful tool for further analysis. In wild-type yeast, fluorescence from the Rnq1-GFP fusion was found in one or more small, intense, cytoplasmic foci. When the fusion protein was expressed in the isogenic hsp104 strain, fluorescence was diffuse. The C-terminal end of Rnq1 (amino acids 153-405 of SEQ ID NO: 50) contained the region rich in glutamine and asparagine residues. Fusion of this region alone to GFP gave an identical result to that seen with the full length Rnq1-GFP fusion. Since the effect of HSP104 deletion upon the coalescence of the Rnq1 fusion was the most dramatic, it was chosen for further analysis.

[0260] Differential centrifugation was employed to determine if the coalescence observed with Rnq1-GFP fusion proteins reflected the behavior of the endogenous Rnq1 protein. Log phase yeast were lysed using a bead beater (Biospec) into 75 mM Tris-Cl (pH7), 200 mM NaCl, 0.5 mM EDTA, 2.5% glycerol, 0.25 mM EDTA, 0.25% Na-deoxycholate, supplemented with protease inhibitors (Boehringer-Mannheim). Lysates were cleared of crude cellular debris by a 15 second 6000 RPM spin in a microcentrifuge (Eppendorf). Non-denatured total cellular lysates were fractionated by high-speed centrifugation into supernatant and pellet fractions using a TLA-100 rotor on an Optima TL ultracentrifuge (Beckman) at 280,000xg (85,000 RPM) for 30 minutes. Protein fractions were resolved by 10% SDS-PAGE and immunoblotted with an α -Rnq1 antibody. Rnq1 remained in the supernatant of a hsp104 strain, but pelleted in the wild-type. Thus, the GFP coalescence is not an artifact of the fusion; the Rnq1 protein itself is sequestered into an insoluble aggregate in an Hsp104-dependent fashion. We also examined the solubility of Rnq1 in several unrelated yeast strains. In four (S288c, YJM436, SK1 and W303) the protein fractionated in the pellet, in two (YJM128, YJM309) it partitioned between the pellet and supernatant fractions, and in two others (33G, 10B-H49) the protein was chiefly recovered in the supernatant fraction. Thus, Rnq1 naturally exists in distinct physical states in different strains.

[0261] C. The insoluble state of Rnq1 is Transmitted by Cytoduction

[0262] The heritability of the known yeast prions is based upon the ability of protein in the prion state to influence other protein of the same sequence to adopt the same state. Because the protein is passed from cell to cell through the cytoplasm, the conformational conversion is heritable, dominant in crosses, and segregates in a non-Mendelian manner. To determine if the insoluble state of Rnq1 is transmissible in this way, we used cytoduction, a well-established tool for the analysis of the [PSI⁺] and [URE3] prion. The karyogamy deficient (kar1-1) strain 10B-H49 (ade2-1, lys1-1, his3-11,15, leu2-3, 112, kar1-1, ura3::KANR) can undergo normal conjugation between a and cells but is unable to fuse its nucleus with its mating partner. Cytoplasmic proteins and organelles are mixed in fused cells, but the haploid progeny that bud from them contain nuclear information from only one of the two parents.

[0263] 10B-H49 shows diffuse expression of Rnq1-GFP, and served as the recipient for the transfer of insoluble Rnq1 from W303 (Mata, his3-11, 15, leu2-3, 112, trp1-1, ura3-1, ade2-1), the donor. After cytoduction, colonies derived from haploid cells that contained the 10B-H49 nuclear genome but had undergone cytoplasmic mixing, as demonstrated by mitochondrial transfer, were selected. Cytoeductants were selected after overnight mating on defined media lacking tryptophan that had glycerol as the sole carbon source. All showed single or multiple cytoplasmic aggregates of Rnq1-GFP—a pattern indistinguishable from that of the W303 parent. Furthermore, density-based centrifugation of protein extracts, performed as above, indicated that cytoduction caused the endogenous Rnq1 protein of the 10B-H49 strain to shift from the soluble to the insoluble fraction. Thus exposure of 10B-H49 cells to the cytoplasm of W303 is sufficient to cause a heritable change in the physical state of Rnq1. Because RNQ1 is a nuclear gene (not transmitted during cytoduction) the protein's insoluble state is not due to polymorphisms in its amino acid sequence, nor to any other trait carried by the W303 genome. Rather, like the Sup35 and Ure2 prions, its altered conformational state is “infectious”, transmissible from one protein to another.

[0264] D. Purified Rnq1 Forms Fibers and Shows Seeded Polymerization

[0265] Both Sup35 and Ure2 have the capacity to form highly ordered amyloid fibers in vitro, as analyzed by the binding of amyloid specific dyes and by electron microscopy. To examine conformational transitions of Rnq1 in vitro, the protein was expressed in *E. coli* and studied as a purified protein. Rnq1 was cloned into pPROEX-HTb (GibcoBRL). The primers 5'-GGA GGA TCC ATG GAT ACG GAT AAG TTA ATC TCAG-3' (SEQ ID NO: 53) and 5'-CC AAG CTT TCA GTA GCG GTT CTG TTG AGA AAA GTTG-3' (SEQ ID NO: 54) were used for PCR in a solution containing 10 mM Tris (pH8.3), 50 mM KCl, 2.5 mM MgCl₂, 2 mM dNTPs, 1 μ M of each primer and 2 U of Taq polymerase; and using genomic 74D DNA as template under the following conditions: incubation at 94° C. for 2 min, followed by 29 cycles of 94° C. for 30 sec, 50° C. for 30 sec, and 72° C. for 90 sec, followed by a final incubation at 72° C. for 10 minutes. The PCR product was then digested and ligated into the BamHI and HindIII sites of pPROEX-HTb (GibcoBRL). The plasmid was electroporated into BL21-DE3 lacIq cells. Transformed bacterial cultures were induced at OD₆₀₀=1 with 1 mM IPTG for four hours at 30 C. The cells were lysed in 8M urea (Rnq1 was purified under denaturing conditions (8M urea) because it had a tendency to form gels during purification in the absence of denaturant), 20 mM Tris-Cl pH8. Protein was purified over a Ni-NTA column (Qiagen) followed by Q-sepharose (Pharmacia). The (His)₆-tag from the vector was cleaved under native conditions (150 mM NaCl, 5 mM KPi) using TEV protease followed by passage of the protease product over a Ni-NTA column to remove uncleaved protein. Protein was methanol precipitated prior to use. Recombinant protein was resuspended in 4M urea, 150 mM NaCl, 5 mM KPi, pH 7.4 at a concentration of 10 μ M. Seeded samples were created by sonication of 1/50 volume of a 10 μ M solution of pre-formed fibers verified by electron microscopy. The protein samples were incubated at room temperature on a wheel rotating at 60 r.p.m.

[0266] To determine if Rnq1 forms amyloids we used Thioflavin T fluorescence. This dye exhibits an increase in fluorescence and a red-shift in the λ_{max} of emission upon

binding to multimeric fibrillar β -sheet structures characteristic of many amyloids, including transthyretin, insulin, β -2 microglobulin and Sup35. Fluorimeter samples were prepared as 3.3 μ M Rnq1, 50 μ M Thioflavin T in buffer. Samples were analyzed on a Jasco FP750 with the following settings; λ_{exc} =409 nm, λ_{emi} =484 nm, bandwidth 10 nm. The acquisition of Thioflavin T binding was sigmoidal (lag phase-six) suggesting a self-seeded process of protein assembly. The addition of 2% preformed fibers to fresh solutions of Rnq1 reduced the lag time—from 6.40.2 hrs to 4.30.2 hrs (n=4).

[0267] The formation of higher ordered structures was confirmed by transmission electron microscopy. For electron microscopy analysis, 5 μ l of a 10 μ M protein solution was placed on a 400 mesh carbon coated EM grid (Ted Pella, Cat. 01822), and allowed to adsorb for 1 minute. The sample was negatively stained with 200 μ l of 2% aqueous uranyl acetate, and wicked dry. Samples were observed in a Philips CM120 transmission electron microscope operating at 120 kV in low dose mode. Micrographs were recorded at a magnification of 45,000 on Kodak SO-163 film. The protein formed fibers with a diameter of 11.3–1.4 nm. This figure is comparable to the reported range for Ure2 (~20 nm) and Sup35 (~17 nm) fibers. The fibers appeared to be branching and the termini were unremarkable. The appearance of the fibers was coincident with the onset of rapid increases in Thioflavin T fluorescence.

[0268] E. Rnq1 Disruption

[0269] [URE3] and [PSI⁺] produce phenotypes that mimic loss-of-function mutations in their protein determinants. To determine the loss of function phenotype of Rnq1, the entire ORF was deleted by homologous recombination in a diploid 74D-694 strain using a kanamycin resistance gene. Strains deleted of the Rnq1 open reading frame were created using the long flanking homology PCR method. Primers 5'-GGT GTC TTG GCC AAT TGC CC-3' (SEQ ID NO: 55) and 5'-GTC GAC CTG CAG CGT ACG CAT TTC AGA TCT TTG CTA TAC-3' (SEQ ID NO: 56) or 5'-CGA GCT CGA ATT CAT CGA TTG ATT CAG TTC GCC TTC TATC-3' (SEQ ID NO: 57) and 5'-CTG TTT TGA AAG GGT CCA CATG-3' (SEQ ID NO: 58) were used to amplify genomic DNA. These PCR products were used as primers for a second round of PCR on plasmid pFA6a, which is described in Wach et al., *Yeast* 13:1065-75 (1994), digested with NotI. The product of the second PCR round was used to transform log-phase yeast cultures. Transformants were selected on YPD containing 200 mg/mL G418 (GibcoBRL). Upon sporulation each tetrad produced four viable colonies, two of which contained the Rnq1 disruption, confirmed by immunoblotting total cellular proteins with an -Rnq1 antibody and PCR analysis of the genomic region. The rnq1 strain had a growth rate comparable to that of wild-type cells on a variety of carbon and nitrogen sources and was competent for mating and sporulation. The strain grew similarly to the wild-type in media with high and low osmolarity, and in assays testing sensitivity to various metals (cadmium, cobalt, copper).

[0270] F. Fusion of Rnq1 (153-405) to Sup35 (124-685)-Nonsense Suppression Phenotype

[0271] The lack of an obvious loss-of-function phenotype was not unexpected, as the two known yeast prions, [URE3] and [PSI⁺] only exhibit phenotypes under unusual selective conditions. However, the absence of a phenotype presented difficulties in determining whether Rnq1 could direct the epigenetic inheritance of a trait. To determine if the prion-like domain of Rnq1 could produce an epigenetic loss-of-function phenotype we asked if it could replace the prion-determining

domain of Sup35. When the wild-type Sup35 translation termination factor enters the prion state the loss-of-function phenotype it produces is nonsense suppression—the readthrough of stop codons. This phenotype can be conveniently assayed in the strain 74D-694 because it contains a UGA stop codon in the ADE1 gene. In [psi⁻] 74D-694 cells, ribosomes efficiently terminate translation at this codon. Cells are therefore unable to grow on media lacking adenine (SD-ade), and colonies appear red on rich media due to the accumulation of a pigmented by-product. In [PSI⁺] strains, sufficient readthrough occurs to support growth on SD-ade and prevent accumulation of the pigment on rich media.

[0272] The coding region for amino acid residues 153-405 of Rnq1 (amino acid residues 153-405 of SEQ ID NO: 50) was substituted for 1-123 of Sup35 and the resulting fusion gene, RMC, was inserted into the genome in place of the endogenous SUP35 gene. RNQ1, SUP35 and its promoter were cloned by amplification of 74D-694 genomic DNA. The RNQ1 open reading frame was cloned using 5'-GGA GGA TCC ATG GAT ACG GAT AAG TTA ATC TCAG-3' (SEQ ID NO: 59) and (A) 5'-GGA CCG CGG GTA GCG GTT CTG TTG AGA AAA GTT GCC-3' (SEQ ID NO: 60). RNQ1 (153-405) was cloned using 5'-GA GGA TCC ATG CCT GAT GAT GAG GAA GAA GAC GAGG-3' (SEQ ID NO: 61) and (A). The SUP35 promoter was cloned using 5'-CG GAA TTC CTC GAG AAG ATA TCC ATC-3' (SEQ ID NO: 62) and 5'-G GGA TCC TGT TGC TAG TGG GCA GA-3' (SEQ ID NO: 63). SUP35 (124-685) was cloned using 5'-GTA CCG CGG ATG TCT TTG AAC GAC TTT CAA AAGC-3' (SEQ ID NO: 64) and 5'-GTG GAG CTC TTA CTC GGC AAT TTT AAC AAT TTT AC-3' (SEQ ID NO: 65) by PCR using the conditions described above in section D.

[0273] The RMC gene replacement was performed as described in Rothstein, 1991. To create the plasmid for pop-in/pop-out replacement in pRS306 (available from ATCC), the SUP35 promoter was ligated into the EcoRI-BamHI site, RNQ1 (153-405) was ligated into the BamHI-SacII site, and SUP35 (124-685) was ligated into the SacII-SacI site. To create the disrupting fragment, this plasmid was linearized with MluI and transformed. Pop-outs were selected on 5-FOA (Diagnostic Chemicals Ltd.) and verified by PCR. The resulting strain, RMC, had a growth rate similar to that of wild-type cells on YPD, although the accumulation of red pigment was not as intense as seen in [psi⁻] strains. RMC strains showed no growth on SD-ade even after 2 weeks of incubation). Thus, the protein encoded by the RMC gene (Rmc) fulfilled the essential translational termination function of Sup35.

[0274] At a low frequency, RMC variants appeared that were white on rich media and grew on SD-ade even more robustly than [PSI⁺] cells did. The frequency at which these variants appeared (~10⁻⁴) was far greater than expected for reversion of the UGA stop codon mutation in ade1-14, and subsequent analysis demonstrated that the allele had not reverted. The suppressor phenotype of these variants was comparable in stability to that of [PSI⁺]. Because Sup35 proteins that lack residues 1-123 are incapable of making such conversions, these observations suggest that the Rnq1 prion-like domain can direct a prion conversion in the Rmc fusion protein.

[0275] Transient over-expression of Sup35 can produce new [PSI⁺] elements, because higher protein concentrations make it more likely that a prion conformation will be achieved. To test whether over-expression of Rmc can pro-

duce heritable suppressing variants, the original, non-suppressing RMC strain was transformed with an expression plasmid for RMC. These transformants showed a greatly elevated frequency of conversion to the suppressor state compared to control strains carrying the plasmid alone. Once a prion conformation is achieved it should be self-perpetuating and normal expression should then be sufficient for maintenance. When the RMC expression plasmid was lost all strains retained the suppressor phenotype. Thus, transient over-expression of Rmc produced a heritable change in the fidelity of translation termination.

[0276] G. Non-Mendelian Segregation of Rmc-Based Suppression Phenotype

[0277] To examine the genetic behavior of the suppressor phenotype in RMC strains, an isogenic mating partner was created from a non-suppressing RMC strain. When this strain was crossed to the original, non-suppressing, RMC strain, neither the diploids nor their haploid meiotic progeny exhibited the suppressor phenotype. However, when this strain was mated to RMC suppressor strains, the resulting diploids all displayed the suppressor phenotype, demonstrating that suppression is dominant. In fourteen tetrads dissected from two different diploids of this cross, all four haploid progeny showed inheritance of the suppression phenotype, instead of the 2:2 segregation expected for a phenotype encoded in the nuclear genome. Following convention, we henceforth refer to the dominant, non-Mendelian suppressor phenotype as [RPS⁺] (for Rnq1 [PSI⁺]-like Suppression) and the non-suppressed phenotype as [rps⁻].

[0278] To determine if the dominant, non-Mendelian [RPS⁺] phenotype arises from the ability of Rmc protein to form a prion, we tested it for two additional unusual genetic behaviors that are not expected for other non-Mendelian genetic elements, such as viruses or mitochondrial genomes. First, it should become recessive and Mendelian in crosses to strains carrying a wild-type Sup35 allele. This is because Sup35 lacks the Rnq1 sequences that would allow it to be incorporated into an [RPS⁺] prion. Wild-type Sup35, therefore, should cover the impaired translation-termination phenotype associated with the [RPS⁺] prion. However, even when this phenotype has disappeared, Rmc protein in the prion state should still convert new Rmc protein to the same state. Therefore, in haploid meiotic progeny of this diploid, the phenotype will reappear in segregants carrying the RMC gene, but not in segregants carrying the SUP35 gene (2:2 segregation).

[0279] Indeed, diploids of a cross between an [RPS⁺] strain and an isogenic strain with a wild-type SUP35 gene did not exhibit a suppressor phenotype. Upon sporulation, suppression reappeared in only two of the four progeny. By PCR genotyping, these strains had the RMC gene at the SUP35 locus. Thus the [RPS⁺] factor had been preserved in the diploid, even though the phenotype had become cryptic.

[0280] Second, maintenance of [RPS⁺] should depend upon continued expression of the Rmc protein. Although [RPS⁺] is maintained in a cryptic state in diploids with a wild-type Sup35 gene, it should not be maintained in their haploid progeny whose only source of translational termination factor is wild-type Sup35. To determine if these progeny harbored the [RPS⁺] element in a cryptic state, they were mated to an [rps⁻] RMC strain whose protein would be converted if [RPS⁺] were still present. When this diploid was sporulated, none of the progeny exhibited the suppressor

phenotype. Thus, the [RPS⁺] element was not maintained in a cryptic state unless the Rmc protein was present.

[0281] H. Curing of [RPS⁺]

[0282] One of the hallmarks of yeast prions is that cells can be readily and reversibly cured of them. [PSI⁺] is curable by several means, including growth on media containing low concentrations of the protein denaturant guanidine hydrochloride and transient over-expression or deletion of the protein remodeling factor HSP104.

[0283] Strains carrying [RPS⁺] were passaged on medium containing 2.5 mM guanidine hydrochloride (GdnHCl) (Fluka) and then plated to YPD and to SD-ade to assay the suppressor phenotype. Cells passaged on GdnHCl no longer displayed the [RPS⁺] phenotype, while cells not treated with GdnHCl retained it. [RPS⁺] was also lost when the HSP104 gene was deleted by homologous recombination, performed using the same strategy as described above in section E, or when HSP104 was over expressed from a multicopy plasmid using the constitutive GPD promoter. Cells that had been cured of [RPS⁺] by over-expression of HSP104 were passaged on YPD medium to isolate strains that had lost the over-expression plasmid. These strains remained [rps⁻]. Thus transient over-expression of HSP104 is sufficient to heritably cure cells of [RPS⁺].

[0284] Finally, we asked if Hsp104-mediated curing was reversible. Cells cured by over-expression of HSP104 were re-transformed with a plasmid bearing a single copy of RMC. To create the single-copy RMC plasmid in pRS316 (available from ATCC) the ClaI-SacI fragment (includes promoter and RMC) from the plasmid used above for the RMC gene replacement was ligated into the ClaI-SacI site. Transformants were then plated onto SD-ade to assess the rate at which they converted to the [RPS⁺] suppressor phenotype. [RPS⁺] was regained at a rate comparable to that seen in the parental RMC strain, indicating that the transient over-expression of HSP104 caused no permanent alteration in susceptibility to [RPS⁺] conversion.

[0285] I. Effect of Endogenous Rnq1 Upon [RPS⁺]

[0286] To determine if [RPS⁺] can act as an independent genetic element, the gene encoding the endogenous Rnq1 protein was deleted in strains carrying the RMC replacement of SUP35 using methods described above. The deletion had no effect upon the maintenance of the [RPS⁺] suppression phenotype. Growth on SD-ade was equally robust in [RPS⁺] and [RPS⁺] rnq1 strains. This indicates that Rmc can behave as an independent prion and is not dependent upon pre-existing Rnq1 in an insoluble state.

[0287] J. Physical State of the Rmc Protein in [RPS⁺] and [rps⁻] Strains

[0288] Finally, we examined the localization of the Rmc fusion protein in the [RPS⁺] and [rps⁻] strains. Both strains were transformed with inducible plasmids that provided Rnq1(153-405)-GFP expression that were constructed as described above in section A. Strains that lacked the endogenous Rnq1 gene were used to prevent the GFP marker from localizing to the endogenous Rnq1 aggregate. Short-term expression of the GFP-fusion protein prevented the formation of new [RPS⁺] elements in the [rps⁻] strain.

[0289] Two distinct patterns of Rmc protein localization were revealed by this assay and these correlated with the phenotypic differences between [RPS⁺] and [rps⁻] strains. In the non-suppressing [rps⁻] strains, the Rnq1(153-405)-GFP label was diffuse. In the suppressing [RPS⁺] strains, fluorescence was punctate, and was excluded from the nucleus. This

punctate pattern was different from that observed with the endogenous Rnq1 aggregates, as Rmc aggregates are numerous and very small.

[0290] Collectively, the foregoing experiments demonstrate that Rnq1, which was identified based on sequence analysis, exhibits prion-like behavior in numerous *in vitro* and *in vivo* assays. The search method used here shows that putative prions can be identified by a directed prion search rather than by the study of a pre-existing phenotype. In addition, this method will be applicable to the identification of prion proteins in many other organisms. Our demonstration that a new prion protein domain can substitute for that of another well-characterized prion, reproducing its phenotypic characteristics and epigenetic mode of inheritance, also provides a crucial tool in the analysis of uncharacterized candidates.

[0291] We have shown that Rnq1 exists in distinct physical states—soluble and insoluble—in unrelated yeast strains. The insoluble state can be transmitted through cytoduction, and once transmitted is stably inherited. When the N-terminal prion-determining region of SUP35 was replaced with the C-terminal domain of RNQ1, the hybrid Rmc protein provided translation termination activity, mimicking the phenotype of [psi⁻] strains. At a low spontaneous frequency, the strain acquired a stable, heritable suppressor phenotype, [RPS⁺], which mimicked the phenotype of [PSI⁺] strains. Suppression was dominant and segregated to meiotic progeny in non-Mendelian ratios. The possibility that this phenotype is caused by an epigenetic factor unrelated to the fusion protein was ruled out by genetic crosses showing that the phenotype is not expressed and can not be transmitted in strains that do not produce the fusion protein. The relationship of the suppression phenotype to protein conformation was further demonstrated by fluorescence localization of the hybrid protein in isogenic [RPS⁺] and [rps⁻] strains. In [RPS⁺] strains, most of the protein is sequestered into small foci and is presumably inhibited in its function in translational termination. Transient over-expression of Rmc greatly increased the frequency of conversion to [RPS⁺].

[0292] It is highly unusual for over-expression of a protein to cause a loss-of-function phenotype. It is even more unusual for phenotypes produced by over-expression to be stable after over-expression has ceased. Yet these properties are shared by the two yeast prion determinants and, to our knowledge, have been uniquely shared by them until now. They are believed to derive from stabilization of an otherwise unstable protein conformation by protein-protein interactions. Proteins in the altered form then have the capacity to recruit new proteins of the same type to the same form. The phenotype associated with this change is, therefore, stably inherited from generation to generation and transferred to mating partners in crosses.

[0293] The ability of amino acid residues 153-405 of Rnq1 (SEQ ID NO: 50) to substitute for the N-terminal domain of Sup35 and recapitulate its prion behavior was by no means predictable. The C-terminal region of Rnq1 (residues 153-405) and the N-terminal region of Sup35 have no primary amino-acid sequence homology—only a similar enrichment in polar amino acids. Reconstituting the epigenetic behavior of a prion requires that the Rmc fusion protein achieve an unusual balance between solubility and aggregation. If the fusion protein is too likely to aggregate, the inactive state will be ubiquitous; if it is too likely to remain soluble, the inactive state will not be stable. To recapitulate the epigenetic behav-

ior of [PSI⁺] the fusion protein must be able to switch from one state to the other and maintain either the inactive or the active state in a manner that is self-perpetuating and highly stable from generation to generation. Even minor variations in the sequence of the N-terminal region of Sup35, including several single amino-acid substitutions and small deletions, can prevent maintenance of the inactive state. And a small internal duplication destabilizes maintenance of the active state. Therefore, the ability of the Rnq1 domain to substitute for the prion domain of Sup35 and to fully recapitulate its epigenetic behavior provides a rigorous test for its capacity to act as a prion and suggests that it has been honed through evolution to serve this function.

[0294] The fusion of prion-determining regions with different functional proteins could be used to create a variety of recombinant proteins whose functions can be switched on or off in a heritable manner, both by nature and by experimental design. The two regions that constitute a prion, a functional domain and an epigenetic modifier of function, are modular and transferable.

Example 8

High-Throughput Assay to Identify Novel Prion-Like Amyloidogenic Sequences

[0295] The procedures described in Example 5 are particularly useful for identifying candidate prion-like sequences based on sequence characteristics and for screening these candidate sequences for useful prion-like properties. The following modification of those procedures provides a high-throughput genetic screen that is particularly useful for identifying sequences having prion-like properties from any set of clones, including a set of uncharacterized clones, such as cDNA or genomic libraries.

[0296] A library of short DNA fragments, such as genomic DNA fragments or cDNAs, is cloned in front of a sequence encoding the C-terminal domain of yeast Sup35 to create a library of CSup35 chimeric constructs of the formula 5'-X-CSup35-3', wherein X is the candidate DNA fragment. Optionally, the 3' end of the construct encodes both the M and C domains of Sup35. This library is transformed into a [psi⁻] strain of yeast that carries Sup35 as a Ura⁺ plasmid (with its chromosomal Sup35 deleted). Transformants are plated onto FOA-containing medium, which will cure the Ura⁺ plasmid so that the only functioning copy of Sup35 will be a fusion construct from the chimeric library.

[0297] Viable transformants are transferred to a selective media to screen for transformants which can suppress nonsense codons in a [PSI⁺]-like manner. For example, if the host cell is a yeast strain carrying a nonsense mutation in the ADE1 gene, the transformants are screened for cells that are viable on a SD-ADE media. Cells that can survive via suppression of nonsense codons are selected for further analysis (e.g., as described in preceding Examples), under the assumption that the library chimera has altered the function of Sup35. By using prion-specific tests such as histological examination for protein aggregates, curing, and Hsp104-dosage alteration, true aggregation-directing protein domains will be identified from original library of DNA constructs. The constructs which display prion-like properties can be used as described herein. Also, such constructs can be isolated and sequenced and used to identify and study the complete genes from which they were derived, to see if the original gene/protein possesses prion properties in its native host. The foregoing assay

also is useful for rapidly identifying fragments and variants of known prion-like proteins (NMSup35, NUre2, PrP, and so on) that retain prion-like properties. The assay, as well as chimeric constructs of the formula 5'-X-CSup35-3' and expression vectors containing such constructs, are considered additional aspects of the present invention.

Example 9

Fiber Assembly Mechanism of the Prion-Determining Region (NM) of Yeast Sup35p

[0298] The investigation of specific protein aggregation is gaining an increasing role in conjunction with increasing numbers of human diseases characterized by altered protein structures, including prion-based encephalopathies, noninfectious neurodegenerative diseases, and systemic amyloidosis. Amyloid protein aggregates are β -sheet rich structures that form fibers in vitro and bind dyes such as CongoRed and ThioflavinT. Strikingly, most amyloids can promote the propagation of their own altered conformations, which is thought to be the basis of protein-mediated infectivity in prion diseases. This feature of protein self-propagation in amyloids may also be critical to disease progression in noninfectious amyloid diseases such as Alzheimer's or Parkinson's disease. A powerful system to study the molecular mechanism of amyloid propagation and specificity is the prion-like phenomenon [PSI⁺] of *Saccharomyces cerevisiae*. Formation of higher ordered Sup35p complexes and the propagation of [PSI⁺] is caused by NM region of Sup35p. In vitro, both full-length Sup35p and NM form amyloid fibers with NM dictating the formation of the fiber axis while the C-terminal region of Sup35p is thought to be located on the periphery of the fibers. Detailed analysis by circular dichroism showed that NM adopts a mainly random coil structure in solution before it changes slowly to a structure that is β -sheet-rich. This conformational conversion was shown to occur simultaneously to the formation of amyloid fibrils.

[0299] In general, amyloid polymerization is considered to be a two-stage process initiated by the formation of a small nucleating seed or protofibril. Seed formation is thought to be oligomerization of soluble protein accompanied by a transition from a predominantly random coil to an amyloidogenic β -sheet conformation. Subsequent to nucleation, the seeds assemble with soluble protein to form the observed amyloid fibrils. The mechanisms for nucleation and fiber assembly are not well understood.

[0300] Strikingly, the secondary structure of all proteins that form amyloid fibrils under physiological conditions is partially random coil in aqueous solutions. Such structure is usually significant for partially unfolded protein as found in folding intermediates. It is possible that this unique "high-energy" structure in solution is the driving force for fiber assembly of such proteins. Thereby, the fibrous aggregates might present the lowest energy conformer of these proteins. As a consequence, interference with their structural state in solution should influence their fiber assembly ability. This has been shown for Alzheimer's β -amyloid peptide, islet amyloid polypeptide, and the artificial peptide DAR16-IV, where changes in the secondary structure dramatically altered the fiber assembly process.

[0301] The following experiments were performed to examine and characterize the folding and association pathway of soluble NM by starting with chemically denatured protein. Similar results were obtained with proteins isolated

under non-denaturing conditions. These studies were facilitated by use of labeled cysteine-substituted NM mutants. A better understanding of the mechanisms of fiber assembly will facilitate manipulations of fiber growth under various conditions.

A. Materials and Methods

[0302] Bacterial Strains and Culture

[0303] Using pEMBL-Sup35p (an *E. coli* plasmid containing the Sup35 protein) as template, DNA encoding NM was amplified by PCR with various linkers for subcloning. For recombinant NM expression, the PCR products were subcloned as NdeI-BamHI fragments into pJC25. For GST-NM fusions, the PCR products were subcloned as BamHI-EcoRI fragments into pGEX-2T (Pharmacia). For site-directed mutagenesis the protocol by Howorka and Bayley, *Biotechniques*, 25:764-766 (1998), was used for a high throughput cysteine scanning mutagenesis. A non-mutagenic primer pair for the β -lactamase gene and a mutagenic primer pair for each respective mutant were employed. In addition to generating a unique NsiI site, we used SphI and NspI sites, which allows introduction of a cysteine codon in front of methionine and isoleucine or after alanine and threonine codons, to increase the number of mutants in our cysteine screen. The fidelity of each construct was confirmed by Sanger sequencing. Protein was expressed in *E. coli* BL21 [DE3] after inducing with 1 mM IPTG (OD_{600nm} of 0.6) at 25° C. for 3 hours.

[0304] Yeast Strains and Culture

[0305] Using pJLI-Sup35pC-Sup35p as a template, DNA encoding each of the respective NM^{CYS} was amplified by PCR with two EcoRI sites for subcloning. To investigate the propagation and maintenance of [PSI⁺] by each NM^{CYS} used, integrative constructs, constructed using the standard pRS series of vectors (available from ATCC), were digested with XbaI and transformed into 74-D694 [PSI⁺] and [psi] strains. Transformants were selected on uracil-deficient (SD-Ura) medium and confirmed by genomic PCR followed by digestion with AatII, which cleaves the HA-tag between NM^{CYS} and Sup35pC. Recombinant excision events were selected on medium containing 5-fluoro-ortotic acid. Only cells that have lost remaining integrative plasmids are able to grow on medium containing 5-fluoro-ortotic acid. Again, replacements were confirmed by PCR followed by digestion with AatII as described above.

[0306] Protein Purification

[0307] NM and each NM^{CYS} were purified after recombinant expression in *E. coli* by chromatography using Q-Sepharose (Pharmacia), hydroxyapatite (BioRad), and Poros HQ (Boehringer Mannheim) as a final step. All purification steps for NM or NM^{CYS} were performed in the presence of 8M urea. GST-NM was purified by chromatography using Glutathione-Sepharose (Boehringer Mannheim), Poros HQ (Boehringer Mannheim), and S-Sepharose (Pharmacia) as a final step. All purification steps for GST-NM were performed in the presence of 50 mM Arginine-HCl. Protein concentrations were determined using the calculated extinction coefficient of 0.90 (NM, NM^{CYS}) or 1.23 (GST-NM) for a 1 mg/ml solution in a 1 cm cuvette at 280 nm.

[0308] Secondary Structure Prediction

[0309] Secondary structure of NM was predicted by using two independent prediction methods, GOR IV and Hierarchical Neural Network. Both methods were provided by Pôle Bio-Informatique Lyonnais.

[0310] Secondary Structure Analysis

[0311] CD spectra were obtained using a Jasco 715 spectropolarimeter equipped with a temperature control unit. All UV spectra were taken with a 0.1 cm pathlength quartz cuvette (Hellma) in 5 mM potassium phosphate (pH 7.4), 150 mM NaCl and respective additives such as osmolytes in certain experiments. Protein concentration varied from 0.5 μ M to 65 μ M. Folding of chemically denatured NM or NM^{CYS} was monitored at 222 nm in time course experiments by diluting protein out of 8M Gdm*Cl (Guanidinium HCl; final concentration 50 mM) in the respective phosphate buffer. Thermal transition of NM or NM^{CYS} was performed with a heating/cooling increment of 0.5° C./min. Spectra were recorded between 200 nm and 250 nm (2 accumulations). In a separate measurement, time courses were recorded for 30 sec at single wavelengths (208 nm and 222 nm) for each temperature and the mean value of each time course was determined. Temperature jump experiments were performed by incubating the sample in a water bath with the respective starting temperature for 30 min. The cuvette was transferred to the spectropolarimeter already set to the final temperature and time courses were taken with a constant wavelength of 222 nm. Settings for wavelength scans: bandwidth, 5 nm; response time, 0.25 sec; speed, 20 nm/min; accumulations, 4. All spectra were buffer-corrected.

[0312] Fluorescent Labeling of NM^{CYS}

[0313] The thiol-reactive fluorescent labels acrylodan and IANBD amide (Molecular Probes) were incubated with NM^{CYS} for 2 hours at 25° C. according to the manufacturer's protocol. Remaining free label was removed by size exclusion chromatography using D-Salt Excellulose desalting columns (Pierce). The labeling efficiencies were determined by visible absorption using the extinction coefficients of 2×10^4 for acrylodan at 391 nm and 2.5×10^4 for IANBD

B. Construction and Analysis of NM Mutants

[0314] To investigate the structural requirements for amyloid fiber assembly, we used yeast Sup35p's NM-region as a model protein. Until recently, fiber assembly kinetics of NM and other amyloid forming proteins have been monitored by binding of dyes such as CongoRed (CR) or ThioflavinT. To gain further insight into NM folding and fiber assembly, a more sensitive method for detecting structural changes, such as that provided by intrinsic fluorescence, was necessary. As NM naturally lacks tryptophan, the only native amino acid with a reasonable environmental-sensitive fluorescence, site-directed mutagenesis could have been employed to artificially introduce tryptophan in NM. However, to improve experimental flexibility we introduced single cysteine substitutions throughout NM. Since NM naturally lacks cysteine, such single point mutations would allow probing of NM folding and assembly in a specific, well defined manner after cross-linking of fluorescent probes to the sulfhydryl-groups of cysteines.

[0315] NM mutants with single cysteine replacements at amino acids throughout NM that were predicted to be in structured regions or that were likely involved in the fiber assembly process were constructed. These included the following fifteen mutants: NM^{S2C}, NM^{I35C}, NM^{Q38C}, NM^{G43C}, NM^{G68C}, NM^{M124C}, NM^{P138C}, NM^{L144C}, NM^{T158C}, NM^{E167C}, NM^{K184C}, NM^{E203C}, NM^{S234C}, and NM^{L238C}. As indicated in table 1 below, three of the fifteen mutants, NM^{I35C}, NM^{Q40C} and NM^{M124C}, were not stably expressed at a sufficiently high protein levels in *E. coli*. All other mutants were purified to homogeneity under denaturing conditions. To confirm that refolded NM attained a native protein structure, a GST-NM fusion protein was purified with thrombin, and GST was removed by binding to Glutathione-Sepharose. A structural comparison of refolded and native NM using far-UV circular dichroism (CD) showed no apparent differences between the two proteins.

TABLE 1

NM Protein	Expression in <i>E. coli</i>	Secondary Structure [0 _{222 nm}]	Fiber assembly (CR-binding)	Fiber morphology (EM)
wild-type (wt)	yes	-2950	yes	smooth fibers up to 35 μ m long
NM				
NM ^{S2C}	yes	as wt	as wt	as wt
NM ^{I35C}	not detectable	—	—	—
NM ^{Q38C}	yes	as wt	as wt	as wt
NM ^{Q40C}	very low, not stable	—	—	—
NM ^{G43C}	yes	-6420	slower assembly rate	short fibers, only few are longer than 1 μ m
NM ^{G68C}	yes	-6250	slower assembly rate	short fibers, only few are longer than 1 μ m
NM ^{M124C}	very low, not stable	—	—	—
NM ^{P138C}	yes	-4570	as wt	as wt
NM ^{L144C}	yes	-4198	as wt	as wt
NM ^{T158C}	yes	as wt	as wt	as wt
NM ^{E167C}	yes	as wt	as wt	as wt
NM ^{K184C}	yes	-4400	as wt	as wt
NM ^{E203C}	yes	-4000	as wt	less smooth, many short fibers

TABLE 1-continued

NM Protein	Expression in <i>E. coli</i>	Secondary Structure [$[\theta]_{222nm}$]	Fiber assembly (CR-binding)	Fiber morphology (EM)
NM ^{S234C}	yes	-6410	slower assembly rate	many short fibers
NM ^{L238C}	yes	-3730	no	no detectable fibers

[0316] To determine the direct influence of individual cysteine replacements on the folding and assembly of NM in vitro, the secondary structure of each NM^{cys} was compared to wild-type NM structure by far-UV CD after refolding. The results are summarized in table 1. Structurally, only NM^{S2C}, NM^{Q38C}, NM^{T158C}, and NM^{E167C} were identical to wild-type NM. All other mutants contained a higher content of secondary structure as indicated by an increased mean residue ellipticity at $[\theta]_{222nm}$. NM and all ^{Nm}cys, with the exception of NM^{L238C}, had identical mean residue ellipticities at $[\theta]_{208nm}$ of $-9000 \text{ degree cm}^2 \text{ dmol}^{-1}$. In contrast, NM^{L238C} had a decreased mean residue ellipticity at $[\theta]_{208nm}$ indicating that this mutant had an aberrant structure in comparison to wild-type NM than the other NM^{cys}.

[0317] Next, fiber assembly of each mutant was performed on a roller drum and compared to wild-type NM assembly kinetics by binding of CongoRed (CR), which shows a spectral shift after interacting with amyloid fibers. Results from these experiments are summarized in table 1. Only NM^{L238C} did not bind CR under all conditions tested. NM^{G43C}, NM^{G68C}, and NM^{S234C} showed slightly altered CR-binding kinetics suggesting slower fiber assembly rates in comparison to wild-type NM.

[0318] Electron microscopy (EM) was used to confirm that NM^{cys} fibers were morphologically identical to wild-type fibers. As indicated in table 1, the electron micrographs showed no apparent differences in fiber density, fiber diameter, or other morphological features in comparison to wild-type NM for NM^{S2C}, NM^{Q3C}, NM^{O138C}, NM^{L144C}, NM^{T158C}, NM^{E167C}, and NM^{K184C}. NM^{L238C} fibers were not detectable by EM, suggesting that the apparent lack of CR-binding of NM^{L238C} was not due to structural differences in fibers that affected CR-binding. Results from CD (secondary structure), CR-binding (fiber assembly kinetics), and EM (fiber morphology) indicate that the NM^{S2C}, NM^{Q38C}, NM^{T158C}, and NM^{E167C} mutants display no apparent differences to wild-type NM with respect to these parameters. To further confirm that the chosen cysteine mutants were not influencing the principal properties of NM, genomic wild-type NM could be replaced by Nm^{cys}.

C. Covalent Binding of Fluorescent Labels to NM^{cys}

[0319] Environmentally sensitive fluorescent probes, such as naphthalene derivatives or benzofurazans, are commonly used to detect conformational changes and assembly processes of proteins. Here, we made use of 6-acryloyl-2-dimethylaminonaphthalene (acrylodan) and N,N'-dimethyl-N-(iodoacetyl)-N'-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)ethylene diamine (IANBD amide) both of which react specifically with free thiol-groups on proteins. Whereas acrylodan is very sensitive to its structural environment, IANBD amide exhibits appreciable fluorescence when linked to buried or unsolvated thiols. Therefore, the latter fluorescence is highly sen-

sitive to changes in the solvation level of the fluorophore as seen in folding events, whereas acrylodan is more powerful for investigating conformational changes of a protein. The specific labeling efficiencies of soluble NM^{cys} were in the range of 0.40 to 0.78 (mol label/mol protein) with unspecific binding below 0.05 mol/mol for both fluorescent probes.

[0320] After covalent binding to NM^{cys}, the influence of the fluorescent labels on fiber assembly was investigated. No differences were found in fiber assembly for 7 mutants (see table 1) in the presence of fluorescent labels in comparison to non-labeled protein as detected by CR-binding. No gross structural changes in assembled fibers were visible by EM for NM^{Q38C}, NM^{P138C}, NM^{L144C}, NM^{T158C}, NM^{E167C}, and NM^{K184C}. In contrast, NM^{S2C} fibers labeled with both acrylodan and IANBD amide appeared rougher with an overall shorter length, although these changes were subtle.

[0321] To determine the incorporation of labeled NM^{cys} into fibers, equal amounts of labeled and non-labeled protein were mixed. The amount of label in the soluble protein fraction was detected over the course of fiber assembly. During the experiment, the label to protein ratio was constant indicating an equal incorporation of labeled and non-labeled protein into fibers. The resulting fibers were monitored for fluorescent emission of the respective label. Both measurements showed that fluorescent-labeled protein was sufficiently incorporated into amyloid fibers without influencing the assembly kinetics or the assembled state for NM^{Q38C}, NM^{P138C}, NM^{L144C}, NM^{T158C}, NM^{E167C}, and NM^{K184C}.

[0322] The foregoing experiments examined the folding process of NM using NM^{cys} mutants that exhibited folding processes and structural characteristics similar to wild-type NM. These results provide a better understanding of the process of NM folding.

Example 10

Kinetic Analysis of Fiber Elongation

[0323] The following experiments were performed to characterize how nuclei mediate the conversion of soluble NM to the amyloid form in the elongation phase of fiber formation.

[0324] Effect of Fluorescent Labeling

[0325] To determine if fluorescent labels themselves affected fiber assembly, mixed assembly reactions were performed with equal quantities of labeled and unlabeled protein of each mutant. The ratio of labeled protein to unlabeled protein that remained in the soluble phase was constant throughout the assembly time course, and the final level of assembly was the same. The fibers formed with each of the labeled NM^{cys} mutants were indistinguishable from unlabeled NM^{cys} fibers in terms of their diameter ($11.5 \pm 1.5 \text{ nm}$) and concentration. Thus, covalent attachment of acrylodan/IANBD amide to cysteines did not influence the assembly of these mutants.

[0326] Fluorescence Assay for Conformational Conversion
[0327] Next, it was investigated which residues of the NM residues are located in positions that would provide a change in fluorescent signal (upon fiber assembly) in conformational conversion reactions (during seeded fiber elongation). For NM^{S2C}, NM^{Q38C}, NM^{T158C}, and NM^{E167C}, cysteine-linked acrylodan showed a blue shift in fluorescence emission maximum (λ_{max}), indicating that the environment of each cysteine substitution changed. To determine if these changes were based on the conformational transitions that are associated with the transition from soluble protein into fibers, fluorescent changes were analyzed for 12 hours in undisturbed, non-seeded reactions. Such reactions depend upon spontaneous nucleation and no NM fibers are detected in this time frame. This experiment revealed that acrylodan fluorescence emission showed a gradual change of λ_{max} during the pre-assembly stage for NM^{S2C} and NM^{Q38C}.

[0328] By many criteria, the N-region of NM has been established as the region responsible for nucleation. Thus, these changes most likely reflect early conformational transitions involved in the first stage of nucleated conformational conversion (NCC). Acrylodan fluorescence emission of NM^{T158C} and NM^{E167C} revealed no significant change after 12 hours in non-seeded samples (Both of these residues are located in the M-region). However, coincident with seeded fiber assembly, solutions of NM^{T158C} and NM^{E167C}-acrylodan showed increased fluorescence intensities accompanied by a blue shift of λ_{max} (NM^{T158C}: 521 nm to 486 nm, FIG. 2A; NM^{E167C}: 528 nm to 502 nm). Thus, acrylodan labels at cysteine 158 and 167 are sensitive to the conformational differences between soluble and fibrous NM.


[0329] Seeded elongation occurs in two steps

[0330] Both NM^{T158C}- and NM^{E167C}-acrylodan (2 μ M each) showed a rate of fiber assembly of $v_{fiber} = 8 \pm 0.4 \times 10^{-4} \mu\text{mol s}^{-1}$ at 25° C. in the presence of seed (4% w/w), at which seed concentration soluble NM is present in excess over the seeding fiber ends by approximately 50,000 fold. This fiber assembly rate was similar to that measured for NM^{wt} by far-UV CD ($3 \times 10^{-4} \mu\text{mol s}^{-1}$) and light scattering ($5 \pm 0.3 \times 10^{-4} \mu\text{mol s}^{-1}$) at identical experimental conditions. To determine the kinetic parameters of fiber assembly it was essential to ensure that both the substrate and the seed were in excess in the reactions. To do this, fiber assembly rates were determined with constant seed concentrations (4% w/w calculated for a 5 μ M protein concentration) and varying soluble protein concentrations. Decreasing the soluble NM concentration 100-fold only decreased fiber assembly rates by a factor of two. Hence, soluble protein is in excess with 4% w/w seed and 5 μ M soluble NM.

[0331] The kinetics of seeded fiber elongation reproducibly showed a lag-phase of 80 ± 10 s at 25° C., then exhibited linear kinetics. The fact that fiber assembly did not begin immediately suggested that an assembly intermediate is formed. Non-fibrous NM is soluble in SDS while fibrous NM shows SDS-resistance. Based on this fact, an assay was developed to detect intermediate complexes, which identifies soluble NM that is associated with seed but still not converted into the fiber state. Seeds were prepared from NM^{K184C}, a cysteine substitution mutant with surface accessible sulfhydryl groups that allow for labeling after fiber formation and that shows a seeding efficiency indistinguishable to that of NM^{wt}, and these NM^{K184C} seeds were biotinylated. Further, NM^{T158C} was labeled with iodo[1-¹⁴C]acetamide. Reactions were started by addition of biotinylated NM^{K184C} seed (50%

(w/w)) to soluble NM^{T158C}-iodo[1-¹⁴C]acetamide and at distinct time points aliquots of the reaction were taken and incubated with Streptavidin-coated Dynabeads. A high ratio of seed to soluble protein was used to ensure that the fiber ends (i.e. the seeds) were saturated with soluble NM, which would therefore allow us the best opportunity of observing short-lived intermediate complexes. The beads were removed at different time points using a magnet and washed with SDS to detect non-converted intermediates. Both the SDS soluble protein and the SDS resistant fiber, which were attached to the beads, were analyzed by scintillation counting. It took 30 seconds to collect the beads. At early time points a substantial fraction (~50%) of the NM assembled with bead-bound seeds was soluble in SDS, at later time points the fraction of SDS-soluble material diminished. In a control experiment, in which the NM^{K184C} seeds were not biotinylated, no radioactivity could be detected attached to the beads. The ability to capture material bound to the seed that had not completely converted, established the formation of a detergent susceptible complex. However, this method did not have sufficient resolving power to analyze kinetic parameters of the assembly process.

[0332] To establish kinetic parameters, it was necessary to precisely discriminate between soluble and seed-bound NM. Therefore a sedimentation assay was developed to detect the disappearance of soluble NM^{T158C}-acrylodan during fiber assembly. The total acrylodan concentration was plotted against the acrylodan concentration in the supernatant, and each measurement was repeated 6 times to estimate the level of variation. In combination with the wavelength shift assay described above, this provided sufficient data to kinetically analyze fiber assembly and develop a model for nucleated fiber elongation. These reactions have several components: two reactants—the seed and the soluble NM, with the soluble NM as the substrate being in excess of the seed, and a catalyst that is not used up as the reaction progresses (the catalyst is the fiber ends, which are bound to by the soluble NM, but the same number of ends are present as the fiber elongates). These components and the fact that these reactions reach steady state kinetics suggest that they can be analyzed with the same mathematical formula that has been used to describe enzyme kinetics—the Michaelis-Menten equation:



[0333] where S is soluble NM, A is assembled protein (seed), SA is bound but not converted intermediate (akin to an enzyme:substrate complex), and AA is converted fiber, which again can act as seed. Importantly, we were unable to discriminate whether seed associates with monomers or oligomers or both. The observed rate of conformational conversion is determined experimentally by k_1 , k_{-1} , k_{conf} , k_1 and k_{-1} represent the rate constants for binding and dissociation, and k_{conf} is the first-order conformational conversion rate. Since the dissociation rate of converted protein from the amyloid fibers is too slow to be detected in our experimental set-up, the back reaction



[0334] is quasi-irreversible and ignored in our model.

[0335] Next, we analyzed our experimental data using a Lineweaver-Burk plot in order to gain more information on

the kinetic parameters of fiber assembly. In these experimental conditions, the Lineweaver-Burk plot yielded a straight line and a protein concentration of $K_m=0.12\pm 0.01 \mu\text{M}$, at which the rate of reaction is equal to one half of the limiting rate (maximum rate). We also calculated a maximal rate of conformational conversion $V_{max}=10\pm 0.3\times 10^{-4} \mu\text{mol s}^{-1}$, the rate constant of conformational conversion of $k_{conf}=5\pm 0.1\times 10^{-3} \text{s}^{-1}$, and a conformational conversion efficiency of $k_{conf}/K_m=42000 \text{M}^{-1} \text{s}^{-1}$, which is equivalent to an enzyme's specificity constant.

[0336] Influences of Temperature on Seeded Fiber Elongation

[0337] The effect of increased temperature on seeded fiber elongation was investigated with NM^{T158C} -acrylodan in the presence of 4% w/w seed. A low temperature optimum of the rate of fiber assembly as seen in the logarithm of NCC velocities plotted against the reciprocal temperature (Arrhenius plot) was found. The sticking probability of soluble protein, which is reflected by k_{conf}/k_{-1} , characterizes the rate at which soluble NM (S) associates with seed (SA) relative to dissociation, i.e., the sticking probability is high if $k_{-1}<k_{conf}$. In these experiments the abnormal temperature dependence with decreasing ratios of k_{conf}/k_{-1} at elevated temperature indicates a significant rate enhancement for the dissociation of the seed-NM (SA) complex in comparison to its conversion into an assembled fiber (AA). At low temperature $k_{-1}\ll k_{conf}$ and k_{conf}/K_m becomes equal to k_1 . Because the dissociation of non-converted, but seed-bound NM, has a high activation energy, k_{-1} becomes predominant at high temperature.

[0338] In order to test this experimentally, the velocities of fiber elongation at 25° C. and 40° C. were measured with a constant soluble NM^{T158C} -acrylodan concentration (2 μM) and increasing seed concentrations. It was confirmed that increasing seed concentrations led to increasing fiber elongation velocities at both temperatures yielding maximal elongation rates above 10% w/w of seed. Therefore, fiber elongation velocities at 12% w/w seed, which should be not-rate limiting seed concentrations for fiber elongation, were plotted against the reciprocal temperature. The plot revealed a temperature dependence of fiber elongation that is consistent with the collision theory of Arrhenius. The Arrhenius plot gives a straight line and its slope is equivalent to the activation energy E_a divided by the gas constant $R=8.3145 \text{J K}^{-1} \text{mol}^{-1}$. Using this equation, the activation energy for fiber elongation was calculated to be $E_a=11.7 \text{0.2 kJ mol}^{-1}$.

[0339] Acquisition of Secondary and Tertiary Structure of Soluble NM

[0340] In order to elucidate the influence of the conformation of soluble NM on the association with seed, we investigated the rate at which secondary, tertiary, and quaternary structures were acquired in soluble material. When NM is first diluted out of denaturants such as urea or guanidinium chloride (GdmCl), it adopts the characteristics of a molecule that is rich in random coil but partially structured (typical for intrinsically unstructured proteins) indistinguishable from that of NM purified under non-denaturing conditions. To analyze whether the rate of this process influences seeded fiber assembly, 6M GdmCl was used to form a homogenous and monomeric population of denatured NM. After dilution into 5 mM sodium phosphate, pH 7.4, 150 mM NaCl, the time course of far-UV Circular Dichroism (CD) changes at 222 nm was monitored. The acquisition of secondary structure reached half maximal amplitude after $24\pm 2 \text{s}$ with a rate-


constant of $k_{gain}^{farUV}=2.1\pm 0.2\times 10^{-2} \text{s}^{-1}$. Thus, the formation of secondary structure is not rate determining for seeded fiber elongation.

[0341] The kinetics of acquisition of NM tertiary structure was investigated by the four fluorescently-labeled NM^{CS} mutants. Changes in tertiary structure of NM upon dilution into buffer from 6M GdmCl were investigated with two different techniques: IANDB-amide labeled protein was investigated by fluorescence emission and acrylodan labeled protein with near-UV CD. The fluorescence emission of IANBD-amide revealed solvent exposure in all four mutants in 6M GdmCl, as expected. A stable IANDB-amide emission signal was reached after dilution into buffer indicative of a higher ordered environment. The time course had a half maximal amplitude at 31.4 s and a rate constant of $k_{gain}^{fluor}=1.6\pm 0.2\times 10^{-2} \text{s}^{-1}$. Similarly, near UV-CD time courses with acrylodan-labeled NM (all four mutants led to the same results) showed a half maximal amplitude after $33\pm 2 \text{s}$ and a rate constant of $k_{gain}^{nearUV}=1.5\pm 0.1\times 10^{-2} \text{s}^{-1}$. Both independent measurements revealed that formation of some tertiary structure is also not rate limiting for seeded fiber assembly under the experimental conditions chosen.

[0342] Quaternary Structure Analysis

[0343] Dilution of NM^{wt} out of denaturant led to the formation of a mixed population of monomers and oligomers. $87\pm 5\%$ of NM was monomeric and the remaining fraction heterogeneously oligomeric with varying molecular masses from tetramers to 30mers. Oligomerization was preceded by a lag phase of approximately 60 seconds after dilution out of denaturant, which may suggest that some acquisition of secondary and tertiary structure is required prior to oligomerisation. Populations of monomers and oligomers were established after a half time of 75 ± 5 seconds and remained constant for 3 hours. Since this steady state was achieved far before spontaneous nucleation (and well before seed was added), NM oligomerisation is not likely to be rate-determining for seeded fiber assembly in our experiments.

[0344] The data suggested the following mechanism for initial structural changes of soluble NM, starting from the denatured state:



[0345] where M_u is the unfolded monomer, M is the random-coil monomer with some structure, and O_x are the oligomers. The rate constant for structural gain of monomeric NM from the denatured state was $k_{gain}=1.5\pm 0.2\times 10^{-2} \text{s}^{-1}$. Remarkably, the rate of oligomerisation and establishment of a steady state distribution of monomers and oligomers showed little dependence on the concentration of NM between 0.7 μM and 46 μM NM. This observation agrees with that of a previous study that NM fiber assembly proceeds via the conversion of oligomers to nuclei with little concentration dependence. Nuclei form by conformational rearrangements of NM within the context of oligomeric intermediates and not by assembly of structurally converted monomers.

Example 11

Bi-Directional Formation of Fibers Composed of the Prion-Determining Region (NM) of Yeast Sup35p

[0346] The following experiments were performed to demonstrate that fibers composed of the NM region of Sup35p are

capable of adding NM protein at both ends of the fiber. This was investigated using a mutant NM protein, in which the lysine residue at position 184 was substituted by cysteine, that was capable of forming fibers labeled with specifically modified gold colloids. Visualization of the gold-labeled fibers allowed determination of the directionality of fiber growth.

A. Determining the Accessibility of Cysteine Residues in Assembled Fibers

[0347] First, the accessibility of cysteine residues was assayed in fibers composed of cysteine-substituted mutant NM (NM^{C35S}) proteins, each of which carried different single cysteine replacements at amino acid residues throughout the NM protein. All NM^{C35S}, described in Example 9 above, that formed fibers were examined. For fiber assembly, NM^{C35S} protein was diluted out of 4M Gdm*Cl 80-fold into 5 mM potassium phosphate (pH 7.4), 150 mM NaCl to yield a final NM^{C35S} protein concentration of 10 μ M. To accelerate the rate of fiber assembly, all NM^{C35S} proteins were incubated on a roller drum (9 rpm) for 12 hours. The resulting fibers were sonicated with a Sonic Dismembrator Model 302 (Artek) using an intermediate tip for 15 seconds. Sonication resulted in small sized fibers that did not reassemble to larger fibers as determined by electron microscopy (EM). Seeding of fiber assembly was performed by addition of 1% (v/v) of the sonicated fibers to soluble NM^{C35S} protein.

[0348] To test the accessibility of cysteines in assembled fibers composed of NM^{C35S} proteins, EZ-link PEO-maleimide-conjugated biotin (Pierce, product number 21901) was added to the assembled fibers and the labeling efficiency of the biotin was assayed. EZ-link PEO-maleimide-conjugated biotin was covalently linked to assembled NM^{C35S} fibers for 2 hours at 25° C. according to the manufacturer's protocol (protocol number 0748). Remaining free biotin was removed by size exclusion chromatography using D-Salt Excellulose desalting columns (Pierce, product number 20450). Labeling efficiency was determined by competing for avidin binding between biotin and [2-(4'-hydroxybenzene)] benzoic acid (HABA). The binding of HABA to avidin results in a specific absorption band at 500 nm. Since biotin displaces the HABA dye due to higher affinity of biotin for avidin, as compared to that of HABA dye for avidin, the binding of HABA to avidin and thus the specific absorption at 500 nm decreases proportionately when biotin is added to the reaction. Results from this assay indicated that fibers composed of either NM^{C35S} proteins in which the lysine residue at position 184 was substituted by a cysteine residue (K184C) or NM^{C35S} proteins in which the serine residue at position 2 was substituted by a cysteine residue (S2C), bound a detectable amount of biotin. S2C fibers had a labeling efficiency of 0.16 mol biotin/mol protein, and K184C fibers exhibited a labeling efficiency of 0.56 mol biotin/mol protein. Thus, the cysteine residue at position 184 is highly accessible and the cysteine residue at position 2 is partially accessible on the surface of assembled fibers.

B. Analysis of Fiber Growth Using EM

[0349] K184C sonicated fibers were tested for their ability to seed fiber assembly of soluble wild-type NM protein. Fiber assembly was performed as described above using sonicated K184C fibers as seeds to assemble soluble wild-type NM protein. The rate of fiber assembly was assayed by CongoRed binding (CR-binding) and fiber morphology was examined

by EM. For EM studies, protein solutions were negatively stained as previously described in Spiess et al., 1987, *Electron Microscopy and Molecular Biology: A Practical Approach*, Oxford Press, p. 147-166. Images were obtained with a CM120 Transmission Electron Microscope (Phillips) with an LaB6 filament, operating at 120 V in low dose mode at a magnification of 4500 \times and recorded on Kodak SO163 film. Results from CR-binding and EM experiments show that K184C fibers are able to seed wild-type NM fiber assembly. The resulting mixed K184C/NM fibers showed no apparent differences in assembly rate or morphology to fibers seeded with sonicated wild-type NM fibers. Similar results were obtained when biotinylated K184C seeds were used for fiber assembly.

[0350] The surface exposure of the cysteine at position 184 in assembled fibers composed of the K184C mutant protein allowed sufficient labeling of fibers with specifically modified gold colloids. Monomaleimido Nanogold™ (Nanoprobes, product number 2020A) with a particle diameter of 1.4 nm was covalently cross-linked to the sulfhydryl group of accessible cysteine residues in sonicated K184C fibers for 18 hours at 4° C. according to the manufacturer's protocol. Remaining free Nanogold™ was removed by a repeated size exclusion chromatography using D-Salt Excellulose desalting columns (Pierce, product number 20450). The extent of labeling was determined by UV/visible absorption using extinction coefficients for Nanogold™ of 2.25×10^5 at 280 nm and 1.12×10^5 at 420 nm. Ratios of optical densities at 280 nm and 420 nm allowed an approximation of the labeling efficiency. These gold-labeled fibers were employed to seed fiber growth of soluble wild-type NM protein.

[0351] To visualize the 104 nm Nanogold™ particles attached to the assembled mixed K184C/NM fibers, we used Goldenhance™ (Nanoprobes) according to the manufacturer's instructions. Briefly, equal volumes of enhancer (Solution A) and activator (Solution B) were combined and incubated for 15 min at room temperature. Initiator (Solution C) was then added at a volume equal to that of enhancer or activator, and the resulting mixture was diluted (1:2) with phosphate buffer (Solution D). The final solution acts as an enhancing reagent by selectively depositing gold onto Nanogold™ particles, thereby providing enlargement of Nanogold™ to give electron-dense enlarged Nanogold™ particles in the electron microscope. For negative staining of gold-labeled fibers, 6 μ l of protein (8 μ M, 1% (w/w) gold labeled seed) were applied to a 400 mesh carbon-coated copper grid (Ted Pella) for 45 seconds. After washing with 100 μ l phosphate buffer, grids were incubated with the final Goldenhance™ enhancing reagent, prepared as described above, for 5 min. After washing with 200 μ l glass-distilled water, negative staining was employed as in Spiess et al., 1987 *Electron Microscopy and Molecular Biology: A Practical Approach*, Oxford Press, p. 147-166. EM results revealed that the gold-labeled K184C regions are located in the middle of the assembled K184C/NM fibers indicating bi-directional fiber assembly with no apparent polarity in the seeds used.

[0352] The foregoing experiments show that fiber assembly of NM proteins occurs at both ends of the fibers. These analyses were performed using K184C, a NM^{C35S} mutant wherein the lysine residue at position 184 has been substituted with a cysteine residue. Experiments by biotin-labeling of the cysteine residues on assembled K184C fibers were carried out to determine accessibility of the cysteines. Since wild-type NM protein does not contain any cysteine residues,

labeling can only occur at position 184. Results show that position 184 is highly accessible in assembled K184C fibers. The ability of specifically modified gold colloids to covalently cross-link the sulfhydryl group of cysteines enabled generation of gold-labeled fibers that can be visualized by EM. Examination of fiber assembly, by taking advantage of the ability of K184C to produce gold-labeled fibers, indicates that fiber growth occurs bi-directionally. It further indicates that fibers with specific modifications and attachments, a single fiber containing modified and unmodified regions, and mixtures of modified and unmodified fibers can be produced.

Example 12

Conducting Nanowires Built by Controlled Self-Assembly of Amyloid Fibers and Selective Metal Deposition

[0353] The following experiments were performed to demonstrate that fibers composed of the NM region of Sup35p can be modified to conduct electricity. This was investigated using a mutant NM protein, in which the lysine residue at position 184 was substituted by cysteine, that was capable of forming fibers labeled with specifically modified gold colloids. These fibers were placed across gold electrodes, and additional metal was deposited by highly specific chemical enhancement of the colloidal gold by reductive deposition of metallic silver and gold from salts. The resulting silver and gold wires were ≈ 100 nm wide. These biotemplated metal wires demonstrated the conductive properties of a solid metal wire, such as low resistance and ohmic behavior.

A. Materials and Methods

[0354] Protein Expression and Purification.

[0355] NM and NM^{K184C} was recombinantly expressed in *Escherichia coli* BL21 [DE3] as described (Scheibel, T., et al., *Curr. Biol.* 11: 366-369 (2001)) and purified by chromatography with Q-Sepharose (Amersham Pharmacia), hydroxyapatite (Bio-Rad), and Poros HQ (Roche Molecular Biochemicals) as a final step. All purification steps were performed in the presence of 8 M urea.

[0356] Fiber Assembly.

[0357] Solutions with protein (NM or NM^{K184C}) concentrations >25 μ M were rotated at 60 rpm to increase turbulence and surface area. At this protein concentration, many seeding events initiate simultaneous fiber assembly, which results in many short fibers (average fiber length from 60 to 200 nm). These short fibers were then used to seed further soluble NM. The polymerization of NM is a two-stage process that starts with the formation of a nucleus that contains protein with a different conformation than that of soluble protein. The nucleus promotes the conformational conversion of the remaining soluble protein into amyloid fibers. When denatured NM is initially diluted into physiological buffers it has the features of an intrinsically unstructured (random coil-rich) protein. After a lag phase, nuclei form and initiate the rapid conversion of soluble NM into β -sheet-rich amyloid. This second stage can be imitated by addition of pre-formed fibers (seed) to soluble NM. Fibers of different average length were generated by changing the ratios of seed to soluble NM (keeping the soluble NM concentration constantly at 5 μ M).

[0358] Analysis of Fiber Structure.

[0359] After fiber assembly, three techniques were used to examine the fibrous state of NM: far-UV CD (far-ultra-violet

circular dichroism), Congo red (CR) binding, and atomic force microscopy (AFM). CD spectra were obtained by using a Jasco (Easton, Md.) 715 spectropolarimeter equipped with a temperature control unit. All spectra were taken with a 0.1-cm pathlength quartz cuvette (Hellma, Forest Hills, N.Y.) in 5 mM potassium phosphate (pH 7.4)/150 mM NaCl (standard buffer). The settings for wavelength scans were 5-nm bandwidth; 0.25-sec response time; speed, 20 nm/min; and four accumulations.

[0360] CR-binding was carried out as described (Glover, J. R., et al. *Cell*, 89: 811-819 (1997)). Proteins were diluted to a final concentration of 1 μ M into standard buffer plus 10 μ M CR and incubated for 1 min at 25° C. before measuring the absorbance at 540 and 477 nm.

[0361] Samples for AFM analysis were placed on freshly cleaved mica attached to 15-mm AFM sample disks (Ted Pella, Redding, Calif.). After 3 min of adsorption at 25° C., disks were rinsed once with buffer and twice with Millipore filtered distilled H₂O. The samples were then allowed to air dry. Contact and tapping-mode imaging were performed on a Digital Instruments (Santa Barbara, Calif.) multimode scanning probe microscope (Veeco, Santa Barbara, Calif.) by using long, thin-leg standard silicon nitride (Si₃N₄) probes for contact mode and standard etched silicon probes for tapping mode.

[0362] Analysis of Fiber Stability.

[0363] To investigate fiber stability at elevated temperatures, NM fibers were incubated in standard buffer for 90 min at 98° C., before assessment by CD, CR binding, and AFM. The stability of the fibers was also tested under other temperatures for varying lengths of time, i.e., several months at 25° C. and after freezing at -20° C. and -80° C. Chemical stability was tested by the addition of high concentrations of salt (2.5 M NaCl) or denaturants [8 M urea or 2 M guanidiniumchloride (Gdm.Cl)] to the standard buffer (5 mM sodium phosphate, pH 6.8) and assessed by CD, CR binding, and AFM. NM fiber stability in strong alkaline or acidic solutions and in organic solvents was tested by immobilizing the fibers on mica, air-drying them, and treating them with NaOH (pH 10), HCl (pH 2), or 100% ethanol for several hours. These conditions were not compatible with CD and CR-binding assessment, therefore only AFM was used.

[0364] Gold Toning.

[0365] Monomaleimido Nanogold (Nanoprobes, Yaphank, N.Y.) with a particle diameter of 1.4 nm was covalently cross-linked to NM^{K184C} fibers as described in Scheibel, T., et al., *Curr. Biol.* 11: 366-369 (2001), incorporated by reference. The Nanogold reagent was dissolved in 0.02 ml isopropanol, then diluted to 0.2 ml with deionized water. The activated Nanogold solution was added to the NM^{K184C} fibers and incubated for 2 hours at 25° C. Unbound gold particles were separated from the NM^{K184C} fibers using gel exclusion chromatography. The Nanogold conjugate was effectively isolated using a Pharmacia Superdex 400HR medium (which fractionate a wide range of molecular weights). The 1.4-nm Nanogold particles were then subjected to "gold toning" (i.e., silver enhancement followed by gold enhancement). In this procedure, the Nanogold particles act as promoters for reducing silver ions from a solution. The Nanogold-labeled fibers are subjected to silver enhancement with LI Silver (Nanoprobes) performed according to the manufacturer's protocol: solutions A (enhancer solution) and B (activator solution) were mixed in a 1:1 ratio and incubated with the fibers at 25° C.). The resulting silver-coated fiber-bound Nanogold par-

ticles were gold-enhanced with GoldEnhance LM (Nano-probes). Enhancement was performed according to the manufacturer's protocol: solutions A-D (A: enhancer; B: activator; C: initiator; D: buffer) were mixed in a 1:1:1:1 ratio and incubated with the fibers at 25° C.). Exposure times varied from 3 min of silver enhancement and 3 min of gold enhancement to 25 min of silver enhancement and 25 min of gold enhancement.

[0366] Electrode Assembly and Visualization.

[0367] Electrodes were prepared on Si₃N₄ membrane substrates as described in Morkved, T. L., et al., *Polymer*, 39: 3871-3875 (1998), incorporated herein by reference. The electrodes were constructed by spinning polymer resist layers onto Si₃N₄ substrates and exposing them to a scanned electron beam. The electron beam demarcated the electrode sites. The exposed polymer was etched away, and gold vapor was applied to fill the resulting gaps. Finally, the remaining polymer was dissolved away, leaving the gold in the pattern inscribed by the electron beam. Typically, gaps between electrodes were 2-10 μm. Transmission electron microscopy (TEM) images of electrodes in the absence and presence of protein fibers were obtained with a CM120 transmission electron microscope (Phillips, FEI, Hillsboro, Oreg.) with a LaB6 filament, operating at 120 kV in low-dose mode at a magnification of ×45,000, and recorded on Kodak SO163 film. Alternatively, samples were imaged by AFM in contact mode. Conductivity measurements were performed as described (Morkved, T. L., et al., *Polymer*, 39: 3871-3875 (1998)). Briefly, conductivity measurements were performed by biasing the sample with a constant voltage from a Hewlett Packard function synthesizer and, using Keithley electrometers, measuring current and voltage across the sample over a range of temperatures.

B. NM Fibers are Highly Stable

[0368] To investigate the feasibility of using NM fibers in building nanoscale devices, fiber stability was first evaluated under extreme conditions such as those that might be encountered in industrial manufacturing processes. NM fibers assembled at physiological pH and room temperature were assayed for stability by three techniques that differentiate between NM in its soluble and amyloid state. Far-UV CD distinguishes the β-sheet-rich secondary structure of NM fibers from the random coil-rich structure of soluble NM. CR exhibits a spectral shift when it intercalates into the cross-pleated β-strands of NM fibers, which is not observed with soluble NM. AFM and EM were used to monitor the maintenance of fiber morphology.

[0369] NM fibers were incubated in standard buffer (5 mM sodium phosphate, pH 6.8) at high and low temperatures, in the absence or presence of high salt (2.5 M NaCl), and in denaturants (8 M urea or 2 M guanidiniumchloride, Gdm.Cl). By all three techniques, fibers were stable in standard buffer after incubation for 90 min at 98° C., for several months at 25° C., and after freezing at -20° and -80° C. (Some shearing of long fibers occurred with repeated cycles of freeze-thawing.) Fibers were completely stable to prolonged incubation in the absence of salt and at 2.5 M salt. They dissociated in <2 h at concentrations of Gdm.Cl >4 M but remained intact in the presence of 2 M Gdm.Cl and 8 M urea.

[0370] To test whether NM fibers can withstand strong alkaline or acidic solutions and incubation in organic solvents, which are incompatible with CD and CR-binding assays, NM fibers were immobilized on mica, imaged by

AFM, incubated with test solutions [NaOH (pH 10), HCl (pH 2), or 100% ethanol], at 25° C. for up to 2 hours and then reimaged. No morphological changes were apparent after any of these treatments. Therefore, NM fibers show unusually high chemical and thermal stability for a biological material.

C. Production of NM Fibers of Variable Lengths

[0371] Studies of the NM amyloid fibers have provided insights into how fibers assemble and how assembly can be controlled (Glover, J. R., et al. *Cell*, 89: 811-819 (1997); Serio, T. R., et al. *Science*, 289: 1317-1321 (2000); Scheibel, T., et al., *Nat. Struct. Biol.*, 8: 958-962 (2001) all of which are incorporated by reference). The rate of fiber formation by purified soluble NM is dramatically increased by the addition of preformed NM fibers, which seed assembly from their ends (DePace, A. H., et al., *Nat. Struct. Biol.*, 9:389396 (2002); Scheibel, T., et al., *Curr. Biol.*, 11: 366-369 (2001)). Pools of fibers with different average lengths were generated by simple manipulation of the assembly conditions. First, short fibers (60-200 nm) were produced by rotating solutions with high NM protein concentrations (>25 μM) at high speeds (60 rpms) to increase turbulence and surface area. These conditions produced short fibers by greatly increasing the efficiency of seeding (such that it dominates over assembly), rather than by simply shearing fibers after they had assembled. Indeed, when preformed fibers were sheared by the much more physically disruptive force of sonication, the resulting fibers had longer average lengths and a much more heterogeneous distribution. The resulting sonicated fibers showed lengths varying from 100 to 500 nm (Scheibel, T., et al., *Curr. Biol.*, 11: 366-369 (2001)).

[0372] The short fibers produced by vigorous rotation of high concentrations of NM were used to seed further soluble NM. By simply changing the ratios of seed to soluble NM and by controlling the assembly temperatures (i.e., for preferred fiber assembly, the temperature was kept constant at 25° C.) fibers of different average length were generated. At seed to soluble NM ratios of 1:1 (wt/wt), fibers showed an average length of 500±100 nm. Increasing the soluble NM concentration increased fiber lengths. At ratios of 1:16 of seed to soluble NM, fibers were ≈5±1 μm long. Ratios of 1:64 led to even longer fibers but these had more variable lengths (10 μm up to several hundred micrometers).

[0373] A remarkable phenomenon that was sometimes observed when long fibers were prepared for microscopy was their alignment next to each other without any external manipulation. This alignment varied with the buffers in which fibers were suspended and the manner in which the surfaces were prepared in a fashion that has not been completely deciphered.

D. NM Fibers are Insulators

[0374] To examine the electrical behavior of the protein fibers, Si₃N₄ membrane substrates were grown on a silicon wafer which allowed for in-plane electrode fabrication, low-temperature transport measurements, and direct visualization by TEM (Morkved, T. L., et al., *Polymer*, 39: 3871-3875 (1998)). The electrodes were constructed by spinning polymer resist layers onto Si₃N₄ substrates and exposing them to a scanned electron beam. The electron beam demarcated the electrode sites. The exposed polymer was etched away, and gold vapor was applied to fill the resulting gaps. Finally, the remaining polymer was dissolved away, leaving the gold in

the pattern inscribed by the electron beam. Typically, gaps between electrodes were 2-10 μm . NM fibers with polydispersed lengths ($>2 \mu\text{m}$) were randomly deposited on the electrodes. Binding of the protein fibers to the electrodes and bridging of the gap between the electrodes were confirmed by AFM. Current (I) and voltage (V) readings were taken as electricity was applied to the electrodes and the I-V curve for bare fibers showed a very high resistance ($R > 10^{14} \Omega$), with no measurable conductivity. Thus, NM amyloid fibers are by themselves good insulators.

E. NM Fibers can be Converted into Conducting Nanowires with Low Ohmic Resistance

[0375] NM fibers were converted to conducting nanowires by a multistep process. A derivative of NM was used that was genetically engineered to contain a cysteine residue that remained accessible after fiber formation (See, for example, Examples 9 and 10 above, and (Scheibel, T., et al., *Curr. Biol.*, 11: 366-369 (2001)). This derivative, NM^{K184C} , assembled in vitro with kinetics that were indistinguishable from those of the wild-type protein and led to fibers with the same physical properties. Monomaleimido Nanogold (Nanoprobes), which has the chemical specificity to form covalent links with the sulfhydryl groups of cysteine residues, was covalently cross-linked to NM^{K184C} fibers. The gold particles had a diameter of 1.4 nm and their distribution along the surface of the NM^{K184C} fibers was confirmed by TEM. Importantly, linking Nanogold covalently to NM fibers affected neither fiber stability nor fiber morphology.

[0376] As the distance between the NM^{K184C} cysteine residues in a fiber is $\approx 3\text{-}5 \text{ nm}$ and the Nanogold particles have a diameter of only 1.4 nm, it was necessary to bridge the particles with metal to gain conductivity. GoldEnhance LM (Nanoprobes) was first used, by which gold ions are deposited from solution onto the preexisting particles of Nanogold, followed by chemical reduction of the gold ions to form metallic gold. This process itself was inefficient in gaining conductivity, because binding and reducing the soluble gold ions did not fill all of the gaps between the covalently linked Nanogold particles as determined by TEM and AFM.

[0377] A different enhancement protocol (gold toning, FIG. 5) proved much more efficient. The Nanogold particles (FIG. 5, number 2) on the labeled fibers (FIG. 5, number 1) acted as promoters for reducing silver ions (FIG. 5, number 3) (LI Silver, Nanoprobes) from a solution. The resulting silver-coated fiber-bound Nanogold particles were then gold-enhanced with GoldEnhance LM (FIG. 5, number 4). This gold-toning technique led to fibers with densely packed gold particles. The gold-toned fibers showed a significant increase in diameter from 9-11 nm (bare fibers; FIG. 6, number 1) to 80-200 nm (labeled fibers; FIG. 6, number 2), with the diameter of the resulting fiber strictly depending on the length of exposure time of both the silver and the gold enhancement solution (longer exposure time = thicker fiber). The diameters of the metal wires varied somewhat with different batches of fibers and gold- and silver-toning solutions but were extremely consistent within reactions, i.e., all were within a 10% range. Gold toning was remarkably specific for fibers that had been covalently labeled with Nanogold particles. When NM^{K184C} fibers that were linked to Nanogold were incubated together with a large excess of unlabeled NM^{K184C} fibers, the toning process was restricted to labeled fibers (FIG. 6). Furthermore, the diameters of the wires were consistent within single experiments with fixed exposure times. There-

fore, controlling the enhancement exposure time controlled the thickness for the resulting gold wires.

[0378] The electrical behavior of NM-templated metallic fibers was assessed by randomly depositing fibers with a length $>2 \mu\text{m}$ and covalently attached Nanogold particles on patterned electrodes, followed by gold toning to form metallic continuous gold nanowires (FIGS. 7-9). Although no background deposition of gold had been detected on unlabeled NM fibers deposited on mica, some gold deposition did occur when enhancement was performed on the Si_3N_4 electrodes. No conductivity was detected in cases where the gold nanowires did not bridge the electrode gap (FIG. 7). In contrast, conductivity was readily detected when single or multiple gold-toned nanowires crossed the gap. I-V curves were linear (FIG. 8), exhibiting ohmic conductivity with low resistance ($R = 86 \Omega$ for fibers with diameters of $\approx 100 \text{ nm}$; this resistance was exhibited in each of six repeated measurements with $<1 \Omega$ variation, and with one to four bridging nanowires). The resistance measurements were stable within tenths of ohms within any given fiber (FIG. 8). Such an ohmic response indicates continuous, metallic connections across the sample. The low resistance is that expected for grain-boundary-dominated transport in a polycrystalline metal. In most cases the current was independent of the voltage scan direction and experiments could be repeated several times with the same pair of electrodes and the same nanowire. Notably, in some instances fibers were vaporized (FIG. 9, number 2) from the electrodes when the voltage was increased after the initial conductivity measurements were finished (FIG. 9). This vaporization is a consequence of Joule heating in which the power delivered to the fiber by the current results in a temperature increase sufficient to vaporize the fiber. The Joule heating power depends not only on the applied voltage but also on fiber resistance, which will vary with fiber length and other factors. Bridging fibers (FIG. 9, number 1) were vaporized and did not reassemble, but non-bridging fibers remained. In such cases conductivity was lost on remeasurement. This loss of conductivity confirmed that the bridging fibers were the active nanowires and demonstrated that they can act as fuses at higher voltages and currents.

[0379] The foregoing experiments demonstrate that NM protein fibers are excellent candidates for nanocircuit construction. They are exceedingly good insulators without metal coating ($R > 10^{14} \Omega$) and have very good electrical conductivity with gold and silver coating ($R = 86 \Omega$) and linear I-V curves. Previously the least resistance achieved with metallized proteinaceous material was of the order of 200 k Ω , $>1,000$ times greater than the resistance for metallized NM fibers (Fritzsche, W., et al. *Appl. Phys. Lett.*, 75: 2854-2856 (1999)).

[0380] The diameter of the wires produced was 80-200 nm, well below the dimensions accessible by standard electronic manufacturing methods. Having achieved the construction of wires with these dimensions, methods to produce even thinner ones are possible. The thickness of these wires was dictated by the relatively large amounts of silver and gold enhancement that were required to fill the gaps between the Nanogold particles attached to cysteine residues (FIGS. 5 and 6). The sizes of these gaps is reduced by introducing additional cysteines into NM (or using other residues), thus providing more frequent binding sites for the gold particles. Smaller gaps between gold particles will require less enhancement to make contacts continuous, and the resulting

wire is thinner. This smaller diameter will allow the manufacture of more intricate circuits and could potentially provide a new model system for quantum confinement and single-electron charging effects when electrons tunnel through restricted pathways (Halperin, W. P., *Rev. Mod. Phys.*, 58: 533-606 (1986); Kastner, M. A., *Rev. Mod. Phys.*, 64: 849-858 (1992); Grabert, H., et al., *Single Charge Tunneling* (Plenum, New York) (1992); Timp, G. L., ed., *Nanotechnology* (Springer, New York) (1999)).

Example 13

Production of Semiconductor Nanowires Built by Controlled Self-Assembly of Amyloid Fibers and Selective Semiconducting Material Deposition

[0381] The following example describes procedures to produce semiconductor nanowires built by controlled self-assembly of amyloid fibrils and selective semiconducting material deposition.

[0382] The Sup35 C terminus (e.g., amino acid 246 to 685) lies externally along the length of Sup35 fibers. Thus by replacing the C terminus with semiconductor binding peptides, and by binding semiconducting materials to those peptides, the fibrils are used to produce continuous self-assembling semiconductor wires.

[0383] Peptides with binding sites specific for different semiconductors are isolated using phage-display technology as described by Whaley et al. (Whaley, et al., *Nature*, 405: 665-668 (2000)) and Mao et al. (Mao et al., *Science*, 303: 213-217 (2004)), both of which are incorporated herein by reference. Amino acid sequences encoding the peptides identified as having semiconductor binding activity are then attached to the C-terminus of Sup35 NM, as a replacement of substitution for all or part of the wild type Sup35p C-terminus, using recombinant DNA techniques. Alternatively, the peptides identified as having semiconductor binding activity are cross-linked to the native amino acid sequence of the NM region of Sup35p (i.e., the C terminus would not be present).

[0384] Subsequently, semiconductor materials such as GaAs, ZnS, CdS, InP and Si are incorporated along the length of NM fibers (using the binding peptides as initial sites of attachment) to produce a continuous semiconductor wire.

[0385] While the present invention has been described in terms of specific embodiments, it is understood that variations and modifications will occur to those in the art, all of which are intended as aspects of the present invention. Accordingly, only such limitations as appear in the claims should be placed on the invention.

SEQUENCE LISTING

```

<160> NUMBER OF SEQ ID NOS: 65

<210> SEQ ID NO 1
<211> LENGTH: 3321
<212> TYPE: DNA
<213> ORGANISM: Saccharomyces cerevisiae
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (739)..(2796)

<400> SEQUENCE: 1

agaaattaaa gctacttaca acaacggtct actacaaatt aagggtccta aaattgtcaa      60
tgacactgaa aagccgaagc caaaaaagag gatcgccatt gaggaatac cgcacgaaga      120
attggagttt gaagaaaatc ccaaccctac ggtagaaaat tgaatatcgt atctgtttat      180
acacacatac atacatttat atttataata agcggttaaaa ttctgcgcaga atatctgtca      240
accacacaaa aatcatacaa cgaatggtat atgcttcatt tctttgtttc gcattagctg      300
cgctatttga ctcaaattat tattttttac taagacgacg cgtcacagtg ttcgagctctg      360
tgtcatttct tttgtaattc tcttaaacca cttcataaag ttgtgaagtt catagcaaaa      420
ttcttccgca aaaagatgaa tcttagttct cagcccacca aaagaggtag atgctaagat      480
catacagaag ttattgtcac ttcttacctt gctcttaaat gtacattaca accgggtatt      540
atatcttaca tcatcgtata atatgatctt tctttatgga gaaaattttt ttttactcog      600
accaaagctc ccattgcttc tgaagagtgt agtgtatatt ggtacatctt ctcttgaaag      660
actccattgt actgtaacaa aaagcggttt cttcatcgac ttgctcgtaa taacatctat      720
atctgcccac tagcaaca atg tcg gat tca aac caa ggc aac aat cag caa      771
Met Ser Asp Ser Asn Gln Gly Asn Asn Gln Gln
1             5             10

aac tac cag caa tac agc cag aac ggt aac caa caa caa ggt aac aac      819

```

-continued

Asn 15	Tyr	Gln	Gln	Tyr	Ser 20	Gln	Asn	Gly	Asn 25	Gln	Gln	Gln	Gly	Asn	Asn	
aga	tac	caa	ggt	tat	caa	gct	tac	aat	gct	caa	gcc	caa	cct	gca	ggt	867
Arg 30	Tyr	Gln	Gly	Tyr	Gln	Ala	Tyr	Asn	Ala	Gln	Ala	Gln	Pro	Ala	Gly	
ggg	tac	tac	caa	aat	tac	caa	ggt	tat	tct	ggg	tac	caa	caa	ggt	ggc	915
Gly 45	Tyr	Tyr	Gln	Asn	Tyr	Gln	Gly	Tyr	Ser	Gly	Tyr	Gln	Gln	Gly	Gly	
tat	caa	cag	tac	aat	ccc	gac	gcc	ggt	tac	cag	caa	cag	tat	aat	cct	963
Tyr 60	Gln	Gln	Tyr	Asn	Pro	Asp	Ala	Gly	Tyr	Gln	Gln	Gln	Tyr	Asn	Pro	
caa	gga	ggc	tat	caa	cag	tac	aat	cct	caa	ggc	ggt	tat	cag	cag	caa	1011
Gln 80	Gly	Gly	Tyr	Gln	Gln	Tyr	Asn	Pro	Gln	Gly	Gly	Tyr	Gln	Gln	Gln	
ttc	aat	cca	caa	ggt	ggc	cgt	gga	aat	tac	aaa	aac	ttc	aac	tac	aat	1059
Phe 95	Asn	Pro	Gln	Gly	Gly	Arg	Gly	Asn	Tyr	Lys	Asn	Phe	Asn	Tyr	Asn	
aac	aat	ttg	caa	gga	tat	caa	gct	ggt	ttc	caa	cca	cag	tct	caa	ggt	1107
Asn 110	Asn	Leu	Gln	Gly	Tyr	Gln	Ala	Gly	Phe	Gln	Pro	Gln	Ser	Gln	Gly	
atg	tct	ttg	aac	gac	ttt	caa	aag	caa	caa	aag	cag	gcc	gct	ccc	aaa	1155
Met 125	Ser	Leu	Asn	Asp	Phe	Gln	Lys	Gln	Gln	Lys	Gln	Ala	Ala	Pro	Lys	
cca	aag	aag	act	ttg	aag	ctt	gtc	tcc	agt	tcc	ggt	atc	aag	ttg	gcc	1203
Pro 140	Lys	Lys	Thr	Leu	Lys	Leu	Val	Ser	Ser	Ser	Gly	Ile	Lys	Leu	Ala	
aat	gct	acc	aag	aag	gtt	ggc	aca	aaa	cct	gcc	gaa	tct	gat	aag	aaa	1251
Asn 160	Ala	Thr	Lys	Lys	Val	Gly	Thr	Lys	Pro	Ala	Glu	Ser	Asp	Lys	Lys	
gag	gaa	gag	aag	tct	gct	gaa	acc	aaa	gaa	cca	act	aaa	gag	cca	aca	1299
Glu 175	Glu	Glu	Lys	Ser	Ala	Glu	Thr	Lys	Glu	Pro	Thr	Lys	Glu	Pro	Thr	
aag	gtc	gaa	gaa	cca	gtt	aaa	aag	gag	gag	aaa	cca	gtc	cag	act	gaa	1347
Lys 190	Val	Glu	Glu	Pro	Val	Lys	Lys	Glu	Glu	Lys	Pro	Val	Gln	Thr	Glu	
gaa	aag	acg	gag	gaa	aaa	tcg	gaa	ctt	cca	aag	gta	gaa	gac	ctt	aaa	1395
Glu 205	Lys	Thr	Glu	Glu	Lys	Ser	Glu	Leu	Pro	Lys	Val	Glu	Asp	Leu	Lys	
atc	tct	gaa	tca	aca	cat	aat	acc	aac	aat	gcc	aat	ggt	acc	agt	gct	1443
Ile 220	Ser	Glu	Ser	Thr	His	Asn	Thr	Asn	Asn	Ala	Asn	Val	Thr	Ser	Ala	
gat	gcc	ttg	atc	aag	gaa	cag	gaa	gaa	gaa	gtg	gat	gac	gaa	ggt	ggt	1491
Asp 240	Ala	Leu	Ile	Lys	Glu	Gln	Glu	Glu	Glu	Val	Asp	Asp	Glu	Val	Val	
aac	gat	atg	ttt	ggt	ggt	aaa	gat	cac	ggt	tct	tta	att	ttc	atg	ggt	1539
Asn 255	Asp	Met	Phe	Gly	Gly	Lys	Asp	His	Val	Ser	Leu	Ile	Phe	Met	Gly	
cat	ggt	gat	gcc	ggt	aaa	tct	act	atg	ggt	ggt	aat	cta	cta	tac	ttg	1587
His 270	Val	Asp	Ala	Gly	Lys	Ser	Thr	Met	Gly	Gly	Asn	Leu	Leu	Tyr	Leu	
act	ggc	tct	gtg	gat	aag	aga	act	att	gag	aaa	tat	gaa	aga	gaa	gcc	1635
Thr 285	Gly	Ser	Val	Asp	Lys	Arg	Thr	Ile	Glu	Lys	Tyr	Glu	Arg	Glu	Ala	
aag	gat	gca	ggc	aga	caa	ggt	tggt	tac	ttg	tca	tggt	gtc	atg	gat	acc	1683
Lys 300	Asp	Ala	Gly	Arg	Gln	Gly	Trp	Tyr	Leu	Ser	Trp	Val	Met	Asp	Thr	
aac	aaa	gaa	gaa	aga	aat	gat	ggt	aag	act	atc	gaa	ggt	ggt	aag	gcc	1731

-continued

Asn Lys Glu Glu Arg	Asn Asp Gly Lys Thr	Ile Glu Val Gly Lys Ala	
320	325	330	
tac ttt gaa act gaa	aaa agg cgt tat acc	ata ttg gat gct cct ggt	1779
Tyr Phe Glu Thr Glu	Lys Arg Arg Tyr Thr	Ile Leu Asp Ala Pro Gly	
335	340	345	
cat aaa atg tac gtt	tcc gag atg atc ggt	ggt gct tct caa gct gat	1827
His Lys Met Tyr Val	Ser Glu Met Ile Gly	Gly Ala Ser Gln Ala Asp	
350	355	360	
gtt ggt gtt ttg gtc	att tcc gcc aga aag	ggt gag tac gaa acc ggt	1875
Val Gly Val Leu Val	Ile Ser Ala Arg Lys	Gly Glu Tyr Glu Thr Gly	
365	370	375	
ttt gag aga ggt ggt	caa act cgt gaa cac	gcc cta ttg gcc aag acc	1923
Phe Glu Arg Gly Gly	Gln Thr Arg Glu His	Ala Leu Leu Ala Lys Thr	
380	385	390 395	
caa ggt gtt aat aag	atg gtt gtc gtc gta	aat aag atg gat gac cca	1971
Gln Gly Val Asn Lys	Met Val Val Val Val	Asn Lys Met Asp Asp Pro	
400	405	410	
acc gtt aac tgg tct	aag gaa cgt tac gac	caa tgt gtg agt aat gtc	2019
Thr Val Asn Trp Ser	Lys Glu Arg Tyr Asp	Gln Cys Val Ser Asn Val	
415	420	425	
agc aat ttc ttg aga	gca att ggt tac aac	att aag aca gac gtt gta	2067
Ser Asn Phe Leu Arg	Ala Ile Gly Tyr Asn	Ile Lys Thr Asp Val Val	
430	435	440	
ttt atg cca gta tcc	ggc tac agt ggt gca	aat ttg aaa gat cac gta	2115
Phe Met Pro Val Ser	Gly Tyr Ser Gly Ala	Asn Leu Lys Asp His Val	
445	450	455	
gat cca aaa gaa tgc	cca tgg tac acc ggc	cca act ctg tta gaa tat	2163
Asp Pro Lys Glu Cys	Pro Trp Tyr Thr Gly	Pro Thr Leu Leu Glu Tyr	
460	465	470 475	
ctg gat aca atg aac	cac gtc gac cgt cac	atc aat gct cca ttc atg	2211
Leu Asp Thr Met Asn	His Val Asp Arg His	Ile Asn Ala Pro Phe Met	
480	485	490	
ttg cct att gcc gct	aag atg aag gat cta	ggt acc atc gtt gaa ggt	2259
Leu Pro Ile Ala Ala	Lys Met Lys Asp Leu	Gly Thr Ile Val Glu Gly	
495	500	505	
aaa att gaa tcc ggt	cat atc aaa aag ggt	caa tcc acc cta ctg atg	2307
Lys Ile Glu Ser Gly	His Ile Lys Lys Gly	Gln Ser Thr Leu Leu Met	
510	515	520	
cct aac aaa acc gct	gtg gaa att caa aat	att tac aac gaa act gaa	2355
Pro Asn Lys Thr Ala	Val Glu Ile Gln Asn	Ile Tyr Asn Glu Thr Glu	
525	530	535	
aat gaa gtt gat atg	gct atg tgt ggt gag	caa gtt aaa cta aga atc	2403
Asn Glu Val Asp Met	Ala Met Cys Gly Glu	Gln Val Lys Leu Arg Ile	
540	545	550 555	
aaa ggt gtt gaa gaa	gaa gac att tca cca	ggt ttt gta cta aca tcg	2451
Lys Gly Val Glu Glu	Glu Asp Ile Ser Pro	Gly Phe Val Leu Thr Ser	
560	565	570	
cca aag aac cct atc	aag agt gtt acc aag	ttt gta gct caa att gct	2499
Pro Lys Asn Pro Ile	Lys Ser Val Thr Lys	Phe Val Ala Gln Ile Ala	
575	580	585	
att gta gaa tta aaa	tct atc ata gca gcc	ggt ttt tca tgt gtt atg	2547
Ile Val Glu Leu Lys	Ser Ile Ile Ala Ala	Gly Phe Ser Cys Val Met	
590	595	600	
cat gtt cat aca gca	att gaa gag gta cat	att gtt aag tta ttg cac	2595
His Val His Thr Ala	Ile Glu Glu Val His	Ile Val Lys Leu Leu His	
605	610	615	
aaa tta gaa aag ggt	acc aac cgt aag tca	aag aaa cca cct gct ttt	2643

-continued

Lys Leu Glu Lys Gly Thr Asn Arg Lys Ser Lys Lys Pro Pro Ala Phe
 620 625 630 635

gct aag aag ggt atg aag gtc atc gct gtt tta gaa act gaa gct cca 2691
 Ala Lys Lys Gly Met Lys Val Ile Ala Val Leu Glu Thr Glu Ala Pro
 640 645 650

gtt tgt gtg gaa act tac caa gat tac cct caa tta ggt aga ttc act 2739
 Val Cys Val Glu Thr Tyr Gln Asp Tyr Pro Gln Leu Gly Arg Phe Thr
 655 660 665

ttg aga gat caa ggt acc aca ata gca att ggt aaa att gtt aaa att 2787
 Leu Arg Asp Gln Gly Thr Thr Ile Ala Ile Gly Lys Ile Val Lys Ile
 670 675 680

gcc gag taa atttcttgca aacataagta aatgcaaaca caataatacc 2836
 Ala Glu
 685

gatcataaag cttttcttc tatattaaaa aacaaggttt aataaagctg ttatatatat 2896

atatatatat atagacgtat aattagttta gttctttttg taccatatac cataaacaag 2956

gtaaacttca cctctcaata tatctagaat ttcataaaaa tatctagcaa ggtttcaact 3016

ccttcaatca cgttttcttc ataacccttc cccggcggtta tttcagaatg tgcaaaatct 3076

attagtgaca tggaactcaa agaaccagtt gtttttttgt cctttgggtcc ttcgctgctt 3136

ccctcggcat catcatcatc atcatcatca ttatcatcat cgtcgtcatc atcgtctata 3196

aaatcatctc gcataagttt gtcaacatca tttagtaatt cccatcgtc cgggtctcct 3256

tcgtaaataa acaaaagact acttgatc attctaactt cttcttctag catagtatta 3316

taaaa 3321

<210> SEQ ID NO 2
 <211> LENGTH: 685
 <212> TYPE: PRT
 <213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 2

Met Ser Asp Ser Asn Gln Gly Asn Asn Gln Gln Asn Tyr Gln Gln Tyr
 1 5 10 15

Ser Gln Asn Gly Asn Gln Gln Gln Gly Asn Asn Arg Tyr Gln Gly Tyr
 20 25 30

Gln Ala Tyr Asn Ala Gln Ala Gln Pro Ala Gly Gly Tyr Tyr Gln Asn
 35 40 45

Tyr Gln Gly Tyr Ser Gly Tyr Gln Gln Gly Gly Tyr Gln Gln Tyr Asn
 50 55 60

Pro Asp Ala Gly Tyr Gln Gln Gln Tyr Asn Pro Gln Gly Gly Tyr Gln
 65 70 75 80

Gln Tyr Asn Pro Gln Gly Gly Tyr Gln Gln Gln Phe Asn Pro Gln Gly
 85 90 95

Gly Arg Gly Asn Tyr Lys Asn Phe Asn Tyr Asn Asn Asn Leu Gln Gly
 100 105 110

Tyr Gln Ala Gly Phe Gln Pro Gln Ser Gln Gly Met Ser Leu Asn Asp
 115 120 125

Phe Gln Lys Gln Gln Lys Gln Ala Ala Pro Lys Pro Lys Lys Thr Leu
 130 135 140

Lys Leu Val Ser Ser Ser Gly Ile Lys Leu Ala Asn Ala Thr Lys Lys
 145 150 155 160

Val Gly Thr Lys Pro Ala Glu Ser Asp Lys Lys Glu Glu Glu Lys Ser

-continued

165	170	175
Ala Glu Thr Lys Glu Pro Thr Lys Glu Pro Thr Lys Val Glu Glu Pro		
180	185	190
Val Lys Lys Glu Glu Lys Pro Val Gln Thr Glu Glu Lys Thr Glu Glu		
195	200	205
Lys Ser Glu Leu Pro Lys Val Glu Asp Leu Lys Ile Ser Glu Ser Thr		
210	215	220
His Asn Thr Asn Asn Ala Asn Val Thr Ser Ala Asp Ala Leu Ile Lys		
225	230	235
Glu Gln Glu Glu Glu Val Asp Asp Glu Val Val Asn Asp Met Phe Gly		
245	250	255
Gly Lys Asp His Val Ser Leu Ile Phe Met Gly His Val Asp Ala Gly		
260	265	270
Lys Ser Thr Met Gly Gly Asn Leu Leu Tyr Leu Thr Gly Ser Val Asp		
275	280	285
Lys Arg Thr Ile Glu Lys Tyr Glu Arg Glu Ala Lys Asp Ala Gly Arg		
290	295	300
Gln Gly Trp Tyr Leu Ser Trp Val Met Asp Thr Asn Lys Glu Glu Arg		
305	310	315
Asn Asp Gly Lys Thr Ile Glu Val Gly Lys Ala Tyr Phe Glu Thr Glu		
325	330	335
Lys Arg Arg Tyr Thr Ile Leu Asp Ala Pro Gly His Lys Met Tyr Val		
340	345	350
Ser Glu Met Ile Gly Gly Ala Ser Gln Ala Asp Val Gly Val Leu Val		
355	360	365
Ile Ser Ala Arg Lys Gly Glu Tyr Glu Thr Gly Phe Glu Arg Gly Gly		
370	375	380
Gln Thr Arg Glu His Ala Leu Leu Ala Lys Thr Gln Gly Val Asn Lys		
385	390	395
Met Val Val Val Val Asn Lys Met Asp Asp Pro Thr Val Asn Trp Ser		
405	410	415
Lys Glu Arg Tyr Asp Gln Cys Val Ser Asn Val Ser Asn Phe Leu Arg		
420	425	430
Ala Ile Gly Tyr Asn Ile Lys Thr Asp Val Val Phe Met Pro Val Ser		
435	440	445
Gly Tyr Ser Gly Ala Asn Leu Lys Asp His Val Asp Pro Lys Glu Cys		
450	455	460
Pro Trp Tyr Thr Gly Pro Thr Leu Leu Glu Tyr Leu Asp Thr Met Asn		
465	470	475
His Val Asp Arg His Ile Asn Ala Pro Phe Met Leu Pro Ile Ala Ala		
485	490	495
Lys Met Lys Asp Leu Gly Thr Ile Val Glu Gly Lys Ile Glu Ser Gly		
500	505	510
His Ile Lys Lys Gly Gln Ser Thr Leu Leu Met Pro Asn Lys Thr Ala		
515	520	525
Val Glu Ile Gln Asn Ile Tyr Asn Glu Thr Glu Asn Glu Val Asp Met		
530	535	540
Ala Met Cys Gly Glu Gln Val Lys Leu Arg Ile Lys Gly Val Glu Glu		
545	550	555
Glu Asp Ile Ser Pro Gly Phe Val Leu Thr Ser Pro Lys Asn Pro Ile		
565	570	575

-continued

Lys Ser Val Thr Lys Phe Val Ala Gln Ile Ala Ile Val Glu Leu Lys
580 585 590

Ser Ile Ile Ala Ala Gly Phe Ser Cys Val Met His Val His Thr Ala
595 600 605

Ile Glu Glu Val His Ile Val Lys Leu Leu His Lys Leu Glu Lys Gly
610 615 620

Thr Asn Arg Lys Ser Lys Lys Pro Pro Ala Phe Ala Lys Lys Gly Met
625 630 635 640

Lys Val Ile Ala Val Leu Glu Thr Glu Ala Pro Val Cys Val Glu Thr
645 650 655

Tyr Gln Asp Tyr Pro Gln Leu Gly Arg Phe Thr Leu Arg Asp Gln Gly
660 665 670

Thr Thr Ile Ala Ile Gly Lys Ile Val Lys Ile Ala Glu
675 680 685

<210> SEQ ID NO 3
<211> LENGTH: 1427
<212> TYPE: DNA
<213> ORGANISM: *Saccharomyces cerevisiae*
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (182)..(1246)

<400> SEQUENCE: 3

ctcgcagggtg aaaagaatag caaaaatctt tccttttcaa acagctcatt tgggaattggt 60

tatagcactg aattgaatcg aagaggaata aagatcccc gtacgaactt ctttattttt 120

agtttttcat tttttggtat tagtcatatt gttttaagct gcaaatgaag ttgtacacca 180

a atg atg aat aac aac ggc aac caa gtg tcg aat ctc tcc aat gcg ctc 229
Met Met Asn Asn Asn Gly Asn Gln Val Ser Asn Leu Ser Asn Ala Leu
1 5 10 15

cgt caa gta aac ata gga aac agg aac agt aat aca acc acc gat caa 277
Arg Gln Val Asn Ile Gly Asn Arg Asn Ser Asn Thr Thr Thr Asp Gln
20 25 30

agt aat ata aat ttt gaa ttt tca aca ggt gta aat aat aat aat aat 325
Ser Asn Ile Asn Phe Glu Phe Ser Thr Gly Val Asn Asn Asn Asn Asn
35 40 45

aac aat agc agt agt aat aac aat aat gtt caa aac aat aac agc ggc 373
Asn Asn Ser Ser Ser Asn Asn Asn Asn Val Gln Asn Asn Asn Ser Gly
50 55 60

cgc aat ggt agc caa aat aat gat aac gag aat aat atc aag aat acc 421
Arg Asn Gly Ser Gln Asn Asn Asp Asn Glu Asn Asn Ile Lys Asn Thr
65 70 75 80

tta gaa caa cat cga caa caa caa cag gca ttt tcg gat atg agt cac 469
Leu Glu Gln His Arg Gln Gln Gln Gln Ala Phe Ser Asp Met Ser His
85 90 95

gtg gag tat tcc aga att aca aaa ttt ttt caa gaa caa cca ctg gag 517
Val Glu Tyr Ser Arg Ile Thr Lys Phe Phe Gln Glu Gln Pro Leu Glu
100 105 110

gga tat acc ctt ttc tct cac agg tct gcg cct aat gga ttc aaa gtt 565
Gly Tyr Thr Leu Phe Ser His Arg Ser Ala Pro Asn Gly Phe Lys Val
115 120 125

gct ata gta cta agt gaa ctt gga ttt cat tat aac aca atc ttc cta 613
Ala Ile Val Leu Ser Glu Leu Gly Phe His Tyr Asn Thr Ile Phe Leu
130 135 140

gat ttc aat ctt ggc gaa cat agg gcc ccc gaa ttt gtg tct gtg aac 661

-continued

Asp Phe Asn Leu Gly Glu His Arg Ala Pro Glu Phe Val Ser Val Asn	
145 150 155 160	
cct aat gca aga gtt cca gct tta atc gat cat ggt atg gac aac ttg	709
Pro Asn Ala Arg Val Pro Ala Leu Ile Asp His Gly Met Asp Asn Leu	
165 170 175	
tct att tgg gaa tca ggg gcg att tta tta cat ttg gta aat aaa tat	757
Ser Ile Trp Glu Ser Gly Ala Ile Leu Leu His Leu Val Asn Lys Tyr	
180 185 190	
tac aaa gag act ggt aat cca tta ctc tgg tcc gat gat tta gct gac	805
Tyr Lys Glu Thr Gly Asn Pro Leu Leu Trp Ser Asp Asp Leu Ala Asp	
195 200 205	
caa tca caa atc aac gca tgg ttg ttc ttc caa acg tca ggg cat gcg	853
Gln Ser Gln Ile Asn Ala Trp Leu Phe Phe Gln Thr Ser Gly His Ala	
210 215 220	
cca atg att gga caa gct tta cat ttc aga tac ttc cat tca caa aag	901
Pro Met Ile Gly Gln Ala Leu His Phe Arg Tyr Phe His Ser Gln Lys	
225 230 235 240	
ata gca agt gct gta gaa aga tat acg gat gag gtt aga aga gtt tac	949
Ile Ala Ser Ala Val Glu Arg Tyr Thr Asp Glu Val Arg Arg Val Tyr	
245 250 255	
ggg gta gtg gag atg gcc ttg gct gaa cgt aga gaa gcg ctg gtg atg	997
Gly Val Val Glu Met Ala Leu Ala Glu Arg Arg Glu Ala Leu Val Met	
260 265 270	
gaa tta gac acg gaa aat gcg gct gca tac tca gct ggt aca aca cca	1045
Glu Leu Asp Thr Glu Asn Ala Ala Ala Tyr Ser Ala Gly Thr Thr Pro	
275 280 285	
atg tca caa agt cgt ttc ttt gat tat ccc gta tgg ctt gta gga gat	1093
Met Ser Gln Ser Arg Phe Phe Asp Tyr Pro Val Trp Leu Val Gly Asp	
290 295 300	
aaa tta act ata gca gat ttg gcc ttt gtc cca tgg aat aat gtc gtg	1141
Lys Leu Thr Ile Ala Asp Leu Ala Phe Val Pro Trp Asn Asn Val Val	
305 310 315 320	
gat aga att ggc att aat atc aaa att gaa ttt cca gaa gtt tac aaa	1189
Asp Arg Ile Gly Ile Asn Ile Lys Ile Glu Phe Pro Glu Val Tyr Lys	
325 330 335	
tgg acg aag cat atg atg aga aga ccc gcg gtc atc aag gca ttg cgt	1237
Trp Thr Lys His Met Met Arg Arg Pro Ala Val Ile Lys Ala Leu Arg	
340 345 350	
ggg gaa tga aggctgcttt aaaaacaaga aagaagaag aaggaggaaa	1286
Gly Glu	
agaaggttat aagggtatgt atataggcag acaaaaagga aaattaagtg caaatataaa	1346
caaaaatgtc atagaagtat ataatagttt tgaattttct gttgcttcta tttattcttt	1406
gttacccecaa ccacagaatt c	1427

<210> SEQ ID NO 4
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 4

Met Met Asn Asn Asn Gly Asn Gln Val Ser Asn Leu Ser Asn Ala Leu	
1 5 10 15	
Arg Gln Val Asn Ile Gly Asn Arg Asn Ser Asn Thr Thr Thr Asp Gln	
20 25 30	
Ser Asn Ile Asn Phe Glu Phe Ser Thr Gly Val Asn Asn Asn Asn Asn	
35 40 45	

-continued

Asn Asn Ser Ser Ser Asn Asn Asn Asn Val Gln Asn Asn Asn Ser Gly
 50 55 60
 Arg Asn Gly Ser Gln Asn Asn Asp Asn Glu Asn Asn Ile Lys Asn Thr
 65 70 75 80
 Leu Glu Gln His Arg Gln Gln Gln Gln Ala Phe Ser Asp Met Ser His
 85 90 95
 Val Glu Tyr Ser Arg Ile Thr Lys Phe Phe Gln Glu Gln Pro Leu Glu
 100 105 110
 Gly Tyr Thr Leu Phe Ser His Arg Ser Ala Pro Asn Gly Phe Lys Val
 115 120 125
 Ala Ile Val Leu Ser Glu Leu Gly Phe His Tyr Asn Thr Ile Phe Leu
 130 135 140
 Asp Phe Asn Leu Gly Glu His Arg Ala Pro Glu Phe Val Ser Val Asn
 145 150 155 160
 Pro Asn Ala Arg Val Pro Ala Leu Ile Asp His Gly Met Asp Asn Leu
 165 170 175
 Ser Ile Trp Glu Ser Gly Ala Ile Leu Leu His Leu Val Asn Lys Tyr
 180 185 190
 Tyr Lys Glu Thr Gly Asn Pro Leu Leu Trp Ser Asp Asp Leu Ala Asp
 195 200 205
 Gln Ser Gln Ile Asn Ala Trp Leu Phe Phe Gln Thr Ser Gly His Ala
 210 215 220
 Pro Met Ile Gly Gln Ala Leu His Phe Arg Tyr Phe His Ser Gln Lys
 225 230 235 240
 Ile Ala Ser Ala Val Glu Arg Tyr Thr Asp Glu Val Arg Arg Val Tyr
 245 250 255
 Gly Val Val Glu Met Ala Leu Ala Glu Arg Arg Glu Ala Leu Val Met
 260 265 270
 Glu Leu Asp Thr Glu Asn Ala Ala Ala Tyr Ser Ala Gly Thr Thr Pro
 275 280 285
 Met Ser Gln Ser Arg Phe Phe Asp Tyr Pro Val Trp Leu Val Gly Asp
 290 295 300
 Lys Leu Thr Ile Ala Asp Leu Ala Phe Val Pro Trp Asn Asn Val Val
 305 310 315 320
 Asp Arg Ile Gly Ile Asn Ile Lys Ile Glu Phe Pro Glu Val Tyr Lys
 325 330 335
 Trp Thr Lys His Met Met Arg Arg Pro Ala Val Ile Lys Ala Leu Arg
 340 345 350

Gly Glu

<210> SEQ ID NO 5
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: FLAG
 peptide

<400> SEQUENCE: 5

Asp Tyr Lys Asp Asp Asp Asp Lys
 1 5

<210> SEQ ID NO 6

-continued

<211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: FLAG
 peptide

<400> SEQUENCE: 6

Asp Tyr Lys Asp Glu Asp Asp Lys
 1 5

<210> SEQ ID NO 7
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Strep
 epitope

<400> SEQUENCE: 7

Ala Trp Arg His Pro Gln Phe Gly Gly
 1 5

<210> SEQ ID NO 8
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 Hemagglutininepitope

<400> SEQUENCE: 8

Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ile Glu Gly Arg
 1 5 10

<210> SEQ ID NO 9
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: myc epitope

<400> SEQUENCE: 9

Glu Gln Lys Leu Leu Ser Glu Glu Asp Leu Asn
 1 5 10

<210> SEQ ID NO 10
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 10

Pro Gln Gly Gly Tyr Gln Gln Tyr Asn
 1 5

<210> SEQ ID NO 11
 <211> LENGTH: 445
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: CUP1
 promoter

<400> SEQUENCE: 11

ccattaccga catttggcg ctatacgtgc atatgttcat gtagtattct gtatttaaaa 60

-continued

```

cacttttgta ttatttttcc tcatatatgt gtataggttt atacggatga ttaattatt 120
acttcaccac cctttatttc aggctgatat cttagccttg ttactagtta gaaaaagaca 180
tttttgctgt cagtcactgt caagagattc ttttgctggc atttcttcta gaagcaaaaa 240
gagcgatgcg tcttttccgc tgaaccgttc cagcaaaaaa gactaccaac gcaatatgga 300
ttgtcagaat catataaaag agaagcaaat aactccttgt cttgtatcaa ttgcattata 360
atatcttctt gttagtgcaa tatcatatag aagtcacga aatagatatt aagaaaaaca 420
aactgtacaa tcaatcaatc aatca 445

```

```

<210> SEQ ID NO 12
<211> LENGTH: 717
<212> TYPE: DNA
<213> ORGANISM: Aequorea victoria

```

```

<400> SEQUENCE: 12

```

```

atgtctaaag gtgaagaatt attcactggg gttgtcccaa ttttggttga attagatggg 60
gatgttaaat gtcacaaatt ttctgtctcc ggtgaaggty aaggatgatc tacttacggg 120
aaattgacct taaaatttat ttgtactact ggtaaattgc cagttccatg gccaacctta 180
gtcactactt tcggttatgg tgttcaatgt tttgctagat acccagatca tatgaaacaa 240
catgactttt tcaagtctgc catgccagaa gggtatgttc aagaaagaac tatttttttc 300
aaagatgacg gtaactacaa gaccagagct gaagtcaagt ttgaaggatga taccttagtt 360
aatagaatcg aattaaaagg tattgatttt aaagaagatg gtaacatttt aggtcacaaa 420
ttggaataca actataaactc tcacaatggt tacatcatgg ctgacaaaaca aaagaatggg 480
atcaaagtta acttcaaaat tagacacaaac attgaagatg gttctgttca attagctgac 540
cattatcaac aaaatactcc aattggatg ggtccagtct tgttaccaga caaccattac 600
ttatccactc aatctgcctt atccaaagat ccaaacgaaa agagagacca catggtcttg 660
ttagaatttg ttactgtctc tggattacc catggtatgg atgaattgta caaataa 717

```

```

<210> SEQ ID NO 13
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: HA
tag-encoding sequence

```

```

<400> SEQUENCE: 13

```

```

taccatagc acgtcccaga ctacgct 27

```

```

<210> SEQ ID NO 14
<211> LENGTH: 645
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: yeast
Sup35Rdelta2-5 encoding sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(645)

```

```

<400> SEQUENCE: 14

```

```

atg tcg gat tca aac caa ggc aac aat cag caa aac tac cag caa tac 48
Met Ser Asp Ser Asn Gln Gly Asn Asn Gln Gln Asn Tyr Gln Gln Tyr

```

-continued

1	5	10	15	
agc cag aac ggt aac caa caa caa ggt aac aac aga tac caa ggt tat				96
Ser Gln Asn Gly Asn Gln Gln Gln Gly Asn Asn Arg Tyr Gln Gly Tyr				
20	25	30		
caa gct tac aat gct caa gcc caa cct gca ggt ggg tac tac caa aat				144
Gln Ala Tyr Asn Ala Gln Ala Gln Pro Ala Gly Gly Tyr Tyr Gln Asn				
35	40	45		
tac caa ggt tat tct ggg tac cca caa ggt ggc cgt gga aat tac aaa				192
Tyr Gln Gly Tyr Ser Gly Tyr Pro Gln Gly Gly Arg Gly Asn Tyr Lys				
50	55	60		
aac ttc aac tac aat aac aat ttg caa gga tat caa gct ggt ttc caa				240
Asn Phe Asn Tyr Asn Asn Asn Leu Gln Gly Tyr Gln Ala Gly Phe Gln				
65	70	75	80	
cca cag tct caa ggt atg tct ttg aac gac ttt caa aag caa caa aag				288
Pro Gln Ser Gln Gly Met Ser Leu Asn Asp Phe Gln Lys Gln Gln Lys				
85	90	95		
cag gcc gct ccc aaa cca aag aag act ttg aag ctt gtc tcc agt tcc				336
Gln Ala Ala Pro Lys Pro Lys Lys Thr Leu Lys Leu Val Ser Ser Ser				
100	105	110		
ggt atc aag ttg gcc aat gct acc aag aag gtt ggc aca aaa cct gcc				384
Gly Ile Lys Leu Ala Asn Ala Thr Lys Lys Val Gly Thr Lys Pro Ala				
115	120	125		
gaa tct gat aag aaa gag gaa gag aag tct gct gaa acc aaa gaa cca				432
Glu Ser Asp Lys Lys Glu Glu Glu Lys Ser Ala Glu Thr Lys Glu Pro				
130	135	140		
act aaa gag cca aca aag gtc gaa gaa cca gtt aaa aag gag gag aaa				480
Thr Lys Glu Pro Thr Lys Val Glu Glu Pro Val Lys Lys Glu Glu Lys				
145	150	155	160	
cca gtc cag act gaa gaa aag acg gag gaa aaa tcg gaa ctt cca aag				528
Pro Val Gln Thr Glu Glu Lys Thr Glu Glu Lys Ser Glu Leu Pro Lys				
165	170	175		
gta gaa gac ctt aaa atc tct gaa tca aca cat aat acc aac aat gcc				576
Val Glu Asp Leu Lys Ile Ser Glu Ser Thr His Asn Thr Asn Asn Ala				
180	185	190		
aat gtt acc agt gct gat gcc ttg atc aag gaa cag gaa gaa gaa gtg				624
Asn Val Thr Ser Ala Asp Ala Leu Ile Lys Glu Gln Glu Glu Glu Val				
195	200	205		
gat gac gaa gtt gtt aac gat				645
Asp Asp Glu Val Val Asn Asp				
210	215			

<210> SEQ ID NO 15
 <211> LENGTH: 215
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: yeast
 Sup35Rdelta2-5 encoding sequence
 <400> SEQUENCE: 15

Met Ser Asp Ser Asn Gln Gly Asn Asn Gln Gln Asn Tyr Gln Gln Tyr				
1	5	10	15	
Ser Gln Asn Gly Asn Gln Gln Gln Gly Asn Asn Arg Tyr Gln Gly Tyr				
20	25	30		
Gln Ala Tyr Asn Ala Gln Ala Gln Pro Ala Gly Gly Tyr Tyr Gln Asn				
35	40	45		
Tyr Gln Gly Tyr Ser Gly Tyr Pro Gln Gly Gly Arg Gly Asn Tyr Lys				
50	55	60		

-continued

Asn Phe Asn Tyr Asn Asn Asn Leu Gln Gly Tyr Gln Ala Gly Phe Gln
 65 70 75 80

Pro Gln Ser Gln Gly Met Ser Leu Asn Asp Phe Gln Lys Gln Gln Lys
 85 90 95

Gln Ala Ala Pro Lys Pro Lys Lys Thr Leu Lys Leu Val Ser Ser Ser
 100 105 110

Gly Ile Lys Leu Ala Asn Ala Thr Lys Lys Val Gly Thr Lys Pro Ala
 115 120 125

Glu Ser Asp Lys Lys Glu Glu Glu Lys Ser Ala Glu Thr Lys Glu Pro
 130 135 140

Thr Lys Glu Pro Thr Lys Val Glu Glu Pro Val Lys Lys Glu Glu Lys
 145 150 155 160

Pro Val Gln Thr Glu Glu Lys Thr Glu Glu Lys Ser Glu Leu Pro Lys
 165 170 175

Val Glu Asp Leu Lys Ile Ser Glu Ser Thr His Asn Thr Asn Asn Ala
 180 185 190

Asn Val Thr Ser Ala Asp Ala Leu Ile Lys Glu Gln Glu Glu Glu Val
 195 200 205

Asp Asp Glu Val Val Asn Asp
 210 215

<210> SEQ ID NO 16
 <211> LENGTH: 813
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: yeast
 Sup35R2E2 encoding sequence
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(813)

<400> SEQUENCE: 16

atg tcg gat tca aac caa ggc aac aat cag caa aac tac cag caa tac 48
 Met Ser Asp Ser Asn Gln Gly Asn Asn Gln Gln Asn Tyr Gln Gln Tyr
 1 5 10 15

agc cag aac ggt aac caa caa caa ggt aac aac aga tac caa ggt tat 96
 Ser Gln Asn Gly Asn Gln Gln Gln Gly Asn Asn Arg Tyr Gln Gly Tyr
 20 25 30

caa gct tac aat gct caa gcc caa cct gca ggt ggg tac tac caa aat 144
 Gln Ala Tyr Asn Ala Gln Ala Gln Pro Ala Gly Gly Tyr Tyr Gln Asn
 35 40 45

tac caa ggt tat tct ggg tac caa caa ggt ggc tat caa cag tac aat 192
 Tyr Gln Gly Tyr Ser Gly Tyr Gln Gln Gly Gly Tyr Gln Gln Tyr Asn
 50 55 60

ccc caa ggt ggc tat caa cag tac aat ccc caa ggt ggc tat caa cag 240
 Pro Gln Gly Gly Tyr Gln Gln Tyr Asn Pro Gln Gly Gly Tyr Gln Gln
 65 70 75 80

tac aat ccc gac gcc ggt tac cag caa cag tat aat cct caa gga ggc 288
 Tyr Asn Pro Asp Ala Gly Tyr Gln Gln Gln Tyr Asn Pro Gln Gly Gly
 85 90 95

tat caa cag tac aat cct caa ggc ggt tat cag cag caa ttc aat cca 336
 Tyr Gln Gln Tyr Asn Pro Gln Gly Gly Tyr Gln Gln Gln Phe Asn Pro
 100 105 110

caa ggt ggc cgt gga aat tac aaa aac ttc aac tac aat aac aat ttg 384
 Gln Gly Gly Arg Gly Asn Tyr Lys Asn Phe Asn Tyr Asn Asn Asn Leu
 115 120 125

-continued

caa gga tat caa gct ggt ttc caa cca cag tct caa ggt atg tct ttg	432
Gln Gly Tyr Gln Ala Gly Phe Gln Pro Gln Ser Gln Gly Met Ser Leu	
130 135 140	
aac gac ttt caa aag caa caa aag cag gcc gct ccc aaa cca aag aag	480
Asn Asp Phe Gln Lys Gln Gln Lys Gln Ala Ala Pro Lys Pro Lys Lys	
145 150 155 160	
act ttg aag ctt gtc tcc agt tcc ggt atc aag ttg gcc aat gct acc	528
Thr Leu Lys Leu Val Ser Ser Ser Gly Ile Lys Leu Ala Asn Ala Thr	
165 170 175	
aag aag gtt ggc aca aaa cct gcc gaa tct gat aag aaa gag gaa gag	576
Lys Lys Val Gly Thr Lys Pro Ala Glu Ser Asp Lys Lys Glu Glu Glu	
180 185 190	
aag tct gct gaa acc aaa gaa cca act aaa gag cca aca aag gtc gaa	624
Lys Ser Ala Glu Thr Lys Glu Pro Thr Lys Glu Pro Thr Lys Val Glu	
195 200 205	
gaa cca gtt aaa aag gag gag aaa cca gtc cag act gaa gaa aag acg	672
Glu Pro Val Lys Lys Glu Glu Lys Pro Val Gln Thr Glu Glu Lys Thr	
210 215 220	
gag gaa aaa tcg gaa ctt cca aag gta gaa gac ctt aaa atc tct gaa	720
Glu Glu Lys Ser Glu Leu Pro Lys Val Glu Asp Leu Lys Ile Ser Glu	
225 230 235 240	
tca aca cat aat acc aac aat gcc aat gtt acc agt gct gat gcc ttg	768
Ser Thr His Asn Thr Asn Asn Ala Asn Val Thr Ser Ala Asp Ala Leu	
245 250 255	
atc aag gaa cag gaa gaa gaa gtg gat gac gaa gtt gtt aac gat	813
Ile Lys Glu Gln Glu Glu Glu Val Asp Asp Glu Val Val Asn Asp	
260 265 270	

<210> SEQ ID NO 17
 <211> LENGTH: 271
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: yeast
 Sup35R2E2 encoding sequence

<400> SEQUENCE: 17

Met Ser Asp Ser Asn Gln Gly Asn Asn Gln Gln Asn Tyr Gln Gln Tyr	
1 5 10 15	
Ser Gln Asn Gly Asn Gln Gln Gln Gly Asn Asn Arg Tyr Gln Gly Tyr	
20 25 30	
Gln Ala Tyr Asn Ala Gln Ala Gln Pro Ala Gly Gly Tyr Tyr Gln Asn	
35 40 45	
Tyr Gln Gly Tyr Ser Gly Tyr Gln Gln Gly Gly Tyr Gln Gln Tyr Asn	
50 55 60	
Pro Gln Gly Gly Tyr Gln Gln Tyr Asn Pro Gln Gly Gly Tyr Gln Gln	
65 70 75 80	
Tyr Asn Pro Asp Ala Gly Tyr Gln Gln Gln Tyr Asn Pro Gln Gly Gly	
85 90 95	
Tyr Gln Gln Tyr Asn Pro Gln Gly Gly Tyr Gln Gln Gln Phe Asn Pro	
100 105 110	
Gln Gly Gly Arg Gly Asn Tyr Lys Asn Phe Asn Tyr Asn Asn Asn Leu	
115 120 125	
Gln Gly Tyr Gln Ala Gly Phe Gln Pro Gln Ser Gln Gly Met Ser Leu	
130 135 140	
Asn Asp Phe Gln Lys Gln Gln Lys Gln Ala Ala Pro Lys Pro Lys Lys	

-continued

145	150	155	160
Thr Leu Lys Leu Val Ser Ser Ser Gly Ile Lys Leu Ala Asn Ala Thr			
165	170	175	
Lys Lys Val Gly Thr Lys Pro Ala Glu Ser Asp Lys Lys Glu Glu Glu			
180	185	190	
Lys Ser Ala Glu Thr Lys Glu Pro Thr Lys Glu Pro Thr Lys Val Glu			
195	200	205	
Glu Pro Val Lys Lys Glu Glu Lys Pro Val Gln Thr Glu Glu Lys Thr			
210	215	220	
Glu Glu Lys Ser Glu Leu Pro Lys Val Glu Asp Leu Lys Ile Ser Glu			
225	230	235	240
Ser Thr His Asn Thr Asn Asn Ala Asn Val Thr Ser Ala Asp Ala Leu			
245	250	255	
Ile Lys Glu Gln Glu Glu Glu Val Asp Asp Glu Val Val Asn Asp			
260	265	270	

<210> SEQ ID NO 18
 <211> LENGTH: 641
 <212> TYPE: DNA
 <213> ORGANISM: MOUSE
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(633)

<400> SEQUENCE: 18

atg tct aaa aag cgg cca aag cct gga ggg tgg aac acc ggt gga agc	48
Met Ser Lys Lys Arg Pro Lys Pro Gly Gly Trp Asn Thr Gly Gly Ser	
1 5 10 15	
cgg tat ccc ggg cag gga agc cct gga ggc aac cgt tac cca cct cag	96
Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly Asn Arg Tyr Pro Pro Gln	
20 25 30	
ggt ggc acc tgg ggg cag ccc cac ggt ggt ggc tgg gga caa ccc cat	144
Gly Gly Thr Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His	
35 40 45	
ggg ggc agc tgg gga caa cct cat ggt ggt agt tgg ggt cag ccc cat	192
Gly Gly Ser Trp Gly Gln Pro His Gly Gly Ser Trp Gly Gln Pro His	
50 55 60	
ggc ggt gga tgg ggc caa gga ggg ggt acc cat aat cag tgg aac aag	240
Gly Gly Gly Trp Gly Gln Gly Gly Gly Thr His Asn Gln Trp Asn Lys	
65 70 75 80	
ccc agc aaa cca aaa acc aac ctc aag cat gtg gca ggg gct gcg gca	288
Pro Ser Lys Pro Lys Thr Asn Leu Lys His Val Ala Gly Ala Ala Ala	
85 90 95	
gct ggg gca gta gtg ggg ggc ctt ggt ggc tac atg ctg ggg agc gcc	336
Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr Met Leu Gly Ser Ala	
100 105 110	
gtg agc agg ccc atg atc cat ttt ggc aac gac tgg gag gac cgc tac	384
Val Ser Arg Pro Met Ile His Phe Gly Asn Asp Trp Glu Asp Arg Tyr	
115 120 125	
tac cgt gaa aac atg tac cgc tac cct aac caa gtg tac tac agg cca	432
Tyr Arg Glu Asn Met Tyr Arg Tyr Pro Asn Gln Val Tyr Tyr Arg Pro	
130 135 140	
gtg gat cag tac agc aac cag aac aac ttc gtg cac gac tgc gtc aat	480
Val Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val His Asp Cys Val Asn	
145 150 155 160	
atc acc atc aag cag cac acg gtc acc acc acc acc aag ggg gag aac	528
Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr Thr Lys Gly Glu Asn	

-continued

```

165             170             175
ttc acc gag acc gat gtg aag atg atg gag cgc gtg gtg gag cag atg   576
Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg Val Val Glu Gln Met
180             185             190

tgc gtc acc cag tac cag aag gag tcc cag gcc tat tac gac ggg aga   624
Cys Val Thr Gln Tyr Gln Lys Glu Ser Gln Ala Tyr Tyr Asp Gly Arg
195             200             205

aga tcc agc tgataacc   641
Arg Ser Ser
210

```

```

<210> SEQ ID NO 19
<211> LENGTH: 211
<212> TYPE: PRT
<213> ORGANISM: MOUSE

```

```

<400> SEQUENCE: 19

```

```

Met Ser Lys Lys Arg Pro Lys Pro Gly Gly Trp Asn Thr Gly Gly Ser
 1             5             10             15
Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly Asn Arg Tyr Pro Pro Gln
20             25             30
Gly Gly Thr Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His
35             40             45
Gly Gly Ser Trp Gly Gln Pro His Gly Gly Ser Trp Gly Gln Pro His
50             55             60
Gly Gly Gly Trp Gly Gln Gly Gly Gly Thr His Asn Gln Trp Asn Lys
65             70             75             80
Pro Ser Lys Pro Lys Thr Asn Leu Lys His Val Ala Gly Ala Ala Ala
85             90             95
Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr Met Leu Gly Ser Ala
100            105            110
Val Ser Arg Pro Met Ile His Phe Gly Asn Asp Trp Glu Asp Arg Tyr
115            120            125
Tyr Arg Glu Asn Met Tyr Arg Tyr Pro Asn Gln Val Tyr Tyr Arg Pro
130            135            140
Val Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val His Asp Cys Val Asn
145            150            155            160
Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr Thr Lys Gly Glu Asn
165            170            175
Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg Val Val Glu Gln Met
180            185            190
Cys Val Thr Gln Tyr Gln Lys Glu Ser Gln Ala Tyr Tyr Asp Gly Arg
195            200            205

Arg Ser Ser
210

```

```

<210> SEQ ID NO 20
<211> LENGTH: 644
<212> TYPE: DNA
<213> ORGANISM: Mesocricetus auratus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(636)

```

```

<400> SEQUENCE: 20

```

```

atg tct aag aag cgg cca aag cct gga ggg tgg aac act ggc gga agc   48

```


-continued

Met	Ser	Lys	Lys	Arg	Pro	Lys	Pro	Gly	Gly	Trp	Asn	Thr	Gly	Gly	Ser		
1				5					10						15		
cga	tac	cct	ggg	cag	ggc	agc	cct	gga	ggc	aac	cgt	tac	cca	cct	cag		96
Arg	Tyr	Pro	Gly	Gln	Gly	Ser	Pro	Gly	Gly	Asn	Arg	Tyr	Pro	Pro	Gln		
20				25						30							
ggg	ggc	ggc	aca	tgg	ggg	caa	ccc	cat	ggt	ggt	ggc	tgg	gga	cag	ccc		144
Gly	Gly	Gly	Thr	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Pro		
35				40					45								
cat	ggt	ggt	ggc	tgg	gga	cag	ccc	cat	ggt	ggt	ggc	tgg	ggt	cag	ccc		192
His	Gly	Gly	Gly	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Pro		
50				55					60								
cat	ggt	ggt	ggc	tgg	ggt	caa	gga	ggt	ggc	acc	cac	aat	cag	tgg	aac		240
His	Gly	Gly	Gly	Trp	Gly	Gln	Gly	Gly	Gly	Thr	His	Asn	Gln	Trp	Asn		
65				70					75					80			
aag	ccc	agt	aag	cca	aaa	acc	aac	atg	aag	cac	atg	gcc	ggc	gct	gct		288
Lys	Pro	Ser	Lys	Pro	Lys	Thr	Asn	Met	Lys	His	Met	Ala	Gly	Ala	Ala		
85				90					95								
gcg	gea	ggg	gcc	gtg	gtg	ggg	ggc	ctt	ggt	ggc	tac	atg	ctg	ggg	agt		336
Ala	Ala	Gly	Ala	Val	Val	Gly	Gly	Leu	Gly	Gly	Tyr	Met	Leu	Gly	Ser		
100				105					110								
gcc	atg	agc	agg	ccc	atg	atg	cat	ttt	ggc	aat	gac	tgg	gag	gac	cgc		384
Ala	Met	Ser	Arg	Pro	Met	Met	His	Phe	Gly	Asn	Asp	Trp	Glu	Asp	Arg		
115				120					125								
tac	tac	cgt	gaa	aac	atg	aac	cgc	tac	cct	aac	caa	gtg	tat	tac	cgg		432
Tyr	Tyr	Arg	Glu	Asn	Met	Asn	Arg	Tyr	Pro	Asn	Gln	Val	Tyr	Tyr	Arg		
130				135					140								
cca	gtg	gac	cag	tac	aac	aac	cag	aac	aac	ttt	gtg	cac	gat	tgt	gtc		480
Pro	Val	Asp	Gln	Tyr	Asn	Asn	Gln	Asn	Asn	Phe	Val	His	Asp	Cys	Val		
145				150					155					160			
aac	atc	acc	atc	aag	cag	cac	aca	gtc	acc	acc	acc	acc	aag	ggg	gag		528
Asn	Ile	Thr	Ile	Lys	Gln	His	Thr	Val	Thr	Thr	Thr	Thr	Lys	Gly	Glu		
165				170					175								
aac	ttc	acg	gag	acc	gac	atc	aag	ata	atg	gag	cgc	gtg	gtg	gag	cag		576
Asn	Phe	Thr	Glu	Thr	Asp	Ile	Lys	Ile	Met	Glu	Arg	Val	Val	Glu	Gln		
180				185					190								
atg	tgt	acc	acc	cag	tat	cag	aag	gag	tcc	cag	gcc	tac	tac	gat	gga		624
Met	Cys	Thr	Thr	Gln	Tyr	Gln	Lys	Glu	Ser	Gln	Ala	Tyr	Tyr	Asp	Gly		
195				200					205								
aga	agg	tcc	agc	tgataacc													644
Arg	Arg	Ser	Ser														
210																	

<210> SEQ ID NO 21
 <211> LENGTH: 212
 <212> TYPE: PRT
 <213> ORGANISM: Mesocricetus auratus

<400> SEQUENCE: 21

Met	Ser	Lys	Lys	Arg	Pro	Lys	Pro	Gly	Gly	Trp	Asn	Thr	Gly	Gly	Ser		
1				5					10						15		
Arg	Tyr	Pro	Gly	Gln	Gly	Ser	Pro	Gly	Gly	Asn	Arg	Tyr	Pro	Pro	Gln		
20				25					30								
Gly	Gly	Gly	Thr	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Pro		
35				40					45								
His	Gly	Gly	Gly	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Pro		
50				55					60								
His	Gly	Gly	Gly	Trp	Gly	Gln	Gly	Gly	Gly	Thr	His	Asn	Gln	Trp	Asn		

-continued

65					70						75					80
Lys	Pro	Ser	Lys	Pro	Lys	Thr	Asn	Met	Lys	His	Met	Ala	Gly	Ala	Ala	
85					90					95						
Ala	Ala	Gly	Ala	Val	Val	Gly	Gly	Leu	Gly	Gly	Tyr	Met	Leu	Gly	Ser	
100					105					110						
Ala	Met	Ser	Arg	Pro	Met	Met	His	Phe	Gly	Asn	Asp	Trp	Glu	Asp	Arg	
115					120					125						
Tyr	Tyr	Arg	Glu	Asn	Met	Asn	Arg	Tyr	Pro	Asn	Gln	Val	Tyr	Tyr	Arg	
130					135					140						
Pro	Val	Asp	Gln	Tyr	Asn	Asn	Gln	Asn	Asn	Phe	Val	His	Asp	Cys	Val	
145					150					155					160	
Asn	Ile	Thr	Ile	Lys	Gln	His	Thr	Val	Thr	Thr	Thr	Lys	Gly	Glu		
165					170					175						
Asn	Phe	Thr	Glu	Thr	Asp	Ile	Lys	Ile	Met	Glu	Arg	Val	Val	Glu	Gln	
180					185					190						
Met	Cys	Thr	Thr	Gln	Tyr	Gln	Lys	Glu	Ser	Gln	Ala	Tyr	Tyr	Asp	Gly	
195					200					205						
Arg	Arg	Ser	Ser													
210																

<210> SEQ ID NO 22
 <211> LENGTH: 780
 <212> TYPE: PRT
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 22

Met	Lys	Lys	Lys	Asp	Asn	Ser	Asp	Asp	Lys	Asp	Asn	Val	Ala	Ser	Gly	
1				5					10					15		
Gly	Tyr	Lys	Asn	Ala	Ala	Asp	Ala	Gly	Ser	Asn	Asn	Ala	Ser	Lys	Lys	
20				25						30						
Ser	Ser	Tyr	Arg	Asn	Trp	Lys	Gly	Gly	Asn	Tyr	Gly	Gly	Tyr	Ser	Tyr	
35				40						45						
Asn	Ser	Asn	Tyr	Asn	Asn	Tyr	Asn	Asn	Tyr	Asn	Asn	Tyr	Asn	Asn	Tyr	
50				55						60						
Asn	Asn	Tyr	Asn	Asn	Tyr	Asn	Lys	Tyr	Asn	Gly	Gly	Tyr	Lys	Ser	Thr	
65				70						75					80	
Tyr	Lys	Ser	Ala	Val	Thr	Asn	Ser	Gly	Thr	Thr	Ser	Ala	Ser	Thr	Thr	
85				90						95						
Ser	Thr	Ser	Asn	Lys	Ser	Asn	Thr	Ser	Ser	Lys	Cys	Ser	Thr	Asp	Cys	
100				105						110						
Lys	Asn	Lys	Gly	Lys	Gly	Asn	Ser	Thr	Gly	Lys	Trp	Lys	Val	Asp	Val	
115				120						125						
Ser	Lys	Lys	Lys	Asn	Ser	Val	Arg	Ser	Ala	Met	Ser	Asn	Ala	Ser	Gly	
130				135						140						
Lys	Ala	Tyr	Asn	Val	Ala	Asp	Cys	Ser	Asp	Lys	Asn	Thr	Val	Lys	Arg	
145				150						155					160	
Ala	Ala	His	Ala	Asp	Ser	Asn	Cys	Met	Ala	Thr	Cys	Val	Thr	Asp	Tyr	
165				170						175						
Ser	Ser	Gly	Ala	Lys	Trp	Ala	Lys	Met	Ala	Ala	Ser	Val	Val	Asp	Arg	
180				185						190						
Arg	Asp	Ser	Ala	Asn	Asp	Thr	Lys	Asp	Ala	Val	Val	Thr	Asp	Val	Ala	
195				200						205						

-continued

Thr Asp Lys Ala Lys Gly Tyr Lys Thr Asp Tyr Val Ser Asp Asn Asp
 210 215 220
 Ser Arg Tyr Lys Val Asp Thr Asp Ser Lys Val Ser Val Lys Ser Ser
 225 230 235 240
 Ser Val Thr Val Ala Val Thr Ser Ser Val Asn Arg Ser Asn Ser Ser
 245 250 255
 Ser Ser Arg Thr Val Val Val Asn Thr Arg Val Asn Asn Arg Asn Ser
 260 265 270
 Gly Lys Val Val Asp Thr Ala Ser Val Arg Ala Lys Ala Asn Val Lys
 275 280 285
 Asp Asp Ala Asp Lys Asn Lys Ser Gly Arg Thr Gly Arg Asp Asp His
 290 295 300
 Lys Asp Lys Ala Asp Asp Ser Cys Val Lys Tyr Met Asn Asp Thr Val
 305 310 315 320
 Lys Tyr Met Ser Lys Thr Val Asp Ser Asn Val Asn Asp Trp Lys Arg
 325 330 335
 Asp Thr Ala Val Gly Gly Ser Asp Ser Arg Val Lys Asp His Asn Arg
 340 345 350
 Ala Tyr Lys Arg Ala Asp Asp Gly Val Asn Thr Asp Ser Ala Tyr Gly
 355 360 365
 Ser Arg Met Asn Lys Thr Asn Arg Lys Gly His Arg Tyr Gly Cys Gly
 370 375 380
 Arg Asn Gly Ala Gly Lys Ser Thr Met Arg Ala Ala Asn Gly Asp Gly
 385 390 395 400
 Asp Lys Asp Thr Arg Thr Cys Val His Lys Gly Gly Asp Asp Val Ser
 405 410 415
 Ala Asp Ser Thr Ser Arg Ala Ala Ala Ser Val Gly Asp Arg Arg Ala
 420 425 430
 Thr Val Gly Ser Ser Gly Gly Trp Lys Met Lys Ala Arg Ala Met Lys
 435 440 445
 Ala Asp Asp Thr Asn His Asp Val Ser Asn Val Lys Trp Tyr His Thr
 450 455 460
 Asp Thr Ser Val Ser His Asp Ser Gly Asp Thr Val Cys Thr Asp His
 465 470 475 480
 Tyr Asn Lys Lys Ala Tyr Tyr Lys Gly Asn Ala Ala Val Lys Ala Lys
 485 490 495
 Ser Tyr Tyr Thr Thr Asp Ser Asn Ala Met Arg Gly Thr Gly Val Lys
 500 505 510
 Ser Asn Thr Arg Ala Val Ala Lys Met Thr Asp Val Thr Ser Tyr Gly
 515 520 525
 Ala Lys Ser Ser His Val Ser Cys Ser Ser Ser Ser Arg Val Ala Cys
 530 535 540
 Gly Asn Gly Ala Gly Lys Ser Thr Lys Thr Gly Val Asn Gly Lys Val
 545 550 555 560
 Lys His Asn Arg Gly Tyr Ala His Ala His Val Asn His Lys Lys Thr
 565 570 575
 Ala Asn Tyr Trp Arg Tyr Gly Asp Asp Arg Val Lys Ser Arg Lys Ser
 580 585 590
 Asp Lys Met Met Thr Lys Asp Asp Asp Gly Arg Gly Lys Arg Ala Ala
 595 600 605
 Val Gly Arg Lys Lys Lys Ser Tyr Val Lys Trp Lys Tyr Trp Lys Lys

-continued

```

610                615                620
Tyr Asn Ser Trp Val Lys Asp Val Val His Gly Lys Val Lys Asp Asp
625                630                635                640
His Ala Ser Arg Gly Gly Tyr Arg Ser Val Thr Lys His Asp Val Gly
645                650                655
Asp Ser Ala Asn His Thr Gly Ser Ser Gly Gly Val Lys Val Val Ala
660                665                670
Gly Ala Met Trp Asn Asn His Val Asp Thr Asn Tyr Asp Arg Asp Ser
675                680                685
Gly Ala Ala Val Ala Arg Asp Trp Ser Gly Gly Val Val Met Ser His
690                695                700
Asn Asn Val Gly Ala Cys Trp Val Asn Gly Lys Met Val Lys Gly Ser
705                710                715                720
Ala Val Asp Ser Lys Asp Gly Gly Asn Ala Asp Ala Val Gly Lys Ala
725                730                735
Ser Asn Ala Lys Ser Val Asp Asp Asp Asp Ser Ala Asn Lys Val Lys
740                745                750
Arg Lys Lys Arg Thr Arg Asn Lys Lys Ala Arg Arg Arg Arg Tyr Trp
755                760                765
Ser Ser Lys Gly Thr Lys Val Asp Thr Asp Asp Asp
770                775                780

```

```

<210> SEQ ID NO 23
<211> LENGTH: 1075
<212> TYPE: PRT
<213> ORGANISM: Saccharomyces cerevisiae

```

```

<400> SEQUENCE: 23

```

```

Met Asp Asn Lys Arg Leu Tyr Asn Gly Asn Leu Ser Asn Ile Pro Glu
1      5      10      15
Val Ile Asp Pro Gly Ile Thr Ile Pro Ile Tyr Glu Glu Asp Ile Arg
20     25     30
Asn Asp Thr Arg Met Asn Thr Asn Ala Arg Ser Val Arg Val Ser Asp
35     40     45
Lys Arg Gly Arg Ser Ser Ser Thr Ser Pro Gln Lys Ile Gly Ser Tyr
50     55     60
Arg Thr Arg Ala Gly Arg Phe Ser Asp Thr Leu Thr Asn Leu Leu Pro
65     70     75     80
Ser Ile Ser Ala Lys Leu His His Ser Lys Lys Ser Thr Pro Val Val
85     90     95
Val Val Pro Pro Thr Ser Ser Thr Pro Asp Ser Leu Asn Ser Thr Thr
100    105    110
Tyr Ala Pro Arg Val Ser Ser Asp Ser Phe Thr Val Ala Thr Pro Leu
115    120    125
Ser Leu Gln Ser Thr Thr Thr Arg Thr Arg Thr Arg Asn Asn Thr Val
130    135    140
Ser Ser Gln Ile Thr Ala Ser Ser Ser Leu Thr Thr Asp Val Gly Asn
145    150    155    160
Ala Thr Ser Ala Asn Ile Trp Ser Ala Asn Ala Glu Ser Asn Thr Ser
165    170    175
Ser Ser Pro Leu Phe Asp Tyr Pro Leu Ala Thr Ser Tyr Phe Glu Pro
180    185    190

```

-continued

Leu	Thr	Arg	Phe	Lys	Ser	Thr	Asp	Asn	Tyr	Thr	Leu	Pro	Gln	Thr	Ala	195	200	205	
Gln	Leu	Asn	Ser	Phe	Leu	Glu	Lys	Asn	Gly	Asn	Pro	Asn	Ile	Trp	Ser	210	215	220	
Ser	Ala	Gly	Asn	Ser	Asn	Thr	Asp	His	Leu	Asn	Thr	Pro	Ile	Val	Asn	225	230	235	240
Arg	Gln	Arg	Ser	Gln	Ser	Gln	Ser	Thr	Thr	Asn	Arg	Val	Tyr	Thr	Asp	245	250	255	
Ala	Pro	Tyr	Tyr	Gln	Gln	Pro	Ala	Gln	Asn	Tyr	Gln	Val	Gln	Val	Pro	260	265	270	
Pro	Arg	Val	Pro	Lys	Ser	Thr	Ser	Ile	Ser	Pro	Val	Ile	Leu	Asp	Asp	275	280	285	
Val	Asp	Pro	Ala	Ser	Ile	Asn	Trp	Ile	Thr	Ala	Asn	Gln	Lys	Val	Pro	290	295	300	
Leu	Val	Asn	Gln	Ile	Ser	Ala	Leu	Leu	Pro	Thr	Asn	Thr	Ile	Ser	Ile	305	310	315	320
Ser	Asn	Val	Phe	Pro	Leu	Gln	Pro	Thr	Gln	Gln	His	Gln	Gln	Asn	Ala	325	330	335	
Val	Asn	Leu	Thr	Ser	Thr	Ser	Leu	Ala	Thr	Leu	Cys	Ser	Gln	Tyr	Gly	340	345	350	
Lys	Val	Leu	Ser	Ala	Arg	Thr	Leu	Arg	Gly	Leu	Asn	Met	Ala	Leu	Val	355	360	365	
Glu	Phe	Ser	Thr	Val	Glu	Ser	Ala	Ile	Cys	Ala	Leu	Glu	Ala	Leu	Gln	370	375	380	
Gly	Lys	Glu	Leu	Ser	Lys	Val	Gly	Ala	Pro	Ser	Thr	Val	Ser	Phe	Ala	385	390	395	400
Arg	Val	Leu	Pro	Met	Tyr	Glu	Gln	Pro	Leu	Asn	Val	Asn	Gly	Phe	Asn	405	410	415	
Asn	Thr	Pro	Lys	Gln	Pro	Leu	Leu	Gln	Glu	Gln	Leu	Asn	His	Gly	Val	420	425	430	
Leu	Asn	Tyr	Gln	Leu	Gln	Gln	Ser	Leu	Gln	Gln	Pro	Glu	Leu	Gln	Gln	435	440	445	
Gln	Pro	Thr	Ser	Phe	Asn	Gln	Pro	Asn	Leu	Thr	Tyr	Cys	Asn	Pro	Thr	450	455	460	
Gln	Asn	Leu	Ser	His	Leu	Gln	Leu	Ser	Ser	Asn	Glu	Asn	Glu	Pro	Tyr	465	470	475	480
Pro	Phe	Pro	Leu	Pro	Pro	Pro	Ser	Leu	Ser	Asp	Ser	Lys	Lys	Asp	Ile	485	490	495	
Leu	His	Thr	Ile	Ser	Ser	Phe	Lys	Leu	Glu	Tyr	Asp	His	Leu	Glu	Leu	500	505	510	
Asn	His	Leu	Leu	Gln	Asn	Ala	Leu	Lys	Asn	Lys	Gly	Val	Ser	Asp	Thr	515	520	525	
Asn	Tyr	Phe	Gly	Pro	Leu	Pro	Glu	His	Asn	Ser	Lys	Val	Pro	Lys	Arg	530	535	540	
Lys	Asp	Thr	Phe	Asp	Ala	Pro	Lys	Leu	Arg	Glu	Leu	Arg	Lys	Gln	Phe	545	550	555	560
Asp	Ser	Asn	Ser	Leu	Ser	Thr	Ile	Glu	Met	Glu	Gln	Leu	Ala	Ile	Val	565	570	575	
Met	Leu	Asp	Gln	Leu	Pro	Glu	Leu	Ser	Ser	Asp	Tyr	Leu	Gly	Asn	Thr	580	585	590	
Val	Ile	Gln	Lys	Leu	Phe	Glu	Asn	Ser	Ser	Asn	Ile	Ile	Arg	Asp	Ile				

-continued

```

Asn Pro Leu Ser Arg Ser Val Ser His Asn Asn Asn Asn Thr Asn
1010                1015                1020

Asn Tyr Asn Asn Asn Asp Asn Asp Asn Asn Asn Asn Asn Asn
1025                1030                1035                1040

Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Ser Asn Asn
1045                1050                1055

Ser Asn Asn Asn Asn Asn Asn Asp Thr Ser Leu Tyr Arg Tyr Arg Ser
1060                1065                1070

Tyr Gly Tyr
1075

```

```

<210> SEQ ID NO 24
<211> LENGTH: 76
<212> TYPE: PRT
<213> ORGANISM: Saccharomyces cerevisiae

```

```

<400> SEQUENCE: 24

```

```

Met Ser Ala Asn Asp Tyr Tyr Gly Gly Thr Ala Gly Lys Ser Tyr Ser
 1                5                10                15

Arg Ser Asn Ser Ser Ala His Asn Lys Thr Arg Gly Tyr Tyr Tyr His
20                25                30

Gly Tyr Tyr Asn Gly Tyr Asn Gly Tyr Asn Gly Tyr Asn Gly Tyr Asn
35                40                45

Gly Tyr Asn Gly Tyr Asn Gly His Val Tyr Val Arg Gly Asn Gly Cys
50                55                60

Ala Ala Cys Ala Ala Cys Cys Cys Thr Met Asp Met
65                70                75

```

```

<210> SEQ ID NO 25
<211> LENGTH: 380
<212> TYPE: PRT
<213> ORGANISM: Saccharomyces cerevisiae

```

```

<400> SEQUENCE: 25

```

```

Met Ser Ser Asp Asp Asn Asp Tyr Gly Asp Asp Lys Thr Thr Thr Val
 1                5                10                15

Lys Lys Asn Lys Ala Gly Ser Gly Thr Ser Asp Ala Ala Ala Ser Ser
20                25                30

Ser Asn Lys Asn Asn Asn Ser Asn Asn Ser Ser Ser Asn Asn Ser Asn
35                40                45

Asp Thr Ser Ser Ser Lys Asp Gly Thr Ala Asn Asp Lys Gly Ser Asn
50                55                60

Asp Thr Lys Asn Lys Lys Ser Ala Thr Ser Ala Asn Ala Asn Ala Asn
65                70                75                80

Ala Ser Ser Ala Gly Ser Gly Trp Thr Met Ser Ser Ser Ser Val Thr
85                90                95

Thr Lys Arg Ser Lys Ala Asp Ser Lys Ser Cys Lys Met Gly Gly Asn
100               105                110

Trp Asp Thr Thr Asp Asn Arg Tyr Gly Lys Tyr Gly Thr Val Thr Asp
115               120                125

Lys Met Lys Asp Ala Thr Gly Arg Ser Arg Gly Gly Ser Lys Ser Ser
130               135                140

Val Asp Val Val Lys Thr His Asp Gly Lys Val Asp Lys Arg Ala Arg
145               150                155                160

```

-continued

Asp Asp Lys Thr Gly Lys Val Gly Gly Gly Asp Val Arg Lys Ser Trp
 165 170 175
 Gly Thr Asp Ala Met Asp Lys Asp Thr Gly Ser Arg Gly Gly Val Thr
 180 185 190
 Tyr Asp Ser Ala Asp Ala Val Asp Arg Val Cys Asn Lys Asp Lys Asp
 195 200 205
 Arg Lys Lys Arg Ala Arg His Met Lys Ser Ser Asn Asn Gly Gly Asn
 210 215 220
 Asn Gly Gly Asn Asn Met Asn Arg Arg Gly Gly Asn Gly Asn Gly Asp
 225 230 235 240
 Asn Met Tyr Asn Met Met Gly Gly Tyr Asn Met Met Asn Ala Met Thr
 245 250 255
 Asp Tyr Tyr Lys Met Tyr Tyr Met Lys Thr Gly Met Asp Tyr Thr Met
 260 265 270
 Tyr Met Met Ala Met Met Met Gly Ala Met Asn Ala Met Thr Asn Asp
 275 280 285
 Ser Asn Ala Thr Gly Ser Ala Ser Asp Ser Asp Asn Asn Lys Ser Asn
 290 295 300
 Asp Val Thr Gly Asn Thr Ser Asn Thr Asp Ser Gly Ser Asn Asn Gly
 305 310 315 320
 Lys Gly Ser Tyr Asn Asp Asp His Asn Ser Gly Tyr Gly Tyr Asn Arg
 325 330 335
 Asp Arg Gly Asp Arg Asp Arg Asn Asp Arg Asp Arg Asp Tyr Asn His
 340 345 350
 Arg Ser Gly Gly Asn His Arg Arg Asn Gly Arg Gly Gly Arg Gly Gly
 355 360 365
 Tyr Asn Arg Arg Asn Asn Gly Tyr His Tyr Asn Arg
 370 375 380

<210> SEQ ID NO 26

<211> LENGTH: 256

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 26

Met Ser Ala Thr His Val Ser Val Val Asp Ala Val His Ala Asp Ala
 1 5 10 15
 Val Ser Ala Ser Ala Ala Asn Asp Val Ser Asn Ala Tyr Gly Ser His
 20 25 30
 Ser Val Asp Tyr Ala His His His Tyr Tyr Gly His Met His Gly Arg
 35 40 45
 Met His His Arg Gly Ser Asn Thr Arg Val Arg Asp Val Ser Asn Gly
 50 55 60
 Gly Met Lys Val Lys Asn Gly Ala Val Ala Ser Ala Ala Lys Ala Val
 65 70 75 80
 His Gly Lys Ser Ala Asn Val Val Tyr Ser Lys Ala Lys Arg Tyr Arg
 85 90 95
 Thr Met Lys Asn Gly Cys Ser Trp Asp Lys Asp Ala Arg Asn Ser Thr
 100 105 110
 Thr Ser Ser Val Asn Thr Arg Asp Asp Gly Thr Gly Ala Ser Val Ala
 115 120 125
 Arg Asn Asn Arg Gly Ser Val Thr Val Arg Asp Asp Asn Arg Arg Ser

-continued

```

130                135                140
Asn Arg Gly Gly Arg Gly Arg Gly Gly Arg Gly Gly Arg Gly Gly Arg
145                150                155                160

Gly Gly Ser Arg Gly Gly Gly Gly Arg Gly Gly Gly Gly Arg Gly Gly
165                170                175

Tyr Gly Gly Tyr Ser Arg Gly Gly Tyr Gly Gly Tyr Ser Arg Gly Gly
180                185                190

Tyr Gly Gly Ser Arg Gly Gly Tyr Asp Ser Arg Gly Gly Tyr Asp Ser
195                200                205

Arg Gly Gly Tyr Ser Arg Gly Gly Tyr Gly Gly Arg Asn Asp Tyr Gly
210                215                220

Arg Gly Ser Tyr Gly Gly Ser Arg Gly Gly Tyr Asp Gly Arg Gly Asp
225                230                235                240

Tyr Gly Arg Asp Ala Tyr Arg Thr Arg Asp Ala Arg Arg Ser Thr Arg
245                250                255

```

```

<210> SEQ ID NO 27
<211> LENGTH: 286
<212> TYPE: PRT
<213> ORGANISM: Saccharomyces cerevisiae

```

```

<400> SEQUENCE: 27

```

```

Met Ser Asp Ile Glu Glu Gly Thr Pro Thr Asn Asn Gly Gln Gln Lys
1      5      10      15

Glu Arg Arg Lys Ile Glu Ile Lys Phe Ile Glu Asn Lys Thr Arg Arg
20     25     30

His Val Thr Phe Ser Lys Arg Lys His Gly Ile Met Lys Lys Ala Phe
35     40     45

Glu Leu Ser Val Leu Thr Gly Thr Gln Val Leu Leu Leu Val Val Ser
50     55     60

Glu Thr Gly Leu Val Tyr Thr Phe Ser Thr Pro Lys Phe Glu Pro Ile
65     70     75     80

Val Thr Gln Gln Glu Gly Arg Asn Leu Ile Gln Ala Cys Leu Asn Ala
85     90     95

Pro Asp Asp Glu Glu Glu Asp Glu Glu Glu Asp Gly Asp Asp Asp Asp
100    105    110

Asp Asp Asp Asp Asp Gly Asn Asp Met Gln Arg Gln Gln Pro Gln Gln
115    120    125

Gln Gln Pro Gln Gln Gln Gln Gln Val Leu Asn Ala His Ala Asn Ser
130    135    140

Leu Gly His Leu Asn Gln Asp Gln Val Pro Ala Gly Ala Leu Lys Gln
145    150    155    160

Glu Val Lys Ser Gln Leu Leu Gly Gly Ala Asn Pro Asn Gln Asn Ser
165    170    175

Met Ile Gln Gln Gln Gln His His Thr Gln Asn Ser Gln Pro Gln Gln
180    185    190

Gln Gln Gln Gln Gln Pro Gln Gln Gln Met Ser Gln Gln Gln Met Ser
195    200    205

Gln His Pro Arg Pro Gln Gln Gly Ile Pro His Pro Gln Gln Ser Gln
210    215    220

Pro Gln Gln Gln Gln Gln Gln Gln Gln Leu Gln Gln Gln Gln Gln
225    230    235    240

```

-continued

Gln Gln Gln Gln Gln Pro Leu Thr Gly Ile His Gln Pro His Gln Gln
 245 250 255

Ala Phe Ala Asn Ala Ala Ser Pro Tyr Leu Asn Ala Glu Gln Asn Ala
 260 265 270

Ala Tyr Gln Gln Tyr Phe Gln Glu Pro Gln Gln Gly Gln Tyr
 275 280 285

<210> SEQ ID NO 28
 <211> LENGTH: 414
 <212> TYPE: PRT
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 28

Met Ala Lys Thr Thr Lys Val Lys Gly Asn Lys Lys Glu Val Lys Ala
 1 5 10 15

Ser Lys Gln Ala Lys Glu Glu Lys Ala Lys Ala Val Ser Ser Ser Ser
 20 25 30

Ser Glu Ser Ser Ser Ser Ser Ser Ser Ser Ser Glu Ser Glu Ser Glu
 35 40 45

Ser Glu Ser Glu Ser Glu Ser Ser Ser Ser Ser Ser Ser Ser Asp Ser
 50 55 60

Glu Ser Ser Ser Ser Ser Ser Ser Asp Ser Glu Ser Glu Ala Glu Thr
 65 70 75 80

Lys Lys Glu Glu Ser Lys Asp Ser Ser Ser Ser Ser Ser Ser Asp Ser Ser
 85 90 95

Ser Asp Glu Glu Glu Glu Glu Lys Glu Glu Thr Lys Lys Glu Glu
 100 105 110

Ser Lys Glu Ser Ser Ser Ser Asp Ser Ser Ser Ser Ser Ser Ser Asp
 115 120 125

Ser Glu Ser Glu Lys Glu Glu Ser Asn Asp Lys Lys Arg Lys Ser Glu
 130 135 140

Asp Ala Glu Glu Glu Glu Asp Glu Glu Ser Ser Asn Lys Lys Gln Lys
 145 150 155 160

Asn Glu Glu Thr Glu Glu Pro Ala Thr Ile Phe Val Gly Arg Leu Ser
 165 170 175

Trp Ser Ile Asp Asp Glu Trp Leu Lys Lys Glu Phe Glu His Ile Gly
 180 185 190

Gly Val Ile Gly Ala Arg Val Ile Tyr Glu Arg Gly Thr Asp Arg Ser
 195 200 205

Arg Gly Tyr Gly Tyr Val Asp Phe Glu Asn Lys Ser Tyr Ala Glu Lys
 210 215 220

Ala Ile Gln Glu Met Gln Gly Lys Glu Ile Asp Gly Arg Pro Ile Asn
 225 230 235 240

Cys Asp Met Ser Thr Ser Lys Pro Ala Gly Asn Asn Asp Arg Ala Lys
 245 250 255

Lys Phe Gly Asp Thr Pro Ser Glu Pro Ser Asp Thr Leu Phe Leu Gly
 260 265 270

Asn Leu Ser Phe Asn Ala Asp Arg Asp Ala Ile Phe Glu Leu Phe Ala
 275 280 285

Lys His Gly Glu Val Val Ser Val Arg Ile Pro Thr His Pro Glu Thr
 290 295 300

Glu Gln Pro Lys Gly Phe Gly Tyr Val Gln Phe Ser Asn Met Glu Asp
 305 310 315 320

-continued

Ala Lys Lys Ala Leu Asp Ala Leu Gln Gly Glu Tyr Ile Asp Asn Arg
 325 330 335

Pro Val Arg Leu Asp Phe Ser Ser Pro Arg Pro Asn Asn Asp Gly Gly
 340 345 350

Arg Gly Gly Ser Arg Gly Phe Gly Gly Arg Gly Gly Gly Arg Gly Gly
 355 360 365

Asn Arg Gly Phe Gly Gly Arg Gly Gly Ala Arg Gly Gly Arg Gly Gly
 370 375 380

Phe Arg Pro Ser Gly Ser Gly Ala Asn Thr Ala Pro Leu Gly Arg Ser
 385 390 395 400

Arg Asn Thr Ala Ser Phe Ala Gly Ser Lys Lys Thr Phe Asp
 405 410

<210> SEQ ID NO 29

<211> LENGTH: 405

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 29

Met Asp Thr Asp Lys Leu Ile Ser Glu Ala Glu Ser His Phe Ser Gln
 1 5 10 15

Gly Asn His Ala Glu Ala Val Ala Lys Leu Thr Ser Ala Ala Gln Ser
 20 25 30

Asn Pro Asn Asp Glu Gln Met Ser Thr Ile Glu Ser Leu Ile Gln Lys
 35 40 45

Ile Ala Gly Tyr Val Met Asp Asn Arg Ser Gly Gly Ser Asp Ala Ser
 50 55 60

Gln Asp Arg Ala Ala Gly Gly Gly Ser Ser Phe Met Asn Thr Leu Met
 65 70 75 80

Ala Asp Ser Lys Gly Ser Ser Gln Thr Gln Leu Gly Lys Leu Ala Leu
 85 90 95

Leu Ala Thr Val Met Thr His Ser Ser Asn Lys Gly Ser Ser Asn Arg
 100 105 110

Gly Phe Asp Val Gly Thr Val Met Ser Met Leu Ser Gly Ser Gly Gly
 115 120 125

Gly Ser Gln Ser Met Gly Ala Ser Gly Leu Ala Ala Leu Ala Ser Gln
 130 135 140

Phe Phe Lys Ser Gly Asn Asn Ser Gln Gly Gln Gly Gln Gly Gln Gly
 145 150 155 160

Gln Gly Gln Gly Gln Gly Gln Gly Gln Gly Gln Gly Ser Phe Thr Ala
 165 170 175

Leu Ala Ser Leu Ala Ser Ser Phe Met Asn Ser Asn Asn Asn Asn Gln
 180 185 190

Gln Gly Gln Asn Gln Ser Ser Gly Gly Ser Ser Phe Gly Ala Leu Ala
 195 200 205

Ser Met Ala Ser Ser Phe Met His Ser Asn Asn Asn Gln Asn Ser Asn
 210 215 220

Asn Ser Gln Gln Gly Tyr Asn Gln Ser Tyr Gln Asn Gly Asn Gln Asn
 225 230 235 240

Ser Gln Gly Tyr Asn Asn Gln Gln Tyr Gln Gly Gly Asn Gly Gly Tyr
 245 250 255

Gln Gln Gln Gln Gly Gln Ser Gly Gly Ala Phe Ser Ser Leu Ala Ser

-continued

260 265 270
 Met Ala Gln Ser Tyr Leu Gly Gly Gly Gln Thr Gln Ser Asn Gln Gln
 275 280 285
 Gln Tyr Asn Gln Gln Gly Gln Asn Asn Gln Gln Gln Tyr Gln Gln Gln
 290 295 300
 Gly Gln Asn Tyr Gln His Gln Gln Gln Gly Gln Gln Gln Gln Gly
 305 310 315 320
 His Ser Ser Ser Phe Ser Ala Leu Ala Ser Met Ala Ser Ser Tyr Leu
 325 330 335
 Gly Asn Asn Ser Asn Ser Asn Ser Ser Tyr Gly Gly Gln Gln Gln Ala
 340 345 350
 Asn Glu Tyr Gly Arg Pro Gln His Asn Gly Gln Gln Gln Ser Asn Glu
 355 360 365
 Tyr Gly Arg Pro Gln Tyr Gly Gly Asn Gln Asn Ser Asn Gly Gln His
 370 375 380
 Glu Ser Phe Asn Phe Ser Gly Asn Phe Ser Gln Gln Asn Asn Asn Gly
 385 390 395 400
 Asn Gln Asn Arg Tyr
 405

<210> SEQ ID NO 30
 <211> LENGTH: 964
 <212> TYPE: PRT
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 30

Met Pro Glu Gln Ala Gln Gln Gly Glu Gln Ser Val Lys Arg Arg Arg
 1 5 10 15
 Val Thr Arg Ala Cys Asp Glu Cys Arg Lys Lys Lys Val Lys Cys Asp
 20 25 30
 Gly Gln Gln Pro Cys Ile His Cys Thr Val Tyr Ser Tyr Glu Cys Thr
 35 40 45
 Tyr Lys Lys Pro Thr Lys Arg Thr Gln Asn Ser Gly Asn Ser Gly Val
 50 55 60
 Leu Thr Leu Gly Asn Val Thr Thr Gly Pro Ser Ser Ser Thr Val Val
 65 70 75 80
 Ala Ala Ala Ala Ser Asn Pro Asn Lys Leu Leu Ser Asn Ile Lys Thr
 85 90 95
 Glu Arg Ala Ile Leu Pro Gly Ala Ser Thr Ile Pro Ala Ser Asn Asn
 100 105 110
 Pro Ser Lys Pro Arg Lys Tyr Lys Thr Lys Ser Thr Arg Leu Gln Ser
 115 120 125
 Lys Ile Asp Arg Tyr Lys Gln Ile Phe Asp Glu Val Phe Pro Gln Leu
 130 135 140
 Pro Asp Ile Asp Asn Leu Asp Ile Pro Val Phe Leu Gln Ile Phe His
 145 150 155 160
 Asn Phe Lys Arg Asp Ser Gln Ser Phe Leu Asp Asp Thr Val Lys Glu
 165 170 175
 Tyr Thr Leu Ile Val Asn Asp Ser Ser Ser Pro Ile Gln Pro Val Leu
 180 185 190
 Ser Ser Asn Ser Lys Asn Ser Thr Pro Asp Glu Phe Leu Pro Asn Met
 195 200 205

-continued

Lys Ser Asp Ser Asn Ser Ala Ser Ser Asn Arg Glu Gln Asp Ser Val
 210 215 220
 Asp Thr Tyr Ser Asn Ile Pro Val Gly Arg Glu Ile Lys Ile Ile Leu
 225 230 235 240
 Pro Pro Lys Ala Ile Ala Leu Gln Phe Val Lys Ser Thr Trp Glu His
 245 250 255
 Cys Cys Val Leu Leu Arg Phe Tyr His Arg Pro Ser Phe Ile Arg Gln
 260 265 270
 Leu Asp Glu Leu Tyr Glu Thr Asp Pro Asn Asn Tyr Thr Ser Lys Gln
 275 280 285
 Met Gln Phe Leu Pro Leu Cys Tyr Ala Ala Ile Ala Val Gly Ala Leu
 290 295 300
 Phe Ser Lys Ser Ile Val Ser Asn Asp Ser Ser Arg Glu Lys Phe Leu
 305 310 315 320
 Gln Asp Glu Gly Tyr Lys Tyr Phe Ile Ala Ala Arg Lys Leu Ile Asp
 325 330 335
 Ile Thr Asn Ala Arg Asp Leu Asn Ser Ile Gln Ala Ile Leu Met Leu
 340 345 350
 Ile Ile Phe Leu Gln Cys Ser Ala Arg Leu Ser Thr Cys Tyr Thr Tyr
 355 360 365
 Ile Gly Val Ala Met Arg Ser Ala Leu Arg Ala Gly Phe His Arg Lys
 370 375 380
 Leu Ser Pro Asn Ser Gly Phe Ser Pro Ile Glu Ile Glu Met Arg Lys
 385 390 395 400
 Arg Leu Phe Tyr Thr Ile Tyr Lys Leu Asp Val Tyr Ile Asn Ala Met
 405 410 415
 Leu Gly Leu Pro Arg Ser Ile Ser Pro Asp Asp Phe Asp Gln Thr Leu
 420 425 430
 Pro Leu Asp Leu Ser Asp Glu Asn Ile Thr Glu Val Ala Tyr Leu Pro
 435 440 445
 Glu Asn Gln His Ser Val Leu Ser Ser Thr Gly Ile Ser Asn Glu His
 450 455 460
 Thr Lys Leu Phe Leu Ile Leu Asn Glu Ile Ile Ser Glu Leu Tyr Pro
 465 470 475 480
 Ile Lys Lys Thr Ser Asn Ile Ile Ser His Glu Thr Val Thr Ser Leu
 485 490 495
 Glu Leu Lys Leu Arg Asn Trp Leu Asp Ser Leu Pro Lys Glu Leu Ile
 500 505 510
 Pro Asn Ala Glu Asn Ile Asp Pro Glu Tyr Glu Arg Ala Asn Arg Leu
 515 520 525
 Leu His Leu Ser Phe Leu His Val Gln Ile Ile Leu Tyr Arg Pro Phe
 530 535 540
 Ile His Tyr Leu Ser Arg Asn Met Asn Ala Glu Asn Val Asp Pro Leu
 545 550 555 560
 Cys Tyr Arg Arg Ala Arg Asn Ser Ile Ala Val Ala Arg Thr Val Ile
 565 570 575
 Lys Leu Ala Lys Glu Met Val Ser Asn Asn Leu Leu Thr Gly Ser Tyr
 580 585 590
 Trp Tyr Ala Cys Tyr Thr Ile Phe Tyr Ser Val Ala Gly Leu Leu Phe
 595 600 605
 Tyr Ile His Glu Ala Gln Leu Pro Asp Lys Asp Ser Ala Arg Glu Tyr

-continued

1	5	10	15
Asp Phe Ser Ser Ala	Pro Asn Ile Lys Gly	Leu Asn Ser His Thr Gln	
20	25	30	
Leu Gln Phe Asp Ala	Asp Ser Arg Val Phe	Val Ser Asp Val Met Ala	
35	40	45	
Lys Asn Ser Lys Gln	Leu Leu Tyr Ala His	Ile Tyr Asn Tyr Leu Ile	
50	55	60	
Lys Asn Asn Tyr Trp	Asn Ser Ala Ala Lys	Phe Leu Ser Glu Ala Asp	
65	70	75	80
Leu Pro Leu Ser Arg	Ile Asn Gly Ser Ala	Ser Gly Gly Lys Thr Ser	
85	90	95	
Leu Asn Ala Ser Leu	Lys Gln Gly Leu Met	Asp Ile Ala Ser Lys Gly	
100	105	110	
Asp Ile Val Ser Glu	Asp Gly Leu Leu Pro	Ser Lys Met Leu Met Asp	
115	120	125	
Ala Asn Asp Thr Phe	Leu Leu Glu Trp Trp	Glu Ile Phe Gln Ser Leu	
130	135	140	
Phe Asn Gly Asp Leu	Glu Ser Gly Tyr Gln	Gln Asp His Asn Pro Leu	
145	150	155	160
Arg Glu Arg Ile Ile	Pro Ile Leu Pro Ala	Asn Ser Lys Ser Asn Met	
165	170	175	
Pro Ser His Phe Ser	Asn Leu Pro Pro Asn	Val Ile Pro Pro Thr Gln	
180	185	190	
Asn Ser Phe Pro Val	Ser Glu Glu Ser Phe	Arg Pro Asn Gly Asp Gly	
195	200	205	
Ser Asn Phe Asn Leu	Asn Asp Pro Thr Asn	Arg Asn Val Ser Glu Arg	
210	215	220	
Phe Leu Ser Arg Thr	Ser Gly Val Tyr Asp	Lys Gln Asn Ser Ala Asn	
225	230	235	240
Phe Ala Pro Asp Thr	Ala Ile Asn Ser Asp	Ile Ala Gly Gln Gln Tyr	
245	250	255	
Ala Thr Ile Asn Leu	His Lys His Phe Asn	Asp Leu Gln Ser Pro Ala	
260	265	270	
Gln Pro Gln Gln Ser	Ser Gln Gln Gln Ile	Gln Gln Pro Gln His Gln	
275	280	285	
Pro Gln His Gln Pro	Gln Gln Gln Gln Gln	Gln Gln Gln Gln Gln Gln	
290	295	300	
Gln Gln Gln Gln Gln	Gln Gln Gln Gln Gln	Gln Gln Gln Gln Gln Gln	
305	310	315	320
Gln Gln Gln His Gln	Gln Gln Gln Gln Thr	Pro Tyr Pro Ile Val Asn	
325	330	335	
Pro Gln Met Val Pro	His Ile Pro Ser Glu	Asn Ser His Ser Thr Gly	
340	345	350	
Leu Met Pro Ser Val	Pro Pro Thr Asn Gln	Gln Phe Asn Ala Gln Thr	
355	360	365	
Gln Ser Ser Met Phe	Ser Asp Gln Gln Arg	Phe Phe Gln Tyr Gln Leu	
370	375	380	
His His Gln Asn Gln	Gly Gln Ala Pro Ser	Phe Gln Gln Ser Gln Ser	
385	390	395	400
Gly Arg Phe Asp Asp	Met Asn Ala Met Lys	Met Phe Phe Gln Gln Gln	
405	410	415	

-continued

Ala Leu Gln Gln Asn Ser Leu Gln Gln Asn Leu Gly Asn Gln Asn Tyr
420 425 430

Gln Ser Asn Thr Arg Asn Asn Thr Ala Glu Glu Thr Thr Pro Thr Asn
435 440 445

Asp Asn Asn Ala Asn Gly Asn Ser Leu Leu Gln Glu His Ile Arg Ala
450 455 460

Arg Phe Asn Lys Met Lys Thr Ile Pro Gln Gln Met Lys Asn Gln Ser
465 470 475 480

Thr Val Ala Asn Pro Val Val Ser Asp Ile Thr Ser Gln Gln Gln Tyr
485 490 495

Met His Met Met Met Gln Arg Met Ala Ala Asn Gln Gln Leu Gln Asn
500 505 510

Ser Ala Phe Pro Pro Asp Thr Asn Arg Ile Ala Pro Ala Asn Asn Thr
515 520 525

Met Pro Leu Gln Pro Gly Asn Met Gly Ser Pro Val Ile Glu Asn Pro
530 535 540

Gly Met Arg Gln Thr Asn Pro Ser Gly Gln Asn Pro Met Ile Asn Met
545 550 555 560

Gln Pro Leu Tyr Gln Asn Val Ser Ser Ala Met His Ala Phe Ala Pro
565 570 575

Gln Gln Gln Phe His Leu Pro Gln His Tyr Lys Thr Asn Thr Ser Val
580 585 590

Pro Gln Asn Asp Ser Thr Ser Val Phe Pro Leu Pro Asn Asn Asn Asn
595 600 605

Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Ser Asn Asn
610 615 620

Ser Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Ser Asn Asn
625 630 635 640

Thr Pro Thr Val Ser Gln Pro Ser Ser Lys Cys Thr Ser Ser Ser Ser
645 650 655

Thr Thr Pro Asn Ile Thr Thr Thr Ile Gln Pro Lys Arg Lys Gln Arg
660 665 670

Val Gly Lys Thr Lys Thr Lys Glu Ser Arg Lys Val Ala Ala Ala Gln
675 680 685

Lys Val Met Lys Ser Lys Lys Leu Glu Gln Asn Gly Asp Ser Ala Ala
690 695 700

Thr Asn Phe Ile Asn Val Thr Pro Lys Asp Ser Gly Gly Lys Gly Thr
705 710 715 720

Val Lys Val Gln Asn Ser Asn Ser Gln Gln Gln Leu Asn Gly Ser Phe
725 730 735

Ser Met Asp Thr Glu Thr Phe Asp Ile Phe Asn Ile Gly Asp Phe Ser
740 745 750

Pro Asp Leu Met Asp Ser
755

<210> SEQ ID NO 32

<211> LENGTH: 750

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 32

Met Thr Ser Val Asn Arg Ser Asn Asn Thr Arg Ser Met Ser Ala Ser

-continued

1	5	10	15
Arg Ser Ala Thr Ser	Arg Val Arg Asn Thr	Thr Ala Asn Ser Ser Asp	
20	25	30	
Val Asn Ser Ser Lys	Arg Asn Ser Asn Ser	Val Tyr Asp Asp Asn Ser	
35	40	45	
Ser Lys Arg Arg Ser	Arg Arg Ser Asp Gly	Lys Asn Asn Asp His Thr	
50	55	60	
Tyr Arg Thr Thr Val	Lys Ser Lys Asn Ser	Arg Tyr Val Ser Ser Ser	
65	70	75	80
Lys Arg Ala Lys Arg	Asn Ser Val Gly Thr	Ser Ser Ala Ser Lys Ser	
85	90	95	
Ser Asn Gly Gly Ser	Ala His Lys Trp Ser	Asn Met Lys Asn Val Ser	
100	105	110	
Asn Ser Ala Val Asp	Ala Gly Ser Asp Ser	Lys Ser Val Gly Gly Arg	
115	120	125	
Lys Ser Asn Asn Ser	Asn Asp Lys Asp Asn	Ser Ala Arg Asp Asp Asn	
130	135	140	
Asn Ser Gly Asn Asn	Asn Asn Asn Asn Asn	His Ser Ser Asn Asn Asn	
145	150	155	160
Asp Asn Asn Asn Asn	Asn Asn Asp Asp Asn	Asn Asn Asn Asn Asn Ser	
165	170	175	
Asn Ser Arg Asp Asn	Asn Asn Asn Ser Asp	Asp Ser Asn Arg Asn Asp	
180	185	190	
Ser Cys Lys Ala Ser	Asn Lys Arg Ser Gly	Ala Lys Tyr Lys Val Val	
195	200	205	
Lys Arg Cys Ser Thr	Asn Ser Thr Thr Lys	Ser Trp Thr Tyr Lys Asn	
210	215	220	
Thr Asp Val Asn Asn	Tyr Val Thr Thr Thr	Ala Ser His Asp Val Gly	
225	230	235	240
Val Tyr Arg Arg Arg	Trp Val Tyr Gly Thr	Thr Asp Val Lys Asn Ser	
245	250	255	
Asn Met Asp Val Cys	Cys Thr His Val Val	Ser Ser Thr Met Ser Asp	
260	265	270	
Ser Lys Tyr Ser Thr	Trp Arg Gly Asp Ser	Arg Met Ala Ala Tyr Ser	
275	280	285	
Ser Asp Trp Lys Ser	Ala His Trp Tyr Thr	Ala Met Lys Tyr Tyr Asn	
290	295	300	
His Gly Lys Tyr Tyr	His Met Ser Thr Val	Asn Thr Ala Val Asn Gly	
305	310	315	320
Lys Ser Val Cys Thr	Thr Ser Tyr Met Val	Asp Asn Tyr Arg Ala Val	
325	330	335	
Arg Asn Asn Gly Asn	Arg Asn Ser Tyr Lys	His Ser Ala Met Ser Ser	
340	345	350	
Asp Asn Val Val Ser	Tyr Lys Gly Asp Ala	Asn Gly Cys Asn Asn Ala	
355	360	365	
Asp Met Val Asn Asp	Lys Tyr Arg His Gly	Ser Ala Ser His Val Gly	
370	375	380	
Gly Lys Asn Ala Lys	Tyr Lys Arg Lys Asp	Lys Lys Arg Lys Lys Ser	
385	390	395	400
Ser Asn Asn Asp Ser	Ser Val Thr Ser Ser	Thr Gly Asn Ser Arg Asn	
405	410	415	

-continued

Asp Asn Asp Asp Asp Met Ser Ser Thr Thr Ser Ser Asp His Asp Ala
 420 425 430

Asn Asp Asp Thr Arg Arg Ser Met Thr Asn Ala Trp Thr Lys Asn Met
 435 440 445

Thr Ser Lys Cys Gly Val Arg Lys His Gly Gly Ala His Trp Tyr Ser
 450 455 460

Cys Lys Ser Ser Ser Asp Val Ser Lys Trp Met Val Lys Arg Ala Trp
 465 470 475 480

Asp Thr Met Val Thr Met Asn Val Val Tyr Asp Asn Thr Ser Asn Ser
 485 490 495

Gly Asp Cys Asp Asp Tyr Asp Lys Ser Ser Asn Gly Gly Cys Trp Gly
 500 505 510

Thr Trp Asp Thr Cys Lys Asn Thr His Ser Ser Ser Asp Asn Gly Lys
 515 520 525

Asp Tyr Met Ala Asp Ser Thr Asp Gly Asp Lys Asp Asn Gly Lys Trp
 530 535 540

Lys Arg Ala Cys Arg Thr Arg Ser Arg Ser Gly Val Arg Asn Asp Tyr
 545 550 555 560

Arg Ser Ser Asn Thr Asn Gly Ser Val Lys Cys Asn His Asn Asn Val
 565 570 575

Gly Ala Ser Asp Ser Ala Arg Ser Asn Asn Thr Asp His Ala Val Ser
 580 585 590

Val Asn Gly Asp Asn His Tyr Val Gly Tyr Lys Lys Arg Ala Asp Tyr
 595 600 605

Thr Cys Asp Lys Asn Gly Ser Ala Ser Tyr Thr Thr Trp Tyr Val Asn
 610 615 620

Ser Asn Asn Thr Asn Asp Asn Asn Tyr Asn Ser Lys Asn Gly Cys Lys
 625 630 635 640

Ser Asp Tyr Asp Lys Thr Thr Tyr Val Asp Ala Thr Ser Trp Arg His
 645 650 655

Ser Ala Arg Lys Ala Asn Arg Arg Ala Cys Thr Thr Arg Arg Lys Ser
 660 665 670

Lys Asp Asn Val Met Ala Ala Thr Arg Gly Thr Arg Tyr Tyr Asn Lys
 675 680 685

Val Arg Thr Gly Asn Val Ala Thr His Asn Thr Trp Arg Thr His Val
 690 695 700

Asp Val Ser Val Met Lys Ala Lys Ser Ala Ser Arg Ser Arg Arg Asn
 705 710 715 720

Tyr Val Val Ser Asp Asp Asp Ala Met Lys Lys Lys Ala Lys Lys Thr
 725 730 735

Ser Thr Arg Val Ser Cys Thr Lys Gly Arg His Cys Thr Asp
 740 745 750

<210> SEQ ID NO 33
 <211> LENGTH: 710
 <212> TYPE: PRT
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 33

Met Asp Asn Lys Arg Tyr Asn Gly Asn Ser Asn Val Asp Gly Thr Tyr
 1 5 10 15
 Asp Arg Asn Asp Thr Arg Met Asn Thr Asn Ala Arg Ser Val Arg Val

-continued

20	25	30
Ser Asp Lys Arg Gly	Arg Ser Ser Ser Thr	Ser Lys Gly Ser Tyr Arg
35	40	45
Thr Arg Ala Gly Arg	Ser Asp Thr Thr Asn	Ser Ser Ala Lys His His
50	55	60
Ser Lys Lys Ser Thr	Val Val Val Val Thr	Ser Ser Thr Asp Ser Asn
65	70	75
Ser Thr Thr Tyr Ala	Arg Val Ser Ser Asp	Ser Thr Val Ala Thr Ser
85	90	95
Ser Thr Thr Thr Arg	Thr Arg Thr Arg Asn	Asn Thr Val Ser Ser Thr
100	105	110
Ala Ser Ser Ser Thr	Thr Asp Val Gly Asn	Ala Thr Ser Ala Asn Trp
115	120	125
Ser Ala Asn Ala Ser	Asn Thr Ser Ser Ser	Asp Tyr Ala Thr Ser Tyr
130	135	140
Thr Arg Lys Ser Thr	Asp Asn Tyr Thr Thr	Ala Asn Ser Lys Asn Gly
145	150	155
Asn Asn Trp Ser Ser	Ala Gly Asn Ser Asn	Thr Asp His Asn Thr Val
165	170	175
Asn Arg Arg Ser Ser	Ser Thr Thr Asn Arg	Val Tyr Thr Asp Ala Tyr
180	185	190
Tyr Ala Asn Tyr Val	Val Arg Val Lys Ser	Thr Ser Ser Val Asp Asp
195	200	205
Val Asp Ala Ser Asn	Trp Thr Ala Asn Lys	Val Val Asn Ser Ala Thr
210	215	220
Asn Thr Ser Ser Asn	Val Thr His Asn Ala	Val Asn Thr Ser Thr Ser
225	230	235
Ala Thr Cys Ser Tyr	Gly Lys Val Ser Ala	Arg Thr Arg Gly Asn Met
245	250	255
Ala Val Ser Thr Val	Ser Ala Cys Ala Ala	Gly Lys Ser Lys Val Gly
260	265	270
Ala Ser Thr Val Ser	Ala Arg Val Met Tyr	Asn Val Asn Gly Asn Asn
275	280	285
Thr Lys Asn His Gly	Val Asn Tyr Ser Thr	Ser Asn Asn Thr Tyr Cys
290	295	300
Asn Thr Asn Ser His	Ser Ser Asn Asn Tyr	Ser Ser Asp Ser Lys Lys
305	310	315
Asp His Thr Ser Ser	Lys Tyr Asp His Asn	His Asn Ala Lys Asn Lys
325	330	335
Gly Val Ser Asp Thr	Asn Tyr Gly His Asn	Ser Lys Val Lys Arg Lys
340	345	350
Asp Thr Asp Ala Lys	Arg Arg Lys Asp Ser	Asn Ser Ser Thr Met Ala
355	360	365
Val Met Asp Ser Ser	Asp Tyr Gly Asn Thr	Val Lys Asn Ser Ser Asn
370	375	380
Arg Asp Met Arg Lys	Cys Asn Lys Tyr Thr	Ser Met Gly Val His Lys
385	390	395
Asn Gly Thr Trp Val	Cys Lys Lys Met Ala	Asn Thr Arg Asn Val Thr
405	410	415
Ser Gly Val Ser Asp	Tyr Cys Thr Asn Asp	Gly Asn Tyr Val Gly Lys
420	425	430

-continued

Gly Trp Asn Ser Ser Val Ser His Trp Thr Val Asn Arg Tyr Gly Ser
 435 440 445
 Arg Ala Val Arg Ala Cys Ala Asp Ser Thr Cys Thr Thr Ser Val Ser
 450 455 460
 Tyr Ala Thr Asp Thr Asn Gly Thr Thr Trp Asp Thr Cys Thr Asn Lys
 465 470 475 480
 Asn Cys Asp Lys Val Asn Lys Asn Val Lys Cys Cys His Lys Gly Ser
 485 490 495
 Thr Val Lys Asn Arg Gly Gly Ala Ser Lys Asn Lys His Ala Asp Gly
 500 505 510
 Ser Ser Asp Ser Asp Gly Asn Tyr Gly Thr Tyr Lys Val Thr Ser Arg
 515 520 525
 Asp Asn Ser Val Arg Asp Ala Thr Lys Arg Asn Ser Asn Asn Ser Arg
 530 535 540
 Val Gly Ser Ser Ala Gly Ser Lys Ser Ser Lys Asn His Arg Lys His
 545 550 555 560
 Gly His Ser Gly Arg Ala Arg Gly Val Ser Val Ser Ser Val Arg Ser
 565 570 575
 Ser Asn Ser Arg His Asn Ser Val Met Asn Asn Ala Gly Thr Ala Asn
 580 585 590
 Asn Ala Met Ser Asn Ser Tyr Asn Asn Val Val Tyr Ser Gly Asn Asn
 595 600 605
 Asn Asn Gly Asn Ser Asn Gly Asp Asn Ser Asp Ser Arg Ala Asn Gly
 610 615 620
 Thr Asn Ser Val Asn Asn Val Ser Asn Asn Asn Asn Asn Tyr Asn Asn
 625 630 635 640
 Ser Gly Tyr Ser Ser Met Asn Ser Arg Ser Val Ser His Asn Asn Asn
 645 650 655
 Asn Asn Thr Asn Asn Tyr Asn Asn Asn Asp Asn Asp Asn Asn Asn Asn
 660 665 670
 Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn Asn
 675 680 685
 Asn Ser Asn Asn Ser Asn Asn Asn Asn Asn Asn Asp Thr Ser Tyr Arg
 690 695 700
 Tyr Arg Ser Tyr Gly Tyr
 705 710

<210> SEQ ID NO 34

<211> LENGTH: 477

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 34

Asp Thr Lys Gly Tyr Asp Asp Asp Ala Ala Thr Asp Gly Lys Lys His
 1 5 10 15
 Arg Arg Tyr Arg Tyr Val Ser Gly Ser Val Ser Gly Lys Arg Trp Thr
 20 25 30
 Asp Gly Val Ser Trp Ser Ser Arg Ser Gly Lys Tyr Lys Asp Lys Asn
 35 40 45
 Ala Gly Ser Asn Ala Asn Ala Thr Ser Ser Gly Ser Thr Asp Ser Ala
 50 55 60
 Val Thr Asp Gly Thr Ser Gly Ala Arg Asn Asn Ser Ser Ser Lys Lys

-continued

65	70	75	80
Lys Asn His Asp Thr 85	Met Gly His Ser Ser 90	Ser Asp Thr Ser Ser Ser 95	
Asn Arg Ser Asn Lys 100	Tyr Thr Gly Val Lys 105	Lys Thr Ser Val Lys Lys 110	
Arg Asn Ser Asn His 115	Val Ser Tyr Tyr Ser 120	Val Lys Asp Lys Asn Cys 125	
Val Thr Lys Ala Ser 130	Lys Asp Val Arg Ser 135	Val Ala Met Gly Asn Thr 140	
Thr Gly Asn Val Lys 145	Asn Asn Ser Thr Thr 150	Thr Gly Asn Gly Asn Asn 155	160
Asn Asn Lys Ser Asn 165	Ser Ser Thr Asn Thr 170	Val Ser Thr Asn Asn Asn 175	
Ser Ala Asn Asn Ala 180	Ala Gly Ser Asn Thr 185	Ser Ala Asn Lys Asn Tyr 190	
Tyr Tyr Lys Asn Asp 195	Ser Ser Gly Tyr Thr 200	Ala Ala Ser Thr Thr Met 205	
Tyr Thr Ala Asn Tyr 210	Thr Ser Asp Asn Thr 215	Asn Ala Thr Gly Met Asn 220	
Thr His Val Asn Asn 225	Asn Asn Asn Asn Ser 230	Asn Asn Ser Ser Asn Ser 235	240
Asn Asn Ser Asn Asn 245	Asn Asn Asn Asn Asn 250	Asn Asn Asn Asn Asn 255	
Asn Asn Asn Asn Asn 260	Asn Asn Asn Val Asn 265	Thr Asn Ala Gly Asn Gly 270	
Asn Asn Asn Arg His 275	Asn Ala Ser Ala Tyr 280	Asn Thr Thr Gly Asp Asn 285	
Gly Ser Tyr Tyr Tyr 290	Thr Thr Asn Asn Asn 295	Tyr Tyr Thr Thr Asn Val 300	
Thr Asn Ala Ser Thr 305	Asn Asn Gly Tyr Ser 310	Thr Ser Ser Thr His Tyr 315	320
Tyr Gly His Thr Ser 325	Ser Ala Ser Ala Ala 330	Ala Gly Ala Thr Gly Thr 335	
Gly Thr Ala Asn Val 340	Val Ser Ser Met His 345	Ala Asn Asn Asn Ser Ala 350	
Ser Ser Ala Thr Ser 355	Thr Ala Tyr Val Tyr 360	Ser Met Asn Val Asn Val 365	
Tyr Tyr Asn Ser Ser 370	Ala Ser Ala Tyr Lys 375	Arg Ala Asn Thr Thr Ser 380	
Asn Thr Asn Ala Ser 385	Gly Ala Thr Ser Thr 390	Asn Ser Gly Thr Met Ser 395	400
Asn Ala Tyr Ala Asn 405	Ser Tyr Thr Ser Val 410	Tyr Tyr Gly Tyr Ala Met 415	
Ala Ser Ala Asn Ser 420	Met Tyr His His His 425	Thr Val Tyr Ala Thr Asn 430	
Met Ser Ser Gly His 435	Thr Ser Thr Gly Ser 440	Asp His His His Tyr Asn 445	
Asp His Lys Asn Ala 450	Met Gly His Ala Asn 455	Asn Asn Asn Thr Asn Asn 460	
Asp Thr Met Asn Asn 465	Asn Thr Asn Thr Ser 470	Thr Thr Thr 475	

-continued

```

<210> SEQ ID NO 35
<211> LENGTH: 454
<212> TYPE: PRT
<213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 35

Met Asp Val Arg Ala Ala Cys Ser Ala Ser Gly Arg Thr Gly Lys Lys
 1          5          10          15

Gly Tyr Ser Tyr Lys Met Ser Asn Ser Gly Gly Ser Ser Ser Gly Gly
20          25          30

Ser Asp Val Gly Ser Thr Asn Gly Ser Asn Arg Ala Lys Asn Thr Asn
35          40          45

Tyr Lys Lys Thr Asn Lys Lys Tyr Lys Ala Thr Asp Lys Ala Asn Asp
50          55          60

Thr Lys Tyr Tyr Ser Asn Asp Lys Lys Ser Lys Arg Ser Ala Asn Ser
65          70          75          80

Met Asn Asp Lys Asp Lys Cys Arg Thr Thr Asn Lys Asp Met Thr Arg
85          90          95

Tyr Asp Ser Lys Ser Lys Val Thr Asn Cys Asp His Lys Ala Ser Ser
100         105         110

His Ser Met Lys Tyr Lys Lys Arg Ser Val Asp Lys Asp His Val Met
115         120         125

Lys Asp Asp Ser Ser Val Lys Ala Ser Lys Met Asn Ser His Asn Tyr
130         135         140

Ser Thr Asn Thr Met Asn Lys Met Asp Val Tyr Thr Lys Ala Asn Met
145         150         155         160

Ala Asn Lys Lys Lys Ser Asp Thr Ser Thr Trp Lys Asn Lys Asn Lys
165         170         175

Ser His Val Ser Tyr Asn Asn Asp Lys Ser Lys Thr Lys Trp Tyr Asn
180         185         190

Asp Ser Asp Asp Asp Asp Asp Asn Asn Val Asn Asn Asn Asp Asn Asn
195         200         205

Asn Asn Asn Lys Asn Asp Asn Asn Asn Asp Asn Asn Asn Asp Thr Ser
210         215         220

Asn Asn Asn Asn Asn Asn Asn Asn Arg Thr Lys Asn Asn Arg Asn Asn
225         230         235         240

Arg Asp Trp Lys Thr Lys Lys Cys Thr Asp Met Asn Asp Lys Arg Asp
245         250         255

Asn Asn Asn Lys Asn Asp Met Ala Arg Asn Asp Asn Lys Asn Tyr Asn
260         265         270

Asn Val Asn Lys Arg Asn His Lys Ser Ser Cys Arg Arg Asp Gly Tyr
275         280         285

Ser Ala Asn Asn Ala Val Asn Ser Thr His Ala Ser Asn Lys Asn Val
290         295         300

Asn Asp Met Asn Asn Asp Thr Tyr Lys Asn Lys Thr Asp Thr Asn Lys
305         310         315         320

Lys Asn Asp Ser Asn Ser Asn Asp Val Thr Arg Lys Lys Arg Lys Thr
325         330         335

Ser Asp Gly Asn Tyr Ser Arg Asn Asn Val Ser Val Ser Arg Ser Lys
340         345         350

Ala Thr Thr Lys Lys Thr Lys Lys Lys Lys Arg Arg Asp Gly Lys Asp

```

-continued

```

355                360                365
Lys Lys Asn Lys Lys Asn Ala Asp Asn Lys Lys Asn Asn Ala Val Thr
370                375                380

Val Ser Val Tyr Asp Ser Asn Lys Val Lys Ser Asn Lys Arg Ser Arg
385                390                395                400

Lys Val Asn Asn Lys Ser Asp Val Val Asn Ser Gly Lys Asp Ser Arg
405                410                415

Val Lys Ser Cys Lys Lys Tyr Ala Asp Asn Asn Thr Lys Ser Asn Asp
420                425                430

Ala Asp Gly Trp Asp Asp Met Asn Trp Val Asp Arg Gly Cys Ala Thr
435                440                445

Thr Arg Trp Arg Ala Lys
450

<210> SEQ ID NO 36
<211> LENGTH: 284
<212> TYPE: PRT
<213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 36
Met Asn Val Thr Ser Lys Asp Gly Asn His Ser Ser Lys Lys Asn Arg
1      5      10      15
Asn Thr Asn Lys Arg His Lys Asn Ala Ser Asn Asp Arg Asp Ser Val
20     25     30
Ser Ser Asn Thr Thr Ser Met Thr Asp Asp Ala Asp Tyr Asn Gly Ala
35     40     45
Ser Arg Thr Lys Asn Asn Ser Asp Ser Asp Arg Ser Asn Asp Thr Lys
50     55     60
Asn Asn Tyr Asn Lys Arg Thr Gly Tyr Asn Tyr Asn Gly Ser Gly Asn
65     70     75     80
Arg Tyr Thr Arg Lys Arg Thr Ala Asn Lys Ala Tyr Ser Asp Asp Asn
85     90     95
Val Lys Asp Asp Asn Asn Thr Lys Lys Ala Ser Arg Ser Ser Gly Arg
100    105   110
Asn Val Asn Thr Arg Asn Lys Ser Lys Ser His Lys Val Lys Asn Asn
115    120   125
Lys Ser Ser Ser Arg Lys Ser Ser Ala Ala Arg Lys Gly Lys Tyr Asn
130    135   140
Ser Asn Ser Asp Ser Thr Thr Arg Lys Val Thr Asp Val Lys Lys Arg
145    150   155   160
Ser Lys Trp His Arg His Asp Lys Lys Met Val Lys Lys Ser Arg Tyr
165    170   175
Arg Lys Arg Met Arg Gly Thr Asp Val Ser Ser Ser Asp Asn Ser Lys
180    185   190
Ser Thr Thr Lys Ser Tyr Val Ser Lys Asn Ser Ala Met Asn Asn Asn
195    200   205
Asp Val Thr Asp Asn Lys Lys Thr Asn Asn Asn Lys Ala Arg Asp Ser
210    215   220
Met His Thr Lys Lys Asp Thr Lys Asp Asp Thr Asp Ser Lys Lys Arg
225    230   235   240
Lys Val Val Thr Asn Asp Asn Ala Ala Met Val Asn Lys Gly Trp Arg
245    250   255

```

-continued

Lys Asn Val Met Met Tyr Lys Lys Ser Gly Asn Met Lys Lys Tyr Arg
 260 265 270

Tyr Trp Thr Cys Tyr Cys Asn Tyr Val Tyr Tyr Arg
 275 280

<210> SEQ ID NO 37
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer
 <400> SEQUENCE: 37

gggaattccc attaccgaca ttggggcgc 29

<210> SEQ ID NO 38
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer
 <400> SEQUENCE: 38

ggggattctg attgattgat tgattgtac 29

<210> SEQ ID NO 39
 <211> LENGTH: 720
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: superbright
 GFP encoding sequence
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(720)
 <400> SEQUENCE: 39

atg gct agc aaa gga gaa gaa ctc ttc act gga gtt gtc cca att ctt 48
 Met Ala Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
 1 5 10 15

gtt gaa tta gat ggt gat gtt aat ggg cac aaa ttt tct gtc agt gga 96
 Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
 20 25 30

gag ggt gaa ggt gat gca aca tac gga aaa ctt acc ctt aaa ttt att 144
 Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
 35 40 45

tgc act act gga aaa cta cct gtt cca tgg cca aca ctt gtc act act 192
 Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
 50 55 60

ttc act tat ggt gtt cag tgc ttt tca aga tac ccg gat cat atg aaa 240
 Phe Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
 65 70 75 80

cgg cat gac ttt ttc aag agt gcc atg ccc gaa ggt tat gta cag gaa 288
 Arg His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
 85 90 95

aga act ata ttt ttc aaa gat gac ggg aac tac aag aca cgt gct gaa 336
 Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
 100 105 110

gtc aag ttt gaa ggt gat acc ctt gtt aat aga atc gag tta aaa ggt 384
 Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
 115 120 125

-continued

```

att gat ttt aaa gaa gat gga aac att ctt ggg cac aaa ttg gaa tac      432
Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
130                135                140

aac tat aac tca cac aat gta tac atc atg gca gac aaa caa aag aat      480
Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145                150                155                160

gga atc aaa gct aac ttc aaa att aga cac aac att gaa gat gga agc      528
Gly Ile Lys Ala Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
165                170                175

gtt caa cta gca gac cat tat caa caa aat act cca att ggc gat ggc      576
Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180                185                190

cct gtc ctt tta cca gac aac cat tac ctg tcc aca caa tct gcc ctt      624
Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
195                200                205

tcg aaa gat ccc aac gaa aag aga gac cac atg gtc ctt ctt gag ttt      672
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
210                215                220

gta aca gct gct ggg att aca cat ggc atg gat gaa cta tac aaa tga      720
Val Thr Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu Tyr Lys
225                230                235

```

<210> SEQ ID NO 40

<211> LENGTH: 239

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: superbright GFP encoding sequence

<400> SEQUENCE: 40

```

Met Ala Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
 1                5                10                15

Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
20                25                30

Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
35                40                45

Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
50                55                60

Phe Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
65                70                75                80

Arg His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
85                90                95

Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
100               105               110

Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
115               120               125

Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
130               135               140

Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145               150               155               160

Gly Ile Lys Ala Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
165               170               175

Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180               185               190

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu

```

-continued

195	200	205	
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe			
210	215	220	
Val Thr Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu Tyr Lys			
225	230	235	

<210> SEQ ID NO 41
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 synthetic primer

<400> SEQUENCE: 41

gaccgcgat ggctagcaaa ggagaag 27

<210> SEQ ID NO 42
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 synthetic primer

<400> SEQUENCE: 42

cctgagctct catttgata gttcatcc 28

<210> SEQ ID NO 43
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 synthetic primer

<400> SEQUENCE: 43

ggaggatcca tggatcggga taagttaatc tcag 34

<210> SEQ ID NO 44
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 synthetic primer

<400> SEQUENCE: 44

ggaccgcggt tagcggttct gttgagaaaa gttgcc 36

<210> SEQ ID NO 45
 <211> LENGTH: 7239
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: vector
 containing chimeric gene

<400> SEQUENCE: 45

gacgaaagg cctcgtgata cgcctatatt tataggttaa tgtcatgata ataatggttt 60

cttaggacgg atcgcttgcc tgtaacttac acgcgcctcg tatcttttaa tgatggaata 120

atttggaat ttactctgtg tttatttatt tttatgtttt gtatttggat tttagaaagt 180

-continued

aaataaagaa	ggtagaagag	ttacggaatg	aagaaaaaaaa	aataaaciaa	ggtttaaaaa	240
atltcaacia	aaagcgtact	ttacatatat	atltattaga	caagaaaagc	agattaaata	300
gatatacatt	cgattaacga	taagtaaaat	gtaaaatcac	aggatlttcg	tgtgtggtct	360
tctacacaga	caagatgaaa	caattcggca	ttaatacctg	agagcaggaa	gagcaagata	420
aaaggtagta	ttgttggtcg	atccccctag	agtctlttac	atcttcggaa	aacaaaaact	480
atlttttctt	taatttcttt	ttttactttc	tatltttaat	ttatatattt	atattaaaa	540
atltaaatta	taattatltt	tatagcacgt	gatgaaaagg	accaggtgg	cactlttcgg	600
ggaaatgtgc	gcggaacccc	tatltgttta	ttlttctaaa	tacattcaaa	tatgtatccg	660
ctcatgagac	aataaccctg	ataaatgctt	caataatatt	gaaaaaggaa	gagtatgagt	720
atltcaacatt	tccgtgtcgc	ccttatltcc	ttltttgctg	cattltgcct	tccgtltttt	780
gctcacccag	aaagcgtggt	gaaagtaaaa	gatgctgaag	atcagttggg	tgcacgagtg	840
ggttacatcg	aactggatct	caacagcgg	aagatccttg	agagtlttcg	ccccgaagaa	900
cgltttccaa	tgatgagcac	ttlttaagtt	ctgctatgtg	gcgcggtatt	atccccgatt	960
gagcggggc	aagagcaact	cggtcggcgc	atacactatt	ctcagaatga	cttggttgag	1020
tactcacccag	tcacagaaaa	gcatcttacg	gatggcatga	cagtaagaga	attatgcagt	1080
gctgccataa	ccatgagtga	taacactgcg	gccaaactac	ttctgacaac	gatcggagga	1140
ccgaaggagc	taaccgcttt	tttgcacaac	atgggggatc	atgtaactcg	ccttgatcgt	1200
tgggaaccgg	agctgaatga	agccatacca	aacgacgagc	gtgacaccac	gatgcctgta	1260
gcaatggcaa	caacgttgcg	caaaactatta	actggcgaac	tacttactct	agcttccccg	1320
caacaattaa	tagactggat	ggaggcggat	aaagttgcag	gaccacttct	gcgctcggcc	1380
cttccggctg	gctggtttat	tgctgataaa	tctggagccg	gtgagcgtgg	gtctcgcggt	1440
atcattgcag	cactggggcc	agatggtaag	ccctccccga	tcgtagtatt	ctacacgacg	1500
gggagtcagg	caactatgga	tgaacgaaat	agacagatcg	ctgagatagg	tgcctcactg	1560
attaagcatt	ggtaactgtc	agaccaagtt	tactcatata	tactttagat	tgatttaaaa	1620
cttcatlttt	aatlttaaa	gatctagggt	aagatccttt	ttgataatct	catgacccaa	1680
atccctaac	gtgagtlttc	gttccactga	gcgtcagacc	ccgtagaaaa	gatcaaagga	1740
tcttcttgag	atcctltttt	tctgcgcgta	atctgctgct	tgcaaacaaa	aaaaccaccg	1800
ctaccagcgg	tgglttggtt	gccggatcaa	gagctaccaa	ctctltttcc	gaaggtaact	1860
ggcttcagca	gagcgcagat	accaaatact	gtccttctag	tgtagccgta	gtaggcccac	1920
cacttcaaga	actctgtagc	accgcctaca	tacctcgtc	tgctaactcct	gtagcagtg	1980
gctgctgcca	gtggcgataa	gtcgtgtcct	accgggttgg	actcaagacg	atagttaccg	2040
gataaggcgc	agcggtcggg	ctgaaacggg	ggttcgtgca	cacagcccag	cttgagcgga	2100
acgacctaca	ccgaactgag	atacctacag	cgtgagctat	gagaaagcgc	cacgcttccc	2160
gaaggagaaa	aggcggacag	gtatccggta	agcggcaggg	tcggaacagg	agagcgcacg	2220
agggagcttc	cagggggaaa	cgctcgtgat	ctlttatagtc	ctgtcgggtt	tcgccacctc	2280
tgacttgagc	gtcgatlttt	gtgatgctcg	tcaggggggc	ggagcctatg	gaaaaacgcc	2340
agcaacgcgg	cctlttttacg	gttccctggc	ttlttgctggc	ctltttgctca	catgttcttt	2400
cctcgcgttat	ccctgatct	tgtggataac	cgtattaccg	cctlttgagtg	agctgatacc	2460

-continued

gctcgcgcga gccgaacgac cgagcgcagc gagtcagtga gcgaggaagc ggaagagcgc	2520
ccaatacgcga aaccgcctct ccccgcgcggt tggccgattc attaatgcag ctggcacgac	2580
aggtttcccg actggaagc gggcagtgag cgcaacgcaa ttaatgtgag ttacctcact	2640
cattaggcac ccagggcttt acactttatg cttccggctc gtatgttggtg tggaaatgtg	2700
agcggataac aatttcacac aggaaacagc tatgacctg attacgcaa gctcggaaat	2760
aaccctcact aaaggaaca aaagctgggt accgggcccc ccctcgaggt cgacggatc	2820
gataagcttg atategaatt ccattaccg acatttgggc gctatacgtg catatgttca	2880
tgtatgtatc tgtatttaaa acacttttgt attatttttc ctcatatag tgtatagggt	2940
tatacggatg atttaattat tacttcaaca ccctttattt caggctgata tcttagcctt	3000
gttactagtt agaaaaagac atttttgctg tcagtcactg tcaagagatt cttttgctgg	3060
catttcttct agaagcaaaa agagcagatgc gtcttttccg ctgaaccgtt ccagcaaaaa	3120
agactaccaaa cgcaatatgg attgtcagaa tcatataaaa gagaagcaaa taactccttg	3180
tcttgatca attgcattat aatatcttct tggtagtgca atatcatata gaagtcateg	3240
aaatagatat taagaaaaac aaactgtaca atcaatcaat caatcaggat ccatggatac	3300
ggataagtta atctcagagg ctgagtctca tttttctcaa ggaaacctag cagaagctgt	3360
tgcgaagttg acatccgcag ctcagtcgaa cccaatgac gagcaaatgt caactattga	3420
atcattaatt caaaaaatcg caggatacgt catggacaac cgtagtgggtg gtagtgacgc	3480
ctcgcaagat cgtgctgctg gtggtgggtc atcttttatg aacactttaa tggcagactc	3540
taagggttct tcccaaacgc aactagggaa actagctttg ttagccacag tgatgacaca	3600
ctcatcaaat aaaggttctt ctaacagagg gtttgacgta gggactgtca tgtcaatgct	3660
aaaggttctt ggcggcggga gccaaagtat ggggtgcttc ggctggtg ccttggett	3720
tcaattcttt aagtcaggta acaattccca aggtcaggga caaggcaag gtcaaggtea	3780
aggteaagga caaggtcaag gtcaagggtc ttttactgct ttggcgtctt tggcttcac	3840
ttcatgaat tccaacaaca ataatcagca aggtcaaaat caaagctccg gtggttctc	3900
ctttggagca ctagcttcta tggcaagttc ttttatgcat tccaataata atcagaactc	3960
caacaatagt caacagggtt ataaccaatc ctatcaaaac ggtaacccaa atagtcaagg	4020
ttacaataat caacagtacc aaggtggcaa cgggtggttac caacaacaac agggacaatc	4080
tgggtggtct ttttctcat tggctccat ggctcaatct tacttaggtg gtggacaaac	4140
tcaatccaac caacagcaat acaatcaaca aggcacaaac aaccagcagc aataccagca	4200
acaaggccaa aactatcagc accaacaaca gggtcagcag cagcaacaag gccactccag	4260
ttcattctca gctttggctt ccattggcaag ttcctacctg ggcaataact ccaattcaaa	4320
ttcaggttat gggggccagc aacaggctaa tgagtatggt agaccacaac acaatggtea	4380
acaacaatct aatgagtacg gaagaccgca atacggcggga aaccagaact ccaatggaca	4440
gcacgaatcc ttaattttt ctggcaactt ttctcaacag aacaataacg gcaaccagaa	4500
ccgtacccc cggtaggcta gcaaaggaga agaactcttc actggagttg tcccaattct	4560
tgttgaaata gatggtgatg ttaatgggca caaattttct gtcagtggag aggggtgaagg	4620
tgatgcaaca tacggaaaac ttacccttaa atttatttgc actactggaa aactacctgt	4680
tccatggcca acacttgtea ctactttcac ttatggtgtt cagtgtcttt caagataccc	4740

-continued

ggatcatatg aaacggcatg actttttcaa gagtgccatg cccgaagggt atgtacagga 4800
aagaactata tttttcaaag atgacgggaa ctacaagaca cgtgctgaag tcaagtttga 4860
aggtgatacc ctgttaata gaatcgagtt aaaaggatt gattttaaag aagatggaaa 4920
cattcttggg cacaaattgg aatacaacta taactcacac aatgtatata tcatggcaga 4980
caaaaaaaaa aatggaatca aagctaactt caaaattaga cacaacattg aagatggaaag 5040
cgttcaacta gcagaccatt atcaacaaaa tactccaatt ggcgatggcc ctgtcctttt 5100
accagacaac cattacctgt ccacacaatc tgccctttcg aaagatccca acgaaaagag 5160
agaccacatg gtccttcttg agtttgtaac agctgctggg attacacatg gcatggatga 5220
actatacaaa tgagagctcc aattcgccct atagtgagtc gtattacaat tcaactggccg 5280
tcgttttaca acgctgtgac tgggaaaacc ctggcggttac ccaacttaat cgccttgag 5340
cacatcccc tttcgcagc tggcgtaata gcgaagaggc ccgaccgat cgccttccc 5400
aacagttgag cagcctgaat ggcgaatggc gcgacgcgcc ctgtagcggc gcattaagcg 5460
cggcgggtgt ggtggttacg cgcagcgtga ccgctacact tgccagcggc ctagcgcggc 5520
ctcctttcgc tttcttccct tctttctcg ccacgttcgc cggctttccc cgtcaagctc 5580
taaatcgggg gctcccttta gggttccgat ttagtgcttt acggcacctc gacccccaaa 5640
aacttgatta gggatgatgt tcaagtagtg ggccatcgcc ctgatagacg gtttttcggc 5700
ctttgacggt ggagtccacg tcttttaata gtggactctt gttccaaact ggaacaacac 5760
tcaaccctat ctcggtctat tcttttgatt tataagggat tttgccgatt tcggcctatt 5820
ggttaaaaaa tgagctgatt taacaaaaat ttaacgcgaa ttttaacaaa atattaacgt 5880
ttacaatttc ctgatcgggt attttctcct tacgcatctg tgcggtatth cacaccgat 5940
agggtaataa ctgatataat taaattgaag ctctaatttg tgagtttagt atacatgcat 6000
ttacttataa tacagttttt tagttttgct ggccgcatct tctcaaatat gcttcccagc 6060
ctgcttttct gtaacgttca cctctacct tagcatccct tccctttgca aatagtctc 6120
ttccaacaat aataatgtca gatcctgtag agaccacatc atccacgggt ctatactgtt 6180
gacccaatgc gtctcccttg tcatctaaac ccacaccggg tgcataatc aaccaatcgt 6240
aaccttcate tcttccacc atgtctctt gagcaataaa gccgataaca aatctttgt 6300
cgctcttcgc aatgtcaaca gtacccttag tatattctcc agtagatagg gagcccttg 6360
atgacaatc tgtaacatc aaaaggcctc taggttctt tgttacttct tctgccgct 6420
gcttcaaac gtaacaata cctgggccc caacaccgtg tgcattcgta atgtctgcc 6480
attctgctat tctgtataca cccgcagagt actgcaattt gactgtatta ccaatgtcag 6540
caaatttct gtcttgaag agtaaaaaat tgtacttggc ggataatgcc tttagcggct 6600
taactgtgcc ctccatgaa aatcagtc agatatccac atgtgttttt agtaacaaa 6660
ttttgggacc taatgcttca actaactcca gtaattcctt ggtggtacga acatccaatg 6720
aagcacacaa gtttgtttgc tttctgtgca tgatattaaa tagcttggca gcaacaggac 6780
taggatgagt agcagcagct tcttatatg tagctttcga catgatttat ctctgttcc 6840
tgagggtttt tgttctgtgc agttgggtta agaatactgg gcaatttcat gtttcttcaa 6900
cactacatat gcgtatatat accaatctaa gtctgtgctc ctctctctgt tcttcttct 6960
gttcggagat taccgaatca aaaaaatttc aaagaaccg aatcaaaaa aagaataaaa 7020

-continued

```

aaaaaaatga tgaattgaat tgaaaagctg tggatggg cactctcagt acaatctgct 7080
ctgatgccgc atagttaagc cagccccgac acccgccaac acccgctgac ggcgcctgac 7140
gggcttgtct gctccccgca tccgcttaca gacaagctgt gaccgtctcc gggagctgca 7200
tgtgtcagag gttttaccgc tcataccgca aacgcgcga 7239

```

```

<210> SEQ ID NO 46
<211> LENGTH: 741
<212> TYPE: PRT
<213> ORGANISM: Pichia pinus

```

```

<400> SEQUENCE: 46

```

```

Met Ser Gln Asp Gln Gln Gln Gln Gln Phe Asn Ala Asn Asn Leu
1          5          10          15
Ala Gly Asn Val Gln Asn Ile Asn Leu Asn Ala Pro Ala Tyr Asp Pro
20          25          30
Ala Val Gln Ser Tyr Ile Pro Asn Thr Ala Gln Ala Phe Val Pro Ser
35          40          45
Ala Gln Pro Tyr Ile Pro Gly Gln Gln Glu Gln Gln Phe Gly Gln Tyr
50          55          60
Gly Gln Gln Gln Gln Asn Tyr Asn Gln Gly Gly Tyr Asn Asn Tyr Asn
65          70          75          80
Asn Arg Gly Gly Tyr Ser Asn Asn Arg Gly Gly Tyr Asn Asn Ser Asn
85          90          95
Arg Gly Gly Tyr Ser Asn Tyr Asn Ser Tyr Asn Thr Asn Ser Asn Gln
100         105         110
Gly Gly Tyr Ser Asn Tyr Asn Asn Asn Tyr Ala Asn Asn Ser Tyr Asn
115         120         125
Asn Asn Asn Asn Tyr Asn Asn Asn Tyr Asn Gln Gly Tyr Asn Asn Tyr
130         135         140
Asn Ser Gln Pro Gln Gly Gln Asp Gln Gln Gln Glu Thr Gly Ser Gly
145         150         155         160
Gln Met Ser Leu Glu Asp Tyr Gln Lys Gln Gln Lys Glu Ser Leu Asn
165         170         175
Lys Leu Asn Thr Lys Pro Lys Lys Val Leu Lys Leu Asn Leu Asn Ser
180         185         190
Ser Thr Val Lys Ala Pro Ile Val Thr Lys Lys Lys Glu Glu Glu Pro
195         200         205
Val Asn Gln Glu Ser Lys Thr Glu Glu Pro Ala Lys Glu Glu Ile Lys
210         215         220
Asn Gln Glu Pro Ala Glu Ala Glu Asn Lys Val Glu Glu Glu Ser Lys
225         230         235         240
Val Glu Ala Pro Thr Ala Ala Lys Pro Val Ser Glu Ser Glu Phe Pro
245         250         255
Ala Ser Thr Pro Lys Thr Glu Ala Lys Ala Ser Lys Glu Val Ala Ala
260         265         270
Ala Ala Ala Ala Leu Lys Lys Glu Val Ser Gln Ala Lys Lys Glu Ser
275         280         285
Asn Val Thr Asn Ala Asp Ala Leu Val Lys Glu Gln Glu Glu Gln Ile
290         295         300
Asp Ala Ser Ile Val Asn Asp Met Phe Gly Gly Lys Asp His Met Ser
305         310         315         320

```

-continued

Ile Ile Phe Met Gly His Val Asp Ala Gly Lys Ser Thr Met Gly Gly
 325 330 335
 Asn Leu Leu Phe Leu Thr Gly Ala Val Asp Lys Arg Thr Val Glu Lys
 340 345 350
 Tyr Glu Arg Glu Ala Lys Asp Ala Gly Arg Gln Gly Trp Tyr Leu Ser
 355 360 365
 Trp Ile Met Asp Thr Asn Lys Glu Glu Arg Asn Asp Gly Lys Thr Ile
 370 375 380
 Glu Val Gly Lys Ser Tyr Phe Glu Thr Asp Lys Arg Arg Tyr Thr Ile
 385 390 395 400
 Leu Asp Ala Pro Gly His Lys Leu Tyr Ile Ser Glu Met Ile Gly Gly
 405 410 415
 Ala Ser Gln Ala Asp Val Gly Val Leu Val Ile Ser Ser Arg Lys Gly
 420 425 430
 Glu Tyr Glu Ala Gly Phe Glu Arg Gly Gly Gln Ser Arg Glu His Ala
 435 440 445
 Ile Leu Ala Lys Thr Gln Gly Val Asn Lys Leu Val Val Val Ile Asn
 450 455 460
 Lys Met Asp Asp Pro Thr Val Asn Trp Ser Lys Glu Arg Tyr Glu Glu
 465 470 475 480
 Cys Thr Thr Lys Leu Ala Met Tyr Leu Lys Gly Val Gly Tyr Gln Lys
 485 490 495
 Gly Asp Val Leu Phe Met Pro Val Ser Gly Tyr Thr Gly Ala Gly Leu
 500 505 510
 Lys Glu Arg Val Ser Gln Lys Asp Ala Pro Trp Tyr Asn Gly Pro Ser
 515 520 525
 Leu Leu Glu Tyr Leu Asp Ser Met Pro Leu Ala Val Arg Lys Ile Asn
 530 535 540
 Asp Pro Phe Met Leu Pro Ile Ser Ser Lys Met Lys Asp Leu Gly Thr
 545 550 555 560
 Val Ile Glu Gly Lys Ile Glu Ser Gly His Val Lys Lys Gly Gln Asn
 565 570 575
 Leu Leu Val Met Pro Asn Lys Thr Gln Val Glu Val Thr Thr Ile Tyr
 580 585 590
 Asn Glu Thr Glu Ala Glu Ala Asp Ser Ala Phe Cys Gly Glu Gln Val
 595 600 605
 Arg Leu Arg Leu Arg Gly Ile Glu Glu Glu Asp Leu Ser Ala Gly Tyr
 610 615 620
 Val Leu Ser Ser Ile Asn His Pro Val Lys Thr Val Thr Arg Phe Glu
 625 630 635 640
 Ala Gln Ile Ala Ile Val Glu Leu Lys Ser Ile Leu Ser Thr Gly Phe
 645 650 655
 Ser Cys Val Met His Val His Thr Ala Ile Glu Glu Val Thr Phe Thr
 660 665 670
 Gln Leu Leu His Asn Leu Gln Lys Gly Thr Asn Arg Arg Ser Lys Lys
 675 680 685
 Ala Pro Ala Phe Ala Lys Gln Gly Met Lys Ile Ile Ala Val Leu Glu
 690 695 700
 Thr Thr Glu Pro Val Cys Ile Glu Ser Tyr Asp Asp Tyr Pro Gln Leu
 705 710 715 720

-continued

Gly Arg Phe Thr Leu Arg Asp Gln Gly Gln Thr Ile Ala Ile Gly Lys
725 730 735

Val Thr Lys Leu Leu
740

<210> SEQ ID NO 47
<211> LENGTH: 715
<212> TYPE: PRT
<213> ORGANISM: Candida albicans

<400> SEQUENCE: 47

Met Ala Asn Ala Ser Leu Asn Gly Asp Gln Ser Lys Gln Gln Gln Gln
1 5 10 15

Gln Gln Gln Gln Gln Gln Gln Gln Asn Tyr Tyr Asn Pro Asn Ala
20 25 30

Ala Gln Ser Phe Val Pro Gln Gly Gly Tyr Gln Gln Phe Gln Gln Phe
35 40 45

Gln Pro Gln Gln Gln Gln Gln Tyr Gly Gly Tyr Asn Gln Tyr Asn
50 55 60

Gln Tyr Gln Gly Gly Tyr Gln Gln Asn Tyr Asn Asn Arg Gly Gly Tyr
65 70 75 80

Gln Gln Gly Tyr Asn Asn Arg Gly Gly Tyr Gln Gln Asn Tyr Asn Asn
85 90 95

Arg Gly Gly Tyr Gln Gly Tyr Asn Gln Asn Gln Gln Tyr Gly Gly Tyr
100 105 110

Gln Gln Tyr Asn Ser Gln Pro Gln Gln Gln Gln Gln Gln Ser Gln
115 120 125

Gly Met Ser Leu Ala Asp Phe Gln Lys Gln Lys Thr Glu Gln Gln Ala
130 135 140

Ser Leu Asn Lys Pro Ala Val Lys Lys Thr Leu Lys Leu Ala Gly Ser
145 150 155 160

Ser Gly Ile Lys Leu Ala Asn Ala Thr Lys Lys Val Asp Thr Thr Ser
165 170 175

Lys Pro Gln Ser Lys Glu Ser Ser Pro Ala Pro Ala Pro Ala Ala Ser
180 185 190

Ala Ser Ala Ser Ala Pro Gln Glu Glu Lys Lys Glu Glu Lys Glu Ala
195 200 205

Ala Ala Ala Thr Pro Ala Ala Ala Pro Glu Thr Lys Lys Glu Thr Ser
210 215 220

Ala Pro Ala Glu Thr Lys Lys Glu Ala Thr Pro Thr Pro Ala Ala Lys
225 230 235 240

Asn Glu Ser Thr Pro Ile Pro Ala Ala Ala Ala Lys Lys Glu Ser Thr
245 250 255

Pro Val Ser Asn Ser Ala Ser Val Ala Thr Ala Asp Ala Leu Val Lys
260 265 270

Glu Gln Glu Asp Glu Ile Asp Glu Glu Val Val Lys Asp Met Phe Gly
275 280 285

Gly Lys Asp His Val Ser Ile Ile Phe Met Gly His Val Asp Ala Gly
290 295 300

Lys Ser Thr Met Gly Gly Asn Ile Leu Tyr Leu Thr Gly Ser Val Asp
305 310 315 320

Lys Arg Thr Val Glu Lys Tyr Glu Arg Glu Ala Lys Asp Ala Gly Arg
325 330 335

-continued

Gln Gly Trp Tyr Leu Ser Trp Val Met Asp Thr Asn Lys Glu Glu Arg
 340 345 350
 Asn Asp Gly Lys Thr Ile Glu Val Gly Lys Ala Tyr Phe Glu Thr Asp
 355 360 365
 Lys Arg Arg Tyr Thr Ile Leu Asp Ala Pro Gly His Lys Met Tyr Val
 370 375 380
 Ser Glu Met Ile Gly Gly Ala Ser Gln Ala Asp Val Gly Ile Leu Val
 385 390 395 400
 Ile Ser Ala Arg Lys Gly Glu Tyr Glu Thr Gly Phe Glu Lys Gly Gly
 405 410 415
 Gln Thr Arg Glu His Ala Leu Leu Ala Lys Thr Gln Gly Val Asn Lys
 420 425 430
 Ile Ile Val Val Val Asn Lys Met Asp Asp Ser Thr Val Gly Trp Ser
 435 440 445
 Lys Glu Arg Tyr Gln Glu Cys Thr Thr Lys Leu Gly Ala Phe Leu Lys
 450 455 460
 Gly Ile Gly Tyr Ala Lys Asp Asp Ile Ile Tyr Met Pro Val Ser Gly
 465 470 475 480
 Tyr Thr Gly Ala Gly Leu Lys Asp Arg Val Asp Pro Lys Asp Cys Pro
 485 490 495
 Trp Tyr Asp Gly Pro Ser Leu Leu Glu Tyr Leu Asp Asn Met Asp Thr
 500 505 510
 Met Asn Arg Lys Ile Asn Gly Pro Phe Met Met Pro Val Ser Gly Lys
 515 520 525
 Met Lys Asp Leu Gly Thr Ile Val Glu Gly Lys Ile Glu Ser Gly His
 530 535 540
 Val Lys Lys Gly Thr Asn Leu Ile Met Met Pro Asn Lys Thr Pro Ile
 545 550 555 560
 Glu Val Leu Thr Ile Phe Asn Glu Thr Glu Gln Glu Cys Asp Thr Ala
 565 570 575
 Phe Ser Gly Glu Gln Val Arg Leu Lys Ile Lys Gly Ile Glu Glu Glu
 580 585 590
 Asp Leu Gln Pro Gly Tyr Val Leu Thr Ser Pro Lys Asn Pro Val Lys
 595 600 605
 Thr Val Thr Arg Phe Glu Ala Gln Ile Ala Ile Val Glu Leu Lys Ser
 610 615 620
 Ile Leu Ser Asn Gly Phe Ser Cys Val Met His Leu His Thr Ala Ile
 625 630 635 640
 Glu Glu Val Lys Phe Ile Glu Leu Lys His Lys Leu Glu Lys Gly Thr
 645 650 655
 Asn Arg Lys Ser Lys Lys Pro Pro Ala Phe Ala Lys Lys Gly Met Lys
 660 665 670
 Ile Ile Ala Ile Leu Glu Val Gly Glu Leu Val Cys Ala Glu Thr Tyr
 675 680 685
 Lys Asp Tyr Pro Gln Leu Gly Arg Phe Thr Leu Arg Asp Gln Gly Thr
 690 695 700
 Thr Ile Ala Ile Gly Lys Ile Thr Lys Leu Leu
 705 710 715

<210> SEQ ID NO 48

<211> LENGTH: 653

-continued

<212> TYPE: DNA

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 48

```
tcgagtttat cattatcaat actcgcatt tcaaagaata cgtaaataat taatagtagt    60
gattttccta actttattta gtcaaaaaat tagcctttta attctgctgt aaccggtaca    120
tgccaaaata gggggcgggt tacacagaat atataacact gatggtgctt ggggtgaacag    180
gtttattcct ggcatccact aaatataatg gagcccgctt ttaagctgg catccagaaa    240
aaaaaagaat ccagcacca aaatattggt ttcttcacca accatcagtt catagggtcca    300
ttctcttagc gcaactacag agaacagggc acaaacaggc aaaaaacggg cacaacctca    360
atggagtgat gcaacctgcc tggagtaaat gatgacacaa ggcaattgac ccacgcatgt    420
atctatctca ttttcttaca ccttctatta ccttctgctc tctctgattt ggaaaaagct    480
gaaaaaaaaag gtttaaacca gttccctgaa attattcccc tacttgacta ataagtatat    540
aaagacggta ggtattgatt gtaattctgt aaatctattt cttaaacttc ttaaattcta    600
cttttatagt tagtcttttt tttagtttta aaacaccaag aacttagttt cga          653
```

<210> SEQ ID NO 49

<211> LENGTH: 7988

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Ure2N-Sup35C integration plasmid

<400> SEQUENCE: 49

```
tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagctcccg gagacggtca    60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg    120
ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc    180
accataccac agcttttcaa ttcaattcat catttttttt ttattctttt ttttgatttc    240
ggtttctttg aaattttttt gattcggtaa tctccgaaca gaaggaagaa cgaaggaagg    300
agcacagact tagattggta tatatacgca tatgtagtgt tgaagaaaca tgaattgcc    360
cagtattcct aacccaactg cacagaacaa aaacctgcag gaaacgaaga taaatcatgt    420
cgaaagctac atataaggaa cgtgctgcta ctcatcctag tcctggtgct gccaaagetat    480
ttaatatcat gcacgaaaag caaacaaact tgtgtgcttc attggatggt cgtaccacca    540
aggaattact ggagttagtt gaagcattag gtcccaaat ttgtttacta aaaacacatg    600
tggatatcct gactgatttt tccatggagg gcacagttaa gccgctaaag gcattatccg    660
ccaagtacaa ttttttactc ttcgaagaca gaaaatttgc tgacattggt aatacagtca    720
aattgcagta ctctgcgggt gtatacagaa tagcagaatg ggcagacatt acgaatgcac    780
acggtgtggt gggcccagggt attgttagcg gtttgaagca ggcggcagaa gaagtaacaa    840
aggaacctag aggccttttg atgttagcag aattgtcatg caagggtccc ctatctactg    900
gagaatatac taagggtact gttgacattg cgaagagcga caaagatttt gttatcggct    960
ttattgctca aagagacatg ggtggaagag atgaaggtta cgattggttg attatgacac    1020
ccggtgtggg tttagatgac aaggagacg cattgggtca acagtataga accgtggatg    1080
atgtgtctc tacaggatct gacattatta ttgttgaag aggactatth gcaaagggaa    1140
```

-continued

gggatgctaa ggtagagggg gaacgttaca gaaaagcagg ctgggaagca tatttgagaa 1200
gatgcgccca gcaaaactaa aaaactgtat tataagtaaa tgcattgata ctaaactcac 1260
aaattagagc ttcaatttaa ttatatcagt tattacccta tgcgggtgta aataccgcac 1320
agatgcgtaa ggagaaaata ccgcatcagg aaattgtaaa cgtaaatatt ttgttaaaat 1380
tcgcgtaaa tttttgttaa atcagctcat ttttaacca ataggccgaa atcggcaaaa 1440
tccctataa atcaaaagaa tagaccgaga tagggttgag tgttgtcca gtttggaaaca 1500
agagtccact attaaagaac gtggactcca acgtcaaagg gcgaaaaacc gtctatcagg 1560
gcgatggccc actacgtgaa ccatcacct aatcaagttt tttgggtcg aggtgcccga 1620
aagcactaaa tcggaacct aaaggagcc cccgatttag agcttgacgg ggaagccgg 1680
cgaacgtggc gagaaaggaa gggaagaaag cgaagagagc gggcgctagg gcgctggcaa 1740
gtgtagcggc cacgctgcgc gtaaccacca caccgcgcgc gcttaatgag ccgctacagg 1800
gcgctgcgag ccattcgcca ttcaggctgc gcaactgttg ggaagggcga tcggtgcggg 1860
cctcttcgct attacgccag ctggcgaaag ggggatgtgc tgcaaggcga ttaagttggg 1920
taacgccagg gttttccag tcacgacgtt gtaaaacgac ggccagtga ttaataacg 1980
actcactata gggcgaattg gagctccacc gcggtgaaaa gagtcagtga gacgacgact 2040
tcaggatctt tgggtttcag gatatgtggc atgaaaatac agaaaaatcc ctctgtgctg 2100
aatcagcttt cgctcgaata ttacgaagaa gaagcagaca gtgattatat ctttataaac 2160
aaattgtatg tgcgttcaag aaccgatcaa aatgtttcag atgcaattga actttatatt 2220
aacaatcctc atctgtcggg tgcgagaaa catcaactga agaaaaatc tttgaaaaga 2280
ttgcagttgt tttataatac tatgctagaa gaagaagtta gaatgatatc aagtagctt 2340
ttgtttatc acgaaggaga cccggagcga tgggaattac taaatgatgt tgacaaactt 2400
atgcgagatg attttataga cgatgatgac gacgatgatg ataatgatga tgatgatgat 2460
gatgatgccg agggaagcag cgaaggacca aaggacaaaa aaacaactgg ttctttgagt 2520
tccatgtcac taatagattt tgcacattct gaaataacgc cggggaaggg ttatgatgaa 2580
aacgtgattg aaggagtga aacctgcta gatattttta tgaaattcta gatatattga 2640
gaggtgaagt ttacctgtt tatggtatat ggtacaaaa gaaactaaact aattatacgt 2700
ctatatatat atatatatat ataacagctt tattaacct tgtttttta tatagaagaa 2760
aatgctttat gatcggattt attgtgttg catttactta tgtttgcaag aatggatcc 2820
ttactcggca attttaacaa ttttaccat tgcattgtg gtacctgat ctctcaaagt 2880
gaatctacct aattgaggg aatcttgta agttccaca caaactggag ctctcagttc 2940
taaaacagcg atgacctca tacccttctt agcaaaagca ggtggttct ttgacttacg 3000
gttggtaacc ttttctaatt tgtgcaata cttacaata tgtacctct caattgctgt 3060
atgaacatgc ataacacatg aaaaaccggc tgctatgata gattttaatt ctacaatagc 3120
aatttgagct acaacttg taacactctt gatagggctt tttggcgatg ttagtacaaa 3180
acctggtgaa atgtcttctt cttcaacacc tttgattctt agtttaactt gctcaccaca 3240
catagccata tcaacttcat tttcagttc gttgtaata ttttgaattt ccacagcggg 3300
tttgttaggc atcagtaggg tggattgacc ctttttgata tgaccggatt caattttacc 3360
ttcaacgatg gtacctagat ccttcatctt agcggcaata ggcaacatga atggagcatt 3420

-continued

gatgtgacgg tcgacgtggt tcattgtatc cagatattct aacagagttg ggccgggtga	3480
ccatgggcat tcttttgat ctacgtgatc tttcaaattt gcaccactgt agccggatac	3540
tggcataaat acaacgtctg tcttaatggt gtaaccaatt gctctcaaga aattgctgac	3600
attactcaca cattggtcgt aacgttcctt agaccagtta acggttgggt catccatctt	3660
atttacgacg acaaccatct tattaacacc ttgggtcttg gccaataggg cgtgttcacg	3720
agtttgacca cctctctcaa aaccggtttc gtactcacc tttctggcgg aaatgaccaa	3780
aacaccaaca tcagcttgag aagcaccacc gatcatctcg gaaacgtaca ttttatgacc	3840
aggagcatcc aatatgggat aacgcctttt ttcagtttca aagtaggcct taccaacttc	3900
gatagtctta ccatcatttc tttctctctt gttggtatcc atgacccatg acaagtacca	3960
accttgtctg cctgcatcct tggtctctct ttcataattc tcaatagttc tcttatccac	4020
agagccagtc aagtatagta gattaccacc catagtagat ttaccggcat caacatgacc	4080
catgaaaatt aaagaaacgt gatctttacc accaaacata gcgtagtctg ggacgtcgta	4140
tgggtagcgg ccgctgttat tgttttgaac attattgtta ttactactgc tattgttatt	4200
attattatta ttacacctg ttgaaaatc aaaatttata ttactttgat cggtggttgt	4260
attactgttc ctgtttccta tgtttacttg acggagcgca ttggagagat tcgacacttg	4320
gttgccgttg ttattcatca tgaattctgt tgctagtggg cagatataga tgttattccg	4380
agcaagtcga tgaagaaacc gctttttggt acagtacaat ggagtctttc aagagaagat	4440
gtaccaatat aactacact cttcagaagc aatgggagct ttggtcgagt gaaaaaaaaa	4500
ttttctccat aaagaaagat catattatac gatgatgtaa gatataatac ccggttgtaa	4560
tgtacattta agagcaaggt aagaagtgac aataacttct gtatgatctt agcatgtacc	4620
tcttttggtg ggctgagaac taagattcat ctttttgcgg aagaattttg ctatgaactt	4680
cacaacttta tgaagtgggt taagagaatt acaaaagaaa tgacacagac tcgaacactg	4740
tgacgcgtcg tcttagtaaa aaataataat ttgagtcaaa tagcgcagct aatgcgaaac	4800
aaagaaatga agcatatacc attcgttgta tgatttttgt gtggttgaca gatattctgc	4860
cgaattttta acgcttatta taaatataaa tgotatgatg tgtgtataaa cagatacgat	4920
attcaatttt ctaccgtagg gttgggattt tttcaaact ccaattcttc gtcgggtatt	4980
tectcaatgg cgatcctctt ttttggttc ggcttttcag tgcattgac aattttagge	5040
accttaattt gtagtagacc gttgttgtaa gtagctttaa tttcttcgtc cttaatgcgt	5100
ggcagcacgg ggaatttaac ggttctctca aacgcacat attttagttc cgtgatcttc	5160
agaatttttt catcaatgcc cactctgtct togatcttac ccttgatgag catctcatga	5220
gaagatggat ggtaatcaat gtggaagcc ctagagttag cacctggtaa cgcaagaaca	5280
actacgtaag tgcctcgggt atcatagaca ttcacttctg gtgaaatgg taagtccatt	5340
ctcgtttcag gcttgatac ttgtaacggg tcaggtattg ggaaggcgc ttgtaggtga	5400
gcgaacgatg aagatttttt ggctaaggt ggtctcgac attcctccag ctgattcaaa	5460
ggtttttctt tgttggttc gccagcttc tctttgggtg cttcagactt atccttctta	5520
tccttttctt ctcccttttc gccctcctgt tcggattttg cttcaatttc tggttcagtg	5580
ccttcatatg gtggaacacc tattaacgcg gtaataaagt cgtttaaact gttgcctgt	5640
tggttggtcc tattattcct gccagtatta caatggtaat atgatggata tcttctcgag	5700

-continued

ggggggcccg	gtaccagct	ttgttccct	ttagtgaggg	ttaattccga	gcttggcgtgta	5760
atcatgggtca	tagctgtttc	ctgtgtgaaa	ttgttatccg	ctcacaattc	cacacaacat	5820
acgagccgga	agcataaagt	gtaaagcctg	gggtgcctaa	tgagtgaggt	aactcacatt	5880
aattgcgttg	cgctcactgc	ccgctttcca	gtcgggaaac	ctgtcgtgcc	agctgcatta	5940
atgaatcggc	caacgcgcgg	ggagaggcgg	tttgcgtatt	gggcgctctt	ccgcttcttc	6000
gctcactgac	tcgctcgcgt	eggctgttcg	gctcgcggca	gcggtatcag	ctcactcaaa	6060
ggcggttaata	cggttatcca	cagaatcagg	ggataacgca	ggaaagaaca	tgtgagcaaa	6120
aggccagcaa	aaggccagga	accgtaaaaa	ggccgcgttg	ctggcgtttt	tccataggct	6180
ccgccccct	gacgagcatc	acaaaaatcg	acgctcaagt	cagaggtggc	gaaaccgac	6240
aggactataa	agataccagg	cgtttcccc	tggaaactcc	ctcgtcgcgt	ctcctgttcc	6300
gaccctgccg	cttaccggat	acctgtccgc	ctttctccct	tcgggaagcg	tgccgctttc	6360
tcatagtctca	cgctgtaggt	atctcagttc	gggtgtagtc	gttcgctcca	agctgggctg	6420
tgtgcacgaa	ccccccgttc	agcccagccg	ctgcgcctta	tccggtaact	atcgtcttga	6480
gtccaaccgg	gtaagacacg	acttatcgcc	actggcagca	gccactggta	acaggattag	6540
cagagcgagg	tatgtaggcg	gtgctacaga	gttcttgaag	tggtggccta	actacggcta	6600
cactagaagg	acagtatttg	gtatctgcgc	tctgctgaag	ccagttacct	tcggaaaaag	6660
agttgtagc	tcttgatccg	gaaacaaac	caccgctggg	agcggtggtt	ttttgtttg	6720
caagcagcag	attacgcgca	gaaaaaagg	atctcaagaa	gatcctttga	tcttttctac	6780
ggggtctgac	gctcagtggg	acgaaaactc	acgttaaggg	atthtggca	tgagattatc	6840
aaaaaggatc	ttcacctaga	tccttttaaa	ttaaaaatga	agttttaaat	caatctaaag	6900
tataatagag	taaacttggt	ctgacagtta	ccaatgctta	atcagtgagg	caactatctc	6960
agcgtatctg	ctatttctgt	catccatagt	tgctgactc	cccgtcgtgt	agataactac	7020
gatacgggag	ggcttaccat	ctggccccag	tgctgcaatg	ataccgcgag	acccacgctc	7080
accggctcca	gatttatcag	caataaacca	gccagccgga	aggcccgagc	gcagaagtgg	7140
tcttgcaact	ttatccgctc	ccatccagtc	tattaattgt	tgccgggaag	ctagagtaag	7200
tagttcgcca	gttaatagtt	tgccgaacct	tggtgccatt	gctacaggca	tcgtggtgtc	7260
acgctcgtcg	tttggatagg	cttcattcag	ctccggttcc	caacgatcaa	ggcgagttae	7320
atgatcccc	atggtgtgca	aaaaagcggg	tagctccttc	ggctctccga	tcgttgtcag	7380
aagtaagtty	gccgcagtg	tatcactcat	ggttatggca	gcactgcata	attctcttac	7440
tgctatgcca	tcgtaagat	gcttttctgt	gactgggtgag	tactcaacca	agtcattctg	7500
agaatagtgt	atgcccggac	cgagttgctc	tgccccggcg	tcaatacggg	ataataccgc	7560
gccacatagc	agaactttaa	aagtgtcat	cattggaaaa	cgttcttcgg	ggcgaaaact	7620
ctcaaggatc	ttaccgctgt	tgagatccag	ttcgatgtaa	cccactcgtg	cacccaactg	7680
atcttcagca	tcttttactt	tcaccagcgt	ttctgggtga	gcaaaaacag	gaaggcaaaa	7740
tgccgcaaaa	aaggaataa	gggcgacacg	gaaatggtga	atactcatac	tcttcctttt	7800
tcaatattat	tgaagcattt	atcaggggta	ttgtctcatg	agcggataca	tatttgaatg	7860
tatttagaaa	aataaacaaa	taggggttcc	gogcacattt	ccccgaaaag	tgccacctga	7920
cgtctaagaa	accattatta	tcatgacatt	aacctataaa	aataggcgta	tcacgagggc	7980

-continued

ctttcgtc

7988

<210> SEQ ID NO 50

<211> LENGTH: 405

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 50

```

Met Asp Thr Asp Lys Leu Ile Ser Glu Ala Glu Ser His Phe Ser Gln
 1           5           10           15
Gly Asn His Ala Glu Ala Val Ala Lys Leu Thr Ser Ala Ala Gln Ser
20           25           30
Asn Pro Asn Asp Glu Gln Met Ser Thr Ile Glu Ser Leu Ile Gln Lys
35           40           45
Ile Ala Gly Tyr Val Met Asp Asn Arg Ser Gly Gly Ser Asp Ala Ser
50           55           60
Gln Asp Arg Ala Ala Gly Gly Gly Ser Ser Phe Met Asn Thr Leu Met
65           70           75           80
Ala Asp Ser Lys Gly Ser Ser Gln Thr Gln Leu Gly Lys Leu Ala Leu
85           90           95
Leu Ala Thr Val Met Thr His Ser Ser Asn Lys Gly Ser Ser Asn Arg
100          105          110
Gly Phe Asp Val Gly Thr Val Met Ser Met Leu Ser Gly Ser Gly Gly
115          120          125
Gly Ser Gln Ser Met Gly Ala Ser Gly Leu Ala Ala Leu Ala Ser Gln
130          135          140
Phe Phe Lys Ser Gly Asn Asn Ser Gln Gly Gln Gly Gln Gly Gln Gly
145          150          155          160
Gln Gly Gln Gly Gln Gly Gln Gly Gln Gly Gln Gly Ser Phe Thr Ala
165          170          175
Leu Ala Ser Leu Ala Ser Ser Phe Met Asn Ser Asn Asn Asn Asn Gln
180          185          190
Gln Gly Gln Asn Gln Ser Ser Gly Gly Ser Ser Phe Gly Ala Leu Ala
195          200          205
Ser Met Ala Ser Ser Phe Met His Ser Asn Asn Asn Gln Asn Ser Asn
210          215          220
Asn Ser Gln Gln Gly Tyr Asn Gln Ser Tyr Gln Asn Gly Asn Gln Asn
225          230          235          240
Ser Gln Gly Tyr Asn Asn Gln Gln Tyr Gln Gly Gly Asn Gly Gly Tyr
245          250          255
Gln Gln Gln Gln Gly Gln Ser Gly Gly Ala Phe Ser Ser Leu Ala Ser
260          265          270
Met Ala Gln Ser Tyr Leu Gly Gly Gly Gln Thr Gln Ser Asn Gln Gln
275          280          285
Gln Tyr Asn Gln Gln Gly Gln Asn Asn Gln Gln Gln Tyr Gln Gln Gln
290          295          300
Gly Gln Asn Tyr Gln His Gln Gln Gln Gly Gln Gln Gln Gln Gly
305          310          315          320
His Ser Ser Ser Phe Ser Ala Leu Ala Ser Met Ala Ser Ser Tyr Leu
325          330          335
Gly Asn Asn Ser Asn Ser Asn Ser Ser Tyr Gly Gly Gln Gln Gln Ala
340          345          350

```

-continued

Asn Glu Tyr Gly Arg Pro Gln His Asn Gly Gln Gln Gln Ser Asn Glu
 355 360 365

Tyr Gly Arg Pro Gln Tyr Gly Gly Asn Gln Asn Ser Asn Gly Gln His
 370 375 380

Glu Ser Phe Asn Phe Ser Gly Asn Phe Ser Gln Gln Asn Asn Asn Gly
 385 390 395 400

Asn Gln Asn Arg Tyr
 405

<210> SEQ ID NO 51
 <211> LENGTH: 128
 <212> TYPE: PRT
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 51

Met Ser Ala Asn Asp Tyr Tyr Gly Gly Thr Ala Gly Glu Lys Ser Gln
 1 5 10 15

Tyr Ser Arg Pro Ser Asn Pro Pro Pro Ser Ser Ala His Gln Asn Lys
 20 25 30

Thr Gln Glu Arg Gly Tyr Pro Pro Gln Gln Gln Gln Tyr Tyr Gln
 35 40 45

Gln Gln Gln Gln His Pro Gly Tyr Tyr Asn Gln Gln Gly Tyr Asn Gln
 50 55 60

Gln Gly Tyr Asn Gln Gln Gly Tyr Asn Gln Gln Gly Tyr Asn Gln Gln
 65 70 75 80

Gly Tyr Asn Gln Gln Gly Tyr Asn Gln Gln Gly His Gln Gln Pro Val
 85 90 95

Tyr Val Gln Gln Gln Pro Pro Gln Arg Gly Asn Glu Gly Cys Leu Ala
 100 105 110

Ala Cys Leu Ala Ala Leu Cys Ile Cys Cys Thr Met Asp Met Leu Phe
 115 120 125

<210> SEQ ID NO 52
 <211> LENGTH: 534
 <212> TYPE: PRT
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 52

Met Ser Ser Asp Glu Glu Asp Phe Asn Asp Ile Tyr Gly Asp Asp Lys
 1 5 10 15

Pro Thr Thr Thr Glu Glu Val Lys Lys Glu Glu Glu Gln Asn Lys Ala
 20 25 30

Gly Ser Gly Thr Ser Gln Leu Asp Gln Leu Ala Ala Leu Gln Ala Leu
 35 40 45

Ser Ser Ser Leu Asn Lys Leu Asn Asn Pro Asn Ser Asn Asn Ser Ser
 50 55 60

Ser Asn Asn Ser Asn Gln Asp Thr Ser Ser Ser Lys Gln Asp Gly Thr
 65 70 75 80

Ala Asn Asp Lys Glu Gly Ser Asn Glu Asp Thr Lys Asn Glu Lys Lys
 85 90 95

Gln Glu Ser Ala Thr Ser Ala Asn Ala Asn Ala Asn Ala Ser Ser Ala
 100 105 110

Gly Pro Ser Gly Leu Pro Trp Glu Gln Leu Gln Gln Thr Met Ser Gln
 115 120 125

-continued

530

<210> SEQ ID NO 53
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 53

ggaggatcca tggatcacga taagttaatc tcag 34

<210> SEQ ID NO 54
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 54

ccaagctttc agtagcgggt ctggtgagaa aagttg 36

<210> SEQ ID NO 55
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 55

ggtgtcttgg ccaattgcc 20

<210> SEQ ID NO 56
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 56

gtcgacctgc agcgtacgca tttcagatct ttgctatac 39

<210> SEQ ID NO 57
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 57

cgagctcgaa ttcacgatt gattcagttc gccttctatc 40

<210> SEQ ID NO 58
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

-continued

<400> SEQUENCE: 58

ctgttttgaa aggggccaca tg

22

<210> SEQ ID NO 59

<211> LENGTH: 34

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 59

ggaggatcca tggatcggga taagttaatc tcag

34

<210> SEQ ID NO 60

<211> LENGTH: 36

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 60

ggaccgcggg tagcggttct gttgagaaaa gttgcc

36

<210> SEQ ID NO 61

<211> LENGTH: 36

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 61

gaggatccat gcctgatgat gaggaagaag acgagg

36

<210> SEQ ID NO 62

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 62

cggaattcct cgagaagata tccatc

26

<210> SEQ ID NO 63

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer

<400> SEQUENCE: 63

gggatcctgt tgctagtggg caga

24

<210> SEQ ID NO 64

<211> LENGTH: 34

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:

-continued

```

synthetic primer
<400> SEQUENCE: 64
gtaccgcgga tgtctttgaa cgactttcaa aagc
34

<210> SEQ ID NO 65
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:
synthetic primer
<400> SEQUENCE: 65
gtggagctct tactcggcaa ttttaacaat ttac
35

```

1. An electrical conductor comprising a fibril having a first location separated from a second location and an electrically conductive material disposed on the fibril between the first location and second location to conduct electricity along the fibril from the first location to the second location.

2. The electrical conductor of claim **1** wherein the fibril comprises polypeptide subunits coalesced into an ordered aggregate.

3-11. (canceled)

12. The electrical conductor of claim **2** wherein the electrical conductor is characterized by a length of 60 nm to 300 μm , and a diameter of 9 nm to 200 nm.

13. The electrical conductor of claim **2** wherein at least one of the polypeptide subunits comprises a SCHAG amino acid sequence.

14. The electrical conductor of claim **13** wherein 90-100% of the polypeptide subunits comprise a SCHAG amino acid sequence.

15. The electrical conductor of claim **13** wherein the SCHAG amino acid sequence includes at least one amino acid residue having a reactive amino acid side chain.

16. The electrical conductor of claim **13** wherein the SCHAG amino acid sequence includes at least one substitution of an amino acid residue having a reactive amino acid side chain.

17. The electrical conductor of claim **15** wherein the reactive amino acid side chain is exposed to the environment of the fibril to permit attachment of electrically conductive material thereto, and wherein the electrically conductive material is attached to the fibril at the reactive amino acid side chain.

18. (canceled)

19. The electrical conductor of claim **13** wherein at least 30% of the SCHAG amino acid sequence comprises asparagine or glutamine residues.

20-25. (canceled)

26. The electrical conductor of claim **1** wherein the electrically conductive material comprises a material selected from the group consisting of a metal atom and a semiconductor material.

27. The electrical conductor of claim **26** wherein the metal atom is selected from the group consisting of gold, silver, nickel, copper, platinum, aluminum, gallium, palladium, iridium, rhodium, tungsten, titanium, zinc, and tin.

28. (canceled)

29. The electrical conductor of claim **27** wherein the fibril is gold-toned.

30. The electrical conductor of claim **29** wherein the fibril is characterized by a resistance range of 0-100 Ω and linear I-V curves.

31. (canceled)

32. A method of making an electrical conductor comprising the steps of: (a) making a fibril with first and second separated locations; and (b) disposing on the fibril an electrically conductive material in an amount effective to conduct electricity along the fibril from the first location to the second location.

33. The method according to claim **32** wherein step (a) comprises providing a solution or suspension of polypeptides that have the ability to coalesce into ordered aggregates, and incubating the solution or suspension under conditions to form fibrils from the polypeptides.

34-35. (canceled)

36. The method according to claim **32** wherein step (b) comprises disposing a substrate on the fibril, and disposing a first electrically conductive material on the substrate.

37. The method according to claim **36** wherein a second electrically conductive material is disposed on the first electrically conductive material.

38-70. (canceled)

71. The method according to claim **32** wherein the electrically conductive material comprises a metal atom or a semiconductor material.

72. The method according to claim **71** wherein the electrical conductor material is a metal atom selected from the group consisting of gold, silver, nickel, copper, palladium, iridium, rhodium, tungsten, titanium, zinc, and tin.

73-75. (canceled)

76. An electrical circuit comprising a source of electricity, one or more circuit elements, and electrical conductors disposed between the source of electricity and the one or more circuit elements, and wherein at least one of the electrical conductors comprises a fibril and an electrically conductive material disposed on the fibril to conduct electricity along the fibril between the source of electricity and circuit element or between two circuit elements.

77-79. (canceled)

* * * * *