



Mast cells promote small bowel cancer in a tumor stage-specific and cytokine-dependent manner

Abdulrahman M. Saadalla^a, Abu Osman^a, Michael F. Gurish^b, Kristen L. Dennis^c, Nichole R. Blatner^c, Abdulmohammad Pezeshki^a, Kelly M. McNaghy^d, Hilde Cheroutre^e, Fotini Gounari^f, and Khashayarsha Khazaie^{a,1}

^aDepartment of Immunology, Mayo Clinic, Rochester, MN 55905; ^bDivision of Rheumatology, Immunology and Allergy, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115; ^cRobert H. Lurie Comprehensive Cancer Center, Northwestern University Feinberg School of Medicine, Chicago, IL 60611; ^dThe Biomedical Research Centre, University of British Columbia, Vancouver, BC, V6T 1Z3 Canada; ^eLa Jolla Institute of Allergy and Immunology, San Diego, CA 92121; and ^fKnapp Center for Lupus Research, Section of Rheumatology, Department of Medicine, University of Chicago, Chicago, IL 60637

Edited by Harvey Cantor, Dana-Farber Cancer Institute, Boston, MA, and approved January 5, 2018 (received for review September 25, 2017)

Mast cells (MCs) are tissue resident sentinels that mature and orchestrate inflammation in response to infection and allergy. While they are also frequently observed in tumors, the contribution of MCs to carcinogenesis remains unclear. Here, we show that sequential oncogenic events in gut epithelia expand different types of MCs in a temporal-, spatial-, and cytokine-dependent manner. The first wave of MCs expands focally in benign adenomatous polyps, which have elevated levels of IL-10, IL-13, and IL-33, and are rich in type-2 innate lymphoid cells (ILC2s). These vanguard MCs adhere to the transformed epithelial cells and express murine mast cell protease 2 (mMCP2; a typical mucosal MC protease) and, to a lesser extent, the connective tissue mast cell (CTMC) protease mMCP6. Persistence of MCs is strictly dependent on T cell-derived IL-10, and their loss in the absence of IL-10-expressing T cells markedly delays small bowel (SB) polyposis. MCs expand profusely in polyposis-prone mice when T cells overexpress IL-10. The frequency of polyp-associated MCs is unaltered in response to broad-spectrum antibiotics, arguing against a microbial component driving their recruitment. Intriguingly, when polyps become invasive, a second wave of mMCP5⁺/mMCP6⁺ CTMCs expands in the tumor stroma and at invasive tumor borders. Ablation of mMCP6 expression attenuates polyposis, but invasive properties of the remaining lesions remain intact. Our findings argue for a multistep process in SB carcinogenesis in which distinct MC subsets, and their elaborated proteases, guide disease progression.

small bowel | cancer | mast cells | ILC2 | inflammation

Type-2 inflammation characterized by the abundance of IL-4, IL-13, IL-15, and IL-33, as well as associated cellular components, has been best studied in connection with asthma, allergy, and arthritis. In these situations, two types of tissue sentinel cells respond to cellular stress signals, or danger-associated molecular patterns, and orchestrate type-2 inflammation. These are the type-2 innate lymphoid cells (ILC2s) that seed tissues during embryonic development and mast cells (MCs), which partially reside in tissues but are also recruited from the bone marrow as progenitors. Communication between these two sentinel cell types can produce protective (1) or pathogenic type-2 inflammation (2, 3). Their activation involves IL-10 (4) and IL-33 (3).

In the gastrointestinal (GI) tract, MCs are classified based on their response to infection with nematodes, including *Trichinella spiralis* (5, 6). Two distinct subsets of MCs are identified by their expression of specific proteases, distribution, and kinetics of expansion and contraction (7, 8). Mucosal mast cells (MMC) expand during the early phase of infection in small bowel (SB) crypts and contract during resolution of inflammation. This contraction coincides with the expansion of connective tissue mast cells (CTMCs) in the stroma and then in the submucosa of the infested SB (5, 6). Based on these and later molecular analyses, expression of the β -chymases murine mast cell protease 1 (mMCP1) and mMCP2 defines lymphocyte-dependent MMCs,

whereas the tetrameric tryptases mMCP6 and mMCP7, the β -chymase mMCP4, and the α -chymase mMCP5 define lymphocyte-independent CTMCs (9, 10).

There is clinical and experimental evidence linking MCs to carcinogenesis (11, 12). The protumorigenic properties of MCs are, in part, attributed to their ability to alter regulatory T cell (Treg) properties (13) as well as to influence lipid metabolism (14). In earlier studies, we reported expansion of MCs during polyposis (11, 15). Here, we evaluate the significance of MCs in mouse models of multistep SB cancer, induced by sequential inactivation of the adenomatous polyposis (APC) gene and PTEN or activation of Kras in gut epithelial cells. Combining these models with T cell-specific IL-10 deficiency allows us to dissect cancer-driven inflammation in the course of tumor initiation and progression. We observe a tumor stage-specific expansion of distinct MC subsets with characteristics of MMCs and CTMCs, and elucidate their contribution to cancer initiation and progression. Our data suggest that maturation and lineage commitment of MCs to MMC and then CTMC subsets is inherently and etiologically intertwined with SB multistage carcinogenesis.

Results

Polyps Develop Type-2 Inflammation. Previously, we showed that type-3 cytokines are critical for polyp growth in the SB and colon

Significance

We show that distinct subsets of mast cells (MCs) expand with sequential oncogenic events in small bowel cancer. Mucosal mast cells (MMCs) previously detected early during *Trichinella spiralis* infection expand in adenomatous polyps in an IL-10-dependent manner. Connective tissue mast cells (CTMCs), earlier shown to expand during the resolution of inflammation following clearance of *T. spiralis*, are independent of IL-10 and associate with the transition of polyps to adenocarcinoma. IL-33 upregulates the CTMC lineage-specific protease murine mast cell protease 6 (mMCP6). Ablation of mMCP6 attenuates tumor growth. Thus, tissue sentinel cells respond to oncogenic events and cellular transformation in effect to help promote cancer. Delineating the types of MCs present at various stages of disease offers actionable cellular targets for therapeutic intervention in disease progression.

Author contributions: A.M.S., A.O., M.F.G., H.C., F.G., and K.K. designed research; A.M.S., A.O., K.L.D., N.R.B., and A.P. performed research; M.F.G., K.L.D., N.R.B., K.M.M., H.C., and F.G. contributed new reagents/analytic tools; A.M.S., A.O., K.L.D., N.R.B., A.P., F.G., and K.K. analyzed data; and A.M.S., K.M.M., and K.K. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

¹To whom correspondence should be addressed. Email: khazaie@mayo.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1716804115/-DCSupplemental.




Fig. 1. Type-2 inflammation in polyps. (A) Quantification of expression of nuclear IL-33 in polyps; immunohistochemistry ($n > 6$). (B) Percentage of ST2-expressing MCs among total MNCs ($n = 9$). (C) Mean fluorescence intensity (MFI) of MC ST2 ($n = 5$). (D and E) Frequencies of T helper 2 cells and ILC2s by immunostaining ($n > 40$). Unpaired t test: **** $P < 0.0001$; *** $P < 0.001$; * $P < 0.005$.

of APC^{Δ468} mice. These mice were born with a hereditary defect in the APC gene rendering them susceptible to polyposis (16); however, somewhat surprisingly, they also exhibited mastocytosis in polyps (11, 13–15) consistent with type-2 inflammation. To test whether expansion of MCs in polyps is indicative of local type-2 inflammation, we measured tissue cytokine levels in the intestines of polyp-ridden mice. IL-10, IL-13, and IL-5, as well as TGF- β , were strongly expressed (Fig. S1). Although we did not detect a significant increase in soluble IL-33, transformed epithelial cells of the polyps displayed strong nuclear IL-33 staining compared with epithelia of the underlying hyperproliferative crypts, and healthy neighboring tissues (Fig. 1A and Fig. S2A–C). IL-33 is a nuclear alarmin and maturation factor for both MCs and ILCs (17). IL-33 functions both as a cytokine and as a nuclear factor that regulates expression of ST2/IL-33 receptor (IL-33R) and its soluble decoy receptor counterpart (18). Therefore, we measured expression of the IL-33R (ST2/IL-33R) by polyp-infiltrating MCs and compared this with the healthy surrounding tissues. Microdissected polyps were enzymatically digested to prepare mononuclear cells (MNCs), which were then subjected to flow cytometric analyses. MCs were identified after negative gating for lineage-specific markers and positive gating for Fc ϵ R1 and cKit. While all MCs expressed ST2/IL-33R, the mean fluorescence intensity of staining, which represents the density of receptors per cell, was significantly higher on polyp-infiltrating MCs compared with MCs infiltrating the healthy surrounding tissues (Fig. 1B and C and Fig. S2D). Up-regulation of ST2 on

polyp-infiltrating MCs is consistent with their response to IL-33. Next, we stained paraffin sections of fixed gut “jelly-rolls” for infiltrating T helper 2 cells and ILC2s. Both cell types expanded significantly within polyps compared with the healthy surrounding tissues (Fig. 1D and E). There were parallel increases in IL-1 β (inflammasome activity) and type-3 cytokines in line with ongoing type-3 inflammation (Fig. S1). The T cell tropic chemokines CXCL9, CXCL10, CCL5, and CCL11 and cytokines IL-2 and IL-15 were also increased (Fig. S1). In conclusion, the microenvironment of SB polyps was strongly supportive of type-2 and type-3 inflammation and of T lymphocyte recruitment.

IL-10 Promotes MC Expansion and Polyp Growth in the SB. In earlier studies, we showed that IL-10 deficiency attenuates SB polyposis (19). We reasoned that the initial attenuation of polyposis in IL-10-deficient APC^{Δ468} mice reflected a role for IL-10 in MC function. Therefore, we tested here whether polyp-infiltrating MCs depend on IL-10 using three strategies: (i) conditional ablation of IL-10 in T cells using CD4Cre (CD4CreIL10^{fl}), (ii) germline ablation of IL-10 (IL-10^{-/-}) (19, 20), and (iii) transgenic overexpression of IL-10 in T cells under the control of the IL-2 gene promoter (IL2pIL10) (21). Mice were aged 4 mo to allow SB polyps to develop.

To better understand the kinetics of the disease, we examined mice at 2, 4, and 7 mo of age. Immune-competent APC^{Δ468} mice had few small polyps and mostly aberrant crypts at 2 mo but multiple visible polyps at 4 mo (Fig. 2A). In contrast, T cell-specific IL-10 ablation or germline IL-10 deficiency delayed SB polyposis, such that there was no significant change in polyp load at 4 mo compared with 2 mo (Fig. 2A). Microscopic polyps of 2-mo-old IL-10-deficient APC^{Δ468} mice and age-matched IL-10-competent APC^{Δ468} mice had comparable densities of intra-epithelial mMCP2⁺ (Fig. 2B and Fig. S3B), but had very few mMCP6⁺ MCs in the stroma or submucosa (Fig. 2C and Fig. S3C and D). However, by 4 mo of age, polyp-infiltrating MCs were absent or barely visible in the IL-10-deficient mice (Fig. 2B and C and Fig. 3B–D), suggesting that IL-10 was needed for MC persistence and expansion in the polyps. By 7 mo of age, the mice




Fig. 2. Kinetics of MC expansion in tumors of IL-10-deficient or -proficient mice. (A) Frequency of polyps in the SB of indicated mice at 2, 4, and 7 mo (m) of age. (B–D) Frequencies of mMCP2⁺ MCs, mMCP6⁺ MCs, and myelin basic protein (MBP)-stained eosinophils in the small intestine polyps of the indicated mice ($n \geq 3$ mice, $n > 7$ fields). Unpaired t test: **** $P < 0.0001$; *** $P < 0.001$; ** $P < 0.005$; * $P < 0.05$.

developed invasive tumors (Fig. S3A). New waves of MCs infiltrated the stroma and the submucosa of the lesions, and notably accumulated in the invasive front of the tumors (Fig. S3C). These MCs had a strong preference to express mMCP6 relative to mMCP2 (Fig. 2B; compare with Fig. 2C). In addition to MCs, eosinophils were detected in and around the polyps, with significantly higher densities in IL-10-deficient than IL-10-competent APC^{Δ468} mice. Eosinophils showed a different distribution in the SB, where they accumulated at the margins of polyps, and relative to the colon, where they gathered in the submucosa beneath the polyps (Fig. S3D).

The outcome of IL-10 deficiency in the SB sharply contrasted with that in the colon. At 4 mo of age, the SB of the IL-10-deficient mice had normal levels of TGF- β and type-3 cytokines IL-17a, IL-23, and IL-22 but increased concentrations of IL-33, IFN- γ , and the T cell tropic chemokines CXCL9 and CXCL10 (Fig. S1). By contrast, IL-10 deficiency is known to increase microbial-initiated type-3 cytokines in the colon (20). While the APC^{Δ468} mice with IL-10-deficient T cells exhibited a striking slowdown in growth of SB polyps in the colon, polyposis was accelerated (Fig. 24). Consistent with the delayed SB polyposis in IL-10-deficient APC^{Δ468} mice, we noted that mitotic activity of normal SB crypt epithelial cells was reduced (Fig. S4A). By contrast, there was enhanced crypt epithelial mitosis in the colon of IL-10-deficient APC^{Δ468} mice (Fig. S4B). It is well known that IL-10 deficiency worsens microbial-instigated inflammation and colonic polyposis (20). This has to do with the indispensable function of IL-10-expressing T cells in suppressing bacteria-induced type-3 inflammation. Our findings suggested a different and potentially contrasting function for IL-10-expressing T cells in the SB. To test how T cell expression of IL-10 alters polyposis and MCs, we overexpressed IL-10 in T cells under the control of the IL-2 gene promoter (IL2pIL10 APC^{Δ468} mice) and examined mice at 4 mo of age. Polyposis in both the small intestine and colon was reduced, with a larger impact on the small intestine (Fig. 3 A and B). The IL2pIL10 APC^{Δ468} mice did not have elevated levels of IL-10 in the SB, but TGF- β was reduced compared with the parental APC^{Δ468} mice (Fig. S1). There was extensive expansion of MCs in the SB of IL2pIL10 APC^{Δ468} mice (Fig. 3 C and D), both inside polyps and spreading beyond the polyps to the neighboring healthy SB tissues (Fig. 3 E and F). Furthermore, there were 40-fold more T cells expressing IL-10 in the SB of IL2pIL10 mice than control APC^{Δ468} mice (Fig. 3 G–I).

Expansion of MCs and increased densities of IL-10-expressing T cells, but no significant increase in soluble IL-10, are in line with T cell dependence of MMCs. This shows that expression of IL-10 by the T cells is critically important for MMC expansion. However, these observations also reflect the complex biology of polyposis and multifaceted functions of IL-10. IL-10 is naturally elevated in polyps; therefore, when polyposis is attenuated, IL-10 levels go down, which neutralizes the contribution of overexpression of IL-10 by IL2pIL10 T cells. Type-1 response and establishment of CD8 T cell memory, as well as antitumor cytotoxicity (19), depend on IL-10-expressing T cells and Tregs (22). Thus, we cannot rule out that IL2pIL10 mice have stronger immune surveillance. Expression of IL-10 by T cells also could help stabilize the anti-inflammatory properties of Tregs, which are protective in polyposis (23). It is also possible that polyp growth requires a threshold MC density above which it does not get any worse. Interestingly, as delineated below, tumor progression is more related to the subtype of expanded MCs rather than overall MC numbers.

IL-10-dependent expansion of MC was independent of microbiota. Only colonic polyps responded to treatment with broad-spectrum antibiotics; SB polyps showed little change (Fig. S5 A and B). Interestingly, while antibiotic treatment did not curtail MC expansion in SB polyps, it did hinder the expansion of eosinophils (Fig. S5 C and D). Thus, expansion of MCs and the associated inflammation driven by IL-10-expressing T cells were independent of microbiota. It is possible that intraepithelial MCs




Fig. 3. IL-10 dependence of polyp-infiltrating mMCP2⁺ MCs. (A and B) Frequencies of polyps in the SB and colon, respectively, of the indicated mice ($n > 10$ mice). (C and D) MC densities in polyps, measured by choracetate esterase (CAE) or mMCP2 staining ($n > 50$ fields). A polyp (E) and neighboring healthy tissue (F) from IL2pIL10APC^{Δ468} SB stained with CAE for MCs are shown. (G) Frequencies of IL-10-expressing CD4⁺ T cells in the SB. (H) Representative immunofluorescence images of Thy1.1 staining to reveal IL-10-Thy1.1 reporter (red) and DAPI (blue) staining. (I) Control isotype IgG1, k stain ($n > 3$ fields, $n > 7$ fields, $n \geq 3$ mice). Unpaired *t* test: **** $P < 0.0001$; *** $P < 0.001$; ** $P < 0.005$. Arrows in E and F show MCs; arrow in H shows IL-10-expressing CD4⁺ T cell. (Magnification: E, F, H, and I, 200 \times .)

differentiate in response to sterile danger signals from the transformed epithelial cells. Indeed, SB polyposis has been reported to be less dependent on microbiota (24), and the growth of polyps in the SB was reported in germfree mice (25).

Altogether, these observations provide evidence for the existence of distinct IL-10-dependent, as well as IL-10-independent, MC populations in the SB with mucosal and connective tissue characteristics, based on expression of MC-specific proteases and localization. The differential kinetics of their expansion are consistent with each population being associated with and potentially contributing to distinct stages of tumor initiation and progression. Expansion of eosinophils in the absence, but not the presence, of IL-10 is consistent with these cells being suppressed by IL-10. In our model, the response of eosinophils to antibiotics, in contrast to MCs, points to microbiota driving their expansion.

The Second Wave of MCs Marks the Polyp-to-Carcinoma Transition.

Next, we modeled the genetic cascade of multistage carcinogenesis to delineate the nature of MC expansion in progressive cancer. We used a modified fatty acid-binding promoter Cre mouse (26) in combination with conditional APC (cAPC) (27) to initiate polyposis in the distal ileum and colon (28). To study the polyp-to-carcinoma transition, we also activated Kras or inactivated PTEN, superimposed onto the loss of function of APC. The resulting TS4CreAPCKras and TS4CreAPCCPTEN mice developed invasive tumors similar to those in older IL-10-deficient APC^{Δ468} mice. By far the majority of MCs in benign polyps were mucosal, adhered to the aberrant crypt epithelia, and expressed mMCP2 (Fig. 4 A and C). MC distribution and proteases changed in invasive lesions, with new waves of mMCP6⁺ and mMCP5⁺ MCs expanding in the stroma and submucosa (Fig. 4 B and D–F).

Submucosal MCs adhered to the smooth muscle at the tumor-invasive border, and preferentially expressed mMCP5 (Fig. 4 D and F). Thus, sequential oncogenic events led to expansion of distinct MC types with characteristic spatial distributions.

Ablation of mMCP6 Attenuates Polyposis but Does Not Hinder the Polyp-to-Carcinoma Transition. The expansion of mMCP6⁺ MCs in invasive tumors prompted us to examine the role of mMCP6 in SB cancer. Earlier studies suggested that expression of mMCP6 by MCs is controlled by IL-33 (29), whereas expression of mMCP2 is stimulated by IL-10 (30). We cultured MC progenitors in the presence of IL-33 or IL-10. MCs that matured with medium containing IL-33 had stronger expression of mMCP6, whereas IL-10 inhibited IL-33 expression, in agreement with earlier findings (29) (Fig. S6). Next, we generated chimeric TS4CrecAPC^cPTEN mice with mMCP6-deficient bone marrow or control *wt* bone marrow. The chimeric mice had reduced frequencies of mMCP6⁺ MCs in SB polyps (Fig. 5A and Fig. S7) and increased numbers of mMCP2⁺ MCs (Fig. 5B). Polyposis was attenuated in mice with mMCP6-deficient bone marrow compared with control mice (Fig. 5C and D). Our findings are consistent with a tumor-promoting role for mMCP6. Interestingly, the few remaining lesions were still invasive, suggesting mMCP6 has a role in promoting polyp growth, but not invasion.

Discussion

MC and Polyposis. In the current study, we find that MCs which expand in benign polyps share characteristics with MMCs. These characteristics include expression of mMCP2 and adherence to gut epithelial cells. These MCs expanded in a tissue environment containing elevated levels of type-2 cytokines and ILC2s. A striking finding was that the expansion of this wave of mMCP2⁺ MCs was IL-10-dependent. IL-10 is generally thought to be an antiinflammatory cytokine. However, in some circumstances, IL-10 may function as a proinflammatory cytokine. Earlier studies suggest that IL-10 is involved in the recruitment and maturation of MMCs (30–32). IL-10 stimulates expression of the MMC-specific proteases mMCP1 and mMCP2 (30, 33), and promotes the influx




Fig. 4. Sequential expansion of distinct MCs in the SB during multistage carcinogenesis. (A) Expression of specific proteases by intraepithelial polyp-infiltrating MCs in mouse models of benign polyposis (TS4CrecAPC) and SB cancer (TS4CrecAPC^cPTEN and TS4CrecAPCKras) ($n > 3$ mice, $n > 6$ fields). (B) Expression of MC-specific proteases by polyp-infiltrating submucosal MCs in the same mice ($n > 3$ mice, $n > 6$ fields). (C) Intraepithelial MCs in a TS4CrecAPC polyp stained for mMCP2⁺. The black arrow points to MCs. (D) Stromal and intraepithelial MCs in a TS4CrecAPCKras tumor stained for mMCP2. Submucosal MCs in the same mouse stained for mMCP6 (E) and for mMCP5 (F) are shown. The arrows point to MCs, and the arrowheads point to invasion. Unpaired *t* test: *** $P < 0.0005$; ** $P < 0.005$; * $P < 0.05$. (Magnification in C–F: 200 \times .)




Fig. 5. Bone marrow chimeric TS4CrecAPC^cPTEN mice reveal the tumor-promoting function of mMCP6. (A and B) Frequencies of mMCP6⁺ and mMCP2⁺ MCs in SB polyps. (C and D) Polyposis in the same mice ($n > 4$ mice). Unpaired *t* test: *** $P < 0.001$; ** $P < 0.005$; * $P < 0.05$.

and development of MMCs in a food allergy model (4). Thus, IL-10 is directly or indirectly a cofactor in the development and maturation of MMCs. The persistence of MMCs in benign polyps required expression of IL-10 by T cells. IL-10 deficiency eliminated MMCs and delayed SB polyposis. Conversely, overexpression of IL-10 by T cells expanded MMCs within polyps and in the neighboring tissues, confirming that IL-10-expressing T cells have a rate-limiting role in the maturation of MMCs in the SB. MCs did appear in microscopic polyps of young mice with IL-10-deficient T cells but were lost with time, suggesting that IL-10 may be needed for the persistence of MMCs. We conclude that IL-10 shapes the immune environment of the SB by changing the density of IL-10-dependent MCs. The coincident loss of MCs in IL-10-deficient mice and delay in polyposis suggest that MCs are required for polyposis in the SB, consistent with our earlier findings (13–15). However, further expansion of MMCs through overexpression of IL-10 did not enhance SB polyposis. This reflects the complex biology of IL-10 and polyposis, since IL-10 also enhances T cell memory and antitumor immune surveillance (19), as well as help recovery of Treg antiinflammatory T cells, and hence protective, functions (23). These simultaneous events complicate interpretation of how expansion of MCs over and above what naturally happens during polyposis contributed to polyp growth. The lack of response of polyp-infiltrating MMCs to broad-spectrum antibiotics strongly argues that MC expansion is not bacterially driven.

MCs and SB Cancer. The ability to generate mice with a spectrum of benign and invasive lesions allowed us to identify different types of MCs with distinct spatiotemporal distributions as dysplasia progressed to cancer. We saw a striking expansion of MCs in the stroma and submucosa of tumors that arose by introduction of secondary oncogenic mutations, Kras, or PTEN. The new wave of MCs was distinct from the MCs detected in benign polyposis by their appearance in invasive lesions, location, and expression of CTMC-specific proteases mMCP6 and mMCP5. Here, we find an intriguing similarity in the sequential expansion of MMCs and CTMCs in SB cancer and the earlier reported sequence of MMC expansion leading to inflammation, followed by CTMC expansion and resolution of inflammation, in *T. spiralis*-infected mice. Interestingly, IL-10 deficiency, while delaying SB polyposis, eventually, by itself, led to cancer progression. This may be due to the absence of CD8 T cell memory in IL-10-deficient mice (19) or to the loss of Treg-dependent antiinflammatory properties (16, 23, 34), absence of cytotoxic CD4 T cells (35), or all of these. Loss of IL-10 eliminated polyp-infiltrating MCs that had MMC characteristics. However, it did not hinder MCs for long. New waves of MCs with CTMC properties vigorously expanded during the polyp-to-carcinoma transition. Thus, IL-10 deficiency in the long term reproduced the same sequel of events as in mouse

models of cancer based on the addition of secondary oncogenic events. The expression of MC proteases, mMCP6, mMCP5, and potentially also mMCP4, contributes to MC promotion of SB cancer. A glimpse into these mechanisms came from deletion of mMCP6 in the bone marrow of APC^{Δ468} mice, which led to attenuation of polyposis. Because of the preferential localization of mMCP5⁺ MCs to the tumor-invasive border, we predict that this protease may have a role in the transition from polyp to carcinoma. The role of mMCP4 also needs further investigation.

Region-Specific Effects of MC Expansion: Location, Location, Location.

Interestingly, our data suggest that the inflammation-promoting properties of MCs are likely to be less important in the bacteria-rich colon. IL-10 is indispensable for Treg suppression of microbial-instigated type-3 inflammation (36–38), and down-regulation of IL-10 in Tregs is a characteristic of the immune pathology of colon cancer (16). We find that the colon of IL-10-deficient mice has exacerbated polyposis despite the loss of MCs. The role of the microbiota in colon polyposis is well established. In earlier studies, we showed that colon polyposis is driven by opportunistic microbiota that accumulate within the polyps. Furthermore, we showed that colon polyposis can be reversed by oral gavage of mice with strains of *Lactobacillus acidophilus* that lack cell surface expression of the bacterial lipoteichoic acid, a Toll-like receptor 2 ligand (28). However, we did not detect substantial changes in SB polyposis. Our observations clearly demonstrate striking differences between the immune environments of the SB and colon, and differential roles for IL-10 in these two sites. Since IL-10 deficiency increased colonic polyposis despite the loss of MCs, we can conclude that MCs are not required for polyp growth in the colon, where microbiota dominate regulation of inflammation.

In Conclusion

Together, our findings argue that type-2 inflammation, ILC2s, and MCs are etiologically linked to multistep SB carcinogenesis. Specifically, spatiotemporally regulated expansion of distinct MC subsets and expression of their elaborated proteases shape a supportive tumor environment as polyps grow and continue to progress to cancer. Targeted interruption of this pathway is likely to prevent or potentially interrupt the carcinogenesis process in the SB. We predict that this pathway is also important in other cancers of the upper GI tract where microbiotas are at low densities and not rate-limiting for inflammation.

Materials and Methods

SB tissue was fixed with formalin and paraffin-embedded, and then sectioned (5 μm) for immunofluorescent staining as reported before. A Leica light microscope mounted with a Zeiss Axiocam 503 camera was used for imaging immunohistochemistry staining. For fluorescent staining, a Zeiss Observer. Z1 microscope mounted with an Axiocam 506 mono camera was used for imaging. Fiji software was used for image analysis. The ImmunoRatio plug-in was used for nuclear IL-33 ratio analysis. Additional information is available online as [Supporting Information](#).

For fluorescence-activated cell sorting (FACS), SB tissue was digested with a mixture of collagenase (C5138-1G; Sigma), hyaluronidase (H6254, 500 mg; Sigma), and DNaseI (D5025-15KU; Sigma), and MNCs were prepared by Percoll gradient centrifugation as previously described. FACS analysis was done on a BD LSRII flow cytometer (BD Biosciences) and an Accuri BD flow cytometer (BD Biosciences), and was analyzed by FlowJo software (TreeStar).

Details of antibodies used and staining conditions are available online as [Supporting Information](#).

The statistical analyses were performed with the use of Prism7 (GraphPad). Unpaired one-tailed *t* tests with the Wells correction were used.

ACKNOWLEDGMENTS. This work was supported by Grants R01CA160436 and R01AI108682 (to K.K. and F.G.).

- Shimokawa C, et al. (2017) Mast cells are crucial for induction of group 2 innate lymphoid cells and clearance of helminth infections. *Immunity* 46:863–874.e4.
- Moretto S, et al. (2017) A mast cell-ILC2-Th9 pathway promotes lung inflammation in cystic fibrosis. *Nat Commun* 8:14017.
- Barnig C, et al. (2013) Lipoxin A4 regulates natural killer cell and type 2 innate lymphoid cell activation in asthma. *Sci Transl Med* 5:174ra26.
- Polukort SH, et al. (2016) IL-10 enhances IgE-mediated mast cell responses and is essential for the development of experimental food allergy in IL-10-deficient mice. *J Immunol* 196:4865–4876.
- Friend DS, et al. (1996) Mast cells that reside at different locations in the jejunum of mice infected with *Trichinella spiralis* exhibit sequential changes in their granule ultrastructure and chymase phenotype. *J Cell Biol* 135:279–290.
- Friend DS, et al. (1998) Reversible expression of tryptases and chymases in the jejunal mast cells of mice infected with *Trichinella spiralis*. *J Immunol* 160:5537–5545.
- Gurish MF, Boyce JA (2002) Mast cell growth, differentiation, and death. *Clin Rev Allergy Immunol* 22:107–118.
- Pejler G, Rönnberg E, Waern I, Wernerson S (2010) Mast cell proteases: Multifaceted regulators of inflammatory disease. *Blood* 115:4981–4990.
- Shin K, et al. (2006) Lymphocyte-independent connective tissue mast cells populate murine synovium. *Arthritis Rheum* 54:2863–2871.
- Dwyer DF, Barrett NA, Austen KF; Immunological Genome Project Consortium (2016) Expression profiling of constitutive mast cells reveals a unique identity within the immune system. *Nat Immunol* 17:878–887.
- Maltby S, Khazaie K, McNagny KM (2009) Mast cells in tumor growth: Angiogenesis, tissue remodelling and immune-modulation. *Biochim Biophys Acta* 1796:19–26.
- Khazaie K, et al. (2011) The significant role of mast cells in cancer. *Cancer Metastasis Rev* 30:45–60.
- Blatner NR, et al. (2010) In colorectal cancer mast cells contribute to systemic regulatory T-cell dysfunction. *Proc Natl Acad Sci USA* 107:6430–6435.
- Cheon EC, et al. (2011) Mast cell 5-lipoxygenase activity promotes intestinal polyposis in APC^{Δ468} mice. *Cancer Res* 71:1627–1636.
- Gounaris E, et al. (2007) Mast cells are an essential hematopoietic component for polyp development. *Proc Natl Acad Sci USA* 104:19977–19982.
- Blatner NR, et al. (2012) Expression of ROR γ t marks a pathogenic regulatory T cell subset in human colon cancer. *Sci Transl Med* 4:164ra159.
- Mirchandani AS, Salmond RJ, Liew FY (2012) Interleukin-33 and the function of innate lymphoid cells. *Trends Immunol* 33:389–396.
- Shao D, et al. (2014) Nuclear IL-33 regulates soluble ST2 receptor and IL-6 expression in primary human arterial endothelial cells and is decreased in idiopathic pulmonary arterial hypertension. *Biochem Biophys Res Commun* 451:8–14.
- Dennis KL, et al. (2015) T-cell expression of IL-10 is essential for tumor immune surveillance in the small intestine. *Cancer Immunol Res* 3:806–814.
- Dennis KL, et al. (2013) Adenomatous polyps are driven by microbe-instigated focal inflammation and are controlled by IL-10-producing T cells. *Cancer Res* 73:5905–5913.
- Hagenbaugh A, et al. (1997) Altered immune responses in interleukin 10 transgenic mice. *J Exp Med* 185:2101–2110.
- Laidlaw BJ, et al. (2015) Production of IL-10 by CD4(+) regulatory T cells during the resolution of infection promotes the maturation of memory CD8(+) T cells. *Nat Immunol* 16:871–879.
- Chung AY, et al. (2014) Oral interleukin-10 alleviates polyposis via neutralization of pathogenic T-regulatory cells. *Cancer Res* 74:5377–5385.
- Tomkovich S, et al. (2017) Locoregional effects of microbiota in a preclinical model of colon carcinogenesis. *Cancer Res* 77:2620–2632.
- Dove WF, et al. (1997) Intestinal neoplasia in the ApcMin mouse: Independence from the microbial and natural killer (beige locus) status. *Cancer Res* 57:812–814.
- Saam JR, Gordon JI (1999) Inducible gene knockouts in the small intestinal and colonic epithelium. *J Biol Chem* 274:38071–38082.
- Gounaris F, et al. (2005) Loss of adenomatous polyposis coli gene function disrupts thymic development. *Nat Immunol* 6:800–809.
- Khazaie K, et al. (2012) Abating colon cancer polyposis by *Lactobacillus acidophilus* deficient in lipoteichoic acid. *Proc Natl Acad Sci USA* 109:10462–10467.
- Kaieda S, et al. (2010) Synovial fibroblasts promote the expression and granule accumulation of tryptase via interleukin-33 and its receptor ST-2 (IL1RL1). *J Biol Chem* 285:21478–21486.
- Ghildyal N, et al. (1992) IL-10 induces transcription of the gene for mouse mast cell protease-1, a serine protease preferentially expressed in mucosal mast cells of *Trichinella spiralis*-infected mice. *J Immunol* 149:2123–2129.
- Xia Z, Ghildyal N, Austen KF, Stevens RL (1996) Post-transcriptional regulation of chymase expression in mast cells. A cytokine-dependent mechanism for controlling the expression of granule neutral proteases of hematopoietic cells. *J Biol Chem* 271:8747–8753.
- Thompson-Snipes L, et al. (1991) Interleukin 10: A novel stimulatory factor for mast cells and their progenitors. *J Exp Med* 173:507–510.
- Ghildyal N, McNeil HP, Gurish MF, Austen KF, Stevens RL (1992) Transcriptional regulation of the mucosal mast cell-specific protease gene, MMCP-2, by interleukin 10 and interleukin 3. *J Biol Chem* 267:8473–8477.
- Gounaris E, et al. (2009) T-regulatory cells shift from a protective anti-inflammatory to a cancer-promoting proinflammatory phenotype in polyposis. *Cancer Res* 69:5490–5497.
- Mucida D, et al. (2013) Transcriptional reprogramming of mature CD4⁺ helper T cells generates distinct MHC class II-restricted cytotoxic T lymphocytes. *Nat Immunol* 14:281–289.
- Chaudhry A, et al. (2011) Interleukin-10 signaling in regulatory T cells is required for suppression of Th17 cell-mediated inflammation. *Immunity* 34:566–578.
- Huber S, et al. (2011) Th17 cells express interleukin-10 receptor and are controlled by Foxp3⁺ and Foxp3⁺ regulatory CD4⁺ T cells in an interleukin-10-dependent manner. *Immunity* 34:554–565.
- Rubtsov YP, et al. (2008) Regulatory T cell-derived interleukin-10 limits inflammation at environmental interfaces. *Immunity* 28:546–558.